<https://helda.helsinki.fi>

Nitrogen resorption in *Acer platanoides* and *Acer saccharum* : influence of light exposure and leaf pigmentation

Duan, Baoli

2014-11

Duan , B , Paquette , A , Juneau , P , Brisson , J , Fontaine , B & Berninger , F A 2014 , '
Nitrogen resorption in *Acer platanoides* and *Acer saccharum* : influence of light exposure
and leaf pigmentation ' , *Acta Physiologiae Plantarum* , vol. 36 , no. 11 , pp. 3039-3050 . <https://doi.org/10.1007/s11738-014-1674-x>

<http://hdl.handle.net/10138/159373>

<https://doi.org/10.1007/s11738-014-1674-x>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

ArticleTitle	Nitrogen resorption in <i>Acer platanoides</i> and <i>Acer saccharum</i> : influence of light exposure and leaf pigmentation	
Article Sub-Title		
Article CopyRight	Franciszek Górski Institute of Plant Physiology, Polish Academy of Sciences, Kraków (This will be the copyright line in the final PDF)	
Journal Name	Acta Physiologiae Plantarum	
Corresponding Author	Family Name	Duan
	Particle	
	Given Name	Baoli
	Suffix	
	Division	Département des sciences biologiques, CP 8888 Succ, Centre-ville
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Division	Key Laboratory of Mountain Surface Processes and Ecological Regulation
	Organization	Institute of Mountain Hazards and Environment, Chinese Academy of Sciences
	Address	Chengdu, 610041, China
Author	Email	duanbl@imde.ac.cn
	Family Name	Paquette
	Particle	
	Given Name	Alain
	Suffix	
	Division	Center for Forest Research (CFR)
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Email	
Author	Family Name	Juneau
	Particle	
	Given Name	Philippe
	Suffix	
	Division	Département des sciences biologiques, CP 8888 Succ, Centre-ville
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Email	
Author	Family Name	Brisson
	Particle	
	Given Name	Jacques
	Suffix	
	Division	Center for Forest Research (CFR)
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Division	Institut de recherche en biologie végétale, Département de sciences biologiques

	Organization	Université de Montréal
	Address	4101 East, Sherbrooke St, Montréal, QC, H1X 2B2, Canada
	Email	
Author	Family Name	Fontaine
	Particle	
	Given Name	Bastien
	Suffix	
	Division	Center for Forest Research (CFR)
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Division	Institut de recherche en biologie végétale, Département de sciences biologiques
	Organization	Université de Montréal
	Address	4101 East, Sherbrooke St, Montréal, QC, H1X 2B2, Canada
	Email	
Author	Family Name	Berninger
	Particle	
	Given Name	Frank
	Suffix	
	Division	Département des sciences biologiques, CP 8888 Succ, Centre-ville
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Division	Center for Forest Research (CFR)
	Organization	Université du Québec à Montréal
	Address	Montréal, QC, H3C 3P8, Canada
	Division	Department of Forest Sciences
	Organization	University of Helsinki
	Address	PL 27, Helsinki, 00014, Finland
	Email	
Schedule	Received	2 January 2014
	Revised	4 August 2014
	Accepted	2 September 2014
Abstract	<p>We investigated the effects of leaf color change in the fall on photosynthetic production and nitrogen resorption. Seedlings of <i>Acer platanoides</i> L. and <i>A. saccharum</i> Marsh. were grown in a shade house for 5 months in either 21 % (intermediate light, M) or 4.9 % (low light, L) of incident irradiance. After this period, a subset of the intermediate-light grown seedlings was transferred to a high-light stress treatment (H). Gas exchange, chlorophyll fluorescence, pigments, antioxidant activity, and nitrogen (N) resorption were examined at three leaf senescence stages during September and October. Our results show that plants of both species produce more anthocyanins in the H treatment. In comparison with plants grown in the L and M treatments, plants of both species in the H treatments had lower chlorophyll, carotenoid and chlorophyll fluorescence parameters (F_v/F_m, Φ_{PSII}, NPQ and ETR) at the third sampling date (October 12–18), and indicating higher levels of photoinhibition in the seedlings exposed to high light. Our results imply that autumn leaf redness is inducible and closely linked to photo-oxidative stress. However, anthocyanins did not enhance antioxidant capacity in red leaves in either species, when exposed to high light. For both species, our results showed a higher N-resorption for high-light stressed plants. We also observed that the number of abscised leaves at the second sampling dates (September 10) was higher than at the third sampling dates. The intra-leaf distribution of anthocyanin, the association between anthocyanin production and the high-light environments, the retention of red leaves, the substantial physiological gain of photosynthetic activity, as well as the links between anthocyanins and increased N resorption led us to assume that one primary role of autumn</p>	

anthocyanin could be to protect the photosynthetic apparatus from photo-oxidative damage as light filters rather than as antioxidant. Another major role is to extend carbon capture and help supply the energy needed for N resorption from senescing leaves in both *A. saccharum* and *A. Platanoides* during high-light stress. Nevertheless, photoprotective capacity of anthocyanins was not able to fully compensate for photoinhibitory stress as the anthocyanins are not optimally located to efficiently reduce light within the leaves.

Keywords (separated by '-') Nitrogen resorption - Anthocyanins - Chlorophyll - Photoprotection

Footnote Information Communicated by S. Renault.

Electronic supplementary material The online version of this article (doi:10.1007/s11738-014-1674-x) contains supplementary material, which is available to authorized users.

Metadata of the article that will be visualized in OnlineAlone

Electronic supplementary
material

Below is the link to the electronic supplementary material.

MOESM1: Supplementary material 1 (DOC 43 kb).

Nitrogen resorption in *Acer platanoides* and *Acer saccharum*: influence of light exposure and leaf pigmentation

Baoli Duan · Alain Paquette · Philippe Juneau ·
Jacques Brisson · Bastien Fontaine ·
Frank Berninger

Received: 2 January 2014 / Revised: 4 August 2014 / Accepted: 2 September 2014
© Franciszek Górski Institute of Plant Physiology, Polish Academy of Sciences, Kraków 2014

Abstract We investigated the effects of leaf color change in the fall on photosynthetic production and nitrogen resorption. Seedlings of *Acer platanoides* L. and *A. saccharum* Marsh. were grown in a shade house for 5 months in either 21 % (intermediate light, M) or 4.9 % (low light, L) of incident irradiance. After this period, a subset of the intermediate-light grown seedlings was transferred to a high-light stress treatment (H). Gas exchange, chlorophyll fluorescence, pigments, antioxidant activity, and nitrogen

(N) resorption were examined at three leaf senescence stages during September and October. Our results show that plants of both species produce more anthocyanins in the H treatment. In comparison with plants grown in the L and M treatments, plants of both species in the H treatments had lower chlorophyll, carotenoid and chlorophyll fluorescence parameters (F_v/F_m , Φ_{PSII} , NPQ and ETR) at the third sampling date (October 12–18), and indicating higher levels of photoinhibition in the seedlings exposed to high light. Our results imply that autumn leaf redness is inducible and closely linked to photo-oxidative stress. However, anthocyanins did not enhance antioxidant capacity in red leaves in either species, when exposed to high light. For both species, our results showed a higher N-resorption for high-light stressed plants. We also observed that the number of abscised leaves at the second sampling dates (September 10) was higher than at the third sampling dates. The intra-leaf distribution of anthocyanin, the association between anthocyanin production and the high-light environments, the retention of red leaves, the substantial physiological gain of photosynthetic activity, as well as the links between anthocyanins and increased N resorption led us to assume that one primary role of autumn anthocyanin could be to protect the photosynthetic apparatus from photo-oxidative damage as light filters rather than as antioxidant. Another major role is to extend carbon capture and help supply the energy needed for N resorption from senescing leaves in both *A. saccharum* and *A. Platanoides* during high-light stress. Nevertheless, photoprotective capacity of anthocyanins was not able to fully compensate for photoinhibitory stress as the anthocyanins are not optimally located to efficiently reduce light within the leaves.

Keywords Nitrogen resorption · Anthocyanins · Chlorophyll · Photoprotection

Communicated by S. Renault.

Electronic supplementary material The online version of this article (doi:10.1007/s11738-014-1674-x) contains supplementary material, which is available to authorized users.

B. Duan · P. Juneau · F. Berninger
Département des sciences biologiques, CP 8888 Succ,
Centre-ville, Université du Québec à Montréal, Montréal,
QC H3C 3P8, Canada
B. Duan (✉)
Key Laboratory of Mountain Surface Processes and Ecological
Regulation, Institute of Mountain Hazards and Environment,
Chinese Academy of Sciences, Chengdu 610041, China
e-mail: duanbl@imde.ac.cn

A. Paquette · J. Brisson · B. Fontaine · F. Berninger
Center for Forest Research (CFR), Université du Québec à
Montréal, Montréal, QC H3C 3P8, Canada

J. Brisson · B. Fontaine
Institut de recherche en biologie végétale, Département de
sciences biologiques, Université de Montréal, 4101 East,
Sherbrooke St, Montréal, QC H1X 2B2, Canada

Present Address:
F. Berninger
Department of Forest Sciences, University of Helsinki, PL 27,
00014 Helsinki, Finland

Abbreviations

A_{sat}	Photosynthetic rates at saturating irradiance
Chl	Chlorophyll
Φ_{PSII}	Effective PSII quantum yield
ETR	Electron transport rate
F_v/F_m	Maximal PSII quantum yield
IC ₅₀	Free radical scavenging activity
NPQ	Non-photochemical quenching
qP	Coefficient of photochemical quenching

Introduction

While the autumn coloration of tree foliage remains a fascinating spectacle every year, the mechanisms and reasons for temperate and boreal deciduous trees fall coloration are still subject to discussion. At present, there seem to be two main hypotheses, focusing on the role of anthocyanins in photoprotection and on coloration as a signal to herbivores that the tree is not a suitable host (Archetti 2009). It has been reported that some plants up-regulate anthocyanins to protect themselves from photoinhibition by reducing excess excitation energy and avoid oxidative damage (Feild et al. 2001; Hughes et al. 2005). Although there is experimental evidence for a photoprotective role of anthocyanins in many plants, there seems to be cases where anthocyanins do not improve photoprotection (Esteban et al. 2008; Zeliou et al. 2009). The reasons for these conflicting results remain unclear.

Nitrogen (N) remobilization from senescing leaves during the autumn is an important plant nutrient conservation mechanism in temperate deciduous forests and resorption efficiencies in deciduous forests are above 50 % (Vergutz et al. 2012). Changes in irradiance can modify N resorption, which requires energy (Field 1983) supplied by photosynthesis (Yasumura et al. 2005). However, it seems N resorption response to different light environments does not show a consistent pattern. Chapin and Moilanen (1991), and May and Killingbeck (1992) found that shading of senescing leaves dramatically reduced resorption efficiency in birches (*Betula papyrifera*) and oaks (*Quercus ilicifolia*). By contrast, Yasumura et al. (2005) reported that growth irradiance did not influence N resorption efficiency in three deciduous woody species (*Fagus crenata*, *Lindera umbellata* and *Magnolia salicifolia*). High irradiances especially, combined with low temperature, are harmful to plant photosynthetic capacity and can ultimately result in photoinhibition and photodamage (Pietrini et al. 2002), which reduces N resorption (Hoch et al. 2003). The resorption protection hypothesis (Hoch et al. 2003) states that anthocyanins of senescing foliage shield photosynthetic

tissues from light stress and enhance nutrient resorption. However, this hypothesis, confirmed in some studies (Hoch et al. 2003; Lee et al. 2003), was rejected in other studies (Feild et al. 2001). The resorption protection role of anthocyanins, therefore, requires further investigation.

Lev-Yadun and Holopainen (2009) discussed that while yellow autumn colors prevail in Europe, reds seem to be more important in Eastern Asia and North America. They argued that the reason for their difference in autumn colors could be a product of adaptation to past climates and herbivore faunas. *Acer platanoides* L. (Norway maple), is a Eurasian tree species that was introduced in North America, while *A. saccharum* Marsh. (Sugar maple) is a widespread native. *A. saccharum* usually has a flaming orange autumn color while the color of *A. Platanoides* is normally yellow. In this study, we characterized the relationship between anthocyanins, leaf senescence, photosynthesis and nutrient resorption during autumn for *A. Platanoides* and *A. saccharum*. During the experiment, trees were exposed to different light levels to induce photoinhibitory stress. Specifically, we hypothesized that nitrogen resorption is less efficient in stressed plants (high light < low light < intermediate light).

Materials and methods

Experimental design and treatments

The experiment was conducted at the Montreal Botanical Garden, Quebec, Canada (45°33.7'N, 073°34.3'W). Results from the same experimental setup on the competitive performance of the two species are reported in Paquette et al. (2012) identifying species' characteristics that would indicate invasiveness. *Acer Platanoides* (Norway maple) and *A. saccharum* (sugar maple) seedlings were raised from seeds collected from mature trees in Montreal for the former, and from the Québec provincial forest nursery for the latter. Seeds were stratified and then sown in humid sand boxes filled with layers of sand and minced leaf litter. Germinated seeds were transferred to 320-mL multi-cell containers and placed at random in their respective light regime for 2 months, at which time they were transferred to larger 6.7 L pots. Germinated seedlings were raised in dynamic shade houses under two light levels: 21 % (intermediate light, M treatment) and 4.9 % (low light, L treatment) of full incident photosynthetic photon flux density (PPFD) (measured on September 9, 2009), mimicking conditions found under forest gaps and closed forest understories, respectively, (see Paquette et al. 2012 for experimental setup and light measurement details). These light levels were obtained by varying the size of roof openings and calibrated using whole-day PPFD

measurements. In a previous experiment, Paquette et al. (2010) demonstrated the inadequacy of homogenous shade-cloth greenhouses for mimicking forest understories. Maximum incident irradiance during a sunny day in September is around $1,600 \mu\text{mol m}^{-2} \text{s}^{-1}$ in Montreal. All seedlings were arranged into four replicated blocks, each comprising the two light treatments assigned at random, and the two species. Thus, each block is then a replicate of the light treatment.

All seedlings were well watered throughout the experiment and fertilized using 15 g Nutricote 20-7-10 type 180 per pot. On August 25, 2009 (5 months after leaf emergence), we took a total of 48 seedlings from the larger experiment and assigned them to the present study on leaf redness. Two seedlings per species and per original light treatment (M and L) were chosen from each of the blocks to be part of the present study. These 32 seedlings remained in their original location for the present experiment. To make our high-light treatment (H), an additional 16 seedlings, 8 per species (two from each block), were taken and moved out from the M treatment (21 % of incident PPFD) and placed about a meter away from the eastern walls of their respective shadehouse block, under high-light conditions (~ 86 % of full sunlight) but still protected against dominant winds. We compared trait means for plants that remained in the low- and intermediate-light treatment with those switched from the intermediate light to the high-light treatment to determine whether high-light stress induce anthocyanin. The experiment was a factorial design of two species and three light environments. There were eight seedlings per species and per light treatment, which were spatially arranged into four replication blocks (two seedlings per treatment in each block). Measurements of gas exchange, chlorophyll fluorescence imaging, pigments, antioxidant activity, and nitrogen resorption were performed at three different sampling dates from the end of August to the end of September. At the time of the first sampling period (August 25–September 5), both species maintained their green colors across all light treatments. At the second sampling period (September 10–October 3), *A. Platanoides* and *A. saccharum* still could be seen as greenish for all the three light treatments, despite leaves of both species starting to turn color in high-light treatment. At third sampling dates (October 12–18), leaves of the *A. Platanoides* and *A. saccharum* had turned already to their respective fall colors.

Pigment determination

For each species and treatment, leaves of four to six plants from three blocks were collected at each leaf sampling dates for the determination of pigment concentrations. Leaf discs were sampled on all trees and were immediately

frozen on dry ice in the field, and subsequently stored at -80°C until analysis. Frozen discs were ground in 100 % acetone with a small amount of quartz sand in a chilled mortar. Chlorophylls (Chl) were determined using a multiwavelength analysis at 470, 645, 662 and 710 nm (Lichtenthaler and Buschmann 2001) with a CARY 300 UV–Visible spectrophotometer. For anthocyanin determination, leaf discs were disrupted in liquid nitrogen and extracted in 1.25 mL of 3 M HCl:H₂O:MeOH (1:3:16 by vol.) using a tissue homogenizer. The concentration of anthocyanins was estimated spectrometrically according to Murray and Hackett (1991).

Gas exchange measurements

Photosynthetic rates (A_{sat}) were determined on September 3 and 4, September 29, and October 15, 2009 with a portable photosynthesis system (GFS-3000, Walz, Effeltrich, Germany) at saturating photon flux density ($1,400 \mu\text{mol m}^{-2} \text{s}^{-1}$) and ambient CO₂ concentration (400 ppm). Leaf temperatures (mean $\pm$ SD) were 25.0 ± 2.9 on September 3 and 4, 15.2 ± 0.1 on September 29, and $10.4 \pm 0.8^\circ\text{C}$ on October 15, respectively.

Chlorophyll fluorescence measurements

Fluorescence measurements were carried out with an IMAGING-PAM chlorophyll fluorometer (Heinz Walz GmbH, Effeltrich, Germany). The instrument uses blue LEDs for measuring actinic and saturation pulse light. Leaves were dark adapted for at least 30 min prior to the measurements to completely reoxidize PSII electron transporters. Fluorescence was measured with relatively weak light pulses ($<1 \mu\text{mol m}^{-2} \text{s}^{-1}$) at a low frequency (1 Hz) for measurement of minimal fluorescence (F_0). Maximal fluorescence yield of a dark-adapted leaf (F_m) was measured during an 800-ms exposure to a photon flux of approximately $2,600 \mu\text{mol m}^{-2} \text{s}^{-1}$. Leaves were then illuminated for 9 min with actinic light ($400 \mu\text{mol m}^{-2} \text{s}^{-1}$) to induce electron transport, and saturating pulses were applied to determine maximum fluorescence of light-adapted leaves (F'_m). When performing a measurement, an area of interest (AOI) with a diameter of 1 cm was selected in the center of the leaf. Maximal PSII quantum yield (F_v/F_m , equivalent to $(F_m - F_0)/F_m$), effective PSII quantum yield (Φ_{PSII}), non-photochemical quenching (NPQ) and coefficient of photochemical quenching (qP) were averaged over the AOI. Estimates of Φ_{PSII} , qP, and NPQ were calculated for each irradiance step using the equations of Genty et al. (1989) and Maxwell and Johnson (2000). Rapid light curve measurements were carried out using 30-s exposures to stepwise increased PPFD (1, 24, 54, 103, 265, 532, 599, 831, 1,029, 1,322, 1,617, 2,001, and

2,603 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Simultaneously, apparent electron transport rate (ETR) values were estimated as $\text{ETR} = \Phi_{\text{PSII}} \times 0.5 \times A \times \text{incident PPFD}$, where 0.5 is a factor that assumes equal distribution of energy between the two photosystems (Björkman and Demmig 1987), A is the computed sample absorbance and PPFD is the actinic light intensity.

Antioxidant activity assay

Antioxidant activity of leaf extracts was assessed by determining their ability to scavenge 1,1-diphenyl-2-picrylhydrazyl (DPPH), a stable free radical. Leaf discs were sampled at each sampling dates as above. Extractions were conducted at 4 °C in the dark with acetic acid: water: methanol (7:23:70, v/v/v). Reaction mixtures containing 0–100 μL leaf extract and 1.5 mL of 18 μM DPPH in MeOH were diluted with MeOH to a final volume of 1.6 mL, vortexed, and then held at room temperature for 30 min, after which the absorbance of the mixtures at 517 nm was measured. Antioxidant activity of the leaf extracts was expressed as an effective concentration for radical scavenging (IC_{50}): the concentration of fresh leaf material (mg mL^{-1}) required to produce a 50 % reduction in A_{517} relative to the control mixture to which only methanol was added (van den Berg and Perkins 2007).

Leaf nitrogen analysis and leaf nitrogen in cell walls

To examine foliar nitrogen (N) resorption patterns, we collected fully expanded young to medium-aged green leaves of four to six different individuals of each species. Red leaves that were still attached to branches just before defoliation were sampled when some leaves were fully senescent, falling readily at a touch. Also, shed leaves were counted at intervals of 6 days. Leaf samples were ground and passed through a 20 mesh screen after being first dried at 70 °C for 36 h. The total concentrations of N were determined by the semi-micro Kjeldahl method (Mitchell 1998). For each treatment, N resorption efficiency was calculated as $(N_g - N_s)/N_s \times 100\%$ in which N_g is the green leaf N concentration and N_s is the senescent leaf N concentration (Sanz-Pérez et al. 2009).

Leaf proteins can be divided into water-soluble, SDS-soluble, and SDS-insoluble fractions. The contents of water-soluble, SDS-soluble, and SDS-insoluble fractions were determined as described by Takashima et al. (2004). About 0.3 g frozen leaf disc was powdered in liquid nitrogen in a mortar with a pestle and homogenized in 1 mL of 100 mM Na-phosphate buffer (pH 7.5) system with 0.4 M sorbitol, 2 mM MgCl_2 , 10 mM NaCl, 5 mM iodoacetate, 1 % (v/v) polyvinylpyrrolidone (PVP), 5 mM

phenylmethylsulfonyl fluoride (PMSF), and 5 mM dithiothreitol (DTT), after water-soluble, SDS-soluble, and SDS-insoluble fractions were isolated, the protein content were determined by the method of McGrath (1972). The detergent-insoluble fraction is the protein in cell walls. The ratio of cell wall proteins to total leaf proteins was also calculated. N content in cell walls was calculated from cell wall proteins with a conversion coefficient (0.16 g N g^{-1} wall proteins) (Feng et al. 2009). The proportion of leaf N allocated to cell walls was calculated as N content in cell walls/total leaf N.

Statistical analysis

The experiment consisted of a factorial design of two species, three light levels. There were eight seedlings per species and per light treatment, which were spatially arranged into four replication blocks (two seedlings per treatment in each block). The average of the seedlings within a replicate block was used as the value of a true replicate in the analysis. To meet the requirement of normal distribution, N concentration were log-transformed before analyses. We performed three-way ANOVA for the effects of light, sampling date and species for each variable to discover differences between species in response to light and sampling date. When analyses revealed sampling date and species interactions, or light and species interactions, or light and sampling date and species interactions for certain variables, two-way ANOVAs for light and sampling date were conducted for each species. Significant differences among treatment means were analyzed using Tukey's multiple comparison post hoc tests. In addition, the effects of light, species and their interactions for N resorption were determined using two-way analysis of variance (ANOVA). Simple linear regression was used to determine the relationships between N resorption and anthocyanin levels in leaves. All statistical analyses were conducted in SPSS (SPSS 11.5 for windows, SPSS Inc., Chicago, IL, USA). P values lower than 5 % were considered as statistically significant.

Results

Effects of light on pigment contents

Compared to intermediate light (M) and low light (L), high-light treatment (H) significantly decreased Chl concentrations (by 38–91 %) in *A. Platanoides* at the second and the third sampling dates (Fig. 1a), and increased anthocyanin concentrations (by 53–95 %) across the three sampling dates (Fig. 1b). For *A. saccharum*, high light-

Fig. 1 Effects of light on chlorophyll (Chl) and anthocyanin concentrations, free radical scavenging activity (IC₅₀) and number of red leaves in *A. platanoides* and *A. saccharum*. Treatments: M, intermediate light (21 % of full light); L, low light (4.9 % of full light); H, plants from intermediate light switched to the high-light

environment. For each species, the values not sharing the same letters are significantly different ($p < 0.05$) according to Tukey's test. Each value is the mean of three replicates, consisting of 4–6 seedlings of each species and treatment, error bars are SE. Figure 1d, measured at two census dates (10/12, 10/31) in the 2009 growing season

induced significant reductions in Chl concentrations compared with the L and M treatment at the second and the third sampling period. Meanwhile, in comparison with both the L and M treatment, there was significant increase of the anthocyanin concentrations at the first and the third sampling dates in *A. saccharum* (Fig. 1b). There were significant date $\times$ light interactions in chlorophyll, and the proportion of red leaves (see Supplementary Appendix 1), indicating that high-light stress had a more pronounced effect on these parameters during the third sampling period. Further, there were significant date $\times$ light $\times$ species interactions on anthocyanins. This interaction suggests that high light was associated with significantly higher anthocyanins in *A. saccharum* during the third measurement period. However, neither light nor sampling dates significantly affected IC₅₀ (Fig. 1c). In addition, the percentage of red leaves numbers also increased faster in *A. saccharum* than in *A. Platanoides*, as indicated by the date $\times$ species interaction (see Supplementary Appendix 1, Fig. 1d).

Effects of light on photosynthetic rates and chlorophyll fluorescence parameters

Compared to both the L and M treatment, the H treatment significantly reduced A_{sat} by 62–71 % in *A. Platanoides* and 36–75 % in *A. saccharum* at the first and second sampling dates (Fig. 2a). For both species, at the second and the third sampling dates, F_v/F_m was lower in seedlings grown under H treatment compared with seedlings under L and M treatments (Fig. 2b). The changing tendency of ETR was compatible with that of Φ_{PSII} (Fig. 2c, e). At the third sampling dates, Φ_{PSII} decreased at high-light stress for both species (Fig. 2e). Compared to M treatment, the dynamic changes in Φ_{PSII} and NPQ in *A. Platanoides* under H treatment with photooxidation seem to involve three stages of alteration: at the first sampling stage, both Φ_{PSII} and NPQ decreased; at the second stage, Φ_{PSII} slightly decreased and NPQ increased; and at the third leaf sampling stage, both parameters decreased. ETR increases with the intensity of the actinic light during the rapid light

Fig. 2 Effects of light on net photosynthesis rate (A), maximal PSII quantum yield (F_v/F_m), electron transport rate (ETR), coefficient of photochemical quenching (q_p), effective PSII quantum yield (Φ_{PSII}), and non-photochemical quenching (NPQ) at $1,400 \mu\text{mol m}^{-2} \text{s}^{-1}$ PPF in *A. platanoides* and *A. saccharum*. Treatments: M, intermediate light (21 % of full light); L, low light (4.9 % of full light); H, plants from intermediate light switched to the high-light environment. For each species, the values not sharing the same letters are significantly different ($p < 0.05$) according to Tukey's test. Each value is the mean of two replicates, consisting of four seedlings of each species and treatment, error bars are SE

curves. It is lower (in some cases) in leaves of the H treatment (Fig. 3). For both species, above $500 \mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD, light-dependent ETR in plants under high-light conditions at the third sampling dates was lowest. Across all light treatments, ETR was higher under L and M treatments than under H treatment.

Effects of light on leaf N and N resorption efficiency

There was a tendency for *A. Platanoides* seedlings exposed to high light to have lower leaf N relative to low light and intermediate light at the third sampling dates. For *A. saccharum*, the H treatment significantly decreased leaf N at the second sampling date relative to M treatment, and at the third sampling date relative to L treatment (Fig. 4a). N concentration in cell wall, fraction of leaf N in cell wall and N resorption were similar in

seedlings under L and M treatments (Fig. 4b–d). For both species, the fraction of leaf N in cell wall of seedlings exposed to the high light increased significantly relative to seedlings under M treatment at the third sampling date (Fig. 4c). Furthermore, the leaf N concentration in cell wall and the fraction of leaf N in cell wall also increased more rapidly in *A. saccharum* than in *A. Platanoides*, as indicated by the date $\times$ species interaction (see Supplementary Appendix 1, Fig. 4). N resorption of seedlings under H treatment increased by 42 and 36 % in comparison with seedlings under M treatments in *A. Platanoides* and *A. saccharum*, respectively (Fig. 4d). The two species differed in the amount of N resorption only at low light. In addition, there were positive relationships between N resorption and the anthocyanin content of leaves in *A. saccharum* ($r^2 = 0.49$, $P = 0.04$ for all treatment data pooled together) (Fig. 5).

Fig. 3 Effects of light on the electron transport rate (ETR)-light response curves in *A. platanoides* and *A. saccharum* at the first sampling dates (September 3, **a, b**), the second sampling dates (September 29, **c, d**), and the third sampling dates (October 18, **e, f**). Exposures (30 s) to stepwise increased photon irradiance (1, 24, 54, 103, 265, 532, 599, 831, 1,029, 1,322, 1,617, 2,001, and 2,603 $\mu\text{mol m}^{-2} \text{s}^{-1}$) were provided. Each value is the mean of two replicates, consisting of four seedlings of each species and treatment, error bars are SE. Treatments: M, intermediate light (21 % of full light); L, low light (4.9 % of full light); H, plants from intermediate light switched to the high-light environment

417 Effects of light on number of leaves shedding
418 In both species, the accumulated proportions of leaves shed
419 in late September and October differed among treatments.
420 For both species in all treatments, leaf shed sharply at
421 second sampling dates (September 10–October 3) (Fig. 6).
422 In contrast, at the third sampling dates (October 12–18),
423 both species shed few additional leaves. In addition, L
424 treatment had a lower leaf abscission rate at the second
425 sampling stage, compared with H treatment (Fig. 6).

426 Discussion

427 As expected, significant increases in anthocyanin con-
428 centrations and reductions in chlorophyll concentrations,
429 net photosynthetic rate, maximal PSII quantum yield (F_v/F_m),
430 effective PSII quantum yield (Φ_{PSII}) were observed
431 in high-light stressed plants relative to plants grown in

intermediate light and in low light. Moreover, the N
432 content in the high-light stressed leaves of *A. saccharum*
433 was also lower. These lower N contents could be the
434 reason for the lower photosynthetic rates in the high-light
435 plants (Zeliou et al. 2009), as the majority of leaf N is
436 associated with the photosynthetic function of the leaf
437 (Feng et al. 2009). We also observed that resorption of N
438 was more efficient under light stress. This is contrary to
439 our initial hypothesis that photoinhibition would reduce
440 leaf N re-translocation. On the other hand, during the third
441 measurement period, high light was associated with sig-
442 nificantly higher anthocyanin concentrations in *A. sac-*
443 *charum* compared to *A. Platanoides*, indicated by the
444 significant interaction between light, species and sampling
445 dates on anthocyanin concentrations. Nevertheless, our
446 results fail to explain how the high-light stressed leaves of
447 *A. Platanoides*, with lower anthocyanin concentrations,
448 are as efficient in translocating N as *A. saccharum*. It is
449

Fig. 4 Effects of light on leaf nitrogen content, nitrogen resorption in *A. platanoides* and *A. saccharum*. Treatments: M, intermediate light (21 % of full light); L, low light (4.9 % of full light); H, plants from intermediate light switched to the high-light environment. For each species, values not sharing the same letters are significantly different

($p < 0.05$) according to Tukey's test. In addition, for N resorption values, the asterisks indicate statistically significant differences between the two species within the same light treatment. Each value is the mean of three replicates, consisting of 4–6 seedlings of each species and treatment, error bars are SE

possible that the weaker production of anthocyanins by *A. Platanoides* during autumn senescence is compensated by alternative radical scavenging capacity (van den Berg and Perkins 2007). Anthocyanins are not obligatory for physiological protection, as illustrated by the many temperate trees functioning successfully with yellow autumn leaves (Lev-Yadun and Holopainen 2009).

In our study, the occurrence of leaf redness was coupled to the light environment (Fig. 1b, d), with the reddest leaves (higher anthocyanin concentrations) occurring in the sunniest treatments (Kozłowski and Pallardy 1997). Chlorophyll levels differed considerably between both treatments and senescing developmental stage. These observations indicate that autumnal anthocyanins accumulate and the simultaneous chlorophyll loss is correlated to photoinhibitory environments (Zeliou et al. 2009). A decrease in dark-adapted F_v/F_m is generally used as a measure of photoinhibition (Björkman and Demmig 1987;

Perron and Juneau 2011). During the second and third sampling dates, the rapid decrease in F_v/F_m of plants under high-light conditions compared to plants under intermediate-light conditions also indicates onset of photodamage to PSII. Non-photochemical quenching (NPQ) rose between the first and the second sampling period, indicating increased excess energy dissipation (Lepeduš et al. 2011), but declined during third sampling date, possibly due to oxidative damage in thylakoid membranes and lower rates of linear electron transport (Gielen et al. 2007). We would expect lower ETR in red leaves than green leaves when exposed to the same incident light intensity as the actual light intensity reaching the chloroplasts of a red leaf is lower due to absorption by anthocyanins (Zeliou et al. 2009). This was the case in our study as we observed that ETR at the third sampling periods when leaves turned red was lowest with respect to other sampling dates. The reduction of the Φ_{PSII} , NPQ and ETR explained the

Fig. 5 Relationships between the N resorption and the anthocyanin content of leaves in *A. saccharum* ($r^2 = 0.49$, $P = 0.04$ for all treatment data pooled together). The values are shown for individual replicates

Fig. 6 Cumulative leaf shedding pattern in *A. platanooides* and *A. saccharum* at the first sampling dates (August 25–September 5), the second sampling dates (September 10–October 3) and the third sampling dates (October 12–18). Treatments: M, intermediate light (21 % of full light); L, low light (4.9 % of full light); H, plants from intermediate light switched to the high-light environment. Each value is the mean of three plants, error bars are SE

reduction in the CO_2 fixation as suggested by Lu and Zhang (1998). We found that exposure to high light speeds up the senescence process, as suggested by the decreased F_v/F_m (Fig. 2b) and the increased accumulation of anthocyanins. This further indicates that autumn leaf redness is inducible and closely linked to photo-oxidative stress, and points to a higher need for the buildup of a photoprotection system. During the third measurement period, CO_2 assimilation was low, while ETR was at reasonably high values, especially in *A. saccharum*. It maybe that high-light stress during leaf senescence stimulates the partitioning of electron flow to pathways other than CO_2 assimilation (Park et al. 1996). These findings suggest that photosynthesis under high-light conditions is limited by the electron utilization capacity, not by the electron transport capacity (Fujiki et al. 2007). This could be due to the low leaf internal conductance in red leaves (Miyazawa and Yahata 2006). Lower leaf internal conductance leads to lower CO_2 concentration at the site of RuBP carboxylation and oxygenation, which results in more electrons being used for RuBP oxygenation (Miyazawa and Yahata 2006), and hence low CO_2 assimilation while still relatively high ETR. Additionally, we observed the leaves of *A. saccharum* exhibited slower leaf senescence under high light than did *A. Platanoides*, as shown by smaller decreases in chlorophyll, Φ_{PSII} , and ETR which might be due to *A. saccharum* seedlings having bigger amounts of anthocyanins.

The role of anthocyanin as antioxidants or as light screens should depend on their localization within plant

tissue (Gould et al. 2002). Although the leaf anatomy in the test plants was not examined, it has been reported that anthocyanins accumulation in maples is mainly in the palisade mesophyll (Lee et al. 2003; van den Berg et al. 2009). This anthocyanins distribution pattern supports a role for light screening. We also observed that *A. saccharum* at greater risk of photoinhibition (low leaf N and limited photosynthetic capacity) during leaf senescence have higher anthocyanin, a pattern also supports the light screen hypothesis. Nevertheless, it appears that in spite of a large amount of de novo anthocyanin synthesis in senescing leaves, photoinhibition occurs in red and yellow maple leaves, when they are exposed to high-light levels. These results indicate that though anthocyanins may function as light screening, the anthocyanins in senescing maple leaves do not efficiently reduce light within the leaves as proposed

by van den Berg et al. (2009) that anthocyanins are not optimal internal light filters. The relative small anthocyanin concentrations, and anthocyanins present in the palisade mesophyll in maples (Ishikura 1973; Lee et al. 2003) than multiple locations throughout the palisade and spongy mesophyll in other wood species (Gould et al. 2002; Hughes et al. 2005; van den Berg et al. 2009) could explain that anthocyanins do not provide a physiologically significant level of photoprotection in maple leaves tested in our study.

Foliar N concentrations during the first and the second sampling periods were well within the 16.0–23.2 mg g⁻¹ optimal range of Kolb and McCormick (1993) for *A. saccharum*. In contrast, N concentrations during third sampling date averaged only 6.0 mg g⁻¹ under high-light conditions, which is far below the concentrations reported for *A. saccharum* seedlings with N-limited growth (Walters and Reich 1997). The low-N leaves are more vulnerable to photoinhibitory risk (Schaberg et al. 2003), owing to the limited photosynthetic capacity. During the processes of leaf senescence, approximately 69–75 % of the N in green leaves was resorbed in the both species. This represents higher resorption efficiency than the mean (50–52 %) reported for many plants (Chapin and Kedrowski 1983; Aerts 1996), but is consistent with previous measurements of resorption in *Acer rubrum* (Grizzard et al. 1976). As shown in Fig. 4d, N resorption efficiency did not differ significantly among leaves under low light and intermediate light in either species. However, N resorption was enhanced in high light.

We also observed a positive correlation between N resorption and anthocyanin concentrations in *A. saccharum* leaves (Fig. 5). One hypothesis is that anthocyanins protect foliar nutrient resorption by reducing oxidative stress and quenching free radicals sequestered in vacuoles during the chlorophyll degradation (Matile et al. 1999), as free radicals may disturb nitrogen and/or phosphorus resorption from leaves into branches (Lee et al. 2003). Accordingly, the antioxidant activity of high-light stressed leaves would be expected to exceed that of shaded ones. However, we observed that the antioxidant activity of leaves in the different light treatments was equal at all sampling periods. The results further suggest that rather than enhancing antioxidant capacity, anthocyanins may serve as a ‘sunscreen’ from excessive light in senescing leaves and reduce the risk of photo-oxidative damage, thus facilitating nutrient recovery as discussed in previous sections (Feild et al. 2001; Hoch et al. 2003). Nevertheless, we can not rule out the possibility that some other factor associated with high light, might be responsible for the increased nitrogen resorption. The results further suggest alternative strategies may be employed by senescing *A. Platanoides* leaves to compensate for less

photoprotective function by anthocyanins. In addition to photoprotection, light intensity could also influence senescence and anthocyanin accumulation through sugar levels/sugar accumulation (Wingler et al. 2006, 2009), and future experiments should address the connection between leaf sugar levels, senescence regulation and photosynthetic protein degradation.

A recent study by Schaberg et al. (2008) demonstrated a relationship between foliar coloration and leaf retention strength and suggested that the orange-red coloration in *A. saccharum* may allow for an extended period of nutrient and sugar translocation compared with yellow leaves. In the present study, more leaves are shed between the first and second than between the second and third sampling period (Fig. 6), further strengthening the hypothesis that the benefits of anthocyanins may contribute to prolonged retention. Additionally, N concentration in the cell wall was nearly constant across different leaf color periods, which might help to maintaining leaves function (Takashima et al. 2004). We considered two possible explanations for red leaves being retained in high-light stressed plants. First, leaves would be retained as long as they have a positive carbon gain (Ackerly 1999). In September the total irradiance in Montreal is ~74 % of the maximum (July values) and in October it is still ~40 % (calculated from irradiation data from Plattsburgh New York about 70 km south). Therefore, the physiological gain of continued photosynthetic activity can be substantial. By increasing photoprotection and maintaining a photosystem during cold nights, trees could increase the length of their photosynthetically active period by a few weeks. This would concord with the relatively high rates of photosynthesis we still observed in the leaves at the second sampling dates (Fig. 2a). Second, we suggest that red leaves are involved in a conservative function as well, increasing N resorption and mean residence time during the third sampling dates, while other green or yellow leaves in maple trees are mainly involved in a photosynthetic function. Indeed extending the useful life of leaves comes at the risk of leaves freezing and dying before the tree could remove the nutrients therein.

Our results showed a higher N-resorption for high-light stressed plants contrasting to the suggestion of Hoch et al. (2003). *A. saccharum* in our study had greater anthocyanin accumulation in autumn leaves than *A. Platanoides*. Nevertheless, during the fall, *A. saccharum* and *A. Platanoides* only recovered different amounts of N at low light while at high light the two species show no difference in absorption. Additionally, it has been reported that differences in nitrogen allocation strategies between early and late successional species are important for nutrient resorption (Hoch et al. 2001). In this sense, the both maple species tested in our study behave similarly during autumn. The

higher resorption rates in our light-stressed trees are more difficult to explain. It could be that anthocyanins are over-compensating for the light stress or it could be that photosynthetic protection is the main function of anthocyanins and that the facilitation of nutrient resorption is of secondary importance. Yet, we hypothesize that anthocyanins' limited protection against photoinhibition, as several sets of data presented here argue some photoprotection could be assumed in autumn leaves. The association between anthocyanin production and the high-light environments, the prolonged retention of red leaves, the substantial physiological gain of continued photosynthetic activity in September at mid-northern latitudes as well as the links between anthocyanins and increased N resorption led us to assume that autumnal anthocyanins protect senescing foliage from photoinhibitory irradiances (although incomplete), and that leaf redness is a mechanism to squeeze a bit more photosynthesis out of the leaves before winter, allowing for the resorption of critical foliar nutrients to occur.

In conclusion, our results suggested that the one primary role of autumn anthocyanin is to protect the photosynthetic apparatus from photo-oxidative damage as light filters rather than as antioxidant, another major role is to extend carbon capture and help provide the energy needed for N resorption from senescing leaves in both *A. saccharum* and *A. Platanoides* during high-light stress. Nevertheless, we have realized that photoprotective capacity of anthocyanins were not able to compensate fully for the photoinhibitory stress, as the anthocyanins are not optimally located to efficiently reduce light within the leaves (van den Berg et al. 2009). Our results not only confirm earlier reports indicating a possible light screening role for leaf anthocyanins, but the data could also be useful for understanding the relationship between anthocyanins, photosynthesis and nutrient resorption during autumn for *A. Platanoides* and *A. saccharum*.

Author contribution Baoli Duan contributed to all the experimental process, conducting the experiment, dealing with the data and writing the paper. Alain Paquette coordinated the study and carried out the interpretation. Philippe Juneau, Jacques Brisson, Bastien Fontaine, and Frank Berninger mainly contributed to the experimental process. All authors have read and approved the final manuscript and have no conflicts of interest in regard to this research or its funding.

Acknowledgments We gratefully acknowledge funding by the Fonds québécois de la recherche sur la nature et les technologies (FQRNT) and the Youth Talent Team Program of the Institute of Mountain Hazards and Environment, CAS (SDSQB-2012-01). We also thank the Montreal Botanical Garden for providing the space necessary for this experiment.

References

- Ackerly DD (1999) Self-shading, carbon gain and leaf dynamics: a test of alternative optimality models. *Oecologia* 119:300–310
- Aerts R (1996) Nutrient resorption from senescing leaves of perennials: are there general patterns? *J Ecol* 84:597–608
- Archetti M (2009) Classification of hypotheses on the evolution of autumn colours. *Oikos* 118:328–333
- Björkman O, Demmig B (1987) Photon yield of O_2 evolution and chlorophyll fluorescence characteristics at 77 K among vascular plants of diverse origins. *Planta* 170:489–504
- Chapin FS, Kedrowski RA (1983) Seasonal changes in nitrogen and phosphorus fractions and autumn retranslocation in evergreen and deciduous taiga trees. *Ecology* 64:376–391
- Chapin FS, Moilanen L (1991) Nutritional controls over nitrogen and phosphorus resorption from Alaskan birch leaves. *Ecology* 72:709–715
- Esteban R, Fernández-Marín B, Becerril JM, García-Plazaola JI (2008) Photoprotective implications of leaf variegation in *E. dens-canis* L. and *P. officinalis* L. *J Plant Physiol* 165:1255–1263
- Feild TS, Lee DW, Holbrook NM (2001) Why leaves turn red in autumn. The role of anthocyanins in senescing leaves of red-osier dogwood. *Plant Physiol* 127:566–574
- Feng YL, Lei YB, Wang RF, Callaway RM, Valiente-Banuet A et al (2009) Evolutionary tradeoffs for nitrogen allocation to photosynthesis versus cell walls in an invasive plant. *PNAS USA* 106:1853–1856
- Field C (1983) Allocating leaf nitrogen for the maximization of carbon gain: leaf age as a control on the allocation program. *Oecologia* 56:341–347
- Fujiki T, Suzue T, Kimoto H, Saino T (2007) Photosynthetic electron transport in *Dunaliella tertiolecta* (Chlorophyceae) measured by fast repetition rate fluorometry: relation to carbon assimilation. *J Plankton Res* 29:199–208
- Genty B, Briantais J-M, Baker NR (1989) The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica et Biophys Acta* 990:87–92
- Gielen B, Löw M, Deckmyn G, Metzger U, Franck F et al (2007) Chronic ozone exposure affects leaf senescence of adult beech trees: a chlorophyll fluorescence approach. *J Exp Bot* 58:785–795
- Gould KS, Vogelmann TC, Han T, Clearwater MJ (2002) Profiles of photosynthesis within red and green leaves of *Quintinia serrata* A. Cunn *Physi Plant* 116:127–133
- Grizzard T, Henderson GS, Clebsch EEC, Reichle DE (1976) Seasonal nutrient dynamics of foliage and litterfall on Walker Branch, a deciduous forest ecosystem. ORNL/TM-5254, Oak Ridge National Laboratory, Oak Ridge
- Hoch WA, Singsaas EL, McCown BH (2003) Resorption protection. Anthocyanins facilitate nutrient recovery in autumn by shielding leaves from potentially damaging light levels. *Plant Physiol* 133:1296–1305
- Hoch WA, Zeldin EL, McCown BH (2001) Physiological significance of anthocyanins during autumnal leaf senescence. *Tree Physiol* 21:1–8
- Hughes NM, Neufeld HS, Burkey KO (2005) Functional role of anthocyanins in high-light winter leaves of the evergreen herb *Galax urceolata*. *New Phytol* 168:575–587
- Ishikura N (1973) The changes in anthocyanin and chlorophyll content during the autumnal reddening of leaves. *Kumamoto J Sci Biol* 11:43–50
- Kolb TE, McCormick LH (1993) Etiology of sugar maple decline in four Pennsylvania stands. *Can J For Res* 23:2395–2402

- 754 Kozłowski TT, Pallardy SD (1997) Physiology of woody plants. Academic Press, New York
- 755 Lee DW, O' Keefe J, Holbrook NM, Feild TS (2003) Pigment dynamics and autumn leaf senescence in a New England deciduous forest, eastern USA. *Ecol Res* 18:677–694
- 756 Lepeduš H, Gača V, Viljevac M, Kovač S, Fulgosi H, Simić D, Jurković V, Cesar V (2011) Changes in photosynthetic performance and antioxidative strategies during maturation of Norway maple (*Acer platanoides* L.) leaves. *Plant Physiol Bioch* 49:368–376
- 757 Lev-Yadun S, Holopainen JK (2009) Why red-dominated autumn leaves in America and yellow-dominated autumn leaves in Northern Europe? *New Phytol* 183:497–501
- 758 Lichtenthaler HK, Buschmann C (2001) Chlorophylls and carotenoids: measurement and characterization by UV-VIS spectroscopy. Current protocols in food analytical chemistry. Wiley, New York, pp 431–438
- 759 Lu C, Zhang J (1998) Modifications in photosystem II photochemistry in senescent leaves of maize plants. *J Exp Bot* 49:1671–1679
- 760 Matile P, Hörtensteiner S, Thomas H (1999) Chlorophyll degradation. *Annu Rev Plant Physiol Plant Mol Biol* 50:67–95
- 761 Maxwell K, Johnson GN (2000) Chlorophyll fluorescence: a practical guide. *J Exp Bot* 51:659–668
- 762 May JD, Killingbeck KT (1992) Effects of preventing nutrient resorption on plant fitness and foliar nutrient dynamics. *Ecology* 73:1868–1878
- 763 McGrath R (1972) Protein measurement by ninhydrin determination of amino acids released by alkaline hydrolysis. *Anal Chem* 49:95–102
- 764 Mitchell AK (1998) Acclimation of Pacific yew (*Taxus brevifolia*) foliage to sun and shade. *Tree Physiol* 18:749–757
- 765 Miyazawa Y, Yahata H (2006) Is the parameter electron transport rate useful as a predictor of photosynthetic carbon assimilation rate? *Bull Inst Trop Agr Kyushu Univ* 29:39–53
- 766 Murray JR, Hackett WP (1991) Dihydroflavonol reductase activity in relation to differential anthocyanin accumulation in juvenile and mature phase *Hedera helix* L. *Plant Physiol* 97:343–351
- 767 Paquette A, Fontaine B, Berninger F, Dubois K, Lechowicz MJ, Messier C, Posada JM, Valladares F, Brisson J (2012) Norway maple displays greater seasonal growth and phenotypic plasticity to light than native sugar maple. *Tree Physiol* 32:1339–1347
- 768 Paquette A, Fontaine B, Messier C, Brisson J (2010) Homogeneous light regime in shade-house experiment overestimates carbon gains in Norway maple seedlings. *J Hortic For* 2:117–121
- 769 Park YI, Chow WS, Osmond CB, Anderson JM (1996) Electron transport to oxygen mitigates against the photoinactivation of Photosystem II in vivo. *Photosynth Res* 50:23–32
- 770 Perron MC, Juneau P (2011) Effect of endocrine disrupters on photosystem II energy fluxes of green algae and cyanobacteria. *Environ Res* 111:520–529
- 771 Pietrini F, Iannelli MA, Massacci A (2002) Anthocyanin accumulation in the illuminated surface of maize leaves enhances protection from photo-inhibitory risks at low temperature, without further limitation to photosynthesis. *Plant, Cell Environ* 25:1251–1259
- 772 Sanz-Pérez V, Pilar Castro-Díez P, Millard P (2009) Effects of drought and shade on nitrogen cycling in the leaves and canopy of Mediterranean *Quercus* seedlings. *Plant Soil* 316:45–56
- 773 Schaberg PG, Murakami PF, Turner MR, Heitz HK, Hawley GJ (2008) Association of red coloration with senescence of sugar maple leaves in autumn. *Trees* 22:573–578
- 774 Schaberg PG, van den Berg AK, Murakami PF, Shane JB, Donnelly JR (2003) Factors influencing red expression in autumn foliage of sugar maple trees. *Tree Physiol* 23:325–333
- 775 Takashima T, Hikosaka K, Hirose T (2004) Photosynthesis or persistence: nitrogen allocation in leaves of evergreen and deciduous *Quercus* species. *Plant, Cell Environ* 27:1047–1054
- 776 van den Berg AK, Perkins TD (2007) Contribution of anthocyanins to the antioxidant capacity of juvenile and senescing sugar maple (*Acer saccharum*) leaves. *Funct Plant Biol* 34:714–719
- 777 van den Berg AK, Vogelmann TC, Perkins TD (2009) Anthocyanin influence on light absorption within juvenile and senescing sugar maple leaves —do anthocyanins function as photoprotective visible light screens? *Funct Plant Biol* 36:793–800
- 778 Vergutz L, Manzoni S, Porporato A, Novais RF, Jackson RB (2012) Global resorption efficiencies and concentrations of carbon and nutrients in leaves of terrestrial plants. *Ecol Monographs* 82:205–220
- 779 Walters MB, Reich PB (1997) Growth of *Acer saccharum* seedlings in deeply shaded understories of northern Wisconsin: effects of nitrogen and water availability. *Can J Forest Res* 27:237–247
- 780 Wingler A, Masclaux-Daubresse C, Fischer AM (2009) Sugars, senescence, and ageing in plants and heterotrophic organisms. *J Exp Bot* 60:1063–1066
- 781 Wingler A, Purdy S, MacLean JA, Pourtau N (2006) The role of sugars in integrating environmental signals during the regulation of leaf senescence. *J Exp Bot* 57:391–399
- 782 Yasumura Y, Onoda Y, Hikosaka K, Hirose T (2005) Nitrogen resorption from leaves under different growth irradiance in three deciduous woody species. *Plant Ecol* 178:29–37
- 783 Zeliou K, Manetas Y, Petropoulou Y (2009) Transient winter leaf reddening in *Cistus creticus* characterizes weak (stress-sensitive) individuals, yet anthocyanins cannot alleviate the adverse effects on photosynthesis. *J Exp Bot* 60:3031–3042

Journal : **11738**

Article : **1674**

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details Required	Author's Response
AQ1	The given sentence seems to be incomplete. 'For both species, a125bove ...' Please check for missing words/phrases and complete the sentence.	