<https://helda.helsinki.fi>

Ultra-high performance liquid chromatographic and mass spectrometric analysis of active vitamin B12 in cells of *Propionibacterium* and fermented cereal matrices

Chamlagain, Bhawani

2015

Chamlagain , B , Edelmann , M , Kariluoto , S , Ollilainen , V & Piironen , V 2015 , ' Ultra-high performance liquid chromatographic and mass spectrometric analysis of active vitamin B12 in cells of *Propionibacterium* and fermented cereal matrices ' , Food Chemistry , vol. 166 , pp. 630-638 . <https://doi.org/10.1016/j.foodchem.2014.06.068>

<http://hdl.handle.net/10138/173426>

<https://doi.org/10.1016/j.foodchem.2014.06.068>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Accepted Manuscript

Analytical Methods

Ultra-high performance liquid chromatographic and mass spectrometric analysis of active vitamin B12 in cells of *Propionibacterium* and fermented cereal matrices

Bhawani Chamlagain, Minnamari Edelmann, Susanna Kariluoto, Velimatti Ollilainen, Vieno Piironen

PII: S0308-8146(14)00954-6
DOI: <http://dx.doi.org/10.1016/j.foodchem.2014.06.068>
Reference: FOCH 16010

To appear in: *Food Chemistry*

Received Date: 7 December 2013
Revised Date: 4 June 2014
Accepted Date: 15 June 2014

Please cite this article as: Chamlagain, B., Edelmann, M., Kariluoto, S., Ollilainen, V., Piironen, V., Ultra-high performance liquid chromatographic and mass spectrometric analysis of active vitamin B12 in cells of *Propionibacterium* and fermented cereal matrices, *Food Chemistry* (2014), doi: <http://dx.doi.org/10.1016/j.foodchem.2014.06.068>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Ultra-high performance liquid chromatographic and mass spectrometric**
2 **analysis of active vitamin B12 in cells of *Propionibacterium* and fermented**
3 **cereal matrices**

4
5 Bhawani Chamlagain*, Minnamari Edelmann, Susanna Kariluoto, Velimatti
6 Ollilainen, Vieno Piironen

7
8
9 Department of Food and Environmental Sciences, P.O. Box 27, University of
10 Helsinki, FI-00014 University of Helsinki, Finland

11
12
13
14
15 *Corresponding author. Tel.: +358 9 19158249; fax: +358 9 19158475; E-mail:
16 bhawani.chamlagain@helsinki.fi

26 **ABSTRACT**

27 A sensitive and selective method is needed to analyse *in situ* produced vitamin
28 B12 in plant-based materials, potential new dietary sources of vitamin B12. A
29 UHPLC/UV method was developed and validated for the determination of human
30 active vitamin B12 in cell extracts of *Propionibacterium freudenreichii* subsp.
31 *shermanii* and after immunoaffinity purification in extracts of cereal matrices
32 fermented by *P. freudenreichii*. An Acquity HSS T3 C18 column resulted in a
33 baseline separation, a calibration curve of excellent linearity and a low limit of
34 detection (0.075 ng/5 µL injection). As confirmed by UHPLC-MS, the active
35 vitamin B12 could be separated from pseudovitamin B12. The recovery of
36 vitamin B12 from purified spiked cereal matrices was good (> 90%; RSD < 5%).
37 A nutritionally relevant amount of active vitamin B12 was produced by *P.*
38 *freudenreichii* in cereal malt matrices (up to 1.9 µg/100 g) in 24 h at 28 °C.

39

40

41 **Keywords:** Vitamin B12; *Propionibacterium freudenreichii*; fermented cereal
42 matrix; UHPLC; UPLC; immunoaffinity purification

43

44

45

46

47

48

49

50

51 1. Introduction

52

53 Vitamin B12, one of the water-soluble vitamins, is needed in humans and
 54 animals for the metabolic functions of two enzymes: methylmalonyl-CoA mutase
 55 and methionine synthase. According to Institute of Medicine (1998), the
 56 recommended dietary allowance (RDA) of vitamin B12 for adults is 2.4 µg/day. It
 57 is only synthesized by a few bacteria and archaea (Martens, Barg, Warren, & Jahn,
 58 2002). The structure of vitamin B12 consists of a corrin ring with a central cobalt
 59 atom complexed to four pyrrole rings, a lower α -ligand and an upper β -ligand.
 60 This vitamin includes three naturally occurring forms (adenosylcobalamin,
 61 methylcobalamin and hydroxycobalamin) and one chemically transformed form
 62 (cyanocobalamin), differing only in the upper ligand. Cyanocobalamin, the most
 63 stable form, is obtained by reacting natural vitamers with cyanide and it is the
 64 major vitamin B12 used in fortified foods, nutritional formula and pharmaceutical
 65 preparations (Ball, 2006). The lower ligand of vitamin B12 is 5,6-
 66 dimethylbenzimidazole (DMBI), which is essential for the binding of the vitamin
 67 to the intrinsic factor for its absorption (Nielsen, Rasmussen, Andersen, Nexø, &
 68 Moestrup, 2012). Corrinoids with lower ligands other than DMBI are found in
 69 biological materials (Watanabe, Yabuta, Tanioka, & Bito, 2013). They are
 70 biologically inactive in humans but have vitamin activity in microorganisms
 71 (Herbert, 1988).

72 Animal-based foods such as meat, fish, eggs and dairy products are the
 73 major contributors of vitamin B12 in the human diet. It is accumulated in animal
 74 tissues or milk, primarily as a result of the gut microbial activities and feed
 75 supplements. However, plant food materials are devoid of vitamin B12 (Ball,

2006; Herbert, 1988). A low level of vitamin B12 detected in some plant products and mushrooms probably originates from contaminating organisms or from vitamin uptake from soil residues (Koyyalamudi, Jeong, Cho, & Pang, 2009). Processed plant-based products (e.g. Tempe) have been found to contain a low level of vitamin B12 synthesized by the contaminating bacteria (Herbert, 1988; Keuth & Bisping, 1994; Mo et al., 2013). However, most of these data were obtained by a non-specific analytical method, i.e. microbiological assay (MBA), which does not differentiate between active forms of vitamin B12 and other corrinoids.

The analysis of vitamin B12 in non-fortified foods is challenging due to its low concentration. The typical contents in animal-based foods milk, cheese, meat and liver have been reported to be 0.9, 2.4, 1–2, and 83 µg/100 g, respectively (Ball, 2006). Natural forms of vitamin B12 are sensitive to light, therefore analytical methods utilise cyanide during the extraction to convert the natural forms into the more stable cyanocobalamin. Vitamin B12 in foods is traditionally determined by MBA using *Lactobacillus delbrueckii* ATCC 7830 as an assay organism, which is also the reference analytical method of AOAC (2006). Although it is a sensitive technique, MBA suffers from poor selectivity and often results in overestimation by 5–30% since the test organism can react to compounds other than vitamin B12 forms, such as analogues and nucleic acids (Ball, 2006). One such inactive corrinoid for humans has been identified as pseudovitamin B12, which is produced by *Lactobacillus reuteri* CRL1098 (Santos et al., 2007). This corrinoid has adenine instead of DMBI as the lower ligand. In meat products, vitamin B12 contents were up to 2.2-fold higher with MBA compared with high performance liquid chromatography (HPLC; Guggisberg,

101 Risse, & Hadorn, 2012), reflecting the poor specificity of MBA. Moreover, MBA
102 is labour intensive and time consuming.

103 HPLC based methods have been developed for the determination of vitamin
104 B12 in fortified foods, infant formula and vitamin supplements (Campos-Giménez,
105 Fontannaz, Trisconi, Kilinc, Gimenez, & Andrieux, 2008; Chen, Wolf,
106 Castanheira, & Sanches-Silva, 2010; Heudi, Kilinç, Fotannaz, & Marley, 2006;
107 Marley, Mackay, & Young, 2009). Some studies have also focused on the
108 quantification of vitamin B12 in non-fortified foods such as meat products
109 (Guggisberg et al., 2012), and milk-based fermented products (Van Wyk & Britz,
110 2010). These HPLC methods were based on UV detection. One HPLC method
111 reported by Pakin, Bergaentzlé, Aoudé-Werner and Hasselmann (2005) used
112 fluorescence detection after vitamin B12 was derivatised into a fluorescent
113 compound, α -ribazole. The latter method was sensitive, but it required lengthy
114 sample preparation time involving extensive sample derivatisation. The
115 introduction of an immunoaffinity sample purification technique has improved the
116 sensitivity of the HPLC measurement in fortified foods and supplements
117 (Campos-Giménez et al., 2008; Heudi et al., 2006; Marley et al., 2009).

118 However, the development of ultra-high performance liquid
119 chromatographic (UHPLC) technology that uses separation column made up of
120 sub-2 μm particles and improved instrumentation has considerably increased the
121 sensitivity and resolution of the chromatographic analysis (Swartz, 2005). To
122 date, one method based on UHPLC with mass spectrometric (MS) detection has
123 been reported for the analysis of vitamin B12 in milk and dairy products (Zironi,
124 Gazzotti, Barbarossa, Devicienti, Scardilli, & Pagliuca, 2013). The enhanced
125 sensitivity offered by the UHPLC technology and the sample clean-up and

126 concentrating potential of vitamin B12-specific commercial immunoaffinity
127 columns have not yet been utilised for analysing vitamin B12 in natural or
128 microbiologically processed foods. Particularly, it is necessary to confirm that the
129 microbiologically synthesized forms are active for humans when biofortification
130 is applied. Up to now, cereal matrices, which account for a major portion of the
131 human diet but contain no vitamin B12, have not been considered for *in situ*
132 enrichment with vitamin B12 using food-grade microorganisms.

133 We are studying *in situ* production of vitamin B12 in cereal-based matrices
134 using food-grade propionibacteria. Therefore, the aim of this study was to develop
135 a sensitive UHPLC/UV method for the analysis of vitamin B12 in cells of
136 *Propionibacterium* and in fermented cereal matrices. The new method was
137 compared with MBA, and the need for immunoaffinity-based sample purification
138 and confirmation of the structure of the synthesized vitamer by UHPLC-MS were
139 explored.

140

141

142 **2. Materials and methods**

143

144 *2.1. Chemicals, materials and reagents*

145 All chemicals and reagents used were of analytical grade. Sodium hydroxide,
146 acetic acid and the vitamin B12 assay medium were obtained from Merck
147 (Darmstadt, Germany); ethanol from Altia (Rajamäki, Finland); acetonitrile
148 (HPLC grade), trifluoroacetic acid (TFA) and sodium cyanide from Sigma-
149 Aldrich (Steinheim, Germany); cyanocobalamin from Supelco (Bellefonte, USA);
150 Tween 80 from Sigma (Aldrich, USA); and α -amylase (EC 232-588-1, A9857-

151 5MU, *Aspergillus oryzae*) was from Sigma-Aldrich (Steinheim, Germany). A
 152 certified reference material BCR 487 (lyophilised pig liver; Institute for Reference
 153 Materials and Measurements, Geel, Belgium) was bought from Sigma-Aldrich.
 154 Water (later, MilliQ water) used for the preparation of reagents and analyses was
 155 obtained from a Milli-Q Plus system (0.22 μm , $\geq 18.2 \text{ M}\Omega \text{ cm}$, Millipore
 156 Corporation, Bedford, MA, USA).

157 Cyanocobalamin stock solution (200 $\mu\text{g/mL}$) was prepared in 25%
 158 ethanol/MilliQ water and the concentration was measured by a spectrophotometer
 159 (Lambda 25 UV/Vis, Perkin Elmer Inc., USA) at 361 nm, as described by Indyk,
 160 Persson, Caselunghe, Moberg, Filonzi and Woollard (2002).

161

162 2.2. Vitamin B12 extraction

163

164 The protocol for vitamin B12 extraction (as cyanocobalamin) from sample
 165 matrices was adopted from Kelleher and Broin (1991) for plasma samples with
 166 some modifications. Cereal matrix (1–5 g), bacterial cell pellet (0.1–0.2 g) or pig
 167 liver (0.1 g) weighed in an extraction tube was vortexed with 10 mL buffer (8.3
 168 mM sodium hydroxide and 20.7 mM acetic acid; pH 4.5) containing 100 μL 1%
 169 sodium cyanide and placed in a boiling water bath for 30 min. The samples were
 170 then ice-cooled and centrifuged (Hermle, Wehingen, Germany) at $6900 \times g$ for 10
 171 min. The supernatant was collected in a fresh tube. The residue pellet was
 172 vortexed once again with 5 mL buffer (pH 6.2, adjusted from the pH 4.5 buffer
 173 with 3% sodium hydroxide) and centrifuged. The supernatants were combined
 174 and the pH was adjusted to 6.2. The extract was then paper filtered ($\varnothing$ 90 mm,
 175 VWR, Leuven, Belgium) and the volume was adjusted to 25 mL with pH 6.2

176 buffer. To ease the filtration of extracts for immunoaffinity purification, the
177 cereal-based samples were treated with 1 mL of α -amylase solution (50 mg/mL)
178 (37 °C; 30 min) to digest the starch before the boiling-water extraction (Marley et
179 al., 2009). All the analytical processes were carried out under subdued light or
180 were protected from direct light. The vitamin B12 content of the cereal matrices
181 was expressed in terms of fresh weight (ng/g matrix) and that of cells per unit
182 volume fermented medium (ng/mL).

183

184 2.3. Immunoaffinity purification for UHPLC quantification

185

186 The sample clean-up in the immunoaffinity column “Easy-Extract®” (R-
187 Biopharma, Glasgow, Scotland) was carried out according to the manufacturer’s
188 instructions. Briefly, the buffer in the immunoaffinity column was drained, after
189 which 10–15 mL of the filtered (0.45 μ m, Pall, Cornwall, UK) sample extract
190 was loaded into the column. The column was washed with 10 mL of MilliQ water
191 and 50 mL of air was applied. Vitamin B12 was then eluted with 3 mL of
192 methanol and the elution was completed with an additional 0.5 mL of methanol.
193 The eluate was evaporated at 50 °C under a stream of nitrogen and the residue
194 was reconstituted in 300 μ L of MilliQ water, which was then syringe filtered (0.2
195 μ m, Pall, Cornwall, UK) to a Waters total recovery UPLC vial.

196

197 2.4. Ultra-high performance liquid chromatography (UHPLC)

198

199 2.4.1. UHPLC system

200

201 The chromatographic analysis was performed on a Waters Acquity UPLC
 202 system (Milford, MA, USA) equipped with a photodiode array detector (PDA;
 203 210–600 nm). In addition, it consisted of a binary solvent manager, autosampler
 204 and column manager. The detection was performed at 361 nm and the absorption
 205 spectra were recorded by the PDA. The autosampler was maintained at 4 °C and
 206 the column was operated at 30 °C. The autosampler injected the sample solutions
 207 (2–15 µL) onto a column via a 20-µL injection loop operated in a partial-loop
 208 mode. Chromatographic data were collected and processed using the Waters
 209 Empower 2 software.

210

211 2.4.2. Method optimisation and validation

212

213 Two reversed-phase C18 columns, high-strength silica T3 (HSS) and
 214 ethylene bridged hybrid (BEH) from Waters (Milford, MA, USA), of similar
 215 dimensions (2.1 x 100 mm; 1.8 µm and 1.7 µm particles, respectively) were
 216 evaluated for the separation of vitamin B12 from sample extracts. A set of flow
 217 rates (0.32–0.50 mL/min) was tested in a linear gradient mode of the mobile
 218 phase consisting of MilliQ water (solvent A) and acetonitrile (solvent B); each
 219 modified with 0.025% TFA. Finally, the following mobile phase gradients
 220 (solvent A: solvent B) were maintained during a 10-min run: 0–0.5 min (95:5);
 221 0.5–5 min (60:40); 5–6 min (60:40); and 6–10 min a linear gradient from 60:40 to
 222 95:5 for equilibration. A calibration curve was produced in each sample set from
 223 six cyanocobalamin standards with the vitamin concentration ranging from
 224 0.015–0.75 ng/µL; each standard solution was injected twice. The
 225 chromatographic method was assessed for the linearity of the calibration curve,

limit of detection (LOD: signal-to-noise ratio, $S/N=3$), and limit of quantitation (LOQ: 3 times the LOD) of vitamin B12 using the cyanocobalamin standard solutions. The resolution (calculated according to the formulae in the US Pharmacopeia; USP) for the separation of cyanocobalamin in sample extracts and the number of theoretical plates (USP) for each column were compared. LOD, resolution and number of theoretical plates were obtained with the Waters Empower 2 software.

2.5. UHPLC-MS/MS analysis

The mobile phase was modified with 0.5% formic acid instead of 0.025% TFA for the UHPLC-MS study to minimise the ion suppression. The MS analysis was carried out in a positive ion mode on an Esquire-LC quadrupole ion trap mass spectrometer with an electrospray ionization (ESI) interface (Bruker Daltonics, Bremen, Germany) and the data were analysed by the LC-MSD Trap version 5.2 (Bruker Daltonics). The scanning was carried out for a range of m/z 900–1400 and the tandem mass spectrometry (MS/MS) was performed for a particular m/z according to the target molecule(s) using helium as the collision gas. The following instrumental settings optimised for cyanocobalamin were used for the MS analysis of cyanocobalamin ($[M+H]^+$ m/z 1356) and pseudovitamin B12 ($[M+H]^+$ m/z 1345) in sample extracts: nebulizer (nitrogen) 50.0 psi, dry gas (nitrogen) 8.0 L/min, dry temperature 300 °C, capillary 4500 V, end plate offset –250 V and trap drive 84.

2.6. Microbiological assay

251

252 The MBA was performed on a 96–well microtiter plate (Corning, NY, USA)
 253 based on Kelleher and Broin (1991) using *L. delbrueckii* ATCC 7830 as an assay
 254 organism and cyanocobalamin as a calibrant. The sample extracts were
 255 appropriately diluted at two levels with the extraction buffer (pH 6.2). A certified
 256 reference material BCR 487 (pig liver) extract served as a reference sample in
 257 each set of samples. A blank extract was used to obtain the actual readings of the
 258 samples. Cyanocobalamin solutions of increasing concentration (0–8 pg/well) and
 259 sample extracts, both 100 µL, were inoculated into the wells of the microtiter plate
 260 (4 wells for each concentration), and 200 µL of freshly prepared sterile filtered
 261 vitamin B12 assay medium (pH 6.2) inoculated with the cryopreserved assay
 262 organism was introduced into each well. The plate was then incubated at an
 263 optimised condition (35 °C; 19 h) and the turbidity was measured with a micro-
 264 plate reader (Multiskan EX; Labsystems, Finland) at 595 nm. A calibration curve
 265 of 8 concentration levels and the amount of vitamin B12 in each well were
 266 obtained automatically by Ascent software version 2.6 (Labsystems).

267

268 2.7. Recovery study

269

270 Since natural cereal matrices do not contain vitamin B12, the unfermented
 271 rye malt matrix (from Section 2.8.) was used as a blank matrix for the recovery
 272 study. The sample matrix (1.5 g) was spiked with cyanocobalamin at two levels
 273 (17.7 ng and 53.1 ng), each level in triplicate, and vitamin B12 was extracted and
 274 analysed by MBA directly and by UHPLC after immunoaffinity purification, as
 275 explained earlier. The repeatability was shown by carrying out two independent

276 experiments on two days. To test the effect of the sample matrix, the recovery was
277 also studied in the extraction buffer with the lower level of spiking (17.7 ng). The
278 recovery was calculated from the measured cyanocobalamin concentration in the
279 spiked samples to the concentration of added cyanocobalamin in the samples.

280

281 2.8. Bacterial culture and fermentation of sample matrices

282

283 Propionibacteria capable of vitamin B12 synthesis and allowed for use in
284 food production are mostly used in cheese manufacturing and in the industrial
285 production of vitamin B12 with genetically engineered strains (Thierry, Deutsch,
286 Falentin, Dalmaso, Cousin, & Jan, 2011). For this study, a strain of
287 *Propionibacterium freudenreichii* subsp. *shermanii* (*P. freudenreichii* ABM 5378)
288 isolated from a cheese starter culture was used for the production of vitamin B12
289 in the sample matrices.

290 Malt matrices: The fermented cereal samples were produced to study the
291 suitability of the developed method; the fermentation conditions used were not
292 optimised for vitamin B12 production. Rye and barley malt flours obtained by
293 milling malted grains (Laihian Mallas, Laihia, Finland) were separately mixed in
294 boiling MilliQ water (33% w/v) and cooked for 2 min. The gelatinised matrices
295 were transferred to glass flasks and then autoclaved at 121 °C for 15 min. The
296 culture preparation and fermentation of the matrices were carried out as described
297 by Kariluoto et al. (2010). Briefly, the cooled matrices were aseptically
298 transferred into pre-sterilised flasks and a culture of *P. freudenreichii* ABM 5378
299 was inoculated. The matrices were allowed to ferment at 28 °C for 24 h. After the
300 incubation, the cell counts (CFU/g) had increased by a log factor in both matrices

301 while pH had decreased from 5.6 to 5.3 in the rye matrix and 5.4 to 5.2 in the
 302 barley matrix. Likewise, an aqueous barley malt extract was fermented with
 303 unidentified strains of bacteria producing lactic and propionic acids and then
 304 concentrated.

305 Supplemented whey permeate (SWP): *P. freudenreichii* ABM 5378 was
 306 grown in the SWP medium anaerobically for 72 h and then aerobically for 96 h at
 307 30 °C (Hugenschmidt, Schwenninger, Gnehm, & Lacroix, 2010). Cheese whey
 308 powder (Valio, Espoo, Finland) was used instead of whey permeate powder, as in
 309 the study of Hugenschmidt et al. (2010), and the medium was supplemented with
 310 5 mg/L cobalt(II) chloride (Sigma-Aldrich). The fermented broth was centrifuged
 311 (12000 × g; 10 min), the resulting cell pellet was re-suspended in 10 mL of pH 7.3
 312 PBS buffer (Oxoid, Hampshire, England) and centrifuged again. The fresh cell
 313 mass obtained was stored at -20 °C until the analysis.

314 2.9. Statistical analysis

315 All the statistical analyses were performed using IBM SPSS 20.0 (IBM
 316 Corporation, NY, USA). Average, standard deviation (SD) and relative standard
 317 deviation (RSD) are reported. One-way analysis of variance (ANOVA) and Tukey
 318 post hoc test were performed to compare between respective UHPLC and MBA
 319 results. A *p* value of < 0.05 was considered statistically significant different.

320

321 3. Results and Discussion

322

323 3.1. UHPLC method optimisation and validation

324

325 Gradient elution of the mobile phase consisting of water and acetonitrile
326 modified with 0.025% TFA at a flow rate of 0.32 mL/min was found to be
327 optimal for vitamin B12 separation on both the HSS T3 and BEH column. A
328 higher flow rate (0.47 mL/min) resulted in a co-elution of interfering compounds
329 in the sample extracts. Owen, Lee, and Grissom (2011) used the same flow rate
330 and the mobile phase gradient for the separation of physiologic cobalamins from
331 samples of cell culture media by a BEH C18 column.. A sharper cyanocobalamin
332 peak was produced by the HSS column than that by the BEH column both for the
333 standard solution and for the sample extracts (Figure 1). The height of the
334 cyanocobalamin peak for an equal volume of the injected sample was about 1.4-
335 fold on the HSS column compared to that on the BEH column. The number of
336 theoretical plates of the HSS column was greater by a factor of 2.5, as evidenced
337 by a narrower and a sharper peak on the HSS column. Furthermore, a higher USP
338 peak resolution (1.80) was obtained with the HSS column than with the BEH
339 column (1.10) under identical elution conditions (Fig. 1). In addition, co-elution
340 of an interfering compound was observed for the extracts of *P. freudenreichii*
341 cells leading to the tailing of the peak on the BEH column, whereas the
342 cyanocobalamin peak was well separated on the HSS column. Therefore, the HSS
343 T3 C18 column was chosen for the validation and the analysis of vitamin B12 in
344 sample matrices.

345

346 **Fig. 1**

347

348 The retention time of cyanocobalamin was stable during the analyses,
349 irrespective of the sample matrices. The 30-day average retention time was 3.27

min with a narrow change from 3.25 to 3.29 min over the study period. The UHPLC analysis was faster with a total run time of 10 min compared with over 30 min that was required for the HPLC methods (Heudi et al., 2006; Marley et al., 2009). An external calibration curve of excellent linearity ($R^2 > 0.999$) was achieved by injecting a series of cyanocobalamin standard solutions (0.015–0.75 ng/ μ L). The instrumental LOD and LOQ were 0.075 ng/*inj* and 0.225 ng/*inj*, respectively, for a 5 μ L injection. Previous HPLC/UV methods resulted in higher LODs at the larger injection volumes: 1.5 ng/*inj* (50–100 μ L) (Guggisberg et al., 2012; Marley et al., 2009) and 3–3.3 ng/*inj* (100 μ L) (Chen et al., 2010; Heudi et al., 2006). The enhanced sensitivity allowed for the analysis with small samples (< 5 g), unlike the bulk quantity needed for HPLC analysis (Campos-Giménez et al., 2008; Guggisberg et al., 2012); thus minimising the matrix-related interference. Moreover, HPLC methods have mostly been developed for fortified foods and supplements for measuring added cyanocobalamin, at levels greater than those normally found in foods.

Table 1

The accuracy of the UHPLC method was studied by determining the recovery of spiked cyanocobalamin from a rye malt matrix. Low levels were chosen to cover concentrations that are possible in the fermented cereal matrices. The added levels were still too low to be detected with UHPLC without purification of the extracts. On the other hand, the usefulness of the additional purification and enrichment on the immunoaffinity columns was demonstrated by the good recoveries at both levels. The mean inter-day recovery of

375 cyanocobalamin by UHPLC/UV was about 95% with a relative standard deviation
 376 (RSD) of < 5.0%, for both levels (Table 1). The cereal matrix did not affect the
 377 recovery of the cyanocobalamin, as evident from a similar recovery obtained in
 378 the extraction buffer. Equivalent recoveries of spiked concentrations were also
 379 obtained by the MBA ($p > 0.05$; Table 1). Recently, the HPLC determination of
 380 vitamin B12 in complex fortified foods was improved with the application of an
 381 immunoaffinity sample purification and enrichment column (Heudi et al., 2006),
 382 and then successfully validated in further studies (Campos-Giménez et al., 2008;
 383 Marley et al., 2009). In these earlier studies, a good recovery was also obtained
 384 for cyanocobalamin; however, much higher spiking concentrations were used.
 385 Interestingly, Marley et al. (2009) found that the affinity of other vitamin B12
 386 forms to antibodies was lower; e.g. hydroxycobalamin was recovered only up to
 387 75%, suggesting the benefit of converting natural forms into cyanocobalamin for
 388 the accurate determination of vitamin B12.

389 For BCR 487, a vitamin B12 content of 876 ng/g of dm (dry matter) was
 390 obtained by the UHPLC analysis, whereas the MBA resulted in a higher
 391 concentration (1056 ± 80 ng/g of dm) (Table 2). The latter was in agreement with
 392 the certified reference value (1120 ± 90 ng/g of dm). The UHPLC determined
 393 content was 83% of the MBA result ($p < 0.05$; Table 2); the difference could be
 394 due to the non-specificity of MBA. The certified vitamin B12 value of BCR 487
 395 is based on the data obtained with the MBA. To our knowledge, this is the first
 396 report of the chromatographically analysed vitamin B12 content of BCR 487. A
 397 locally lyophilised pig liver sample analysed with HPLC after immunoaffinity
 398 purification had a vitamin B12 content of only $597 \square 656$ ng/g (Guggisberg et al.,
 399 2012). Distribution of corrinoids in pig liver has not been studied in detail.

400 However, Kelly, Gruner, Furlong and Sykes (2006) investigated corrinoids in
401 lamb tissues, including the liver, by HPLC. Contrary to previous reports that
402 corrinoids in the lamb liver were exclusively vitamin B12, these authors found
403 quite a marked fraction of the total corrinoids (up to 30%) as vitamin B12
404 analogues.

405

406 **Table 2**

407

408 *3.2. UHPLC analysis of vitamin B12 in cells and fermented cereal matrices*

409

410 Figure 2 shows example chromatograms with and without the
411 immunoaffinity clean-up for extracts of cells, pig liver, spiked rye malt matrix and
412 fermented rye malt matrix. Without the immunoaffinity purification, a higher
413 chromatographic background was obtained for the pig liver and the fermented
414 malt matrices while a cleaner chromatogram with a well-separated
415 cyanocobalamin peak (retaining at 3.27 min) was observed for the cell extract.
416 Besides the matrix effect on separation, another disadvantage was that the
417 concentration of cyanocobalamin in these extracts was low or below the LOD
418 (0.075 ng/5 μ L injection). The immunoaffinity purified extracts produced a well-
419 separated peak with none or little background interference, demonstrating the
420 usefulness of the immunoaffinity clean-up for the analysis of low levels of
421 vitamin B12 in the complex matrices. Guggisberg et al. (2012) reported its
422 applicability for the analysis of vitamin B12 in meat and meat products.
423 Nevertheless, analysis of the matrices such as microbial cells, containing a higher
424 level of vitamin B12, did not require the immunoaffinity purification. Therefore,

especially with the immunoaffinity purification this method could also be applicable for other matrices. Separate validation for each material however is needed.

Fig. 2

The corrinoid in the cell extract, the immunoaffinity purified fermented malt matrices and the pig liver eluted at the retention time of the cyanocobalamin standard (Fig. 2) and had a PDA spectrum (210–600 nm) identical to that of cyanocobalamin (Fig. 3). Furthermore, the UHPLC-MS/MS analysis revealed that the fragmentation profile of the corrinoid for the m/z of 1356, $[M+H]^+$ of cyanocobalamin, was similar to that of the cyanocobalamin standard, as shown in the example mass spectra (Figs. 4A; 4B). The following major fragmentation products of cyanocobalamin, as explained by Carkeet et al. (2006) and shown in Figure 4A, were observed: m/z 1210 $[M+H-DMBI]^+$; m/z 1124 $[M+H-DMBI-CN-Co]^+$; m/z 997 $[M+H-DMBI-sugar-PO_3]^+$ and m/z 912 $[M+H-DMBI-sugar-PO_3-CN-Co]^+$. These observations confirmed that the lower ligand of the corrinoid in the sample extracts was DMBI as in cyanocobalamin, which is critical for the binding of the B12 vitamer to the intrinsic factor to make it bioavailable for humans (Nielsen, Rasmussen, Andersen, Nexø, & Moestrup, 2012).

Fig. 3

Fig. 4

450 The UHPLC analysis of *P. freudenreichii* cells obtained from the fermented
 451 SWP accounted for a vitamin B12 production of 620 ng/mL. In a study by
 452 Hugenschmidt et al. (2010), 37 screened *P. freudenreichii* strains produced
 453 vitamin B12 of approximately 100 to 2500 ng/mL in SWP with added cobalt and
 454 DMBI, showing an inter-strain diversity of vitamin B12 synthesis. In our study,
 455 the medium was not supplemented with DMBI. Thus, DMBI had to be
 456 synthesized by the bacteria from the substrate in the SWP, which might have
 457 limited the vitamin B12 production.

458 To the best of our knowledge, we are the first to report vitamin B12 *in situ*
 459 production in cereal matrices with propionibacteria and its analysis by UHPLC.
 460 The vitamin B12 content of the *P. freudenreichii* fermented rye and barley malt
 461 matrices measured with the UHPLC was 19.3 ng/g and 15.7 ng/g of fresh weight,
 462 respectively (Table 2). On the other hand, the UHPLC analysis of the fermented
 463 malt extract after the immunoaffinity purification revealed a peak in the
 464 chromatogram at 3.22 min (Fig. 3A), i.e., before the retention time of
 465 cyanocobalamin (3.27 min) from the standard solution and the immunoaffinity
 466 purified pig liver extract (Figs. 3B; 3C). Its PDA spectrum (210–600 nm) was,
 467 however, similar to that of cyanocobalamin. This peak, which was not seen in the
 468 chromatograms of any other analysed samples, was suggested to be of
 469 pseudovitamin B12 with adenine as its lower ligand. It is a corrinoid known to be
 470 present in biological materials (Watanabe et al., 2013) and has been recently
 471 characterised in *L. reuteri* CRL1098 (Santos et al., 2007). When the sample
 472 extract was analysed by MS/MS for an m/z of 1356, $[M+H]^+$ of cyanocobalamin,
 473 no fragment ions characteristic of cyanocobalamin were observed. However, a
 474 fragmentation profile identical to the cyanocobalamin standard was obtained by

the MS/MS for an m/z of 1345, $[M+H]^+$ of pseudovitamin B12 (Fig. 4C: m/z 1210
 $[M+H\text{-adenine}]^+$; m/z 1124 $[M+H\text{-adenine-CN-Co}]^+$; m/z 997
 $[M+H\text{-adenine-sugar-PO}_3]^+$ and m/z 912 $[M+H\text{-adenine-sugar-PO}_3\text{-CN-Co}]^+$.
 The similar fragment ions obtained for the m/z of 1345 with that for the m/z of
 1356 confirmed that the molecule had a structure otherwise similar to
 cyanocobalamin, but a different lower ligand: adenine in place of DMBI.

3.3. Comparison between UHPLC analysis and MBA

The MBA results of the *P. freudenreichii* cells were similar to the UHPLC
 derived results, whereas analysis of the certified reference pig liver (BCR 487)
 resulted in a 17% higher vitamin B12 content by the MBA (Table 2). The vitamin
 B12 contents determined by the UHPLC were 67% and 63% of the MBA results
 for the fermented rye and barley matrices, respectively (Table 2). For the malt
 extract fermented with bacteria producing lactic and propionic acids no detectable
 level of cyanocobalamin was found by the UHPLC method in contrast to 1200
 ng/g of vitamin B12 measured by the MBA. These sample-specific differences of
 the vitamin B12 contents when measured by the MBA or UHPLC method could
 be due to the inherent drawbacks associated with the MBA.

The MBA has been the golden standard method allowing for the
 determination of low levels of vitamin B12 in foods and biological materials.
 However, it has been reported to give higher vitamin B12 results than HPLC in
 biological and fermented materials. The higher results are possible, as the assay
 organism also responds to corrinoids that are inactive for humans, e.g.
 pseudovitamin B12 and nucleic acids (Ball, 2006; Herbert, 1988; Indyk et al.,

2002). These inactive corrinoids were reported in ovine tissues (Kelly et al., 2006) and may also be synthesized by microorganisms (Herbert, 1988), including propionibacteria. Quesada-Chanto et al. (1998) analysed vitamin B12 in the cells of three strains of propionibacteria by an HPLC method and two spectrophotometric methods. Vitamin B12 was only detected in cells of *P. freudenreichii* with HPLC, whereas vitamin B12 analogues with retention time different than authentic vitamin B12 were observed in the cells of other two strains. However, one strain produced authentic vitamin B12 when the culture medium was supplemented with DMBI. On the other hand, some bacteria produce only inactive corrinoids even in the presence of DMBI. *L. reuteri* CRL1098 produced pseudovitamin B12 exclusively in a growth medium supplemented with and without DMBI under anaerobic conditions (Santos et al., 2007).

Campos-Giménez et al. (2008) noticed a 20–30% higher vitamin B12 content in milk-based infant cereals and soy-based fortified formula by MBA compared to HPLC. Poor selectivity of the MBA was further shown in the analysis of spirulina tablets and shellfish where the MBA results were up to 6–8-fold higher than concentrations determined by a specific intrinsic factor based chemiluminescence method (Watanabe, Takenaka, Abe, Tamura, & Nakano, 1998). The majority of the measured vitamin B12 may thus have been inactive corrinoids.

520

521 4. Conclusion

522

523 The developed UHPLC/UV method enabled sensitive and specific
524 quantitation of the vitamin B12 content in microbial cells, and in fermented cereal

525 matrices after purification on immunoaffinity columns. This method allowed for
 526 the selective determination of bioactive vitamin B12, thus eliminating the
 527 possibility of measuring inactive corrinoids as in the MBA. One such analogue,
 528 pseudovitamin B12, was identified and confirmed in a malt extract fermented by
 529 bacteria producing lactic and propionic acids. A nutritionally relevant amount of
 530 active vitamin B12 was produced by *P. freudenreichii* in rye and barley malt
 531 matrices. The LC-MS is a useful tool for the identification of *in situ* produced
 532 bioactive vitamin B12 from inactive corrinoids in food matrices or microbial
 533 materials.

534

535 **Acknowledgements**

536 The authors would like to thank Miikka Olin for his skilful technical
 537 assistance on the UHPLC and MS analyses. Dr. Matti Korhola is acknowledged
 538 for the microbiological analyses. The study was funded by the Research
 539 Foundation of the University of Helsinki; the Finnish Graduate School on Applied
 540 Sciences: Bioengineering, Food & Nutrition, Environment (ABS); and the
 541 Academy of Finland Project No. 257333 (Natural fortification of foods: Microbial
 542 *in situ* synthesis of vitamin B12 and folate in cereal matrix).

543 **References**

- 544 AOAC official method 986.23 (2006). Cobalamin in milk-based infant formula.
 545 AOAC International.
- 546 Ball, G.F.M. (2006). Vitamins in foods: Analysis, bioavailability and stability.
 547 Boca Raton: CRC Press.

- 548 Campos-Giménez, E., Fontannaz, P., Trisconi, M.-J., Kilinc, T., Gimenez, C., &
549 Andrieux, P. (2008). Determination of vitamin B12 in food products by
550 liquid chromatography/UV detection with immunoaffinity extraction: Single-
551 laboratory validation. *Journal of AOAC International*, 91(4), 786-793.
- 552 Carkeet, C., Dueker, S. R., Lango, J., Buchholz, B. A., Miller, J. W., Green, R., et
553 al. (2006). Human vitamin B12 absorption measurement by accelerator mass
554 spectrometry using specifically labeled ^{14}C -cobalamin. *Proceedings of the*
555 *National Academy of Sciences of the United States of America*, 103(15),
556 5694-5699.
- 557 Chen, P., Wolf, W. R., Castanheira, I., & Sanches-Silva, A. (2010). A LC/UV/Vis
558 method for determination of cyanocobalamin (VB12) in multivitamin dietary
559 supplements with on-line sample clean-up. *Analytical Methods*, 2(8), 1171-
560 1175.
- 561 Guggisberg, D., Risse, M. C., & Hadorn, R. (2012). Determination of vitamin B12
562 in meat products by RP-HPLC after enrichment and purification on an
563 immunoaffinity column. *Meat Science*, 90(2), 279-283.
- 564 Heudi, O., Kiliç, T., Fontannaz, P., & Marley, E. (2006). Determination of
565 vitamin B12 in food products and in premixes by reversed-phase high
566 performance liquid chromatography and immunoaffinity extraction. *Journal*
567 *of Chromatography A*, 1101(1-2), 63-68.
- 568 Herbert, V. (1988). Vitamin B12: Plant sources, requirements, and assay.
569 *American Journal of Clinical Nutrition*, 48(3), 852-858.

- 570 Hugenschmidt, S., Schwenninger, S. M., Gnehm, N., & Lacroix, C. (2010).
571 Screening of a natural biodiversity of lactic and propionic acid bacteria for
572 folate and vitamin B12 production in supplemented whey permeate.
573 *International Dairy Journal*, 20(12), 852-857.
- 574 Indyk, H. E., Persson, B. S., Caselunghe, M. C. B., Moberg, A., Filonzi, E. L., &
575 Woollard, D. C. (2002). Determination of vitamin B12 in milk products and
576 selected foods by optical biosensor protein-binding assay: Method
577 comparison. *Journal of AOAC International*, 85(1), 72-81.
- 578 Institute of Medicine (1998). Dietary reference intakes for thiamine, riboflavin,
579 niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline.
580 Washington: The National Academy Press.
- 581 Kariluoto, S., Edelmann, M., Herranen, M., Lampi, A.-M., Shmelev, A.,
582 Salovaara, H., et al. (2010). Production of folate by bacteria isolated from oat
583 bran. *International Journal of Food Microbiology*, 143(1-2), 41-47.
- 584 Kelleher, B. P., & Broin, S. D. O. (1991). Microbiological assay for vitamin B12
585 performed in 96-well microtitre plates. *Journal of Clinical Pathology*, 44(7),
586 592-595.
- 587 Kelly, R. J., Gruner, T. M., Furlong, J. M., & Sykes, A. R. (2006). Analysis of
588 corrinoids in ovine tissues. *Biomedical Chromatography*, 20(8), 806-814.
- 589 Keuth, S., & Bisping, B. (1994). Vitamin B12 production by *Citrobacter freundii*
590 or *Klebsiella pneumoniae* during tempeh fermentation and proof of

- 591 enterotoxin absence by PCR. *Applied and Environmental Microbiology*,
592 60(5), 1495-1499.
- 593 Koyyalamudi, S. R., Jeong, S.-H., Cho, K. Y., & Pang, G. (2009). Vitamin B12 is
594 the active corrinoid produced in cultivated white button mushrooms
595 (*Agaricus bisporus*). *Journal of Agricultural and Food Chemistry*, 57(14),
596 6327-6333.
- 597 Marley, E. C., Mackay, E., & Young, G. (2009). Characterisation of vitamin B12
598 immunoaffinity columns and method development for determination of
599 vitamin B12 in a range of foods, juices and pharmaceutical products using
600 immunoaffinity clean-up and high performance liquid chromatography with
601 UV detection. *Food Additives and Contaminants: Part A* 26(3), 282-288.
- 602 Martens, J.-H., Barg, H., Warren, M. J., & Jahn, D. (2002). Microbial production
603 of vitamin B12. *Applied Microbiology and Biotechnology*, 58(3), 275-285.
- 604 Mo, H., Kariluoto, S., Piironen, V., Zhu, Y., Sanders, M. G., Vincken, J.-P., et al.
605 (2013). Effect of soybean processing on content and bioaccessibility of folate,
606 vitamin B12 and isoflavones in tofu and tempe. *Food Chemistry*, 141(3),
607 2418-2425.
- 608 Nielsen, M. J., Rasmussen, M. R., Andersen, C. B. F., Nexø, E., & Moestrup, S. K.
609 (2012). Vitamin B12 transport from food to the body's cells—a sophisticated,
610 multistep pathway. *Nature Reviews Gastroenterology and Hepatology*, 9(6),
611 345-354.

- 612 Owen, S. C., Lee, M., & Grissom, C. B. (2011). Ultra-performance liquid
613 chromatographic separation and mass spectrometric quantitation of
614 physiologic cobalamins. *Journal of Chromatographic Science*, 49(3), 228-
615 233.
- 616 Pakin, C., Bergaentzlé, M., Aoudé-Werner, D., & Hasselmann, C. (2005). α -
617 Ribazole, a fluorescent marker for the liquid chromatographic determination
618 of vitamin B12 in foodstuffs. *Journal of Chromatography A*, 1081(2), 182-
619 189.
- 620 Quesada-Chanto, A., Schmid-Meyer, A. C., Schroeder, A. G., Fächter, A.,
621 Carvalho-Jonas, M. F., Koehntopp, P. I., et al. (1998). Comparison of
622 methods for determination of vitamin B12 in microbial material.
623 *Biotechnology Techniques*, 12(1), 75-77.
- 624 Santos, F., Vera, J. L., Lamosa, P., de Valdez, G. F., de Vos, W. M., Santos, H., et
625 al. (2007). Pseudovitamin B12 is the corrinoid produced by *Lactobacillus*
626 *reuteri* CRL1098 under anaerobic conditions. *FEBS Letters*, 581(25), 4865-
627 4870.
- 628 Swartz, M. E. (2005). UPLC™: An introduction and review. *Journal of Liquid*
629 *Chromatography and Related Technologies*, 28(7-8), 1253-1263.
- 630 Thierry, A., Deutsch, S.-M., Falentin, H., Dalmaso, M., Cousin, F. J., & Jan, G.
631 (2011). New insights into physiology and metabolism of *Propionibacterium*
632 *freudenreichii*. *International Journal of Food Microbiology*, 149(1), 19-27.

- 633 Van Wyk, J., & Britz, T. J. (2010). A rapid HPLC method for the extraction and
634 quantification of vitamin B12 in dairy products and cultures of
635 *Propionibacterium freudenreichii*. *Dairy Science and Technology*, 90(5),
636 509-520.
- 637 Watanabe, F., Takenaka, S., Abe, K., Tamura, Y., & Nakano, Y. (1998).
638 Comparison of a microbiological assay and a fully automated
639 chemiluminescent system for the determination of vitamin B12 in food.
640 *Journal of Agricultural and Food Chemistry*, 46(4), 1433-1436.
- 641 Watanabe, F., Yabuta, Y., Tanioka, Y., & Bito, T. (2013). Biologically active
642 vitamin B12 compounds in foods for preventing deficiency among
643 vegetarians and elderly subjects. *Journal of Agricultural and Food Chemistry*,
644 61(28), 6769-6775.
- 645 Zironi, E., Gazzotti, T., Barbarossa, A., Devicienti, C., Scardilli, M., & Pagliuca,
646 G. (2013). Development and validation of a method using ultra performance
647 liquid chromatography coupled with tandem mass spectrometry for
648 determination of vitamin B12 concentrations in milk and dairy products.
649 *Journal of Dairy Science*, 96 (5), 2832-2836.

650 **Figure captions:**

651

652 **Fig. 1.** Chromatograms showing the separation characteristics of the HSS T3 and
653 BEH C18 columns (2.1 mm x 100 mm, 1.8 and 1.7 μm particles) for
654 cyanocobalamin from standard solution (0.15 ng/ μL), an immunoaffinity purified
655 extract of BCR 487 (lyophilised pig liver) and an extract of *P. freudenreichii*
656 cells. The injection volume, flow rate, column temperature and mobile phase were
657 identical. 1 = cyanocobalamin peak on HSS column, 2 = cyanocobalamin peak on
658 BEH column.

659

660 **Fig. 2.** Chromatograms of unpurified and immunoaffinity purified extracts of
661 BCR 487 pig liver, cyanocobalamin-spiked control rye malt matrix, fermented rye
662 malt matrix and cells of *P. freudenreichii*. The cyanocobalamin concentration
663 (ng/ μL) of the purified extract of pig liver, spiked matrix, fermented rye malt
664 matrix and cells were 0.166, 0.034, 0.037, and 0.180, respectively. The injection
665 volume was 10 μL . 1 = cyanocobalamin peak (retention time: 3.27 min).

666

667 **Fig. 3.** Chromatograms of immunoaffinity purified extracts of malt extract
668 fermented with bacteria producing lactic and propionic acids (A), *P.*
669 *freudenreichii* fermented rye malt matrix (B) and cyanocobalamin standard
670 solution (C). t_R = retention time.

671

672 **Fig. 4.** UHPLC-MS/MS spectra of an m/z $[\text{M}+\text{H}]^+$ of 1356 for the
673 cyanocobalamin standard (A) and the corrinoid in the immunoaffinity purified
674 extract of the fermented barley malt matrix (B), and of an m/z $[\text{M}+\text{H}]^+$ of 1345

675 (pseudovitamin B12) for immunoaffinity purified malt extract fermented with
676 bacteria producing lactic and propionic acids (C).

677

678

679 **Table captions:**

680

681 **Table 1.** Recovery of cyanocobalamin by UHPLC/UV from immunoaffinity
682 purified extracts of the spiked unfermented rye malt matrix (1.5 g) and the
683 microbiological assay (MBA) of the extracts without purification (n = 3; for both
684 days).

685

686 **Table 2.** Vitamin B12 contents of certified reference material BCR 487
687 (lyophilised pig liver), *P. freudenreichii* fermented cereal matrices (28 °C; 24 h),
688 cell mass from *P. freudenreichii* fermented supplemented whey permeate (SWP)
689 (30 °C; 168 h) and concentrated malt extract fermented with bacteria producing
690 lactic and propionic acids determined by UHPLC and MBA.

Sample	Spiked amount (ng)	Recovery, % (RSD, %)			
		Day 1		Day 2	
		UHPLC	MBA	UHPLC	MBA
	0.0	ND	ND	ND	ND
Rye malt matrix	17.7	97.2 (3.1) ^a	95.0 (0.9) ^a	93.8 (5.3) ^a	98.2 (4.3) ^a
	53.1	96.8 (2.4) ^a	94.8 (3.7) ^a	91.2 (2.5) ^a	96.1 (5.7) ^a
Extraction Buffer	17.7	94.6 (1.0) ^a	92.7 (5.7) ^a	97.4 (1.9)	NA

ND = Not detected

NA = Not analysed

Values with dissimilar superscript letters (a and b) along each row indicate significant difference ($p < 0.05$).

Sample	UHPLC (ng/g)	MBA (ng/g)	UHPLC/MBA, %
BCR 487 pig liver (n = 8)	876 ($\pm$ 49) ^{a*}	1056 ($\pm$ 80) ^{b*}	82.9
Fermented rye malt matrix (n = 2 $\times$ 2) [‡]	19.3 ($\pm$ 1.5) ^a	28.8 ($\pm$ 2.4) ^b	67.0
Fermented barley malt matrix (n = 2 $\times$ 2) [‡]	15.7 ($\pm$ 2.4) ^a	25.0 ($\pm$ 2.3) ^b	62.8
Cell mass (from fermented SWP) (n = 3)	620 ($\pm$ 90) ^{a#} (ng/mL)	650 ($\pm$ 60) ^{a#} (ng/mL)	95.4
Concentrated fermented malt extract (n = 3)	ND	1196 $\pm$ 56	-

Values are means $\pm$ SD

Values with different superscript letters (a and b) along the row indicate significant difference ($p < 0.05$)

*Dry matter basis

[‡]Biological replicates = 2 and analytical replicates = 2

[#]Expressed per unit volume fermented medium

ND = Not detected

SWP = supplemented whey permeate

The certified value for BCR 487 was 1120 $\pm$ 90 ng/g of dry matter.

- A sensitive UHPLC/UV method was developed for the analysis of vitamin B12.
- The method was suitable for analysing microbial cells and fermented cereal matrices.
- Immunoaffinity purification enabled analysis of vitamin B12 in fermented cereal matrices.
- Active vitamin B12 was produced in cells and cereal matrices by *P. freudenreichii*.
- Pseudovitamin B12 could be separated and further identified with UHPLC-MS.