<https://helda.helsinki.fi>

Application of the Red List Index as an indicator of habitat change

Juslen, Aino Kaisa

2016-03

Juslen , A K , Pykälä , J , Kuusela , S , Kaila , L J , Kullberg , J B , Mattila , J A , Muona , J E , Saari , S E & Cardoso , P 2016 , ' Application of the Red List Index as an indicator of habitat change ' , Biodiversity and Conservation , vol. 25 , no. 3 , pp. 569-585 . <https://doi.org/10.1007/s10531-016-1075-0>

<http://hdl.handle.net/10138/173992>

<https://doi.org/10.1007/s10531-016-1075-0>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Application of the Red List Index as an indicator of habitat change

Aino Juslén^{a*}, Juha Pykälä^b, Saija Kuusela^b, Lauri Kaila^a, Jaakko Kullberg^a, Jaakko Mattila^a, Jyrki Muona^a, Sanna Saari^a, Pedro Cardoso^a.

^a Finnish Museum of Natural History, University of Helsinki, P.O. Box 17, FI-00014 University of Helsinki

^b Finnish Environment Institute, P.O. Box 140, FI-00251 Helsinki

Corresponding author:

Aino Juslén: aino.juslen@helsinki.fi; tel. +358 50 3109703, Finnish Museum of Natural History, University of Helsinki, P.O. Box 17, FI-00014 University of Helsinki

Abstract

For the first time ever, the International Union for Conservation of Nature Red List Index for habitat types was calculated for an entire country, Finland. The RLIs were based on species threat assessments from 2000 and 2010 and included habitat definitions for all 10 131 species of 12 organism groups. The RLIs were bootstrapped to track statistically significant changes. The RLI changes of species grouped by habitats were negative for all habitat types except for forests and rural biotopes which showed a stable trend. Trends of beetles and true bugs were positive in rural and forest habitats. Other 16 observed trends of species group and habitat combinations were negative. Several trends observed were in accordance with studies focusing on particular taxa and habitats, and drivers for their change. This study demonstrates the usefulness of the RLI as a tool for observing habitat change based on species threat assessment data.

Keywords: biodiversity indicator, biodiversity loss, habitat, Finland, RLI, threatened species

1. Introduction

The 2010 conference of the parties to the Convention on Biological Diversity (CBD) in Aichi, Japan declared the 2010–2020 decade as a Decade on Biodiversity. Twenty biodiversity targets were set to be met by the year 2020 (Tittensor et al. 2014). Among these, target 12 says “By 2020 the extinction of known threatened species has been prevented and their conservation status, particularly of those most in decline, has been improved and sustained”. The IUCN Red List of Threatened Species is the most widely used information source on the extinction risk of species (Rodrigues et al. 2006; Mace et al. 2008; but see Cardoso et al. 2011, 2012). The IUCN Red List Index (RLI) (Butchart et al. 2004, 2007), which reflects overall changes in IUCN Red List status over time of a group of taxa, was agreed by the parties to the CBD to be used as an overall index of change, to quantify to what extent target 12 is being met.

The RLI uses weight scores based on the Red List status of each of the assessed species. These scores range from 0 (Least Concern) to Extinct/Extinct in the Wild (5). Summing these scores across all species and relating them to the worst-case scenario - all species extinct - gives us an indication of how biodiversity is doing. Importantly, the RLI is based on true improvements or deteriorations in the status of species, i.e. “genuine changes”. It excludes category changes resulting from, e.g., new knowledge (Butchart et al. 2007). The RLI approach helps to develop a better understanding of which taxa, regions or ecosystems are declining or improving. The aim is to provide policy makers, stakeholders, conservation practitioners and the general public with sound knowledge of biodiversity status and change, and tools with which to make informed decisions.

At a global level, the IUCN Red List Index has been calculated for birds (Butchart et al. 2004; Hoffman et al. 2010), mammals (Hoffmann et al. 2010, 2011), amphibians (Hoffman et al. 2010), corals (Butchart et al. 2010), and cycads (The Millenium Development Goals Report 2015). An ongoing project is heading to present a sampled Red List Index (SRLI, Baillie et al. 2008) of plants (Brummitt et al. 2015) and efforts towards a SRLI of butterflies (Lewis and Senior 2011) and Odonata are made (Clausnitzer et al. 2009). At a regional and national level, RLIs or SRLIs have been presented for certain groups (Lopez et al. 2011; Szabo et al. 2012; Moreno Saiz et al. 2015; Woinarski et al. 2015) or multiple species groups (Gärdenfors 2010; Juslén et al. 2013; Rondinini et al. 2014).

A parallel set of criteria was proposed to be applied to ecosystems in lieu of species, with much the same objectives, the IUCN Red List of Ecosystems (RLE, Rodríguez et al. 2011). This has not been widely

adopted as of yet, either at global or regional scales. National assessments of threatened habitat types have been carried out, for example in Finland (Raunio et al. 2008; Kontula and Raunio 2009). Kontula and Raunio (2009) even presented a procedure for assigning IUCN Red List categories for habitat types. However, this assessment has been carried out only once in Finland, and temporal trends cannot be presented as of yet. Until repeated assessments of risk of collapse of particular ecosystem types are available using the Red List of Ecosystems approach, it will not be possible to produce a Red List Index for different ecosystems using the RLE approach. However, as a proxy for ecosystem or habitat change, it is possible to calculate RLIs for sets of species characteristic of particular ecosystem or habitat types. Butchart et al. (2004) has already used such an approach for birds. In practice, any index based on species trends that includes additional information such as habitat types can be used to perceive trends on species groups other than taxonomic. Besides the RLI, we can mention the Living Planet Index (LPI), which is based on population trends of vertebrates from around the world and that has been used in multiple ways, including for quantifying habitat trends (Loh et al. 2005; Collen et al. 2009). The LPI does however require much more information than the RLI, hence its focus on vertebrates.

Here we propose and develop the first national RLI applied to ecosystem level, using Finnish species and their habitats as an example. The approach is intended to complement both the taxon-based RLI and the ecosystem-based RLE, bridging the gap between the two.

2. Material and methods

2.1 Species data

There are approximately 45 000 known species in Finland, and about 21 400 of these had adequate data for threat assessments both in 2000 and 2010 (Rassi et al. 2001, 2010). The present study is based on 10 131 taxa assessed in both years, (Table 1), as we restricted the analyses to species groups well covered in both assessments: beetles (3 384 species), butterflies & moths (below denoted as butterflies) (2 247),

lichens (1 392), vascular plants (1 197), bryophytes (873), true bugs (463), birds (237), polypores (220), mammals (57), dragonflies and damselflies (51) and herptiles (10).

As a part of the method, back casting was used to identify the species with genuine threat category changes. The 2000 Red List categories were adjusted retrospectively based on current information and taxonomy when needed. The RLI calculations include only category changes due to genuine changes in species statuses (Butchart et al. 2007). Back casting was performed already for species groups other than Lepidoptera by Juslén et al. (2013). The reasons for any category change are listed in Rassi et al. 2010 for the species in threatened categories regionally extinct (RE), critically endangered (CR), endangered (EN), vulnerable (VU), near-threatened (NT) and data deficient (DD). The working documentation lists reasons for the Least Concern (LC) species. Any challenging back casting cases were separately discussed with experts of the group in question. Regarding Lepidoptera, LK and JK have made the back casting purposely for the study now presented. Altogether 529 genuine changes were found in the 12 groups studied (Table 1).

2.2. Habitat data

The habitats for species listed in the Finnish Red Data Book (i.e. for those categorized as RE, CR, EN, VU, and NT) were published by Rassi et al. (2010). For LC species we followed the unpublished habitat classification listed at the threat assessment documentation or other working documentation produced by expert groups during two years (except beetles and butterflies, for which no classification was produced previously).

The habitat classification categories were: forests, mires, aquatic habitats, shores, rock outcrops (including erratic boulders), alpine heaths and meadows above tree-level, and rural biotopes and cultural habitats. Definitions of the habitats are given in table 2, and more detailed subcategorizations are published in Rassi et al. 2010. These differ from the standard classifications by IUCN (<http://www.iucnredlist.org/technical-documents/classification-schemes/habitats-classification-scheme-ver3>) in two ways (see also Tables 2 and 5). First, mires were separated from other aquatic habitats due to their exceptional extension in Finland and importance for many Finnish species. Second, marine intertidal and coastal areas were merged due to the difficulty in separating them given the characteristics of Finnish geology and marine hydrology.

The habitat classification for Least Concern beetles and butterflies was conducted in this study. Habitats of the Least Concern species of Coleoptera were based on published sources (Koch 1989a, 1989b, 1992) and checked by Jaakko Mattila and Jyrki Muona. Besides own expertise, we used a database consisting of 670 000 observations of beetles in Finland. This database is not public, but the Finnish Coleoptera Atlas based on the database has been published (The Finnish Expert Group on Coleoptera 2010). The habitats of the least concern species of Lepidoptera were defined by experts Lauri Kaila and Jaakko Kullberg, who also had a database of Lepidoptera of 1.600 000 observations supporting their work (Hyönteistietokanta).

Additionally, a few missing habitats for the other ten groups of organisms were obtained with the help of the Finnish expert groups of species. The whole habitat classification data per species is given in Appendix 1.

Often species occur and establish sustaining populations on several habitat types. Yet, one habitat could always be pointed out by experts as the primary habitat type. This might be the original habitat of the species, for instance, *Thymus serpyllum* is classified to forests, as its original main habitat in Finland is esker forests (Hämet-Ahti et al. 1998), although it nowadays also occurs on sandy riverbanks and sometimes on sandy road banks. Or it might be the habitat where the species occurs in higher abundance. For high-mobility animals, that may occur in different habitats seasonally or during their life cycles, e.g. birds, the primary habitat was the preferential nesting habitat. Habitats of holomethabolic insects were defined according to the larvae preference, as most of their life-cycle is spent on this stage.

2.3. The Red List Index for habitats

Based on the red-list status of species occupying each habitat, we calculated the RLI for habitats.

The RLI value was calculated by multiplying the number of taxa in each red-list category by the category weight (0 for LC, 1 for NT, 2 for VU, 3 for EN, 4 for CR, 5 for RE/EX). These products were summed and then divided by the number of taxa multiplied by the maximum weight 5 ("maximum possible denominator"). To obtain the RLI value, this sum is subtracted from 1. The index value varies between 0 and 1 (Butchart et al. 2007). The lower the value, the closer the set of taxa is heading towards extinction. If the value is 0 all the taxa are (regionally) extinct. If the value is 1 all the taxa are assessed as Least Concern. The

instructions for national and regional use by Bubba et al. (2009) exclude the species that have been assessed as Extinct (EX) in the earlier assessment. We calculated the RLIs including the taxa assessed as Regionally Extinct (RE) in 2000, as some of these taxa were rediscovered in Finland during the observed period (see also Juslén et al. 2013).

2.4. Statistical analysis

We conducted independent analyses with different species groupings by taxon, by habitat and a combination of these. For each group of species in the three groupings we calculated three values: RLI 2000, RLI 2010 and the change between the years (i.e. $RLI_{2010} - RLI_{2000}$). A simple arithmetic analysis would not show whether the group indices were statistically different or the change between the years was significantly different from a null hypothesis of no change. We therefore resampled all the values with non-parametric bootstrapping. For each group, species were randomly sampled with replacement until the original number of species was attained. For each of the 10,000 resampling events the RLI 2000, RLI 2010 and the respective differences were calculated. The confidence limits ($\alpha = 0.05$) of the RLI values per group and year were the 2.5 and 97.5 percentiles of the respective 10,000 randomizations. The change between the years was considered statistically significant if more than 95% of the randomization values had the same sign (either increase or decrease) as the true values. Statistics were performed using the R 3.1.2 statistical environment (R Core Team 2014).

3. Results

The number of taxa in different primary habitats was 4 031 in forests, 513 in mires, 633 in aquatic habitats, 1 257 in shores, 969 in rock outcrops, 411 in alpine heaths and meadows, and 2 317 in rural biotopes (Table 3).

The RLI value for all Finnish species combined was 0.882 in 2000 and 0.879 in 2010. The minor changes observed against Juslén et al. (2013) were due to the inclusion of Lepidoptera in the dataset. The new bootstrap analyses showed that dragonflies, true bugs and beetles were statistically less threatened than the other groups, whose confidence limits mostly overlap (Fig. 1). The RLI changes between the years

were significantly negative for bryophytes, lichens, vascular plants, butterflies and birds and positive for beetles and true bugs (Table 4). Dragonflies, herptiles, mammals and polypores show no significant trend.

Alpine habitats followed by rock outcrops present the most threatened species on average, with aquatic habitats, forests and mires hosting the least threatened (Fig. 2). The RLI changes between the years were significantly negative for all habitat types except forests and rural biotopes, which show no significant trends (Table 5).

Significant RLI trends between 2000 and 2010 were found for 20 combinations of groups of organisms with primary habitats (Table 6; Appendix 2). The trends of beetles and true bugs were positive in rural and forest habitats, otherwise observed trends were all negative. Trends of bryophytes were negative in six habitats and of vascular plants in five. Negative trends were also recovered in two habitats for both birds and lichens, and in one habitat for butterflies. In dragonflies and damselflies, polypores, herptiles and mammals no positive or negative trends were observed (Appendix 2).

4. Discussion

This study demonstrates that it is useful to calculate the RLI for species grouped by habitat, in addition to the usual taxonomic grouping. Several trends were revealed in accordance with published studies focusing on particular taxa and habitats. In general, more negative trends were found, with positive trends being possibly due to the effects of climate warming on several insect species that are expanding northwards. Few scientific papers analyzing reasons for population changes among the Finnish threatened species other than birds exist. Only in one habitat type (forests) several papers focused on recent trends in threatened species were available, such as the simulation study by Fedrowitz et al. (2012) showing continuous decrease of threatened epiphytic lichens. Our main findings, grouped by habitats, are elaborated in the table 5 with likely drivers and references with supporting notes.

We suggest that the habitat-based RLI may show a different, complementary view to the ecosystem-based RLE. Even though some habitats may not be improving, their constituent species may show positive trends

191 due to other factors such as the climate change. The habitat-based RLI clearly bridges the gap between the
192 taxon-based RLI and the RLE.

193 The RLI has been used in multiple ways, usually to evaluate the impact of contrasting policies on the threat
194 status of different taxonomic groups. Hoffmann et al. (2011) used it to attempt to quantify the impact of
195 conservation efforts on the extinction risk of two groups of mammals. Young et al. (2014) quantified the
196 impact of a conservation organisation's programmes on extinction risk of a set of species. Visconti et al.
197 (2015) used the RLI for projecting the likely impact of different policy decisions. Moreno Saiz et al. (2015)
198 tested it as a tool to assess the success of national conservation policies.

200 The latter authors recommended using various indicators as basis for planning regional conservation
201 measures and evaluating their success. However, they also listed several challenges in using and
202 interpreting the RLI. Above all, they recognize it is a summary statistic, which may mask the individual
203 patterns under a global trend. For example, if 10 species increase and 10 decrease in their status the index
204 will reveal the exact same value as if no species change at all, although these are quite different situations.
205 Researchers and stakeholders should therefore always search for individual species that may be at odds
206 with the general trend of the group and try to understand why this might happen. Although this is also
207 verified in the present study, our results show the RLI to be useful for evaluating species trends in different
208 habitat types.

209 As mentioned, besides the RLI other indices can be disaggregated into different groups so that different
210 aspects of biodiversity change can be studied. These might be taxonomic groups (the subject of most RLI
211 studies), habitat types (the subject of this study), or many other. Dividing species into functional groups
212 may be a particularly useful way of using the RLI, as function is related with ecosystem services and thus
213 trends in particular groups may reveal or even precede changes in services, many of them critical for
214 human well-being.

References

- Albrecht A, Mattila K, Rinne V, Rintala T, Söderman G (2010) Hemiptera. In: Rassi P, Hyvärinen E, Juslén A, Mannerkoski I. (eds) The 2010 red list of Finnish species. Ympäristöministeriö and Suomen ympäristökeskus, Helsinki, pp 397–416
- Baillie JEM, Collen B, Amin R, Akçakaya HR, Butchart SHM, Brummitt NA, Meagher TR, Ram M, Hilton-Taylor C, Mace GM (2008) Toward monitoring global biodiversity. *Conservation Letters* 1: 18–26
- Brommer J, Lehikoinen A, Valkama J (2012) The Breeding Ranges of Central European and Arctic Bird Species Move Poleward. *PLoS One* (Article: e43648). DOI: 10.1371/journal.pone.0043648
- Brummitt N, Bachman SP, Aletrari E, Chadburn H, Griffiths-Lee J, Lutz M, Moat J, Rivers MC, Syfert MM, Nic Lughadha EM (2015) The sampled red list index for plants, phase II: ground-truthing specimen-based conservation assessments. *Philos T Roy Soc B* 370 (Article: 20140015). DOI: 10.1098/rstb.2014.0015
- Bubb PJ, Butchart SHM, Collen B, Dublin H, Kapos V, Pollock C, Stuart SN, Vié JC (2009) IUCN Red List index—guidance for national and regional use. IUCN, Gland
- Butchart SHM, Stattersfield AJ, Bennun LA, Shutes SM, Akçakaya HR, Baillie JEM, Stuart SN, Hilton-Taylor C, Mace GM (2004) Measuring global trends in the status of biodiversity: Red List Indices for birds. *PLOS Biology* (Article: e383). DOI:10.1371/journal.pbio.0020383
- Butchart SHM, Akçakaya HR, Chanson J, Baillie JEM, Collen B, Quader S, Turner WR, Amin R, Stuart SN, Hilton-Taylor C (2007) Improvements to the Red List index. *PLOS One* (Article: e140). DOI: 10.1371/journal.pone.0000140.
- Butchart SHM et al (2010) Global biodiversity: indicators of recent declines. *Science* 328: 1164–1168

- 246 Cardoso P, Borges PAV, Triantis KA, Ferrández MA, Martín JL (2011) Adapting the IUCN Red List criteria
247 for invertebrates. *Biol Conserv* 144: 2432–2440
248
- 249 Cardoso P, Borges PAV, Triantis KA, Ferrández MA, Martín JL (2012) The underrepresentation and
250 misrepresentation of invertebrates in the IUCN Red List. *Biol Conserv* 149: 147–148
251
- 252 Clausnitzer V et al. (2009) Odonata enter the biodiversity crisis debate: The first global assessment of an
253 insect group. *Biol Conserv* 142: 1864–1869
254
- 255 Collen B, Loh J, McRae L, Whitmee S, Amin R, Baillie JEM (2009) Monitoring change in vertebrate
256 abundance: the Living Planet Index. *Conservation Biology* 23: 317–327
257
- 258 Fedrowitz K, Kuusinen M, Snäll T (2012) Metapopulation dynamics and future persistence of epiphytic
259 cyanolichens in a European boreal forest ecosystem. *J Appl Ecol* 49: 493–502
260
- 261 The Finnish Expert Group on Coleoptera (2010) *The Finnish Expert Group on Coleoptera 2010. Atlas of the*
262 *Beetles of Finland.* <http://koivu.luomus.fi/elaintiede/kovakuoriaiset/catlas1.htm>.
263
- 264 Gärdenfors U (ed.) (2010) *Rödlistade arter i Sverige 2010 - the 2010 red list of Swedish species.*
265 *ArtDatabanken, Uppsala*
266
- 267 Hanski I (2000) Extinction debt and species credit in boreal forests: modelling the consequences of
268 different approaches to biodiversity conservation. *Ann Zool Fenn* 37: 271–280
269
- 270 Hanski I (2005) *The shrinking world: Ecological consequences of habitat loss.* International Ecology
271 Institute, Oldendorf
272
- 273 Hautala H, Laaka-Lindberg S, Vanha-Majamaa I (2011) Effects of retention felling on epixylic species in
274 boreal spruce forests in southern Finland. *Restor Ecol* 19: 418–429

275

276 Hedenås H, Hedström P (2007) Conservation of epiphytic lichens: Significance of remnant aspen (*Populus*
277 *tremula*) trees in clear-cuts. Biol Conserv 135: 388–395

278

279 Heino J, Virtanen R, Vuori KM, Saastamoinen J, Ohtonen A, Muotka T (2005) Spring bryophytes in
280 forested landscapes: land use effects on bryophyte species richness, community structure and persistence.
281 Biol Conserv 124: 539–545

282

283 Hoffman M et al. (2010) The impact of conservation on the status of the world's vertebrates. Science 330:
284 1503–1509

285

286 Hoffman M, Belant JL, Chanson JS, Cox NA, Lamoreux J, Rodrigues ASL, Schipper J, Stuart SN (2011)
287 The changing fates of the world's mammals. Philos T Roy Soc B 366: 2598–2610

288

289 Hyvärinen E, Kouki J, Martikainen P (2006) Fire and green-tree retention in conservation of red-listed and
290 rare deadwood-dependent beetles in Finnish boreal forests. Conserv Biol 20: 1710–1719

291 Hyönteistietokanta (<http://hyonteiset.luomus.fi/insects/main/EntDatabase.html>) [Entomological database].
292 Finnish Museum of Natural History, Helsinki, 27.10.2015

293 Hämet-Ahti L, Suominen J, Ulvinen T, Uotila P (eds.) (1998) Retkeilykasvio, 4. uudistettu painos,
294 Luonnontieteellisen keskusmuseon kasvimuseo, Helsinki.

295

296 Ikonen I (2011) Integrated coastal planning and management in southwest Finland. In: Reihmanis J (ed.)
297 Nordic-Baltic-Belarus solutions in farming for biodiversity. Latvijas Dabas Fonds, Riga, pp 34–41

298

299 Ilmonen J, Mykrä H, Virtanen R, Paasivirta L, Muotka T (2012) Responses of spring macroinvertebrate and
300 bryophyte communities to habitat modification: community composition, species richness, and red-listed
301 species. Freshwater Science 31: 657–667

302

- 303 Johansson P (2008) Consequences of disturbance on epiphytic lichens in boreal and near boreal forests.
304 Biol Conserv 141: 1933–1944
305
- 306 Juslén A, Hyvärinen E, Virtanen LK (2013) Application of the Red-List Index at a national level for multiple
307 species groups. Conserv Biol 27: 398–406
308
- 309 Jutila H (2001) How does grazing by cattle modify the vegetation of coastal grasslands along the Baltic
310 Sea? Ann Bot Fenn 38: 181–200
311
- 312 Koch K (1989a) Die Käfer Mitteleuropas. Ökologie. Band 1. Krefeld, Goecke & Evers
313
- 314 Koch K (1989b) Die Käfer Mitteleuropas. Ökologie. Band 2. Krefeld, Goecke & Evers
315
- 316 Koch K (1992) Die Käfer Mitteleuropas. Ökologie. Band 3. Krefeld, Goecke & Evers
317
- 318 Kontula T, Raunio A (2009) New method and criteria for national assessments of threatened habitat types.
319 Biodivers Conserv 18: 3861–3876
320
- 321 Kouki J, Arnold K, Martikainen P (2004) Long-term persistence of aspen – a key host for many threatened
322 species – is endangered in old-growth conservation areas in Finland. J Nat Conserv 12: 41–52
323
- 324 Laaksonen T, Lehtikoinen A (2013) Population trends in boreal birds: Continuing declines in agricultural,
325 northern, and long-distance migrant species. Biol Conserv 168: 99–107
326
- 327 Lewis OT, Senior MJM (2011) Assessing conservation status and trends for the world's butterflies: the
328 Sampled Red List Index approach. J Insect Conserv 15: 121–128
329

- 330 Loh J, Green RE, Ricketts T, Lamoreux JF, Jenkins M, et al. (2005) The Living Planet Index: using species
331 population time series to track trends in biodiversity. *Philosophical Transactions of the Royal Society of*
332 *London B*, 360: 289-295
- 333
- 334 Lopez L (2011) Estado de las Aves del Paraguay. Guyra Paraguay, BirdLife International. Asunción,
335 Paraguay.
- 336
- 337 Mace GM, Collar NJ, Gaston KJ, Hilton-Taylor C, Akçakaya HR, Leader-Williams N, Milner-Gulland EJ,
338 Stuart SN (2008) Quantification of extinction risk: IUCN's system for classifying threatened species.
339 *Conserv Biol* 22: 1424–1442
- 340
- 341 Martikainen P (2001) Conservation of threatened saproxylic beetles: significance of retained aspen *Populus*
342 *tremula* on clearcut areas. *Ecol Bull* 49: 205–218
- 343
- 344 The Millenium Development Goals Report 2015. United Nations, New York.
- 345
- 346 Moreno Saiz JC, Domínguez Lozano F, Marrero Gómez M, Bañares Baudet Á (2015) Application of the
347 Red List Index for conservation assessment of Spanish vascular plants. *Conserv Biol* 29: 910–919
- 348
- 349 Olofsson J, Oksanen L (2005) Effects of reindeer density on vascular plant diversity on North Scandinavian
350 mountains. *Rangifer* 25: 5–18
- 351
- 352 Penttilä R, Siitonen J, Kuusinen M (2004) Polypore diversity in a managed and old-growth boreal *Picea*
353 *abies* forests in southern Finland. *Biol Conserv* 117: 271–273
- 354
- 355 Pykälä J (2000) Mitigating human effects on European biodiversity through traditional animal husbandry.
356 *Conserv Biol* 14: 705–712
- 357
- 358 Pykälä J (2004) Effects of new forestry practices on rare epiphytic macrolichens. *Conserv Biol* 18: 831–838

359

360 Pöyry J, Luoto M, Heikkinen RK, Kuussaari M, Saarinen K (2009) Species traits explain recent range shifts
361 of Finnish butterflies. *Glob Change Biol* 15: 732–743

362

363 Pöysä H, Rintala J, Lehikoinen A, Väisänen, R.A., 2012. The importance of hunting pressure, habitat
364 preference and life history for populations trends of breeding waterbirds in Finland. *Eur J Wildlife Res* 59:
365 245–256

366

367 Raatikainen KM, Heikkinen RK, Pykälä J (2007) Impacts of local and regional factors on vegetation of
368 boreal semi-natural grasslands. *Plant Ecol* 189: 155–173

369

370 Rassi P, Alanen A, Kanerva T, Mannerkoski I (eds) (2001) Suomen lajien uhanalaisuus 2000.
371 Ympäristöministeriö & Suomen ympäristökeskus, Helsinki

372

373 Rassi P, Hyvärinen E, Juslén A, Mannerkoski I (eds) (2010) Suomen lajien uhanalaisuus—Punainen kirja
374 2010. The 2010 red list of Finnish species. Ympäristöministeriö ja Suomen ympäristökeskus, Helsinki

375

376 Raunio A, Schulman A, Kontula T, (eds) (2008) Assessment of threatened habitat types in Finland – part 1:
377 Results and basis for assessment (in Finnish). *Suomen ympäristö* 8: 1–264

378

379 R Core Team (2014) R: A language and environment for statistical computing. R Foundation for Statistical
380 Computing, Vienna, Austria. Available at <http://www.R-project.org/>

381

382 Reinikainen A, Mäkipää R, Vanha-Majamaa I, Hotanen JP (eds) (2000) Kasvit muuttuvassa
383 metsäluonnossa. Metsäntutkimuslaitos & Kustannusosakeyhtiö Tammi, Helsinki

384

385 Rodrigues ASL, Pilgrim JD, Lamoreux JF, Hoffmann M, Brooks TM (2006) The value of the IUCN Red List
386 for conservation. *Trends Ecol Evol* 21: 71–76

- Rodríguez JP et al (2011) Establishing IUCN Red List criteria for threatened ecosystems. *Conserv Biol* 25: 21–29
- Rondinini, C., Battistoni, A., Teofili, C. 2014. Lo stato della Biodiversità in Italia: l'applicazione dell'approccio Sampled Red List e Red List Index
- Szabo JK, Butchart SHM, Possingham HP, Garnett ST (2012) Adapting global biodiversity indicators to the national scale: a red list index for Australian birds. *Biol Conserv* 148: 61–68
- Tikkanen OP, Martikainen P, Hyvärinen E, Junninen K, Kouki J (2006) Red-listed boreal forest species of Finland: associations with forest structure, tree species, and decaying wood. *Ann Zool Fenn* 43: 373–383
- Tittensor DP et al. (2014) A mid-term analysis of progress toward international biodiversity targets. *Science* 346: 241–244
- Toivanen T, Kotiaho JS (2007) Mimicking natural disturbances of boreal forests: the effects of controlled burning and creating dead wood on beetle diversity. *Biodivers Conserv* 16: 3193–3211
- Virkkala R, Korhonen KT, Haapanen R, Aapala K (2000) Protected forests and mires in forest and mire vegetation zones in Finland based on the 8th National Forest Inventory (In Finnish with an English summary). *The Finnish Environment* 395: 1–49
- Virkkala R, Heikkinen RK, Leikola N, Luoto M (2008) Projected large-scale range reductions of northern-boreal land bird species due to climate change. *Biol Cons* 141: 1343–1353
- Virkkala R, Rajasärkkä A (2011) Climate change affects populations of northern birds in boreal protected areas. *Biological Letters* 7: 395–398
- Virkkala R, Rajasärkkä A (2012) Preserving species populations in the boreal zone in a changing climate: contrasting trends of bird species groups in a protected area network. *Nature Conserv* 3: 1–20

Visconti P et al (2015) Projecting global biodiversity indicators under future development scenarios.
Conserv Lett DOI: 10.1111/conl.12159

Woinarski JCZ, Burbidgec AA, Harrison PL (2015) Ongoing unraveling of a continental fauna: Decline and
extinction of Australian mammals since European settlement. PNAS 112: 4531–4540

Young RP, Hudson MA, Terry AMR, Jones CG, Lewis RE, Tatayah V, Zuël N, Butchart SHM (2014)
Accounting for conservation: Using the IUCN Red List Index to evaluate the impact of a conservation
organization. Biol Conserv 180: 84–96

Figure captions

Figure 1. The RLI trends between 2000 and 2010 showing the confidence limits for RLI values of each
group of organisms.

Figure 2. The RLI trends between 2000 and 2010 showing the confidence limits for RLI values of each
primary habitat.

455

Figure 1.

456

457

458

459

460

461

462

463

464

465

466

467

468

469

Figure 2.

Table 1. Number of species known in Finland (Total) by organism groups included in our study, number of taxa included in the red-list assessment of 2010, number of RE, CR, EN, VU, and DD taxa together in 2010, number of taxa excluded from the study because they were Data Deficient or not assessed in 2000 as not having an established population, number of taxa included in the present study and those that genuinely changed red-list category between 2000 and 2010.

Organism group	Total	Assessed (% total)	RE, CR, EN, VU, NT, or DD (% assessed)	Excluded as Data Deficient or other reasons (% assessed)	Included (% assessed)	Genuinely changed (% included)
Beetles (Coleoptera)	3 697	3 416 (92.4)	737 (21.6)	32 (0.9)	3 384 (99.1)	138 (4.1)
Birds (Aves)	249	241 (96.8)	89 (36.9)	4 (1.7)	237 (98.3)	66 (27.8)
Bryophytes (Bryophyta, Marchantiophyta and Anthoceroophyta)	906	896 (98.9)	364 (40.6)	23 (2.6)	873 (97.4)	35 (4.0)
Butterflies (Lepidoptera)	2 576	2 313 (89.8)	707 (30.6)	66 (2.9)	2 247 (97.1)	130 (5.8)
Dragonflies and damselflies (Odonata)	55	52 (94.5)	1 (1.9)	1 (1.9)	51 (98.1)	1 (2.0)
Herptiles (Reptilia and Amphibia)	12	10 (83.3)	3 (30.0)	0	10 (100)	1 (10.0)
Lichens (Lichenes)	1 832	1 545 (84.3)	686 (44.4)	153 (9.9)	1 392 (90.1)	59 (4.2)
Mammals (Mammalia)	72	59 (81.9)	22 (37.3)	2 (3.4)	57 (96.6)	4 (7.0)
Polypores (Aphyllorphorales and Heterobasidiomycetes)	237	225 (94.9)	95 (42.2)	5 (2.2)	220 (97.8)	9 (4.1)
True bugs (Heteroptera)	506	469 (92.7)	64 (13.6)	6 (1.3)	463 (98.7)	19 (4.1)
Vascular plants (Tracheophyta)	ca. 3 550	1 206 (40.0)	334 (27.7)	9 (0.7)	1 197 (99.3)	67 (5.6)
All species	ca. 13 692	10 432 (76.2)	3 102 (29.7)	304 (2.9)	10 131 (97.1)	529 (5.2)

Table 2. Habitat classification used in Finnish Red Data Books 2001 and 2010 (Rassi et al. 2001; 2010) and corresponding IUCN Habitat classes.

Habitat	Additional explanation	Corresponding IUCN habitat
Alpine	Alpine heaths and meadows above tree-level	Native grassland
Aquatic habitats	Baltic Sea, lakes and ponds, small ponds, rivers, brooks and streams, rapids, spring complexes	Wetlands
Mires	Rich fens, fens, pine mires, spruce mires	Wetlands (subcategory: bogs, marshes, swamps, fens, peatlands)
Forests	Heath forests, herb-rich forests, mountain birch forests	Forests
Rock outcrops	Rock outcrops, including erratic boulders	Inland rocky areas
Rural biotopes and cultural habitats	Seminatural grasslands, wooded pastures and pollard meadows, ditches, arable land, parks, yards, gardens, roadsides, railway embankments, buildings	Artificial
Shores	Shores of the Baltic Sea, lake shores and river banks	Marine/Intertidal and Marine Coastal/Supratidal

Table 3. The number of taxa in different primary habitats used in the study.

Organism group	Alpine heaths and meadows	Aquatic habitats	Forests	Mires	Rock outcrops	Rural biotopes and cultural habitats	Shores	All habitats
Beetles (Coleoptera)	889	285	1559	72	1	33	545	3 384
Birds (Aves)	36	56	78	20	2	20	25	237
Bryophytes (Bryophyta, Marchantiophyta and Anthoceroophyta)	81	83	138	123	269	108	71	873
Butterflies (Lepidoptera)	688	0	1 143	137	27	54	198	2 247
Dragonflies and damselflies (Odonata)	0	46	0	5	0	0	0	51
Herptiles (Reptilia and Amphibia)	1	2	5	0	0	0	2	10
Lichens (Lichenes)	57	3	537	17	600	79	99	1 392
Mammals (Mammalia)	12	7	32	1	0	2	3	57
Polypores (Aphylophorales and Heterobasidiomycetes)	15	0	198	0	0	0	7	220
True bugs (Heteroptera)	191	44	138	9	1	2	78	463
Vascular plants (Tracheophyta)	346	107	203	129	69	114	229	1 197
All species	2 316	633	4 031	513	969	412	1257	10 131

Table 4. The RLI in 2000 and 2010 and respective change in different groups of organisms and statistical significance of this change.

Group	RLI 2000	RLI 2010	Change	p-value
Beetles	0.905	0.909	0.003	<0.001
Birds	0.877	0.854	-0.023	0.012
Bryophytes	0.824	0.816	-0.008	<0.001
Butterflies	0.881	0.878	-0.004	0.005
Dragonflies	0.984	0.988	0.004	0.372
Herptiles	0.859	0.879	0.020	0.342
Lichens	0.840	0.831	-0.009	<0.001
Mammals	0.814	0.807	-0.007	0.224
Polypores	0.846	0.849	0.004	0.144
True bugs	0.945	0.953	0.008	0.001
Vascular Plants	0.894	0.884	-0.010	<0.001

Table 5. The RLI changes between 2000 and 2010 in different primary habitats (Finnish Red Data Book classification and IUCN Habitat classification) and the statistical significance, statistically significant changes in different organism groups and habitat combinations; and the likely drivers behind the RLI trends shown in the study with supporting notes and references.

Table 6. The changes of RLI for 11 groups of organisms in different primary habitats between 2000 and 2010. Statistically significant combinations are marked with asterisks (* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$).

Group	Alpine	Aquatic	Forests	Mires	Rock	Rural	Shores
Beetles	0	-0.003	0.006***	-0.003	0	0.004*	0
Birds	-0.04	-0.028	-0.002	-0.08*	0	0	-0.048*
Bryophytes	-0.02***	-0.012**	-0.007**	-0.008**	-0.003*	0	-0.02***
Butterflies	-0.015	0	-0.001	-0.007	0.007	-0.003	-0.016**
Dragonflies	0	0.004	0	0	0	0	0
Herptiles	0	0	0	0	0	0	0.101
Lichens	-0.003	0	-0.017***	-0.012	-0.004***	-0.007	0
Mammals	0	0	-0.006	0	0	-0.017	0
Polypores	0	0	0.003	0	0	0.013	0
True bugs	0	0.005	0.013***	0	0	0.008*	0
Vascular Plants	-0.019***	-0.006*	-0.006	-0.014***	-0.006	-0.009***	-0.014***

* $p < 0.05$

** $p < 0.01$

*** $p < 0.001$

Appendix 1.

The species included in the study, their main habitats and the IUCN threat classification in 2000 (backcasted) and 2010.

Appendix 2.

RLI values for all combinations of taxonomic groups and habitat types (Appendix) are available online. The authors are solely responsible for the content and functionality of these materials. Queries (other than absence of the material) should be directed to the corresponding author.