<https://helda.helsinki.fi>

Psychological well-being is associated with better nutrient intakes in heterogeneous older populations

Jyväkorpi, S. K.

2017-02

Jyväkorpi , S K , Pitkala , K H , Björkman , M P , Strandberg , T E , Soini , H & Suominen , M
H 2017 , ' Psychological well-being is associated with better nutrient intakes in
heterogeneous older populations ' , European Geriatric Medicine , vol. 8 , no. 1 , pp. 6-7 . <https://doi.org/10.1016/j.eurger.2016.09.007>

<http://hdl.handle.net/10138/236419>

<https://doi.org/10.1016/j.eurger.2016.09.007>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com/en

Letter to the editor

Psychological well-being is associated with better nutrient intakes in heterogeneous older populations

Introduction

Nutrition is essential for the overall well-being of an older individual, and psychological well-being (PWB) is an important dimension of quality of life [1,2]. Nutritional status is known to associate with better PWB among long-term residents with dementia, but less is known about these relationships in general older population [3]. We analyzed PWB, frailty risk, and dietary intake in various groups of older people.

Methods

Our cross-sectional study combined four datasets of nutritional studies: (1) healthy home-dwelling older people who participated in the Nutrition Education and Cooking Class (NC) follow-up study ($n = 54$) [4], (2) older men from the Helsinki Businessmen Study (HBS) ($n = 66$) [5], (3) home-dwelling older people with signs of frailty and screened for the Porvoo Sarcopenia and Nutrition Trial (PSNT) ($n = 179$) [6], and (4) institutionalized older residents of assisted living facilities (ALF) from the Helsinki metropolitan area

($n = 266$) [7]. The recruitment, eligibility and participant characteristics have been reported elsewhere [4–8].

Nutritional status was assessed using the Mini Nutritional Assessment (MNA) [9]. Energy and nutrient intakes were retrieved from 1 to 3 day food diaries. PWB questionnaire included six validated questions about (1) life satisfaction (yes/no), (2) feeling needed (yes/no), (3) having plans for the future (yes/no), (4) having zest for life (yes/no), (5) feeling depressed (seldom or never/sometimes/often or always), and (6) suffering from loneliness (seldom or never/sometimes/often or always) [10]. In the PWB score each question represented 0 ('no' in questions 1–4, 'often or always' in question 5 or 6), 0.5 ('sometimes' in question 5 or 6), or 1 ('yes' in questions 1–4, 'seldom or never' in question 5 or 6). The score is created by dividing the total score by the number of questions the participant had answered. Thus, a score of 1 represented the best PWB and 0 the poorest.

Results

The mean age of the participants was 82 years, and 70% ($n = 565$) were women. The PWB was good (score ≥ 0.80) in 52% ($n = 295$), moderate (score = 0.40–0.79) in 39% ($n = 219$), and poor in 9% ($n = 51$) of the participants. Those with lower PWB scores were older, had more risk factors of frailty, were more likely institutionalized, and had the lowest mean MNA-scores (Table 1).

Table 1
 Characteristics of older people in heterogeneous populations and their dietary energy, protein and micronutrient intakes according to their Psychological Well-Being (PWB)-scores.

	PWB poor (score: 0–0.39) $n = 51$	PWB moderate (score: 0.4–0.79) $n = 219$	PWB good (score: 0.8–1) $n = 295$	P-value ^a
Age, mean (SD)	84 (7)	83 (7)	80 (8)	<0.001
Place of residence, %				
Home	33	62	72	<0.001 ^b
Institution	67	38	28	
MNA, mean (SD)	20 (4)	22 (4)	23 (3)	<0.001
Energy, kcal (SD)	1606 (417)	1653 (428)	1717 (461)	0.04
Protein, g (SD)	61 (22)	65 (20)	71 (23)	<0.001
Total fat, g (SD)	59 (23)	61 (21)	65 (26)	0.012
SFA	24 (11)	24 (9)	26 (10)	0.116
MUFA	17 (8)	16 (8)	18 (12)	0.130
PUFA	10 (7)	12 (9)	14 (9)	<0.001
Vitamin A, μg (SD)	568 (371)	846 (1130)	947 (1263)	0.039
Vitamin D, μg (SD)	8 (7)	9 (7)	9 (7)	0.093
Vitamin E, mg (SD)	7 (4)	7 (4)	9 (5)	<0.001
Thiamine, mg (SD)	1.2 (0.5)	1.2 (0.4)	1.3 (0.4)	0.20
Vitamin C, mg (SD)	96 (55)	102 (62)	106 (65)	0.25
Folate, μg (SD)	224 (86)	239 (110)	256 (110)	0.018
Iron, mg (SD)	9 (3)	10 (4)	10 (3)	0.06
Zinc, mg (SD)	9 (3)	10 (3)	11 (3)	0.007

PWB: psychological well-being; SD: standard deviation; MNA: Mini-Nutritional Assessment [10]; SFA: saturated fatty acid; MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids.

^a Tested for linearity.

^b Tested by χ^2 test.

A higher PWB score was associated with higher energy, protein, total fat, polyunsaturated fatty acids, folate, vitamin A and vitamin E, iron and zinc intakes (Table 1).

Conclusion

Higher PWB scores and lower risk of frailty were linearly associated with MNA-scores and key nutrient intakes in heterogeneous groups of older people. In earlier studies MNA-status has been associated with higher PWB scores and the present results extend PWB benefits to key nutrient intakes [3]. Consequently, nutrition interventions may have an important role to enhance the quality of life of older people.

Disclosure of interest

SKJ: reports having received lecturing fees from pharmaceutical company Actavis.

KHP: Dr Pitkälä reports having received lecturing fees from pharmaceutical companies (including Lundbeck, Orion), and having participated in clinical trials funded by pharmaceutical companies.

MPB: Dr Björkman has received lecture fees from Valio Ltd. and Valio Ltd. also provided the nutritional supplements used in the Porvoo sarcopenia & nutrition trial that was started after the baseline assessments reported in this study.

TES: reports of having various educational and consultative cooperation with several companies including Nutricia, Abbott, Amgen, Merck, Pfizer, Novartis, NovoNordisk, minor amount of stock in OrionPharma, President of the European Union Geriatric Medicine Society, which has cooperation also with nutrition industry.

HS: reports no conflict of interest.

MHS: reports having received lecture fees from Nutricia and Verman Ltd.

Acknowledgements

This study was supported by The Finland's Slot Machine Association, Helsinki University Hospital EVO-funding, Oulu University Hospital EVO funding, Helsinki City, Konung Gustav V:s och Drottning Victorias Frimurarestiftelse, Yrjö Jahnsson Foundation. The sponsors did not have any role in the study design, analysis or interpretation of data, nor in writing the report or the decision to submit this article. The authors were independent researchers not associated with the funders.

References

- [1] Suominen MH, Jyväkorpi SK, Pitkälä KH, Finne-Soveri H, Hakala P, Mannisto S, et al. Nutritional guidelines for older people in Finland. *J Nutr Health Aging* 2014;18:861–7.
- [2] WHO. WHOQOL-Bref. Introduction. Administration, scoring and generic version of the assessment. Field Trial Version December 1996. In: Rapley M, editor. *Quality of life research. A critical introduction*. London: Sage Publications; 2003.
- [3] Muurinen S, Savikko N, Soini H, Suominen MM, Pitkälä K. Nutrition and psychological well-being among long-term care residents with dementia. *J Nutr Health Aging* 2015;19(2):178–82.
- [4] Jyväkorpi SK, Pitkälä KH, Kautiainen H, Puranen TM, Laakkonen ML, Suominen MH. Nutrition education and cooking classes improve diet quality, nutrient intake, and psychological well-being of home-dwelling older people – a pilot study. *J Aging Res Clin Pract* 2014;3:120–4.
- [5] Strandberg TE, Stenholm S, Strandberg AY, Salomaa VV, Pitkälä KH, Tilvis RS. “Obesity paradox” frailty, disability, and mortality in older men: a prospective, longitudinal cohort study. *Am J Epidemiol* 2013;178(9):1452–60.
- [6] Björkman MP, Suominen MH, Pitkälä KH, Finne-Soveri HU, Tilvis RS. Porvoo sarcopenia and nutrition trial: effects of protein supplementation on functional performance in home-dwelling sarcopenic older people – study protocol for a randomized controlled trial. *Trials* 2013;14:387.
- [7] Vikstedt T, Suominen MH, Joki A, Muurinen S, Soini H, Pitkälä KH. Nutritional status, energy, protein and micronutrient intake of older service house residents. *J Am Med Dir Assoc* 2011;12(4):302–7.
- [8] Jyväkorpi SK, Pitkälä KH, Puranen TM, Björkman MP, Kautiainen H, Strandberg TE, et al. Low protein and micronutrient intakes in heterogeneous older population samples. *Arch Gerontol Geriatr* 2015;15:30022–4.
- [9] Routasalo PE, Tilvis RS, Kautiainen H, Pitkälä KH. Effects of psychological group rehabilitation on social functioning, loneliness and well-being of lonely, older people: randomized controlled trial. *J Adv Nurs* 2009;65:297–305.
- [10] Vellas B, Guigoz Y, Garry PJ, Nourhashemi F, Bennahum D, Launque S, et al. The Mini Nutritional Assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition* 1999;15(2):116–22.

S.K. Jyväkorpi^{a,*}, K.H. Pitkälä^a, M.P. Björkman^a, T.E. Strandberg^{b,c}, H. Soini^{a,d}, M.H. Suominen^a

^aUniversity of Helsinki, Department of General Practice and Primary Health Care, Finland

^bUniversity of Helsinki, Clinicum, and Helsinki University Hospital, Helsinki, Finland

^cUniversity of Oulu, Center for Life Course Health Research, Oulu, Finland

^dDepartment of Social Services and Health Care, Developmental and Operational Support, Helsinki City, Finland

*Corresponding author at: Tukholmankatu 8 B, 00014 University of Helsinki, Finland. Tel.: +358 504920970
E-mail address: satu.jyvakorpi@gery.fi (S.K. Jyväkorpi)

Available online 27 October 2016