<https://helda.helsinki.fi>

Comment to Hansson, S. et al. (2017) : "Competition for the fish - fish extraction from the Baltic Sea by humans, aquatic mammals, and birds", with special reference to cormorants, perch, and pikeperch

Heikinheimo, Outi

2018

Heikinheimo , O , Lehtonen , H & Lehikoinen , A 2018 , ' Comment to Hansson, S. et al. (2017) : "Competition for the fish - fish extraction from the Baltic Sea by humans, aquatic mammals, and birds", with special reference to cormorants, perch, and pikeperch ' , ICES Journal of Marine Science , vol. 75 , no. 5 , pp. 1832-1836 . <https://doi.org/10.1093/icesjms/fsy054>

<http://hdl.handle.net/10138/259794>
<https://doi.org/10.1093/icesjms/fsy054>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 Comment to Hansson, S. *et al.* (2017): “Competition for the fish – fish extraction from the
2 Baltic Sea by humans, aquatic mammals, and birds”, with special reference to cormorants,
3 perch and pikeperch

4 Outi Heikinheimo¹⁾, Hannu Lehtonen²⁾ and Aleksi Lehikoinen³⁾

5 ¹⁾Natural Resources Institute Finland (Luke), Latokartanonkaari 9, FI-00790 Helsinki,
6 Finland

7 ²⁾University of Helsinki, Department of Environmental Sciences, P.O. Box 65, FI-00014
8 University of Helsinki, Finland

9 ³⁾ The Helsinki Lab of Ornithology, Finnish Museum of Natural History, University of
10 Helsinki, P. O. Box 17, FI-00014 University of Helsinki, Finland

11 Abstract

12 Hansson *et al.* (2017) concluded that competition between fisheries and piscivorous
13 mammals and birds exists in the Baltic Sea, based on the estimation of biomass of the fish
14 species consumed in the ICES subdivisions. We compared their results to the data and
15 scientific knowledge from the coastal waters of Finland and show that local differences in
16 fisheries, fish assemblages and abundance of predators should be taken into account to
17 reliably assess potential competition. Hansson *et al.* (2017) did not include the piscivorous
18 fish in their analysis, but these may be the most important predators. In the Archipelago Sea,
19 for instance, the consumption by fish predators is considerably larger than that of cormorants.

20

21 Introduction

22 Hansson *et al.* (2017) compared the estimated fish consumption by birds and mammals to
23 fisheries catches and concluded that competition for some important species, e.g. perch
24 (*Perca fluviatilis*) and whitefish (*Coregonus lavaretus*), is likely.

25 However, it is questionable whether this kind of analysis can tell us anything about
26 competition between predators and fisheries. Our main concerns are the following:

27 1) Hansson *et al.* (2017) compared the catches of fishing and predation in the scale of ICES
28 subdivisions, but locally the situation largely differs in different areas and habitats.

29 2) Hansson *et al.* (2017) ignored the natural year-class fluctuations which are common in the
30 coastal fish stocks and largely determine the ups and downs in the abundance.

31 3) Fishing and natural predation were paralleled even if the predation and fishing are directed
32 to different size classes, and the predation rate depends on the abundance of each prey species
33 (functional response).

34 4) Predation by piscivorous fish was not taken into account although the diet largely overlaps
35 with that of fish-eating birds.

36 1. Comparison of fisheries catches and predation

37 Hansson *et al.* (2017) stated that the cormorants and seals in some subdivisions consumed
38 twice as much perch as caught in fisheries, and this indicated competition. We compared the
39 perch and pikeperch (*Sander lucioperca*) catches on the Finnish coast and the amount
40 consumed by the local great cormorant (*Phalacrocorax carbo sinensis*) population (P.
41 Rusanen, Finnish Environment Institute) to the results of Hansson *et al.* (2017) (Fig. 1). Most
42 of the perch and pikeperch catches came from the Finnish coastal areas, even though the

43 catches in the particular year 2010 were exceptionally low (Commercial fisheries statistics,
44 Natural Resources Institute Finland, Fig. 2). On the contrary, on the basis of the estimates of
45 Hansson *et al.* (2017), most of the consumption by cormorants took place in other parts of the
46 subdivisions. This is partly due to higher estimated food consumption rate by Hansson *et al.*
47 (2017) (500 g daily consumption was assumed even for small chicks), but also to the fact that
48 there were more cormorants in other parts of the subdivisions than in the Finnish coast. The
49 low fisheries catches in other areas, compared to those in the Finnish coast, are most probably
50 an indication of low fishing effort, poorly reported recreational catches or weak fish stocks.
51 We cannot see there any evidence of competition.

52 Hansson *et al.* (2017) calculated the consumption of prey fish species by predators based on
53 local diet studies and used the results to estimate the consumption in the whole ICES
54 subdivision. However, cormorants utilize the prey species that are abundant, most easily
55 available and of suitable size, and thus the diet varies between years, areas and colonies, or
56 even between weeks in the same breeding season (Salmi *et al.* 2015). For instance, Hansson
57 *et al.* (2017) used the average diet of cormorants in the Finnish Archipelago Sea (share of
58 perch 33%, pikeperch 6%, Salmi *et al.*, 2015) to estimate the amount of perch and pikeperch
59 consumed by cormorants in the ICES Subdivision 29, which extends to the coast of Sweden
60 and Estonia. Certainly not all coastal waters of the Subdivision 29 are such suitable habitats
61 for perch and pikeperch as the Archipelago Sea.

62

63

64 Fig. 1. Comparison of perch (upper panel) and pikeperch (lower panel) fisheries catches and
 65 consumption by cormorants in the ICES subdivisions 29, 30 and 32 according to Hansson *et*
 66 *al.* (2017), and corresponding values in the Finnish coast within each area. The proportions of
 67 perch and pikeperch in the diet of cormorants by Hansson *et al.* (2017) were also used for the
 68 Finnish coast.

69

2. Year-class fluctuations of perch and pikeperch

Hansson *et al.* (2017) stated: “Exploitative competition between fisheries and wildlife occurs if the catch/consumption of a fish species by one group has adverse effects on another consumer group. Field observations of decreased abundance of a fish species in response to fisheries and/or predation by wildlife imply exploitative competition.” In fact, decreased fish catches in coastal waters are frequently observed as a consequence of natural year class fluctuations, due to temperatures affecting the reproduction success of e.g. perch and pikeperch (Böhling *et al.*, 1991; Lappalainen *et al.*, 1996; Heikinheimo *et al.*, 2014). It is obvious that sometimes weak year classes may affect the catches simultaneously with an increase of a predator population, but such a correlation (e.g. Vetemaa *et al.*, 2010) is not a sufficient evidence of a negative impact of the predator (Heikinheimo *et al.*, 2016). To study such an impact, the effect of temperature and other potential factors on annual variation in fish stocks should be disclosed.

Hansson *et al.* (2017) stated that the commercial perch catch in the Finnish Archipelago Sea decreased by about 50% from 1998 to 2011, and Salmi *et al.* (2015) proposed that this was caused by predation by cormorants. In fact, the decrease occurred from the end of 1990s to 2009, caused by strong year classes in the beginning of 1990s, and the weak year classes from 2003 onwards (Auvinen and Heikinheimo, 2017), but the catches then rose and almost reached the 1998 level in 2012 and 2014 (Fig. 2). The catches per unit of effort in gillnet fishing show the same development (Commercial Fisheries Statistics, Natural Resources Institute Finland). The predation by cormorants is directed to smaller perch size classes than fisheries (Salmi *et al.*, 2015), about half of which are males that never grow to the sizes mainly taken by fisheries (Heikinheimo and Lehtonen, 2016). Moreover, there was no change in the mortality of perch compared to earlier periods without cormorants (Heikinheimo and Lehtonen, 2016).

Fig. 2. Commercial perch catches in the Archipelago Sea, Finland (ICES rectangles 49H1, 49H2, 50H1) and the number of breeding cormorant pairs in 1980–2016 (Finnish Environment Institute, P. Rusanen).

According to Hansson *et al.* (2017), Östman *et al.* (2012) reported about 80% lower catch of perch in an area with cormorant colonies (Mönsterås) compared to a reference area that had no colonies within 50 km (Vinö). In time series analyses they found negative association between perch abundance and cormorant numbers in 1995–2009. A longer time series of the gillnet monitoring catches, 1995–2011 (Andersson, 2012), shows that the perch catches per unit of effort (CPUEs) were higher in the reference area during the whole period and the fluctuations were wide but rather synchronous in both areas (significant positive correlation between ln-transformed values, $R^2 = 0.25$, $p = 0.039$). There seems to be negative correlation between the number of breeding cormorants and perch CPUEs both in Mönsterås and in the reference area Vinö, but both are not significant (Mönsterås $R^2 = 0.07$, $p = 0.31$, Vinö $R^2 =$

110 0.03, $p = 0.52$, ln-transformed values) (Fig. 3). Thus there is no evidence of cormorant effect
 111 but rather of synchronous year class fluctuation of perch.

112

113 Fig. 3. Perch catches per gillnet day in Mönsterås (average of three fishing sites) and Vinö
 114 (reference area) based on the data by Andersson (2012), and the number of breeding
 115 cormorant pairs in the Mönsterås area (data by T. Larsson, T. M. Johansson, Länsstyrelsen
 116 Kalmar län).

117

118 3. Are fishing and natural predation comparable?

119 Comparing fisheries catches and fish consumption by predators does not tell us about
 120 competition. The predation rate on a prey species depends on its density in the environment
 121 as well as on the densities of other potential prey. The estimates of potential fisheries catch
 122 loss caused by predation on young fish (Östman *et al.*, 2013; Salmi *et al.*, 2015) largely
 123 depend on the assumed rate of other natural mortality. In the case of the pikeperch in the

124 Archipelago Sea, the other mortality exceeded the mortality caused by cormorants at all
 125 alternative assumptions (Heikinheimo *et al.*, 2016).

126 The natural predation mostly targets individuals that are easiest to catch, i.e. fish in bad
 127 condition, sick or unable to avoid predation for some other causes (Huckstorf *et al.*, 2009).

128 Also slow-growing individuals have a higher probability to be caught because of being a
 129 longer time in the suitable size for predators (Craig *et al.*, 2006). Therefore the mortality
 130 caused by predators may not be additive, i.e. the predators take individuals that have a higher
 131 probability of mortality in the first place (Hilborn and Walters, 1992). Fishing, on the
 132 contrary, mainly takes actively moving individuals and is size-selective, taking the fast-
 133 growing individuals as soon as they reach the catchable size (Conover and Munch, 2002).

134

135 4. Food consumption of piscivorous fish

136 Hansson *et al.* (2017) ignore an important group of predators: the piscivorous fish. We
 137 calculated the fish consumption of the pike (*Esox lucius*) population in the Archipelago Sea
 138 (ICES rectangles 49H1, 49H2, 50H1), based on annual catches in 2007–2015 and food
 139 consumption (Heikinheimo and Korhonen, 1996) (Supplementary Table S1).

140 The total range of the estimated food consumption, calculated from minimum and maximum
 141 catches, was 700–3800 tonnes annually, including only the size classes recruited to fisheries
 142 (Supplementary Table S1). Salmi *et al.* (2015) estimated the fish consumption of cormorants
 143 in the same area at 679–835 tonnes in 2010 and Heikinheimo *et al.* (2016) at 576–704 tonnes
 144 in 2009–2010. Thus, the consumption of the pike population is at a minimum on the same
 145 level, or manifold compared to that of cormorants, and the prey species and sizes are largely
 146 the same as those of cormorants (Eklöv and Hamrin, 1989). The food consumption of the

pikeperch population (ages ≥ 5) is on the same level as that of pike, 1000–4300 tonnes, based on the stock assessment by Heikinheimo *et al.* (2014) and food consumption (Vehanen *et al.*, 1998) (Supplementary Fig. S1). We can conclude that in the Archipelago Sea the piscivorous fish are far more important as predators than the cormorants.

Cormorants utilize mostly smaller fish than do the fisheries, and thus the effect in the fish community can be expected to be very similar to that of fish predation. Predator fish are generally considered an important part of the ecosystem, for instance counteracting extreme fluctuations in the prey fish stocks (Pauly *et al.*, 1998).

Hansson *et al.* (2017) with their article aim at “supporting a more informed debate on resource competition between wildlife and fisheries”. In our opinion, this kind of coarse analysis, ignoring local differences in fish abundance, fisheries and predation, tends to rather aggravate the conflicts.

Supplementary data

The following supplementary material is available at ICESJMS online: Estimation of the food consumption of pike and pikeperch populations in the Archipelago Sea.

References

- Andersson, S. 2012. Kustfiskövervakning i recipienten för Mönsterås Bruk 2011. Linnéuniversitetet. Institutionen för naturvetenskap. Rapport 2012:3. 14 pp. (In Swedish). http://www.lansstyrelsen.se/kalmar/SiteCollectionDocuments/Sv/miljo-och-klimat/Kalmarfiske_2011_lev.pdf

- 169 Auvinen, H., and Heikinheimo, O. 2017. Merialueen ahven. *In* Kalakantojen tila vuonna
 170 2016 sekä ennuste vuosille 2017 ja 2018, pp. 73–83. Ed. by J. Raitaniemi, and K. Manninen.
 171 Luonnonvara- ja biotalouden tutkimus 77/2017. Natural Resources Institute Finland,
 172 Helsinki. (In Finnish). <http://urn.fi/URN:ISBN:978-952-326-504-2>.
 173
- 174 Böhling, P., Hudd, R., Lehtonen, H., Karås, P., Neuman, E., and Thoresson, G. 1991.
 175 Variations in year-class strength of different perch (*Perca fluviatilis*) populations in the Baltic
 176 Sea with special reference to temperature and pollution. Canadian Journal of Fisheries and
 177 Aquatic Sciences, 48: 1181–1187.
- 178 Conover, D.O., and Munch, S.B. 2002. Sustaining fisheries yields over evolutionary time
 179 scales. Science, 297: 94–96.
- 180 Craig, J.K., Burke, B.J., Crowder, L.B., and Rice, J.A. 2006. Prey growth and size-dependent
 181 predation in juvenile estuarine fishes: experimental and model analyses. Ecology, 87: 2366–
 182 2377.
- 183 Eklöv, P., and Hamrin, S. F. 1989. Predatory efficiency and prey selection: interactions
 184 between pike *Esox lucius*, perch *Perca fluviatilis* and rudd *Scardinius erythrophthalmus*.
 185 Oikos, 56: 149–156.
- 186 Hansson, S., Bergström, U., Bonsdorff, E., Härkönen, T., Jepsen, N., Kautsky, L.,
 187 Lundström, K. et al. 2017. Competition for the fish – fish extraction from the Baltic Sea by
 188 humans, aquatic mammals, and birds. ICES Journal of Marine Science,
 189 doi:10.1093/icesjms/fsx207
- 190 Heikinheimo, O., and Korhonen, A.P. 1996. Food consumption of northern pike (*Esox lucius*
 191 L.), estimated with a bioenergetics model. Ecology of Freshwater Fish, 5: 37–47.

- 192 Heikinheimo, O., Pekcan-Hekim, Z., and Raitaniemi, J. 2014. Spawning stock – recruitment
193 relationship in pikeperch, *Sander lucioperca*, in the Baltic Sea, with temperature as
194 environmental effect. Fisheries Research, 155: 1–9. [doi:10.1016/j.fishres.2014.02.015](https://doi.org/10.1016/j.fishres.2014.02.015)
- 195 Heikinheimo, O., and Lehtonen, H. 2016. Comment to article by Salmi, J. A. et al., 2015:
196 Perch (*Perca fluviatilis*) and pikeperch (*Sander lucioperca*) in the diet of the great cormorant
197 (*Phalacrocorax carbo*) and effects on catches in the Archipelago Sea, Southwest coast of
198 Finland. Fisheries Research, 164: 26–34. Fisheries Research, 179: 354–357. doi:
199 10.1016/j.fishres.2016.01.020
- 200 Heikinheimo, O., Rusanen, P., and Korhonen, K. 2016. Estimating the mortality caused by
201 great cormorant predation on fish stocks: pikeperch in the Archipelago Sea, northern Baltic
202 Sea, as an example. Canadian Journal of Fisheries and Aquatic Sciences 73: 84–93. doi:
203 10.1139/cjfas-2015-0033
- 204 Hilborn, R., and Walters, C. J. 1992. Quantitative fisheries stock assessment. Choice,
205 dynamics and uncertainty. Chapman and Hall, London. 570 pp.
- 206 Huckstorf, V., Lewin, W.-C., Mehner, T., and Wolter, C. 2009. Performance level and
207 efficiency of two differing predator-avoidance strategies depend on nutritional state of the
208 prey fish. Behavioral Ecology and Sociobiology, 63: 1735–1742. doi 10.1007/s00265-009-
209 0790-x
- 210 Lappalainen, J., Lehtonen, H., Böhling, P., and Erm, V., 1996. Covariation in yearclass
211 strength of perch, *Perca fluviatilis* L. and pikeperch *Stizostedion lucioperca* (L.). Annales.
212 Zoologici Fennici 33: 421–426.
- 213 Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., and Torres, F. Jr. 1998. Fishing down
214 marine food webs. Science, 279: 860–863.

- 215 Salmi, J. A., Auvinen, H., Raitaniemi, J., Kurkilahti, M., Lilja, J., and Maikola, R. 2015.
216 Perch (*Perca fluviatilis*) and pikeperch (*Sander lucioperca*) in the diet of the great cormorant
217 (*Phalacrocorax carbo*) and effects on catches in the Archipelago Sea, Southwest coast of
218 Finland. Fisheries Research, 164: 26–34.
- 219 Vehanen, T., Hyvärinen, P., and Huusko, A. 1998. Food consumption and prey orientation of
220 piscivorous brown trout (*Salmo trutta*) and pikeperch (*Stizostedion lucioperca*) in a large
221 regulated lake. Journal of Applied Ichthyology, 14: 15–22.
- 222 Vetemaa, M., Eschbaum, R., Albert, A., Saks, L., Verliin, A., Jürgens, K., Kesler, M. *et al.*,
223 2010. Changes in fish stocks in an Estonian estuary: overfishing by cormorants? ICES
224 Journal of Marine Science, 67: 1972–1979.
- 225 Östman, Ö., Bergenius, M., Boström, M. K., and Lunneryd, S.-G. 2012. Do cormorant
226 colonies affect local fish communities in the Baltic Sea? Canadian Journal of Fisheries and
227 Aquatic Sciences, 69: 1047–1055. doi: 10.1139/f2012-042
- 228 Östman, Ö., Boström, M.K., Bergström, U., Anderson, J., and Lunneryd, S.-V., 2013.
229 Estimating competition between wildlife and human – a case of cormorants and coastal
230 fisheries in the Baltic Sea. PLOS ONE, 8. [doi:10.1371/journal.pone.0083763](https://doi.org/10.1371/journal.pone.0083763).

232 Supplement: Estimation of the food consumption of pike and pikeperch populations in the
233 Archipelago Sea

234

235 It is relatively straightforward to calculate the biomass of the piscivorous fish species targeted
236 by fisheries, based on the fisheries catches and estimated fishing and natural mortality, using
237 the equation

238
$$C = F / (F + M) * (1 - \exp(-Zt)) * B,$$

239 where C = annual fisheries catch, F = the instantaneous rate of fishing mortality per year; M
240 = the instantaneous rate of natural mortality per year, $Z = F + M$; t = time in years; B = the
241 biomass of the catchable stock.

242 On the basis of the biomass and food consumption estimates of the given species the total
243 consumption of the population can be estimated.

244 The average fisheries catch of pike in the Archipelago Sea was 354 tonnes (range 203–485
245 tonnes) in the years 2007–2015, and the instantaneous annual fishing mortality was assumed
246 at between 0.5 and 0.8, and natural mortality at 0.1. The annual food consumption was
247 estimated at three- to fourfold the biomass (Heikinheimo and Korhonen, 1996), which gives
248 about 1300–2800 tonnes on the average (Table S1).

249

250 Table S1. Food consumption of northern pike (*Esox lucius*) in the Archipelago Sea (ICES rectangles
251 49H1, 49H2 and 50H1), based on mean, minimum and maximum catches in 2007–2015 (Finnish
252 fisheries statistics, Natural Resources Institute Finland). The rate of natural mortality was assumed at
253 0.1. Biomass in the middle of the year ($t = 0.5$).

Catch (tonnes)		Fishing mortality	Biomass (tonnes)	Food consumption (tonnes)	
				3*biomass	4*biomass
Mean	354	0.5	697	2092	2790
Min.	203	0.5	400	1200	1600
Max	485	0.5	956	2867	3822
Mean	354	0.8	428	1284	1712
Min.	203	0.8	245	736	982
Max.	485	0.8	586	1759	2345

254

255 To estimate the food consumption of pikeperch, the number of fish in each age group, based
256 on the stock assessment (see Heikinheimo *et al.* 2014) and the individual food consumption
257 of the pikeperch in Lake Oulujärvi (Vehanen *et al.* 1998) by age were used. For ages >6 we
258 used the food consumption at age 6 because the growth of pikeperch is slower in the
259 Archipelago Sea than in Lake Oulujärvi (Fig. S1).

Fig. S1. Food consumption (tonnes) of the pikeperch (*Sander lucioperca*) population (ages ≥ 5) in the Archipelago Sea (ICES rectangles 49H1, 49H2 and 50H1) in 2000–2014, based on the updated stock assessment (Heikinheimo et al. 2014) and food consumption of pikeperch by age (Vehanen *et al.* 1998). Food consumption at age 6 was used for all older age groups due to slower growth in the Archipelago Sea.

References

- Heikinheimo, O., and Korhonen, A.P. 1996. Food consumption of northern pike (*Esox lucius* L.), estimated with a bioenergetics model. *Ecology of Freshwater Fish*, 5: 37–47.
- Heikinheimo, O., Pekcan-Hekim, Z., and Raitaniemi, J. 2014. Spawning stock – recruitment relationship in pikeperch, *Sander lucioperca*, in the Baltic Sea, with temperature as environmental effect. *Fisheries Research*, 155: 1–9. [doi:10.1016/j.fishres.2014.02.015](https://doi.org/10.1016/j.fishres.2014.02.015)
- Vehanen, T., Hyvärinen, P., and Huusko, A. 1998. Food consumption and prey orientation of piscivorous brown trout (*Salmo trutta*) and pikeperch (*Stizostedion lucioperca*) in a large regulated lake. *Journal of Applied Ichthyology*, 14: 15–22.