<https://helda.helsinki.fi>

Imaging Classification and Treatment of Spontaneous Intracranial Fusiform and Dissecting Aneurysms

Jahromi, Behnam Rezai

2017-11

Jahromi , B R , Jaaskelainen , J E & Niemela , M 2017 , ' Imaging Classification and Treatment of Spontaneous Intracranial Fusiform and Dissecting Aneurysms ' , World Neurosurgery , vol. 107 , pp. 1039-1039 . <https://doi.org/10.1016/j.wneu.2017.07.090>

<http://hdl.handle.net/10138/298153>

<https://doi.org/10.1016/j.wneu.2017.07.090>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Imaging Classification and Treatment of Spontaneous Intracranial Fusiform and Dissecting Aneurysms

LETTER:

We read with great interest the article by Zhang et al.¹ titled “Endovascular Treatment of Spontaneous Intracranial Fusiform and Dissecting Aneurysms: Outcomes Related to Imaging Classification of 309 Cases.” We appreciate the authors’ attempt to try to classify an unusual and diverse (rather black spot) pathology, i.e., various forms of dilations of intracranial artery segments, in a single setting. Unlike Zhang et al., we do not include intracranial artery wall dissections within the category of intracranial aneurysms.

Most intracranial aneurysms (IAs) are saccular pouches that develop during life on the branching sites of intracranial arteries. The “true” fusiform IAs, as we understand them, spontaneous, nontraumatic dilations of intracranial artery segments, are unusual in neurosurgical IA databases.^{2,3} The dissection of intracranial artery wall, separation of the artery wall layers by intraluminal blood, whether spontaneous or traumatic, also may cause dilation of the external surface, but it is a much quicker and different process than the slow formation of the “true” fusiform IAs.^{4,5} How often our “true” fusiform IAs are “healed” end stages of dissection, whether clinical or incidental, remains to be shown.

The elongation and tortuosity of the artery trunk are other processes to be considered.^{6,7} Dolichoectasia is used to characterize arteries throughout the human body with significant elongation and distension. In IAs, dolichoectasia is seen most often in the vertebrobasilar complex, occasionally in the internal carotid artery. Dolichoectasia may associate to aortic aneurysms.⁸ Dilation, elongation, and tortuosity of artery trunks also may be associated with a variety of predisposing genes.

Very large and international databases should be combed for dilated, elongated, or tortuous intracranial arteries to make sense of one of the remaining microsurgical/endovascular black spots. We suggest that the intracranial artery wall dissections, with or without fusiform dilation of the artery wall, are kept separate from the “true” fusiform IAs and dolichoectasias, as the 2 latter do not seem to include the process of arterial wall layer dissection.

Behnam Rezai Jahromi¹, Juha E. Jääskeläinen², Mika Niemelä¹

From the ¹Department of Neurosurgery, Helsinki University Hospital and University of Helsinki, Helsinki; and ²Neurosurgery of KUH NeuroCenter, Kuopio University Hospital, Kuopio, Finland
To whom correspondence should be addressed: Behnam Rezai Jahromi, M.D.
[E-mail: behnam.rezai@hus.fi]

<http://dx.doi.org/10.1016/j.wneu.2017.07.090>.

REFERENCES

1. Zhang Y, Tian Z, Sui B, Wang Y, Liu J, Li M, et al. Endovascular treatment of spontaneous intracranial fusiform and dissecting aneurysms: outcomes related to imaging classification of 309 cases. *World Neurosurg.* 2017;98:444-455.
2. Jahromi BR, Frösen J, Hernesniemi J. Commentary: bypass surgery for the treatment of dolichoectatic basilar trunk aneurysms: a work in progress. *Neurosurgery.* 2016;79:E514-E516.
3. Lawton MT, Abla AA, Rutledge WC, Benet A, Zador Z, Rayz VL, et al. Bypass surgery for the treatment of dolichoectatic basilar trunk aneurysms: a work in progress. *Neurosurgery.* 2015;79:83-99.
4. Nasr DM, Brinjikji W, Rouchaud A, Kadirvel R, Flemming KD, Kallmes DF. Imaging characteristics of growing and ruptured vertebrobasilar non-saccular and dolichoectatic aneurysms. *Stroke.* 2016;47:106-112.
5. Mizutani T. A fatal, chronically growing basilar artery: a new type of dissecting aneurysm. *J Neurosurg.* 1996;84:962-971.
6. Drake CG, Peerless SJ, Hernesniemi J. *Surgery of Vertebrobasilar Aneurysms: London, Ontario Experience on 1767 Patients.* New York: Springer Science & Business Media; 2012.
7. Awad IA, Barrow DL, eds. *Giant Intracranial Aneurysms.* Park Ridge, IL: AANS; 1995.
8. Pico F, Labreuche J, Cohen A, Touboul PJ, Amarenco P. Intracranial arterial dolichoectasia is associated with enlarged descending thoracic aorta. *Neurology.* 2004;63:2016-2021.