<https://helda.helsinki.fi>

Plasticity in a changing world: Behavioural responses to human perturbations

van Baaren, Joan

2018-06

van Baaren , J & Candolin , U 2018 , ' Plasticity in a changing world: Behavioural responses to human perturbations ' , Current opinions in insect science , vol. 27 , pp. 21-25 . <https://doi.org/10.1016/j.cois.2018.02.003>

<http://hdl.handle.net/10138/299852>

<https://doi.org/10.1016/j.cois.2018.02.003>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1
2
3
4
5
6
7

8 Title: Plasticity in a changing world: Behavioural responses to human perturbations

9

10 Joan van Baaren¹ and Ulrika Candolin²

11

12 1. UMR-CNRS 6553 ECOBIO. Université de Rennes. Campus de Beaulieu. Avenue du Gal Leclerc. 35042 Rennes
13 cedex. France. Joan.van-baaren@univ-rennes1.fr

14

15 2. Department of Biosciences, University of Helsinki, PO Box 65, 00014 Helsinki, Finland. ulrika.candolin@helsinki.fi

16

17

18

19

20 Corresponding author

21 Joan van Baaren

22 UMR-CNRS 6553 ECOBIO

23 Université de Rennes

24 Campus de Beaulieu

25 Avenue du Gal Leclerc

26 35042 Rennes cedex

27 France

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

Abstract

Most insect species are affected by Human Induced Rapid Environmental Changes (HIREC). Multiple responses to HIREC are observed in insects, such as modifications of their morphology, physiology, behavioural strategies or phenology. Most of the responses involve phenotypic plasticity rather than genetic evolution. Here, we review the involvement of behavioural plasticity in foraging, reproduction, habitat choice and dispersal; and how behavioural plasticity modifies social behavior and inter-specific interactions. Although important, behavioural plasticity is rarely sufficient to cope with HIREC. An increasing number of studies find species to respond maladaptively or insufficiently to various anthropogenic disturbances, and less often is large degree of plasticity linked to success.

53 Introduction

54 Most insect species are affected by Human Induced Rapid Environmental Changes (HIREC, defined by Sih et al. 2011)[1],
55 which include variable threats like climate change, habitat fragmentation, habitat loss, human harvesting, and pollution.
56 The species can respond to HIREC, as to any other environmental changes, through plasticity, genetic evolution or
57 dispersal. HIREC generally impact negatively the species. However some species cope well with HIREC and their
58 populations increase to a point where they become themselves a threat to other species. Multiple responses to HIREC
59 are observed in insects, such as modifications of their morphology (eg, size, wing area), physiology (eg, immune
60 response, metabolic rate), behavioural strategies or phenology [2]. Hendry et al. (2008) [3] used meta-analysis dedicated
61 to animals and found most of the responses to involve phenotypic plasticity rather than genetic evolution. Moreover,
62 when genetic changes or shifts in demography, distribution or phenology occurred, these were generally preceded by
63 a modification involving phenotypic plasticity [4]. Thus, behavioural plasticity appears important in explaining variation
64 in the success of species to resist HIREC [5] (Figure 1). The behavioural responses can be either maladaptive, such as the
65 incapacity to detect new predators that can precede species' decline [5], or be adaptive and improve fitness, such as
66 the finding of new host plant species by *Drosophila suzukii* when invading new areas [6]. In this chapter, we examine
67 how behavioural plasticity is involved in insect responses to HIREC during foraging, reproduction, habitat choice and
68 dispersal; and how behavioural plasticity modifies social behavior and inter-specific interactions. We also examine if this
69 plasticity is sufficient to respond adaptively to HIREC or not.

70 Foraging behavior

71 The Optimal Foraging Theory (OFT) predicts that generalist species should turn to specialist strategies when resources
72 become rare, such as after habitat changes due to HIREC. Species that show plasticity in their temporal or spatial
73 resource use, such as aphid parasitoids [7], may have a higher ability to resist HIREC. The main plastic response of
74 foraging insects to changes in their habitats is to modify or enlarge their diet choice. Evans & Moustakas (2017) [8]
75 showed with a model that if predators (or species at high trophic level), shift among prey species when the preferred
76 prey becomes unavailable because of climate change, they survive longer. This modification of diet has been observed
77 in phytophagous insects, and sometimes improves resistance to climate change, as shown by Raffa et al. (2016) [9] for
78 bark beetles, and sometimes not, as shown by Pol et al. (2017) [10] for ants. In some species, like the invasive *Drosophila*
79 *suzukii* that shows extreme plasticity in diet choice, including more than 30 plant species, diet breadth is probably
80 responsible for their success [6]. The capacity of this species to use transgenerational medication (preference of

81 oviposition in fruit containing an entomotoxic substance) contribute to its success [11]. However, *D. suzukii* also
82 oviposits on plants that prevent the development of their larvae. Such traps could represent a cost of extreme plasticity
83 [6].

84 Another solution to climatic stress is to selectively feed on particular types of food. The tropical butterfly *Bicyclus*
85 *anyana*, decreases its activity on hot days and increases its intake of polyphenols (antioxidant) in order to maintain
86 elevated antioxidant levels, which may confer fitness benefits by up-regulating endogenous antioxidant defenses. In
87 grasshoppers, individuals shift their nutrient intake between carbohydrates and proteins depending on temperature
88 and predation stress: indeed, it was shown that a chronic risk of predation (i.e. a chronic stress), induces an elevated
89 metabolism and then a change in nutrient demand and resource consumption and that a higher temperature stands to
90 exacerbate this stress and this diet modification. Growth or reproduction are protein-demanding whereas to cope with
91 stress, a diet balanced towards carbohydrates is more appropriate [12].

92 Reproductive behaviour

93 Changes in the environment can influence the ability of individuals to attract and locate mates, as well as their mate
94 choice behaviour. Such changes can alter the number and quality of offspring produced, as well as lead to a demographic
95 Allee effect, and, hence, influence population dynamics [13]. Traffic noise, for instance, interferes with the ability of
96 female field crickets *Gryllus bimaculatus* to locate singing males during mate searching, which can influence their
97 reproductive success [14]. To counteract negative effects of noise on mate location, individuals may alter their mate
98 attraction or mate location behaviour. For example, male grasshoppers *Chortippus biguttulus* elevate the frequency of
99 their courtship song so that it can be distinguished from traffic noise [15].

100 Another common environmental change influencing reproductive behaviours is the increased use of artificial light at
101 night. Male glow-worms *Lampyris noctiluca*, for instance, are less able to locate glowing females under artificial light
102 [16], while females of the Australian black field cricket *Teleogryllus commodus* become less selective in their mate choice
103 [17]. Increased light levels influence also sex pheromone production. For example, females of a noctuid moth *Mamestra*
104 *brassica* reduce their sex pheromone production when light levels are high [18].

105 Changes in habitat structure are common causes of altered reproductive behaviour. An example is the degradation of
106 tropical cloud forests, which has caused an Afrotropical butterfly *Salamis parhassus* to shift its mate search behaviour

107 from perching to patrolling [19]. Rising temperature and climate change influence in turn the production and reception
108 of sex pheromones [20], as well as the choice of oviposition site [21].

109 Choice of micro-habitat

110 The choice of micro-habitat in response to habitat and climate change has received much attention. For example,
111 Burdick et al. (2015) [22] showed that aphids modify their feeding location under increased UV radiation by feeding on
112 the underside of leaves, as this protects against radiation. Alford et al. (2017) [23] showed that aphid parasitoids
113 increase their thermal tolerance and, hence, resistance to climate stress by manipulating the settling place on
114 parasitized aphids.

115 Activity

116 As mentioned previously, Beaulieu et al. (2015) [24] recorded a decrease in daily activity of the tropical butterfly on hot
117 days. Physical activity elevates metabolic rate and the production of reactive oxygen species (ROS) and reduced activity
118 consequently limits these effects. The ant *Iridomyrmex purpureus* (Formicidae) employs a range of strategies to
119 overcome high temperature stress, such as adjusting time of foraging to the colder hours of the day, climbing grass
120 blades to cool down, and foraging only within shaded areas [25].

121 The reaction to thermal stress varies among species. For example, Kruse et al. (2008) [26] showed that spiders increase
122 their activity at high temperatures, whereas carabid beetles decrease their activity. Since both are predators on the
123 same prey (flies), these opposite reactions modify the predation rate on the prey.

124 At the seasonal level, Tougeron et al. (2017) [27] observed a change in overwintering strategy following winter warming
125 in aphid parasitoid wasps, from diapause to active state. This plasticity allows individuals to increase their progeny
126 production. The mechanism behind the change is a decrease in responsiveness to environmental signals, rather than a
127 genetic loss of diapause.

128 Dispersion

129 Poethke et al. (2010) [28] showed with a model that any persistent deterioration of the environment that decreases
130 expected fitness can induce dispersal behaviour in philopatric organisms. The model was developed for aphids
131 responding to declining population viability due to increased predation pressure, but the model can be applied to any
132 environment deterioration that reduces population viability.

133 Interspecific interactions

134 Interactions between species depend on local conditions and, hence, are sensitive to environmental change [29, 30].
 135 Pesticides, for instance, shift the competitive relationship between two thrips species so that an intrinsically inferior
 136 species *Frankliniella occidentalis* displaces an intrinsically superior competitor *Thrips tabaci* [31]. Climate change and
 137 increases in temperature are similarly altering species interactions. Ground beetles, for instance, attack more mobile
 138 prey at higher temperatures [32], while the parasitoid *Aphidius rhopalosiphi* attack fewer aphid hosts [33].
 139 The invasion of alien species can profoundly alter species interactions and thereby the dynamics of populations [34].
 140 When the mosquito *Aedes albopictus* - a vector of Zika, dengue, and chikungunya viruses - invaded North America, it
 141 displaced the resident *Ae. aegypti* through interspecific matings that sterilized the resident species [35]. However,
 142 invaders can as well have positive effects on ecosystems. When insect pollinators were introduced to the Galapagos
 143 Islands, these made more visits to plants than the native species, which improved the stability of the interaction network
 144 [36].
 145 The loss of species can similarly alter species interactions. When large herbivores were lost from the African savanna,
 146 *Acacia* trees invested less in food and shelter rewards for mutualistic ants, which defend the trees against large
 147 herbivores and insect pests. A non-mutualistic ant could then occupy the trees, which caused the trees to suffer
 148 increased attack by insect pests [37].
 149 Social behaviour
 150 Social behaviour is an important determinant of fitness in social insects. Thus, alteration of social behaviour because of
 151 environmental change can have profound demographic consequences [38, 39]. For instance, higher temperature
 152 induces a switch from solitary to social lifestyle in sweet bees *Halictus rubicundus*, which increases the number of
 153 pollinators, which in turn could mitigate the current pollinator crisis [40]. On the other hand, a rise in temperature can
 154 as well disrupt social interactions. For example, higher temperature increases variation in worker size in the ant
 155 *Temnothorax nylanderi*, probably through reduced ability of nestmate workers to regulate larval development [41].
 156 Changes in the distribution and abundance of resources are common causes of altered frequency or type of social
 157 interaction. Changed shelter availability, for instance, influences the formation of social aggregations in the maritime
 158 earwig *Anisolabis maritima* [42]. In the pharaoh's ant *Monomorium pharaonic*, changed resource distribution influences
 159 pheromone deposition along trails to food sources, which allows ants to communicate and reach adaptive collective
 160 decision in changing environments [43].
 161 Is behavioural plasticity sufficient to cope with human perturbation?

162 In this review, we have shown that behaviourally plastic response to HIREC are common. However, whether the plastic
163 responses are sufficient to prevent population decline and extinction is poorly known. The research field is still in its
164 infancy, and examples are scarce for insects. Some species are able to adjust their behaviour in an adaptive manner to
165 changes, such as bark beetles that use multiple, integrated sensory modalities to adjust their choice of host trees to
166 local conditions [9]. However, an increasing number of studies find species to respond maladaptively or insufficiently to
167 various anthropogenic disturbances. For instance, plasticity in feeding behaviour of the harvester ant *Pogonomyrmex*
168 *mendozanus* is insufficient for coping with deteriorating food quality [9]. Similarly, most species have limited potential
169 to adjust behaviourally to extreme temperatures and many species are moving towards the poles to escape climate
170 change [25, 44].

171 Ecological traps can worsen survival prospects in changing environments. For instance, jewel beetles (*Julodimorpha*
172 *bakewellii*) copulate with beer bottles because the texture of the bottles resembles that of females [45], while mayflies
173 oviposit on asphalt because the surface reflects polarized light in a similar manner as water bodies [46].

174 Behavioural responses that are insufficient but still improve survival could facilitate evolutionary adaptation by
175 providing more time for genetic changes [47]. This is especially likely in species with short generation time, as suggested
176 for many pest and disease vectors [48].

177 Conclusions and perspectives

178 Behavioural plasticity is often not sufficient to cope with HIREC. One explanation is proposed by Sih (2013) [5]: if novel
179 items have been poor options in the evolutionary history of a species, this could explain neophobic behaviours, while if
180 novel items have been beneficial in the past, this could explain neophilic behaviours, which can help species to respond
181 to HIREC. Because HIREC often results in novel conditions, not experienced in the recent evolutionary past, the
182 responses may often be maladaptive and result in population decline. In rare cases, human induced changes can have
183 a positive effect on an ecosystem's stability. For instance, in the Galapagos Islands, an introduced pollinators visited
184 more plants than their native or endemics counterparts, which increased ecosystem stability [36, 49].

185 Two fields of research emerge from this review that need more attention: (1) from a fundamental point of view, more
186 research is needed on the effects of transgenerational plasticity on behavioural responses to environmental change, ie,
187 when environments experienced by the parents influence the responses of offspring. Transgenerational plasticity is
188 increasingly found to be involved in adjustments to variable environments, but the degree to which insects adjust to
189 HIREC through transgenerational plasticity in behaviour is a largely unexplored topic. (2) From an applied point of view,

190 the field of Conservation Behavior should receive more attention. This is a young discipline that investigates how the
191 knowledge of animal behaviour can be taken in consideration in actions for preventing the loss of biodiversity [50].

192

193

194

195 Acknowledgements

196 Thanks to Valérie Briand for bibliographic assistance. We would like to thank the section editors Eric Wajnberg and
197 Emmanuel Desouhant for the invitation to contribute to the Behavioural ecology section 2018. This work was supported
198 by the CNRS (Centre National de Recherche Scientifique) and by the University of Rennes 1 for Joan van Baaren, and by
199 the Academy of Finland grant 277667 to Ulrika Candolin.

200

201 References

202 [1] Sih A, Ferrari MCO, Harris DJ: Evolution and behavioural responses to human-induced rapid environmental change.

203 *Evol Appl* 2011, 4:367-387.

204 [2] Sulmon C, van Baaren J, Cabello-Hurtado F, Gouesbet G, Hennion F, Mony C, Renault D, Bormans M, El Amrani A,

205 Wiegand C, Gérard C: Abiotic stressors and stress responses: What commonalities appear between species

206 across biological organization levels? *Environ Pollut* 2015, 202:66-77.

207 [3] Hendry AP, Farrugia TJ, Kinnison MT: Human influences on rates of phenotypic change in wild animal populations.

208 *Mol Ecol* 2008, 17:20-29.

209 [4] Beever EA, Hall LE, Varner J, Loosen AE, Dunham JB, Gahl MK, Smith FA, Lawler JJ: Behavioral flexibility as a

210 mechanism for coping with climate change. *Front Ecol Environ* 2017, 15(6): 299–308.

211 **This review explores in different taxa how behavioural plasticity can be a mechanism to cope with climate

212 change. It also states the limits of plastic responses and the implications for species management and conservation.

213 [5] Sih A: Understanding variation in behavioural responses to human-induced rapid environmental change: A

214 conceptual overview. *Anim Behav* 2013, 85: 1077-1088.

215 [6] Poyet M, Le Roux V, Gibert P, Meirland A, Prévost G, Eslin P, Chabrierie O: The wide potential trophic niche of the

216 Asiatic fruit fly *Drosophila suzukii*: The key of its invasion success in temperate Europe?. *PLoS One* 2015,

217 10:e0142785.

218 [7] Eoche-Bosy D, Outreman Y, Andrade TO, Krespi L, van Baaren J: Seasonal variations of host resources influence
 219 foraging strategy in parasitoids. *Entomol Exp Appl* 2016, 161:11-19.

220 [8] Evans MR and Moustakas A: Plasticity in foraging behavior as a possible response to climate change. *Ecol Inform*
 221 2017, <https://doi.org/10.1016/j.ecoinf.2017.08.001>.

222 [9] Raffa KF, Andersson MN, Schlyter F: Host selection by bark beetles: playing the odds in a high stakes game. Pine
 223 bark beetles. *Adv Insect Physiol* 2016, 50:1-74.

224 * This paper on host selection by bark beetles shows how behavioural plasticity during host selection process can
 225 help these insects to cope with different HIREC. This is one of the rare studies exploring how behavioural plasticity can
 226 be enough to resist to climate and habitat changes.

227 [10] Pol RG, Vargas GA, Marone L: Behavioural flexibility does not prevent numerical declines of harvester ants under
 228 intense livestock grazing. *Ecol Entomol* 2015, 42:283-293.

229 * This study explains why ant behavioural plasticity in foraging is insufficient to cope with habitat changes
 230 (decrease in the quality and availability of seeds available after grazing).

231 [11] Poyet M, Eslin P, Chabrerie O, Prud'homme SM, Desouhant E, Gibert P : The invasive pest *Drosophila suzukii* uses
 232 trans-generational medication to resist parasitoid attack. *Sc Reports* 2017, 7 Article Number: 43696.

233 [12] Schmitz OJ, Rosenblatt AE, Smylie M: Temperature dependence of predation stress and the nutritional ecology of
 234 a generalist herbivore. *Ecology* 2016, 97:3119-3130.

235

236 [13] Shaw AK, Kokko H: Mate finding, Allee effects and selection for sex-biased dispersal. *J Anim Ecol* 2014,
 237 83: 1256-1267. [14] Schmidt R, Morrison A, Kunc HP: Sexy voices - no choices: Male song in noise fails to attract
 238 females. *Anim Behav* 2014, 94:55-59.

239 [15] Lampe U, Schmoll T, Franzke A, Reinhold K: Staying tuned: Grasshoppers from noisy roadside habitats produce
 240 courtship signals with elevated frequency components. *Funct Ecol* 2012, 26:1348-1354.

241 [16] Bird S, Parker J: Low levels of light pollution may block the ability of male glow-worms (*Lampyrus noctiluca* L.) to
 242 locate females. *J Insect Conserv* 2014, 18:737-743.

243 [17] Botha LM, Joenes TM, Hopkins GR: Effects of lifetime exposure to artificial light at night on cricket (*Teleogryllus*
 244 *commodus*) courtship and mating behaviour. *Anim Behav* 2017, 129:181-188.

245 * The study exemplifies how chronic exposure to bright light at night – a growing problem in our human altered
 246 world – can affect mate choice and reproductive behaviour in crickets.

247 [18] Van Van Geffen KG, Groot AT, Van Grunsven RHA, Donners M, Berendse F, Veenendaal EM: Artificial night lighting
 248 disrupts sex pheromone in a noctuid moth. *Ecol Entomol* 2015, 40:401-408.

249 [19] Bonte D, van Dyck H: Mate-locating behaviour, habitat-use, and flight morphology relative to rainforest
 250 disturbance in an Afrotropical butterfly. *Biol J Linn Soc* 2009, 96:830–839.

251 [20] Boullis A, Detrain C, Francis F, Verheggen FJ: Will climate change affect insect pheromonal communication? *Curr*
 252 *Opin Insect Sci* 2016, 17:87–91

253 [21] Sentis A, Ramon-Portugal F, Brodeur J, Hemptinne JL: The smell of change: warming affects species interactions
 254 mediated by chemical information. *Global Change Biology* 2015, 21:3586–3594.

255 [22] Burdick SC, Prischmann-voldseth DA, Harmon JP: Density and distribution of soybean aphid, *Aphis glycines*
 256 *Matsumura* (Hemiptera : Aphididae) in response to UV radiation. *Popul Ecol* 2015, 57:457–466.

257 [23] Alford L, Kishani Farahani H, Pierre JS, Burel F, van Baaren J: Why is there no impact of the host species on the cold
 258 tolerance of a generalist parasitoid? *J Insect Physiol* 2017, 103:71-77.

259 [24] Beaulieu M, Gillen E, Hahn S, Pape JM, Fischer K : Behavioural antioxidant strategies to cope with high
 260 temperatures: a study in a tropical butterfly. *Anim Behav* 2015, 109: 89-99.

261 [25] Andrew NR, Hart RA, Jung M-P, Hemmings Z, Terblanche JS: Can temperate insect take the heat? A case study of
 262 the physiological and behavioural responses in a common ant, *Iridomyrmex purpureus* (Formicidae), with
 263 potential climate change. *J Insect Physiol* 2013, 59:870-880.

264 [26] Kruse PD, Toft S, Sunderland KD: Temperature and prey capture: opposite relationships in two predator taxa. *Ecol*
 265 *Entomol* 2008, 33:305–312.

266 [27] Tougeron K, Le Lann C, Brodeur J, van Baaren J: Are aphid parasitoids from mild winter climates losing their winter
 267 diapause? *Oecologia* 2017, 183:619-629.

268 * This study shows how aphid parasitoids modify their overwintering strategies following climate change in areas
 269 with mild winters. Behavioural plasticity allows a modification in the thresholds of sensitivity to the stimuli inducing
 270 diapause and activity during the whole winter.

271 [28] Poethke HJ, Weisser WW, Hovestadt T: Predator-induced dispersal and the evolution of conditional dispersal in
 272 correlated environments. *Am Nat* 2010, 175:577-586.

273 [29] Hoover SER, Tylianakis JM: Species Interactions. In: *Behavioural Responses to a Changing World*. Edited by
274 Candolin U, Wong BBM. Oxford University Press; 2012:129-142.

275 [30] Valiente-Banuet A: Beyond species loss: the extinction of ecological interactions in a changing world. *Funct Ecol*
276 2015, 29:299–307

277 [31] Zhao X, Reitz SR, Yuan H, Lei Z, Paini DR, GaoY: Pesticide-mediated interspecific competition between local and
278 invasive thrips pests. *Sci Rep* 2017, 7:40512

279 [32] Vucic-Pestic O, Ehnes RB, Rall BC, Brose U: Warming up the system: higher predator feeding rates but lower
280 energetic efficiencies. *Glob Change Biol* 2011, 17:1301-1310.

281 [33] Le Lann C, Lodi M, Ellers J: Thermal change alters the outcome of behavioural interactions between antagonistic
282 partners. *Ecol Entomol* 2014, 9:578-588

283 [34] Grether GF, Peiman KS, Tobias JA, Robinson BW: Causes and consequences of behavioral interference between
284 species. *Trends Ecol Evol* 2017, 32:760-772.

285 ** A review that emphasises the importance of considering aggressive and reproductive interference between
286 species when assessing the influence of environmental change on the distribution and evolution of species.

287 [35] Bargielowski IE, Lounibos LP: (2016) Satyrization and satyrization-resistance in competitive displacements of
288 invasive mosquito species. *Insect Sci* 2016, 23:162–174.

289 [36] Traveset A, Heleno R, Chamarro S, Vargas P, McMullen CK, Castro-Urgal R, Nogales M, Herrera HW, Olesen JM:
290 Invaders of pollination networks in the Galápagos Islands: emergence of novel communities. *Proc R Soc B* 2013,
291 280:20123040

292 [37] Palmer TM, Stanton ML, Young TP, Goheen JR, Pringle RM, Karban R: Breakdown of an ant-plant mutualism follows
293 the loss of large herbivores from an African savanna. *Science* 2008, 319:192-195

294 [38] Blumstein DT: Social behaviour. In *Behavioural responses to a changing world: mechanisms and consequences*.
295 Edited by Candolin U, Wong BBM. Oxford University Press; 2012:119-128.

296 [39] Schwander T, Lo N, Beekman M, Oldroyd BP, Keller L: Nature versus nurture in social insect caste differentiation.
297 *Trends Ecol Evol* 2010, 25:275–282.

298 [40] Schurch R, Accleton C, Field J: Consequences of a warming climate for social organisation in sweat bees. *Behav*
299 *Ecol Sociobiol* 2016, 70:1131–1139.

300 * The study shows how sociality can be plastic and sensitive to changes in local conditions, such as climate change.

301 Sweet bees, switch from solitary to social lifestyle, when temperature increases, which increases worker number.

302 [41] Molet M, Péronnet R, Couette S, Canovas C, Doums C: Effect of temperature and social environment on worker
303 size in the ant *Temnothorax nylanderi*. *J Therm Biol* 2017, 67:22–29.

304 [42] Hack NL, Iyengar VK: Big wigs and small wigs: Time, sex, size and shelter affect cohabitation in the maritime
305 earwig (*Anisolabis maritima*). *PLoS ONE* 2017, 12:e0185754.

306 [43] Czaczkes TJ, Heinze J: Ants adjust their pheromone deposition to a changing environment and their probability
307 of making errors. *Proc R Soc B* 2015, 282:20150679.

308 [44] Gunderson AR, Stillman JH: Plasticity in thermal tolerance has limited potential to buffer ectotherms from global
309 warming. *Proc R Soc B* 2015, 282: 20150401.

310 [45] Gwynne DT, Rentz DCF: Beetles on the bottle – Male Buprestids mistake stubbies for females (Coleoptera). *J Austr*
311 *Entomol Soc* 1983, 22:79–80.

312 [46] Kriska G, Horvath G, Andrikovics S: Why do mayflies lay their eggs en masse on dry asphalt roads? Water-imitating
313 polarized light reflected from asphalt attracts Ephemeroptera. *J Exp Biol* 1998, 201:2273–2286.

314 [47] Tuomainen U, Candolin U: Behavioural responses to human-induced environmental change. *Biol Rev* 2011,
315 86:640–657.

316 [48] Hoffman AA: Rapid adaptation of invertebrate pests to climatic stress? *Curr Opin Insect Sci* 2017, 21:7-13.

317 [49] Wong BBM, Candolin U: Behavioral responses to changing environments. *Behav Ecol* 2015, 26:665-673.

318 [50] Buchholz R: Behavioural biology: an effective and relevant conservation tool. *Trends Ecol Evol* 2007, 22:401-407.

319

320

321

322

323

324

325

326

327

328

329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352

Figure legend

Figure 1 (adapted from Sih 2013 [5]). Extended reaction norms. Within a range of normal past conditions, animals might show optimal reaction norms that match environmental optima reasonably well. In condition A, just outside of the range of past conditions, organisms might simply extend their reaction norms; however, beyond some threshold (condition B), lack of past selection might allow the maintenance of genetic variation in reaction norms, some of which might come close to matching even a very different optimum in a novel condition that is well outside the range of past conditions. The stars represent the optimal trait in each environment.