<https://helda.helsinki.fi>

Hantavirus maintenance and transmission in reservoir host populations

Forbes, Kristian M.

2018-02

Forbes , K M , Sironen , T & Plyusnin , A 2018 , ' Hantavirus maintenance and transmission in reservoir host populations ' , Current opinion in virology , vol. 28 , pp. 1-6 . <https://doi.org/10.1016/j.coviro.2017.09.003>

<http://hdl.handle.net/10138/300721>

<https://doi.org/10.1016/j.coviro.2017.09.003>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Hantavirus maintenance and transmission in reservoir host populations

Kristian M Forbes^{1,2}, Tarja Sironen^{1,3} and Alexander Plyusnin¹

Hantaviruses are primarily hosted by mammalian species of the orders Rodentia, Eulipotyphla and Chiroptera. Spillover to humans is common, and understanding hantavirus maintenance and transmission in reservoir host populations is important for efforts to curtail human disease. Recent field research challenges traditional phases of virus shedding kinetics derived from laboratory rodent infection experiments. Organ infection sites in non-rodent hosts suggest similar transmission routes to rodents, but require direct assessment. Further advances have also been made in understanding virus persistence (and fadeouts) in fluctuating host populations, as well as occupational, recreational and environmental risk factors associated with spillover to humans. However, despite relevance for both intra-species and inter-species transmission, our understanding of the longevity of hantaviruses in natural environments remains limited.

Addresses

¹ Department of Virology, University of Helsinki, Haartmaninkatu 3, Helsinki FI-00290, Finland

² Centre for Infectious Disease Dynamics and Department of Biology, The Pennsylvania State University, Millennium Science Complex, State College, PA 16802, United States

³ Department of Veterinary Biosciences, University of Helsinki, Agnes Sjöbergin katu 2, Helsinki FI-00014, Finland

Corresponding author: Forbes, Kristian M (kristian.forbes@helsinki.fi)

Current Opinion in Virology 2018, 28:1–6

This review comes from a themed issue on **Emerging viruses: intraspecies transmission**

Edited by **Sander Herfst** and **Martin Ludlow**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 13th November 2017

<http://dx.doi.org/10.1016/j.coviro.2017.09.003>

1879-6257/© 2017 Elsevier B.V. All rights reserved.

Introduction

Hantaviruses (family *Bunyaviridae*) are enveloped RNA viruses with a negative sense tri-segmented genome, encoding RNA-dependent RNA-polymerase, two surface glycoproteins, nucleocapsid protein N, and in some cases, a non-structural protein [1]. The latest taxonomical proposal lists 48 hantavirus species [2], which are present on every continent except Australia and Antarctica. These are hosted by mammalian species of the orders Rodentia,

Eulipotyphla and Chiroptera [3,4], with insect hosts also recently identified [5••].

Each hantavirus is associated with one or few closely related reservoir host species in which they cause mostly asymptomatic and persistent (probably life-long) infections. Although several host-switches — establishment in a secondary host species with onward transmission — have been described among wildlife species [6,7], it is a widely shared opinion that co-evolution (co-diversification) of hantaviruses with their respective reservoir hosts is the prevailing trend. Frequent spillover infections to humans, however, have established hantaviruses as an important group of zoonotic disease agents [8].

In order to devise strategies to mitigate and prevent human hantavirus infections, it is necessary to understand how these viruses are maintained and transmitted in their reservoir host species [9••]. Here we summarise current knowledge on this topic, and draw attention to recent advances and important gaps in understanding. Virus spillover to humans is also discussed, which is tightly linked to reservoir host population dynamics and is a consequence of mechanisms that facilitate intraspecies transmission.

Mechanisms of transmission

The majority of knowledge pertaining to hantavirus transmission comes from a small number of well-studied rodent systems; particularly Puumala virus (PUUV) in bank voles (*Myodes glareolus*) and Sin Nombre virus (SNV) in deer mice (*Peromyscus maniculatus*). Most rodent hosts shed virus in their saliva, urine and faeces, with acquisition by conspecifics occurring via the inhalation of aerosolized virus in the environment (see [Figure 1](#)), and through intense contact, such as biting, grooming and sharing food resources (although these routes are usually difficult to demonstrate) [10–13,14••]. SNV and Andes virus (ANDV) are commonly shed in saliva, but conversely, have only rarely been identified in urine and faeces [15–17], suggesting that intense contact is primarily responsible for intraspecies transmission.

Across all systems, the relative importance of different transmission routes, including sex-biases and seasonal variation, remain outstanding questions. For example, aggressive encounters may be important for transmission during reproductive seasons, especially between males [13]. No direct evidence is available regarding virus shedding or transmission routes for shrew-borne, mole-borne

Figure 1

Synthetic depiction of a hantavirus transmission setting. A common route of transmission for many hantaviruses is the inhalation of aerosolized virus, which is shed into the environment in the saliva and excreta of infected rodents (depicted in red). Several factors are known or expected to influence the amount and persistence of infectious hantaviruses once outside the host **(a)**, and the risk of infection in conspecific rodents **(b)** and people **(c)**.

or bat-borne hantaviruses. However, the few studies to assess virus distribution in host tissues have identified similar patterns to rodents, suggesting comparable transmission routes; high viral loads in kidneys particularly support the notion of virus shedding in urine [18,19*,20*]. These assumptions nevertheless require direct validation. As habitat and behavioural characteristics of host species vary, especially across taxonomic orders, viruses are likely to experience some disparity in selection pressures acting on transmission.

Rodent viral loads and shedding in saliva and excreta peak approximately 2–4 weeks following initial infection, and then decrease over subsequent months despite the persistence of viral RNA in host tissues [12,13,21]. Consequently, hantavirus infections are often characterised by an initial acute infectious phase, followed by chronic infection with occasional or no virus shedding. However, this assumption is largely based on laboratory infection experiments, and is challenged by monitoring studies of naturally infected wild bank voles. These have found that

after an initial and comparable peak in shedding, a substantial proportion of voles continue to shed and transmit PUUV, probably throughout their lifetime [14*,22]. Inherent differences between natural and laboratory settings may be responsible for these contrasting results. In the laboratory, for instance, hosts are usually provided with food *ad libitum*, whereas in the wild, finite energy resources may be traded-offs among immune defences and other pertinent processes, impairing the hosts' ability to control the infection [23].

Virus persistence outside the host

Indirect inhalation is favoured as the dominant mode of intraspecies transmission for many hantaviruses due to the ease with which the virus can pass between rodents (for example between cages up to 1.5 m apart for PUUV [24]) and to people, and the longevity of infectious virus once shed into the environment (Figure 1). Clearly for viruses that are transmitted via indirect contact, the longevity of infectiousness once outside the host will

be selected for, and may also interact with other evolutionarily pliable traits such as virulence.

Live vole exposure experiments have demonstrated that PUUV remains infectious at room temperature for 12–15 days once outside the host [25]. Environmental conditions are expected to play an important role in the stability of hantaviruses, with cell culture experiments showing that cool and damp conditions are able to substantially prolong infectiousness [25,26]. Other environmental variables, such as UV radiation, are likely to have the opposite effect. However, the influence of environmental conditions on virus stability remains to be assessed in natural settings and with other hantavirus species.

Heterogeneity in intraspecies transmission

Several factors are known to create individual-level heterogeneity in hantavirus infection risk. Maternal antibodies confer resistance to hantavirus infections in juvenile rodents for up to 80 days [15,22,27,28], while older individuals tend to be disproportionately infected, reflecting their accumulated exposure risk and the persistent nature of infections [13,29]. Male rodents also typically display higher infection prevalence than females [13,28,30], which may result from increased virus exposure due to both aggressive encounters and greater spatial movement. Research with bank voles suggests that resistance/tolerance genes and co-infections by other parasites can create additional heterogeneity in susceptibility [31–33]. Co-infections are particularly noteworthy, as their ability to influence infection dynamics in other host-parasite systems is increasingly recognised [34–36], but their effects on hantaviruses susceptibility and shedding dynamics have so far received little research attention.

The population-level effect of density-dependent transmission has been extensively studied, with mixed results [30]. This isn't surprising given that many studies rely on antibody detection to determine infection status, which usually fails to distinguish between new and old infections (but see [37]). Further confounding the identification of density-dependence is seasonal turnover in population age structure. Research with bank voles has shown that highest hantavirus prevalence typically occurs in early spring, when old, overwintered rodents predominate [29,38]. The introduction of susceptible juveniles through breeding during spring and summer then dilutes the proportion of infected individuals, despite a potential increase in the transmission rate [39].

Extending to a community level, hantavirus transmission is likely to decrease as species richness increases due to mechanisms which reduce contact between naïve and infected hosts or virus in the environment [40]. For example, high species diversity may reduce the abundance of reservoir hosts, or increase predator density, which encourages rodents to seek protective areas and

thereby restricts their movement and virus exposure opportunities. Evidence for this effect comes from studies with SNV in deer mice [41–43], and PUUV in bank voles [44,45]. However, the precise mechanism by which species diversity lowers infection prevalence is often not known (although density is sometimes accounted for).

Landscape effects on hantavirus transmission dynamics present an intriguing line of enquiry, with strong implications for spillover into humans (discussed below). Preliminary evidence suggests that transmission may be greatest in non-optimal host habitats, such as peri-domestic settings for deer mice and young forest stands for bank voles [13,46]; although patterns appear more complex at a micro-habitat scale [47]. Recent studies have examined the persistence of hantaviruses in fluctuating rodent populations, identifying regular fadeouts of PUUV, Dobrava-Belgrade virus (DOBV) and SNV [47,48,49]. Indeed, it was estimated that a sustained population density of at least 17 deer mice/ha is required for the invasion and persistence of SNV [48]. These studies highlight the meta-population structure in which hantaviruses exist, and emphasise the need for research to investigate mechanisms of connectivity that facilitate virus reintroduction to subpopulations (see for example [50]).

Spillover infections

As hantaviruses are shed into the environment, exposure by non-reservoir host species is an invariable consequence. While spillover to sympatric rodents has been identified [51], human infection is by far the most extensively studied form of spillover. All currently known human hantaviruses infections are derived from rodents, with disease severity ranging from asymptomatic to fatal, and the case fatality rate and proportion of asymptomatic infections largely determined by the virus type and human genetic predispositions [52]. These infections are considered dead-end events for the virus, due to an absence of human-to-human transmission. One exception is known; Andes virus (ANDV), which can be occasionally transmitted among people [53]. While the details of how this occurs are not fully understood, ANDV seems to display unique anti-inflammatory properties and resist host salivary antiviral mechanisms [54,55].

Spatio-temporal variation in human infection risk is due to variation in exposure, reservoir host abundance, and virus stability (discussed above). People presumably become infected following inhalation of or direct contact with infectious virus present in wildlife host excreta and saliva. Certain occupations and activities increase the likelihood of virus exposure, including working in forestry and farming, and working with hay and fire wood, cleaning barns and forest cabins, and entering rodent infested buildings (Figure 1) [13,56,57]. As a result, agricultural practices and landscape changes play a significant role in determining the overall exposure risk [58,59]. Recently,

contact with pet rats has been identified as a source of Seoul hantavirus infections [60].

In northern Europe, where specialist predation is likely to regulate vole abundance [61], bank vole density is a reasonable proxy for human infection risk [62]. Meanwhile in temperate Europe, masting induced by warm summers is an important driver of hantavirus epidemics (mostly due to PUUV) [63]. A unique dataset from Central China covering half a century revealed that two climatic variables, temperature and rainfall, play key roles in reservoir host dynamics, and the rate of transmission and risk of Hantaan virus disease outbreaks [64]. Climate variables, in particular temperature, precipitation and rainfall, have similarly been highlighted as drivers of hantavirus cardiopulmonary syndrome outbreaks and nephropathia epidemica infections in other areas [58,65]. Climate not only affects reservoir host dynamics, but also human activities that increase exposure risk, such as seasonal agricultural activities [59] and time spent in the countryside [58,66].

Conclusions

A key purpose of this review is to draw attention to recent developments pertaining to the maintenance and transmission of hantaviruses in their reservoir host populations. As outlined above, several advances have been made, which have in particular increased our understanding of shedding kinetics in wild rodent populations [14], heterogeneity in transmission dynamics [29,38], and occupational, recreational and environmental risk factors for human infection [57,59,64]. While the bank vole-Puumala virus system in northern Europe is by far the most extensively studied host-hantavirus relationship, recent studies on DOBV and SNV in fluctuating rodent populations have shed important light on the persistence and fadeouts of these hantaviruses in their reservoir host populations [48,49].

Despite these advances, significant knowledge gaps remain. Probably most glaring is that almost nothing is known regarding the transmission of hantaviruses in non-rodent hosts. Indeed, relatively little information is available for most rodent-borne hantaviruses. Future research is clearly required to address these deficiencies, and to also assess potential associations between lesser studied hantaviruses and human disease. Other notable areas for further research include seasonal variation in the relative importance of different transmission routes, the effects of environmental variables on virus longevity outside the host, the impact of concurrent infections by other parasites and pathogens on hantavirus susceptibility and shedding, and mechanisms of connectedness between meta-populations that facilitate virus reintroduction.

As the loss of natural habitats continues — due to factors such as land use changes and climate change — so too will

the movement of wildlife into urban and peri-urban environments [67]. Coupled with human occupations and recreations that encroach on natural wildlife habitat and increase the likelihood of exposure, hantavirus infections will remain a global threat to public health; including viruses currently known to science and potentially also those yet to be identified [68]. Understanding how hantaviruses are maintained and transmitted within populations of their reservoir hosts is critical to devising effective intervention strategies [9]. While recent advances have been made towards this endeavour, many important questions remain.

Conflicts of interest

None.

Acknowledgements

We are very grateful to Paula Ahola for drawing the figure. KMF is financially supported by the Finnish Cultural Foundation.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Plyusnin A, Beaty BJ, Elliott RM, Goldbach R, Kormelink R, Lundkvist Å, Schmaljohn CS, Tesh RB: **Bunyaviridae**. In *Virus Taxonomy: Classification and Nomenclature of Viruses. Ninth Report of the International Committee on Taxonomy of Viruses*. Edited by King AMQ, Lefkowitz EJ, Adams MJ, Carstens EB. Elsevier; 2012:725–741.
2. Adams MJ, Lefkowitz EJ, King AMQ, Harrach B, Harrison RL, Knowles NJ, Kropinski AM, Krupovic M, Kuhn JH, Mushegian AR *et al.*: **Changes to taxonomy and the International Code of Virus Classification and Nomenclature ratified by the International Committee on Taxonomy of Viruses**. *Arch Virol* 2017, **162**:2505–2538.
3. Plyusnin A, Sironen T: **Evolution of hantaviruses: co-speciation with reservoir hosts for more than 100 MYR**. *Virus Res* 2014, **187**:22–26.
4. Castel G, Tordo N, Plyusnin A: **Estimation of main diversification time-points of hantaviruses using phylogenetic analyses of complete genomes**. *Virus Res* 2017, **233**:60–69.
5. Shi M, Lin XD, Tian JH, Chen LJ, Chen X, Li CX, Qin XC, Li J, Cao JP, Eden JS *et al.*: **Redefining the invertebrate RNA virosphere**. *Nature* 2016, **540**:539–543.
- Reports the discovery of 1445 novel RNA viruses originating from over 220 invertebrate species. Among these are ancestral hantaviruses.
6. Nemirov K, Vapalahti O, Lundkvist A, Vasilenko V, Golovljova I, Plyusnina A, Niemimaa J, Laakkonen J, Henttonen H, Vaheri A *et al.*: **Isolation and characterization of Dobrava hantavirus carried by the striped field mouse (*Apodemus agrarius*) in Estonia**. *J Gen Virol* 1999, **80**:371–379.
7. Vapalahti O, Lundkvist Å, Fedorov V, Conroy CJ, Hirvonen S, Plyusnina A, Nemirov K, Fredga K, Cook JA, Niemimaa J *et al.*: **Isolation and characterization of a hantavirus from *Lemmus sibiricus*: evidence for host switch during hantavirus evolution**. *J Virol* 1999, **73**:5586–5592.
8. Vaheri A, Strandin T, Hepojoki J, Sironen T, Henttonen H, Mäkelä S, Mustonen J: **Uncovering the mysteries of hantavirus infections 2013**. *Nat Rev Microbiol* 2013, **11**:539–550.
9. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO: **Pathways to zoonotic spillover**. *Nature Rev Microbiol* 2017, **15**:502–510.

Recent review article that presents a synthetic framework for understanding the spillover of zoonotic pathogens. Discusses ecological, epidemiological and behavioral determinants of pathogen exposure, and within-human factors that affect susceptibility.

10. Nuzum EO, Rossi CA, Stephenson EH, LeDuc JW: **Aerosol transmission of Hantaan and related viruses to laboratory rats.** *Am J Trop Med Hyg* 1988, **38**:636-640.
 11. Klein SL, Bird BH, Glass GE: **Sex differences in immune responses and viral shedding following Seoul virus infection in Norway rats.** *Am J Trop Med Hyg* 2001, **65**:57-63.
 12. Hardestam J, Karlsson M, Falk KI, Olsson G, Klingström J, Lundkvist A: **Puumala hantavirus excretion kinetics in bank voles (*Myodes glareolus*).** *Emerg Infect Dis* 2008, **14**:1209-1215.
 13. Carver S, Mills JN, Parmenter CA, Parmenter RR, Richardson KS, Harris RL, Douglass RJ, Kuenzi AJ, Luis AD: **Toward a mechanistic understanding of environmentally forced zoonotic disease emergence: Sin Nombre hantavirus.** *Bioscience* 2015, **65**:651-666.
 14. Voutilainen L, Sironen T, Tonteri E, Bäck AT, Razzauti M, Karlsson M, Wahlström M, Niemimaa J, Henttonen H, Lundkvist Å: **Life-long shedding of Puumala hantavirus in wild bank voles (*Myodes glareolus*).** *J Gen Virol* 2015, **96**:1238-1247.
- A capture-mark-recapture study of PUUV shedding and viraemia in naturally infected wild bank voles. Kinetics differed from results in laboratory settings, and life-long virus shedding was proposed as a means for hantaviruses to persist over bottlenecks in host abundance.
15. Botten J, Mirowsky W, Ye C, Gottlieb K, Saavedra M, Ponce L, Hjelle B: **Shedding and intracage transmission of Sin Nombre hantavirus in the deer mouse (*Peromyscus maniculatus*) model.** *J Virol* 2002, **76**:7587-7594.
 16. Padula P, Figueroa R, Navarrete M, Pizarro E, Cadiz R, Bellomo C, Jofre C, Zor L, Rodriguez E, Murúa R: **Transmission study of Andes hantavirus infection in wild Sigmodontine rodents.** *J Virol* 2004, **78**:11972-11979.
 17. Safronetz D, Drebot MA, Artsob H, Cote T, Makowski K, Lindsay LR: **Sin Nombre virus shedding patterns in naturally infected deer mice (*Peromyscus maniculatus*) in relation to duration of infection.** *Vector Borne Zoonotic Dis* 2008, **8**:97-100.
 18. Gu SH, Hejduk J, Markowski J, Kang HJ, Markowski M, Polatyńska M, Sikorska B, Liberski PP, Yanagihara R: **Co-circulation of soricid- and talpid-borne hantaviruses in Poland.** *Infect Genet Evol* 2014, **28**:296-303.
 19. Witkowski PT, Drexler JF, Kallies R, Ličková M, Bokorová S, Mananga GD, Szemes T, Leroy EM, Krüger DH, Drosten C *et al.*: **Phylogenetic analysis of a newfound bat-borne hantavirus supports a laurasiatherian host association for ancestral mammalian hantaviruses.** *Infect Genet Evol* 2016, **41**:113-119.
- Reports the third bat-borne hantavirus from Africa, and is the first study of virus distribution in bat tissues. Infection sites resembled those of chronically infected rodents.
20. Lee SH, Kim WK, No JS, Kim JA, Kim JI, Gu SH, Kim HC, Klein TA, Park MS, Song JW: **Dynamic circulation and genetic exchange of a shrew-borne hantavirus, Imjin virus, in the Republic of Korea.** *Sci Rep* 2017, **7** <http://dx.doi.org/10.1038/srep44369>.
- First detailed study of the epidemiology of a non-rodent hosted hantavirus. Identified higher prevalence in male than female, and in older than younger shrews. Described the tissue distribution of Imjin virus, which was similar to rodent-borne hantaviruses.
21. Fulhorst CF, Milazzo ML, Duno G, Salas RA: **Experimental infection of the *Sigmodon alstoni* cotton rat with Caño Delgadito virus, a South American hantavirus.** *Am J Trop Med Hyg* 2002, **67**:107-111.
 22. Bernshtein AD, Apekina NS, Mikhaoliva TV, Myasnikov YA, Khlyap LA, Korotkov YS, Gavrilovskaya IN: **Dynamics of Puumala hantavirus infection in naturally infected bank voles (*Clethrionomys glareolus*).** *Arch Virol* 1999, **144**:2415-2428.
 23. Forbes KM, Mappes T, Sironen T, Strandin T, Stuart P, Meri S, Vapalahti O, Henttonen H, Huitu O: **Food limitation constrains host immune responses to nematode infections.** *Biol Lett* 2016, **12** <http://dx.doi.org/10.1098/rsbl.2016.0471>.
 24. Gavrilovskaya IN, Apekina NS, Bernshtein AD, Demina VT, Okulova NM, Myasnikov YA, Chumakov MP: **Pathogenesis of hemorrhagic fever with renal syndrome virus infection and mode of horizontal transmission of hantavirus in bank voles.** *Arch Virol* 1990, **Suppl. 1**:57-62.
 25. Kallio ER, Klingström J, Gustafsson E, Manni T, Vaheri A, Henttonen H, Vapalahti O, Lundkvist A: **Prolonged survival of Puumala hantavirus outside the host: evidence for indirect transmission via the environment.** *J Gen Virol* 2006, **87**:2127-2134.
 26. Hardestam J, Simon N, Hedlund KO, Vaheri A, Klingström J, Lundkvist Å: **Ex vivo stability of the rodent-borne Hantaan virus in comparison to that of arthropod-borne members of the Bunyaviridae family.** *Appl Environ Microbiol* 2007, **73**:2547-2551.
 27. Kallio ER, Poikonen A, Vaheri A, Vapalahti O, Henttonen H, Koskela E, Mappes T: **Maternal antibodies postpone hantavirus infection and enhance individual breeding success.** *Proc Biol Sci* 2006, **273**:2771-2776.
 28. Kallio ER, Henttonen H, Koskela E, Lundkvist Å, Mappes T, Vapalahti O: **Maternal antibodies contribute to sex-biased difference in hantavirus transmission dynamics.** *Biol Lett* 2013, **9** <http://dx.doi.org/10.1098/rsbl.2013.0887>.
 29. Voutilainen L, Kallio ER, Niemimaa J, Vapalahti O, Henttonen H: **Temporal dynamics of Puumala hantavirus infection in cyclic populations of bank voles.** *Sci Rep* 2016, **6** <http://dx.doi.org/10.1038/srep21323>.
- Describes a long-term monitoring study of Puumala virus transmission in wild bank voles, which identified seasonal transmission patterns that reflect population turnover.
30. Khalil H, Hörnfeldt B, Evander M, Magnusson M, Olsson G, Ecke F: **Dynamics and drivers of hantavirus prevalence in rodent populations.** *Vector Borne Zoonotic Dis* 2014, **14**:537-551.
 31. Deter J, Bryja J, Chaval Y, Gala M, Henttonen H, Laakkonen J, Voutilainen L, Vapalahti O, Vaheri A, Salvador AB *et al.*: **Association between the DQA MHC class II gene and Puumala virus infection in *Myodes glareolus*, the bank vole.** *Infect Genet Evol* 2008, **8**:450-458.
 32. Guivier E, Galan M, Salvador AR, Xuéreb A, Chaval Y, Olsson GE, Essbauer S, Henttonen H, Voutilainen L, Cosson JF *et al.*: **Tnf- α expression and promoter sequences reflect the balance of tolerance/resistance to Puumala hantavirus infection in European bank vole populations.** *Infect Genet Evol* 2010, **10**:1208-1217.
 33. Salvador AR, Guivier E, Xuéreb A, Chaval Y, Cadet P, Pouille ML, Sironen T, Voutilainen L, Henttonen H, Cosson JF *et al.*: **Concomitant influence of helminth infection and landscape on the distribution of Puumala hantavirus in its reservoir, *Myodes glareolus*.** *BMC Microbiol* 2011, **11** <http://dx.doi.org/10.1186/1471-2180-11-30>.
 34. Forbes KM, Henttonen H, Hirvelä-Koski V, Kipar A, Mappes T, Stuart P, Huitu O: **Food provisioning alters infection dynamics in populations of a wild rodent.** *Proc Biol Sci* 2015, **282** <http://dx.doi.org/10.1098/rspb.2015.1939>.
 35. Griffiths EC, Fairlie-Clarke K, Allen JE, Metcalf CJ, Graham AL: **Bottom-up regulation of malaria population dynamics in mice co-infected with lung-migratory nematodes.** *Ecol Lett* 2015, **18**:1387-1396.
 36. Antwis RE, Griffiths SM, Harrison XA, Aranega-Bou P, Arce A, Bettridge AS, Brailsford FL, de Menezes A, Devaynes A, Forbes KM *et al.*: **Fifty important research questions in microbial ecology.** *FEMS Microbiol Ecol* 2017, **93** <http://dx.doi.org/10.1093/femsec/fix044>.
 37. Borremans B, Hens N, Beutels P, Leirs H, Reijnders J: **Estimating time of infection using prior serological and individual information can greatly improve incidence estimation of human and wildlife infections.** *PLoS Comput Biol* 2016, **12** <http://dx.doi.org/10.1371/journal.pcbi.1004882>.
 38. Reil D, Rosenfeld UM, Imholt C, Schmidt S, Ulrich RG, Eccard JA, Jacob J: **Puumala hantavirus infections in bank vole populations: host and virus dynamics in Central Europe.** *BMC Ecol* 2017, **17** <http://dx.doi.org/10.1186/s12898-017-0118-z>.

39. Boone JD, McGwire KC, Otteson EW, DeBaca RS, Kuhn EA, St Jeor SC: **Infection dynamics of Sin Nombre virus after a widespread decline in host populations.** *Am J Trop Med Hyg* 2002, **67**:310-318.
40. Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, Hudson P, Jolles A, Jones KE, Mitchell CE *et al.*: **Impacts of biodiversity on the emergence and transmission of infectious diseases.** *Nature* 2010, **468**:647-652.
41. Clay CA, Lehmer EM, St Jeor S, Dearing MD: **Sin Nombre virus and rodent species diversity: a test of the dilution and amplification hypotheses.** *PLoS ONE* 2009, **4** <http://dx.doi.org/10.1371/journal.pone.0006467>.
42. Dizney LJ, Ruedas LA: **Increased host species diversity and decreased prevalence of Sin Nombre virus.** *Emerg Infect Dis* 2009, **15**:1012-1018.
43. Dearing MD, Clay C, Lehmer E, Dizney L: **The roles of community diversity and contact rates on pathogen prevalence.** *J Mammal* 2015, **96**:29-36.
44. Khalil H, Ecker F, Evander M, Magnusson M, Hörnfeldt B: **Declining ecosystem health and the dilution effect.** *Sci Rep* 2016, **6** <http://dx.doi.org/10.1038/srep31314>.
45. Ecker F, Angeler DG, Magnusson M, Khalil H, Hörnfeldt B: **Dampening of population cycles in voles affects small mammal community structure, decreases diversity, and increases prevalence of a zoonotic disease.** *Ecol Evol* 2017, **7**:5331-5342.
46. Voutilainen L, Savola S, Kallio ER, Laakkonen J, Vaheri A, Vapalahti O, Henttonen H: **Environmental change and disease dynamics: effects of intensive forest management on Puumala hantavirus infection in boreal bank vole populations.** *PLoS ONE* 2012, **7** <http://dx.doi.org/10.1371/journal.pone.0039452>.
47. Khalil H, Olsson G, Magnusson M, Evander M, Hörnfeldt B, Ecker F: **Spatial prediction and validation of zoonotic hazard through micro-habitat properties: where does Puumala hantavirus hole-up?** *BMC Infect Dis* 2017, **17** <http://dx.doi.org/10.1186/s12879-017-2618-z>.
48. Luis AD, Douglass RJ, Mills JN, Björnstad ON: **Environmental fluctuations lead to predictability in Sin Nombre hantavirus outbreaks.** *Ecology* 2015, **96**:1691-1701.
Uses a stage-structured host-pathogen model parameterized and cross-validated by long-term monitoring of deer mice to estimate that a sustained abundance of at least 17 mice/ha is required for SNV invasion. Density fluctuations in study population's led to sporadic virus fadeout.
49. Guzzetta G, Tagliapietra V, Perkins SE, Haufler H, Poletti P, Rizzoli A: **Population dynamics of wild rodents induce stochastic fadeouts of a zoonotic pathogen.** *J Anim Ecol* 2017, **86**:451-459.
50. Streicker DG, Winternitz JC, Satterfield DA, Condori-Condori RE, Brooks A, Tello C, Recuenco S, Velasco-Villa A, Altizer S, Valderrama W: **Host-pathogen evolutionary signatures reveal dynamics and future invasions of vampire bat rabies.** *Proc Natl Acad Sci U S A* 2016, **13**:10926-10931.
51. Forbes KM, Voutilainen L, Jääskeläinen A, Sironen T, Kinnunen PM, Stuart P, Vapalahti O, Henttonen H, Huitu O: **Serological survey of rodent-borne viruses in Finnish field voles.** *Vector Borne Zoonotic Dis* 2014, **14**:78-83.
52. Avič-upanc T, Saksida A, Korva M: **Hantavirus infections.** *Clin Microbiol Infect* 2015 <http://dx.doi.org/10.1111/1469-0691.12291>.
53. Martinez-Valdebenito C, Calvo M, Vial C, Mansilla R, Marco C, Palma RE, Vial P, Valdivieso F, Mertz G, Ferrés M: **Person-to-person household and nosocomial transmission of Andes hantavirus, southern Chile, 2011.** *Emerg Infect Dis* 2014, **20**:1637-1644.
54. Hardestam J, Lundkvist Å, Klingström J: **Sensitivity of Andes hantavirus to antiviral effect of human saliva.** *Emerg Infect Dis* 2009, **15**:1140-1142.
55. Cinica V, Dalrymple NA, Roth E, Nasonov A, Mackow ER: **An innate immunity-regulating virulence determinant is uniquely encoded by the Andes virus nucleocapsid protein.** *MBio* 2014, **5** <http://dx.doi.org/10.1128/mBio.01088-13>.
56. Makary P, Kanerva M, Ollgren J, Virtanen MJ, Vapalahti O, Lyytikäinen O: **Disease burden of Puumala virus infections, 1995–2008.** *Epidemiol Infect* 2010, **138**:1484-1492.
57. de St Maurice A, Ervin E, Schumacher M, Yaglom H, VinHatton E, Melman S, Komatsu K, House J, Peterson D, Buttke D *et al.*: **Exposure characteristics of hantavirus pulmonary syndrome patients, United States, 1993–2015.** *Emerg Infect Dis* 2017, **23**:733-739.
Summarises the demographics and rodent exposure settings for more than 600 human hantavirus infections, revealing increased risks for certain occupations and populations.
58. Prist PR, D Andrea PS, Metzger JP: **Landscape, climate and hantavirus cardiopulmonary syndrome outbreaks.** *Ecohealth* 2017 <http://dx.doi.org/10.1007/s10393-017-1255-8>.
59. Tian H, Yu P, Björnstad ON, Cazelles B, Yang J, Tan H, Huang S, Cui Y, Dong L, Ma C *et al.*: **Anthropogenically driven environmental changes shift the ecological dynamics of hemorrhagic fever with renal syndrome.** *PLoS Pathog* 2017, **13** <http://dx.doi.org/10.1371/journal.ppat.1006198>.
A three-decade-long field study on hantavirus infections coupled with epidemiological data, reporting changing dynamics linked to environmental fluctuations at the human-wildlife interface. Modelling demonstrated seasonality in transmission generated from cycles of agricultural phenology and rodent behavior driven by the rainy seasons.
60. Duggan JM, Close R, McCann L, Wright D, Keys M, McCarthy N, Mannes T, Walsh A, Charlett A, Brooks TJG: **A seroprevalence study to determine the frequency of hantavirus infection in people exposed to wild and pet fancy rats in England.** *Epidemiol Infect* 2017 <http://dx.doi.org/10.1017/S0950268817001480>.
61. Korpela K, Helle P, Henttonen H, Korpimäki E, Koskela E, Ovaskainen O, Pietiäinen H, Sundell J, Valkama J, Huitu O: **Predator-vole interactions in Northern Europe: the role of small mustelids revised.** *Proc Biol Sci* 2014, **281** <http://dx.doi.org/10.1098/rspb.2014.2119>.
62. Kallio ER, Begon M, Henttonen H, Koskela E, Mappes T, Vaheri A, Vapalahti O: **Cyclic hantavirus epidemics in humans – predicted by rodent host dynamics.** *Epidemics* 2009, **1**:101-107.
63. Tersago K, Verhagen R, Vapalahti O, Heyman P, Ducoffre G, Leirs H: **Hantavirus outbreak in Western Europe: reservoir host infection dynamics related to human disease patterns.** *Epidemiol Infect* 2011, **139**:381-390.
64. Tian H, Yu P, Cazelles B, Xu L, Tan H, Yang J, Huang S, Xu B, Cai J, Ma C *et al.*: **Interannual cycles of Hantaan virus outbreaks at the human-animal interface in Central China are controlled by temperature and rainfall.** *Proc Natl Acad Sci U S A* 2017, **114**:8041-8046.
A unique field surveillance dataset from Central China, covering half a century, combining rodent density data, epidemiological surveillance and climate variables. Highlights the impact of El Niño/Southern Oscillation on epidemic patterns.
65. Khalil H, Olsson G, Ecker F, Evander M, Hjertqvist M, Magnusson M, Lofvenius MO, Hörnfeldt B: **The importance of bank vole density and rainy winters in predicting Nephropathia Epidemica incidence in Northern Sweden.** *PLOS ONE* 2014, **9** <http://dx.doi.org/10.1371/journal.pone.0111663>.
66. Sane J, Ollgren J, Makary P, Vapalahti O, Kuusi M, Lyytikäinen O: **Regional differences in long-term cycles and seasonality of Puumala virus infections, Finland, 1995–2014.** *Epidemiol Infect* 2016, **144**:2883-2888.
67. Wood JLN, Leach M, Waldman L, Macgregor H, Fooks AR, Jones KE, Restif O, Dechmann D, Hayman DT, Baker KS *et al.*: **A framework for the study of zoonotic disease emergence and its drivers: spillover of bat pathogens as a case study.** *Philos Trans R Soc B* 2012, **367**:2881-2892.
68. Holmes EC, Zhang YZ: **The evolution and emergence of hantaviruses.** *Curr Opin Virol* 2015, **19**:27-33.