Accepted Manuscript

Expansion of the clinical spectrum of frontometaphyseal dysplasia 2 caused by the recurrent mutation p.Pro485Leu in MAP3K7

Alice Costantini, Carina Wallgren-Pettersson, Outi Mäkitie

PII: S1769-7212(18)30030-2

DOI: [10.1016/j.ejmg.2018.04.004](https://doi.org/10.1016/j.ejmg.2018.04.004)

Reference: EJMG 3445

To appear in: *European Journal of Medical Genetics*

Received Date: 12 January 2018

Revised Date: 14 March 2018

Accepted Date: 10 April 2018

Please cite this article as: A. Costantini, C. Wallgren-Pettersson, O. Mäkitie, Expansion of the clinical spectrum of frontometaphyseal dysplasia 2 caused by the recurrent mutation p.Pro485Leu in MAP3K7, *European Journal of Medical Genetics* (2018), doi: 10.1016/j.ejmg.2018.04.004.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Expansion of the clinical spectrum of frontometaphyseal dysplasia 2 caused by the recurrent mutation p.Pro485Leu in MAP3K7

Alice Costantini¹, Carina Wallgren-Pettersson^{2,3}, Outi Mäkitie^{1, 2, 4, 5}

¹Department of Molecular Medicine and Surgery and Center for Molecular Medicine, Karolinska Institutet, Stockholm, Sweden

²The Folkhälsan Institute of Genetics, Biomedicum Helsinki, Finland

³Department of Medical and Clinical Genetics, University of Helsinki, Helsinki, Finland

⁴Children's Hospital, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁵Department of Clinical Genetics, Karolinska University Hospital, Stockholm, Sweden

Corresponding author: Alice Costantini, Department of Molecular Medicine and

Surgery, Center for Molecular Medicine (CMM) L8:02, Karolinska University

Hospital Solna, 171 76 Stockholm, Sweden

E-mail address: alice.costantini@ki.se

Conflicts of interest

Alice Costantini, Carina Wallgren-Pettersson and Outi Mäkitie declare no conflict of interest.

Supplemental data are included in the submission.

ABSTRACT

Frontometaphyseal dysplasia 2 (FMD2) is a skeletal dysplasia with supraorbital hyperostosis combined with undermodeling of the bones, joint contractures and some extraskeletal features. It is caused by heterozygous mutations in *MAP3K7*, encoding the Mitogen-Activated Protein 3-Kinase 7. MAP3K7 is activated by TGF- β and plays an important role in osteogenesis. Less than 20 patients with FMD2 and *MAP3K7* mutations have been described thus far. The majority of the patients harbor a recurrent missense mutation, NM_003188.3: c.1454C>T [NP_003179.1: p.(Pro485Leu)], which leads to a more severe phenotype than mutations in other domains. Here we describe an additional patient with FMD2 caused by the recurrent c.1454C>T *MAP3K7* mutation, identified as a *de novo* variant by whole-genome sequencing. The 17-year-old boy has the characteristic skeletal and facial features of FMD2. However, some novel features were also observed, including growth retardation and spina bifida occulta. In line with other patients harboring the same mutation he also showed keloid scars and had no intellectual disability. This report expands the clinical spectrum of FMD2 caused by the recurrent c.1454C>T [p.(Pro485Leu)] mutation in *MAP3K7*.

Key words: MAP3K7; TAK1; frontometaphyseal dysplasia 2; skeletal dysplasia; growth retardation

41 1 INTRODUCTION

42 Skeletal dysplasias constitute a large and heterogeneous group of disorders characterized by a
43 broad phenotypic and genetic variability (Bonafe et al., 2015). Short stature is the main hallmark
44 but many other features, including deformities and extra-skeletal manifestations, may also be
45 present. Abnormalities in signaling pathways crucial for the growth plate, i.e. Notch, WNT, FGF,
46 Hedgehog, and BMP pathways, are common causes of skeletal dysplasias (Geister and Camper,
47 2015).

48 Recently, heterozygous disease-causing mutations in *MAP3K7*, encoding the Mitogen-Activated
49 Protein Kinase 7 (also known as TAK1), were described in Frontometaphyseal dysplasia 2 (FMD2)
50 in less than 20 patients (Wade et al., 2016; Wade et al., 2017). This kinase is stimulated by TGF- β
51 and BMPs and plays an important role in the NF- κ B signaling and is consequently a key regulator
52 of osteogenesis (Thouverey and Caverzasio, 2012).

53 Phenotypically, patients with FMD2 are very similar to patients with frontometaphyseal dysplasia
54 1 (FMD1), caused instead by mutations in *FLNA* (Robertson et al., 2003). The shared clinical
55 characteristics include sclerosis of the skull and long bones, prominent supraorbital ridges and
56 abnormal fingers. Some patients also present with extra-skeletal manifestations, such as hearing
57 loss, urogenital problems and joint contractures (Wade et al., 2016). In addition, patients with
58 FMD2 also seem to have a predisposition to keloid scars, especially if they are females (Wade et
59 al., 2017).

60 In some aspects, patients with FMD2 also resemble patients with cardiospondylocarpofacial
61 syndrome, which is also caused by *MAP3K7* mutations (Le Goff et al., 2016). However, this

syndrome is characterized by distinct phenotypic features such as growth retardation, brachydactyly, short limbs and joint laxity (Le Goff et al., 2016).

Here we describe a 17-year-old male with FMD2 caused by the recurrent missense mutation NM_003188.3: c.1454C>T [NP_003179.1: p.(Pro485Leu)] in *MAP3K7* and presenting with several novel phenotypic manifestations.

2 CLINICAL REPORT

2.1 Clinical manifestations

The index patient and his healthy parents were recruited at the Folkhälsan Department of Medical Genetics to a research project exploring genetic causes of skeletal dysplasia. The study protocol was approved by the Research Ethics Board at Helsinki University Hospital and informed consents were obtained from the participants prior to the study.

The patient, a boy born to non-consanguineous Finnish parents, presented at birth with tetralogy of Fallot, a clavicular fracture, ulnar deviation of the wrists, contractures of the elbows, pes metatarsovarus, and hydronephrosis due to ureteral reflux (Table 1). The skull was somewhat dolichocephalic with a sloping forehead, the fontanelle was large, and vertebrae C3 and C4 had reduced height. The fingers and toes were long, with broad distal phalanges of the fingers. His birth weight, length and head circumference (3680g / 51cm / 35.5 cm) were normal, and he received 9 Apgar points.

His slight dysmorphic features included hypertelorism, down-slanting palpebral fissures, broad nasal root, full cheeks, grooved philtrum, small mouth, a H-shaped impression in the chin, and

single palmar creases and camptodactyly in both hands. Hemangiomas were present on the forehead and the occiput

and the neck. He was treated with von Rosen's cast for hip dysplasia and joint contractures and underwent corrective cardiac surgery at the age of 6 months.

On examination at age 1 year, in addition to the above-mentioned dysmorphic features, he was noted to have colobomas of the nostrils. Freeman-Sheldon syndrome was suspected. At the age of 3 years, he was observed to have a submucosal cleft palate and a hearing deficit requiring hearing aids.

At 6 years, the boy was found to have a craniocervical malformation (a sharp angle between the clivus and the dens, and caudal displacement of the conus) in combination with Chiari I-malformation. He underwent occipito-cervical fusion to stabilize his neck (Fig. 1). Decompression of the foramen magnum was performed during the same operation because of the Chiari I-malformation. He was found to have spina bifida occulta in several vertebrae and bilateral Sprengel deformities. The Halo bracing, used for stabilization, caused severe keloid formation in his back (Fig. 1D). He also underwent surgery for unilateral cryptorchidism. His orbital ridges were increasingly prominent (Fig. 1A-B) and a corrective surgery was performed at 11 years to flatten the protruding bony parts of the forehead.

The patient had always been sensitive to light but initial examinations revealed no etiology. At the age of 6, he was found to have subepithelial corneal scarring (leucoma corneae), astigmatism, and amblyopia.

At 7 years, he had short stature (-1.5 SD) (Supplemental Fig. S1), scoliosis (Fig. 1C), slightly weak voice, and mild atopy, and was prone to infections. Intellectual development and school performance were normal.

His growth, which decelerated already starting from 3-4 years of age when scoliosis was not present, has progressed slowly despite normal pubertal development (Supplemental Fig. S1). By age 17 years, his scoliosis has significantly progressed (Cobb's angle 42°) and may partly explain the lack of pubertal growth spurt and his severe height deficit (148 cm; -4.3 SDS). He has flexion contractures in the knees, hips and elbows and his feet are deformed and rigid.

2.2 Genetic findings

In order to identify the genetic cause in our index patient, we performed whole-genome sequencing (WGS) on the trio. Pair-end sequencing (2x150 bp) was performed at the SciLifeLab (Stockholm) on the HiSeqX instrument (Illumina), with an average coverage of 30X. Read alignment to the human genome assembly (GRCh37) was performed with the Burrow-Wheeler Aligner (BWA); data processing and variant calling were carried out using Genome Analysis Toolkit (GATK). Variants were annotated using Variant Effect Predictor (VEP) and data analysed using GEMINI. Candidate gene variants were filtered using a MAF < 0.001 in GnomAD and SweGen databases and assuming autosomal recessive/compound heterozygous inheritance pattern or a *de novo* variant (Ameur et al., 2017). Furthermore, we chose the impact severity of the variant to be different than "low" in GEMINI. Findings were validated by Sanger sequencing.

Whole-genome sequencing in our trio led to the identification of two heterozygous candidate genetic variants in the index patient, both identified as *de novo*: 1) a recurrent missense variant NM_003188.3: c.1454C>T [NP_003179.1: p.(Pro485Leu)] in MAP3K7 (Supplemental Fig. S2A) and

2) a novel missense variant NM_020120.3: c.152C>T [NP_064505.1: p.(Leu51Pro)] in *UGGT1*, encoding the UDP-Glucose Glycoprotein Glucosyltransferase 1. *UGGT1* is involved in the quality control of protein folding in the endoplasmic reticulum (Arnold et al., 2000). However, mutations in this gene have not been associated with any disease yet. On the other hand, mutations in *MAP3K7* have been described in two skeletal diseases: FMD2 and cardiospondylocarpofacial syndrome. Based on previous studies and phenotypic correlations, the variant in *MAP3K7* was regarded as the cause of our patient's disorder, confirming a diagnosis of FMD2. This finding, which was also validated with Sanger sequencing (Supplemental Fig. S2B-S2D), allowed us to re-evaluate our patient's clinical features and compare with previously reported patients with the same recurrent p.(Pro485Leu) mutation in *MAP3K7*.

3. DISCUSSION

Our patient had many features in common with the previously reported patients, including prominent supraorbital ridges, hypertelorism, down-slanting palpebral fissures, broad nasal bridge, full cheeks, micrognathia, hydronephrosis, cleft palate, hearing loss, scoliosis, ulnar deviation of the hands, camptodactyly, wrist contractures, long fingers, structural cardiac defect and keloid scars (Table 1, Fig. 1) (Basart et al., 2015; Morava et al., 2003; Wade et al., 2016; Wade et al., 2017). Radiologically, he had a craniocervical malformation, Chiari I-malformation as well as progressive scoliosis requiring surgery, and dislocated and abnormally shaped radial heads.

He did not have valgus but varus deformity of the feet and he had normal intellect. Moreover, he had some features not previously noted in patients with the same mutation: significant growth retardation, only partly explained by scoliosis, resulting in severe short stature (height -4.3 SDS at 17 years) (Supplemental Fig. S1), Chiari I malformation, Sprengel deformity, spina bifida occulta,

single palmar creases, pes cavus, and amblyopia. Amblyopia is common in the general population and it might be a coincidental finding in our patient. Furthermore, Chiari I malformation, Sprengel deformity and single palmar creases are similar to contractures and skeletal deformities described in other individuals with FMD (Basart et al., 2015; Robertson et al., 2003; Wade et al., 2016; Wade et al., 2017). In line with this, spina bifida occulta is part of the cervical spine dysraphism in FMD but it has never been reported hitherto in patients with *MAP3K7* mutations. Finally, leucoma corneae and astigmatism were observed previously in one patient with a *MAP3K7* mutation (Basart et al., 2015), suggesting that this feature may be a part of the phenotypic spectrum of FMD2.

The majority of the patients hitherto reported as having FMD2 harbor the same recurrent mutation as the one identified in our patient, c.1454C>T [p.(Pro485Leu)]. This mutation affects the TAB2 domain of the protein (Table 1; Supplemental Fig.S3) (Wade et al., 2016; Wade et al., 2017). Only 3 patients have been described as having other missense mutations (Wade et al., 2016; Wade et al., 2017). These mutations cluster closer to the N-terminal, in the tyrosine kinase domain, and give rise to milder phenotypes than the ones caused by the recurrent mutation (Table 1; Supplemental Fig.S3). Although the majority of features in our patient overlap with the ones previously described in other FMD2 patients, his phenotype also includes some novel features (Table 1). For example, short stature has not been previously reported in FMD2. However, short stature is one of the hallmarks of patients with cardiospondylocarpofacial syndrome, which is also caused by *MAP3K7* mutations (Le Goff et al., 2016). Furthermore, our patient has additional hand and foot deformities as well as spina bifida occulta, vision problems, undescended testis and hemangiomas that have not previously been described in other patients with FMD2.

In summary, our detailed report corroborates many of the clinical features previously described in patients with FMD2 caused by the recurrent *MAP3K7* mutation c.1454C>T [p.(Pro485Leu)] but also describes some novel or unusual characteristics, in particular growth retardation, which may suggest a partial overlap between FMD2 and cardiospondylocarpofacial syndrome.

ACKNOWLEDGMENTS

The authors thank the Sigrid Jusélius Foundation, the Association Française contre les Myopathies, the Finska Läkaresällskapet, the Folkhälsan Institute of Genetics, the Medicinska understödsföreningen Liv och Hälsa, the Academy of Finland, the Swedish Research Foundation, the Novo Nordisk Foundation and Stockholm County Council (ALF) for funding. The authors would also like to acknowledge support from the National Genomics Infrastructure, NGI, and UPPMAX for providing assistance in massive parallel sequence analysis and computational infrastructure.

184 **WEB RESOURCES**185 Burrow-Wheeler Aligner: <http://bio-bwa.sourceforge.net>186 GATK: <https://software.broadinstitute.org/gatk/>187 VEP: <http://www.ensembl.org/info/docs/tools/vep/index.html>188 GEMINI: <https://gemini.readthedocs.io/en/latest/>189 GnomAD: <http://gnomad.broadinstitute.org>190 SweGen: <https://swefreq.nbis.se>

191

192

193 REFERENCES

- 194 Ameer, A., Dahlberg, J., Olason, P., Vezzi, F., Karlsson, R., Martin, M., Viklund, J., Kahari, A.K.,
 195 Lundin, P., Che, H., Thutkawkorapin, J., Einfeldt, J., Lampa, S., Dahlberg, M., Hagberg, J.,
 196 Jareborg, N., Liljedahl, U., Jonasson, I., Johansson, A., Feuk, L., Lundeberg, J., Syvanen, A.C.,
 197 Lundin, S., Nilsson, D., Nystedt, B., Magnusson, P.K., Gyllensten, U., 2017. SweGen: a whole-
 198 genome data resource of genetic variability in a cross-section of the Swedish population.
 199 *European Journal of Human Genetics* 25(11), 1253-1260.
- 200 Arnold, S.M., Fessler, L.I., Fessler, J.H., Kaufman, R.J., 2000. Two homologues encoding human
 201 UDP-glucose:glycoprotein glucosyltransferase differ in mRNA expression and enzymatic
 202 activity. *Biochemistry* 39(9), 2149-2163.
- 203 Basart, H., van de Kar, A., Ades, L., Cho, T.J., Carter, E., Maas, S.M., Wilson, L.C., van der Horst,
 204 C.M., Wade, E.M., Robertson, S.P., Hennekam, R.C., 2015. Frontometaphyseal dysplasia and
 205 keloid formation without FLNA mutations. *American journal of medical genetics. Part A*
 206 167(6), 1215-1222.
- 207 Bonafe, L., Cormier-Daire, V., Hall, C., Lachman, R., Mortier, G., Mundlos, S., Nishimura, G.,
 208 Sangiorgi, L., Savarirayan, R., Sillence, D., Spranger, J., Superti-Furga, A., Warman, M., Unger, S.,
 209 2015. Nosology and classification of genetic skeletal disorders: 2015 revision. *American*
 210 *journal of medical genetics. Part A* 167A(12), 2869-2892.
- 211 Geister, K.A., Camper, S.A., 2015. Advances in Skeletal Dysplasia Genetics. *Annu Rev Genomics*
 212 *Hum Genet* 16, 199-227.
- 213 Le Goff, C., Rogers, C., Le Goff, W., Pinto, G., Bonnet, D., Chrabieh, M., Alibeu, O., Nistchke, P.,
 214 Munnich, A., Picard, C., Cormier-Daire, V., 2016. Heterozygous Mutations in MAP3K7,
 215 Encoding TGF-beta-Activated Kinase 1, Cause Cardiospondylocarpofacial Syndrome.
 216 *American journal of human genetics* 99(2), 407-413.
- 217 Morava, E., Illes, T., Weisenbach, J., Kartesz, J., Kosztolanyi, G., 2003. Clinical and genetic
 218 heterogeneity in frontometaphyseal dysplasia: severe progressive scoliosis in two families.
 219 *American journal of medical genetics. Part A* 116A(3), 272-277.
- 220 Robertson, S.P., Twigg, S.R.F., Sutherland-Smith, A.J., Biancalana, V., Gorlin, R.J., Horn, D.,
 221 Kenwright, S.J., Kim, C.A., Morava, E., Newbury-Ecob, R., Orstavik, K.H., Quarrell, O.W.J.,
 222 Schwartz, C.E., Shears, D.J., Suri, M., Kendrick-Jones, J., Wilkie, A.O.M., Co, O.-S.D.C., 2003.
 223 Localized mutations in the gene encoding the cytoskeletal protein filamin A cause diverse
 224 malformations in humans. *Nature genetics* 33(4), 487-491.
- 225 Thouverey, C., Caverzasio, J., 2012. The p38alpha MAPK positively regulates osteoblast
 226 function and postnatal bone acquisition. *Cell Mol Life Sci* 69(18), 3115-3125.
- 227 Wade, E.M., Daniel, P.B., Jenkins, Z.A., McInerney-Leo, A., Leo, P., Morgan, T., Addor, M.C., Ades,
 228 L.C., Bertola, D., Bohring, A., Carter, E., Cho, T.J., Duba, H.C., Fletcher, E., Kim, C.A., Krakow, D.,
 229 Morava, E., Neuhaus, T., Superti-Furga, A., Veenstra-Knol, I., Wiczorek, D., Wilson, L.C.,
 230 Hennekam, R.C., Sutherland-Smith, A.J., Strom, T.M., Wilkie, A.O., Brown, M.A., Duncan, E.L.,
 231 Markie, D.M., Robertson, S.P., 2016. Mutations in MAP3K7 that Alter the Activity of the TAK1
 232 Signaling Complex Cause Frontometaphyseal Dysplasia. *American journal of human genetics*
 233 99(2), 392-406.
- 234 Wade, E.M., Jenkins, Z.A., Daniel, P.B., Morgan, T., Addor, M.C., Ades, L.C., Bertola, D., Bohring,
 235 A., Carter, E., Cho, T.J., de Geus, C.M., Duba, H.C., Fletcher, E., Hadzsiev, K., Hennekam, R.C.M.,
 236 Kim, C.A., Krakow, D., Morava, E., Neuhaus, T., Sillence, D., Superti-Furga, A., Veenstra-Knol,
 237 H.E., Wiczorek, D., Wilson, L.C., Markie, D.M., Robertson, S.P., 2017. Autosomal dominant

238 frontometaphyseal dysplasia: Delineation of the clinical phenotype. American journal of
239 medical genetics. Part A.
240

241

242 **FIGURE LEGENDS**

243 **Figure 1.** At 11 years the patient had significant supraorbital hyperostosis (A), as seen also in the
244 preoperative CT at 11 years (B). He required hearing aids for hearing deficit, had atopic eczema
245 and abnormal scapulae (A). He underwent occipito-cervical fusion to stabilize the neck at 6 years
246 and had by 8 years developed significant scoliosis (C). Halo bracing resulted in keloid formation in
247 the back (D). Radiograph of the pelvis and hips at 3 years (E) showed dysplastic hips and coxa
248 valga. Hand radiograph at 16 years (F) showed undertubulation of the digital bones. Cavovarus
249 deformities were seen in radiographs (G) and clinically (H, I).

250

Table 1. Summary of the clinical features in the 19 thus far reported index patients with FMD2 due to *MAP3K7* mutations. Mutations are given as amino acid changes. The novel or unusual clinical features in our patient are marked in blue text. Colours indicate features that were present (yellow), absent (dark blue) or unknown (grey) in the patients. F= female; M=male; D= deceased.

[illegible]