<https://helda.helsinki.fi>

Improving forecasts of arctic-alpine refugia persistence with landscape-scale variables

Niskanen, Annina K. J.

2017-02

Niskanen , A K J , Heikkinen , R K , Mod , H K , Väre , H & Luoto , M 2017 , ' Improving forecasts of arctic-alpine refugia persistence with landscape-scale variables ' , Geografiska Annaler. Series A. Physical Geography , vol. 99 , no. 1 , pp. 2-14 . <https://doi.org/10.1080/04353676.2016.1256746>

<http://hdl.handle.net/10138/307291>

<https://doi.org/10.1080/04353676.2016.1256746>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **Improving forecasts of arctic-alpine refugia persistence with landscape-scale variables**

2
3 Annina K.J. Niskanen¹, Risto K. Heikkinen², Heidi K. Mod¹, Miska Luoto¹

4 ¹Department of Geosciences and Geography, P.O. Box 64 (Gustaf Hållströmin katu 2a) , FI-00014 University of Helsinki,
5 Finland; ² Finnish Environment Institute, Natural Environment Centre, P.O. Box 140, FI-00251, Helsinki, Finland

6
7 Niskanen A.K.J., Heikkinen R.K., Mod H.K. and Luoto M., 20xx.. Improving forecasts of arctic-alpine refugia
8 persistence with landscape-scale variables. *Geografiska Annaler, Series A: Physical Geography*.

9 **ABSTRACT**

10 Refugia, sites preserving conditions reminiscent of suitable climates, are projected to be crucial for species in a changing
11 climate, particularly at high latitudes. However, the knowledge of current locations of high-latitude refugia and
12 particularly their ability to retain suitable conditions under future climatic changes is limited. Occurrences of refugia have
13 previously been mainly assessed and modelled based solely on climatic features, with insufficient attention being paid to
14 potentially important landscape-scale factors. Here, climate-only models and ‘full’ models incorporating topo-edaphic
15 landscape-scale variables (radiation, soil moisture and calcareousness) were developed and compared for 111 arctic-
16 alpine plant species in Northern Fennoscandia. This was done for both current and future climates to determine cells with
17 resilient climatic suitability harbouring refugia. Our results show that topographic and edaphic landscape-scale predictors
18 both significantly improve models of arctic-alpine species distributions and alter projections of refugia occurrence.
19 Predictions of species-climate models ignore landscape-scale ecological processes and may thus provide inaccurate
20 estimates of extinction risk and forecasts of refugia where species can persist under a changing climate.

21 *Keywords:* climate change; refugia; landscape-scale factor

22 **Introduction**

23 Previous research has shown the importance of refugia for species survival during past environmental changes (Taberlet,
24 1998; Birks & Willis, 2008; Svenning *et al.*, 2008). Refugia are sites that provide environmental conditions deviant from
25 the regional average, thereby supporting species persistence in a changing climate (Médail & Diadema, 2009; Keppel *et*
26 *al.*, 2012). Due to the rate and magnitude of global change, refugia are also likely to be among the major mechanisms
27 assisting species persistence in the future (Reside *et al.*, 2013) and should thus be acknowledged in climate change–based
28 conservation planning. A challenge still remains in the credible identification of present-day refugia across landscapes,
29 and particularly in assessing their capability to retain suitable conditions under future climatic changes. Spatial

information regarding future refugia is particularly important at high latitudes, as they are among the most susceptible environments to a changing climate (Root *et al.*, 2003; Ashcroft, 2010). However, little is known about which high-latitude landscapes and species are most vulnerable to projected future climatic trends. This knowledge gap calls for increased attention on robust forecasts of how the suitable areas for the species might change, which environmental factors are most critical for developing the most reliable models of this change, and how well potential future refugia are discernible from a given landscape.

Species distribution models (SDMs) are a commonly used tool to forecast the spatial changes in realized or potential species distributions (Guisan & Thuiller, 2005; Araújo & Peterson, 2012). SDMs have also been shown to complement and be spatially correlated with traditionally used phylogeographic estimates of past refugia distribution while providing less subjective and spatially more explicit predictions (Waltari *et al.*, 2007). However, it is increasingly recognized that SDMs often require both climatic variables and non-climatic variables describing landscape-scale or local environmental conditions to maximize predictive ability (Franklin, 1995; Austin & Van Niel, 2011a; Beauregard & de Blois, 2014). Such integrative modelling studies have not, however, been carried out for arctic-alpine species or their refugia in high-latitude regions. This is a shortcoming as the inclusion of landscape-scale factors into SDMs developed for high-latitude refugia may be of particular significance and help identify locations able to preserve environmental conditions reminiscent of the current climate within the surrounding changing matrix, thus leading to more robust predictions of future refugia (Austin & Van Niel, 2011a; Hodd *et al.*, 2014).

A frequently applied paradigm in the study of global change impacts on biota involves models of the distributional changes of species – or the areas environmentally suitable for them – as spatially explicit functions of projected change (Guisan & Thuiller, 2005; Botkin *et al.*, 2007). Similarly, previous studies using spatial models of predicted species distributions have mainly approached refugia from the viewpoint of a single species (Hugall *et al.*, 2002; Austin & Van Niel, 2011a), past distributions (Fløjgaard *et al.*, 2009) or coarser spatial scales (Hodd *et al.*, 2014). Here, we investigate whether the addition of landscape-scale variables to climate-only SDMs (1) leads to different interpretations of how projected changes in climate may affect the distributions of 111 arctic-alpine plant species, and (2) modifies predictions of in situ refugia for these species in N-Fennoscandia. We base these analyses on high-quality species and environmental data, and predict species distributions under contemporary climate as well as two projected future climates i.e. Representative Concentration Pathway (RCP) scenarios: the moderate RCP 4.5 (Smith & Wigley, 2006; Clarke *et al.*, 2007; Wise *et al.*, 2009) and extreme RCP 8.5 scenarios (Riahi *et al.*, 2007).

58 **Materials and methods**

59 *Study area*

60 Located in Northern Fennoscandia between 67°N and 70°N, the sub-arctic climate of our study region is driven by its
61 position at the edge of the Eurasian continent, the Polar Front, the warm North Atlantic current, and proximity to the
62 Scandes Mountains (Fig. 1) (Tikkanen, 2005; Aalto *et al.*, 2014b). The area is characterized by varying climatic,
63 topographic and geologic gradients (Oksanen & Virtanen, 1995). Average July temperatures range from 6.1 °C to 15.2 °C
64 and mean annual precipitation from 449 mm to 600 mm (1981 – 2010 means) (Pirinen *et al.*, 2012). Elevation ranges
65 from 72 to 1365 m.a.s.l. The vegetation gradient varies from spruce (*Picea abies*) and Scots pine (*Pinus sylvestris*) forests
66 in the south to mountain birch (*Betula pubescens* subsp. *czerepanovii*) in the north, with shrub-dominated tundra-like
67 vegetation above the tree-line (Sormunen *et al.*, 2011).

68 Preliminary data analysis indicated that parts of our study area are predicted to have a future climate with
69 no current equivalent within the study region. To reduce errors from predicting species responses to climatic conditions
70 without current equivalents, and as arctic-alpine species are generally expected to move northwards as the climate warms,
71 we used data from the whole area for training the models, but excluded the non-analogue southern- and easternmost
72 regions of our study area from model projections (dashed line in Fig. 1) (Sætersdal *et al.*, 1998; Heikkinen *et al.*, 2006).

73 *Species data*

74 Species occurrence data for 1341 1 km² cells were collected by botanists and volunteers, and refined using scientific
75 literature and herbaria. The sampling cells cover all important vegetation zones and ecosystems present in the study area,
76 ranging from taiga to treeless tundra (Kurtto & Lampinen, 1999). Only arctic-alpine species (defined according to
77 biogeographic distribution by professional botanists; see le Roux *et al.* (2012)) with an occurrence in a minimum of eight
78 cells within our study area were included in the subsequent analysis ($n=111$).

79 *Selection of predictors*

80 We used an extensive grid of environmental data matching the resolution of the species data (1 km²) to investigate key
81 refugia predictors. We chose to demonstrate our methodology using a set of climate and landscape-scale terrain and
82 geological variables (Table 1) known to be important for plant species (Skov & Svenning, 2004; Austin & Van Niel,
83 2011a; Dobrowski, 2011; Scherrer & Körner, 2011). The chosen direct and resource variable predictors (Austin & Smith,
84 1990) represent growing conditions, surface temperature, light, moisture and soil conditions considered critical for high-
85 latitude vegetation.

86 The climate data, comprising of observations from 61 stations in Northern Fennoscandia, were acquired
87 from the national observation networks of Finland (Finnish Meteorological Institute), Sweden (Swedish Meteorological
88 and Hydrological Institute) and Norway (Norwegian Meteorological Institute), and modelled to a 1 km² resolution grid
89 (see Aalto *et al.*, 2014). Three climatic predictors with a temporal coverage of 1981–2010 were initially included in the
90 analyses: growing degree days (GDD3; annual accumulated daily temperature sum above 3°C, i.e. growing conditions),
91 temperature of the coldest quarter (TCQ; overwintering conditions) and water balance (WAB; moisture conditions).
92 However, due to ecologically erroneous projections resulting from the counter effects of different climatic variables
93 caused by variable multicollinearity, (namely strong correlations of GDD3 with TCQ and WAB [Spearman's rank
94 correlation coefficients, $r_s = -0.88$ and $r_s = -0.87$, respectively]), only GDD3, which correlates the most with arctic-alpine
95 species diversity in the study area ($r_s = -0.75$), was ultimately included in the final models.

96 Landscape features exert a strong influence through numerous geomorphological, hydrological,
97 geological and biological processes (Moore *et al.*, 1991), thus controlling the growing conditions experienced by plants
98 (Guisan *et al.*, 1998; Ackerly *et al.*, 2010; Scherrer & Körner, 2011). Here, three topo-edaphic landscape-scale predictors
99 were used: (1) incoming potential solar *radiation* representing surface temperature conditions (McCune & Keon, 2002);
100 (2) digital elevation model-derived topographic wetness index (*TWI*) representing the availability of soil moisture from
101 upslope areas (Beven & Kirkby, 1979); and (3) *calcareousness* representing the proportion of calcareous, nutrient-rich
102 bedrock in a given 1 km² grid cell (Dubuis *et al.*, 2013). In contrast to GDD3, the relative effects of radiation, fluvial
103 processes (e.g. *TWI*) and soil properties (e.g. *calcareousness*) should remain more constant in the future (Reside *et al.*,
104 2013). Radiation and *TWI* were based on an Aster -derived digital elevation model (NASA Land Processes Distributed
105 Active Archive Center (2013); spatial resolution 30 m²). Radiation was calculated using the ArcView 3.2 Solar analyst
106 extension which uses digital elevation models to generate radiation data by accounting for latitude, elevation, slope angle,
107 slope aspect, shadows cast by surrounding topography, daily and seasonal shifts in solar angle, and atmospheric
108 attenuation (Fu & Rich, 1999). Due to the landscape scale of this study, only one latitude value (here, 69°) for the whole
109 digital elevation model was used. *TWI*, the most often used surrogate for soil moisture, was calculated using a Python
110 script written by Prasad Pathak that accounts for slope and local upslope contributing area (Beven & Kirkby, 1979; Esri,
111 2013). *Calcareousness* was reclassified from a digital database (Geological Survey of Finland 2010; spatial resolution 20
112 m²). The landscape predictors were resampled by spatial averaging to a 1 km² resolution to match the species and climate
113 data.

115 Forecasts for the locations of refugia may vary between climate scenarios, thus making it imperative to consider
 116 predictions based on multiple scenarios (Reside *et al.*, 2013). Four sets of scenarios containing different emission,
 117 concentration and land-use trajectories, have been developed (Van Vuuren *et al.*, 2011). These scenario sets are referred
 118 to as Representative Concentration Pathways (RCPs) and they correspond to different greenhouse gas (GHG)
 119 concentration trajectories (Moss *et al.*, 2010). Here we explore the implications of two possible future pathways, RCP 4.5
 120 and RCP 8.5 (where the number refers to radiative forcing in watts/m² by the year 2100). RCP 4.5 is an intermediate
 121 mitigation scenario where GHG emissions are stabilized at low to medium levels (Van Vuuren *et al.*, 2011). RCP 8.5, the
 122 unabated ‘business-as-usual’ scenario, represents the highest GHG emission profile (Riahi *et al.*, 2011; Van Vuuren *et*
 123 *al.*, 2011).

124 *Statistical analyses*

125 Observed species distributions were related to the set of predictors using five statistical modelling techniques. These
 126 included two regression methods: generalized linear modeling (GLM) (McCullagh & Nelder, 1989) and generalized
 127 additive modeling (GAM) (Hastie & Tibshirani, 1990) and three machine-learning methods: generalized boosting
 128 methods (GBM, also known as boosted regression trees (BRT)) (Elith *et al.*, 2008), random forest (RF) (Breiman, 2001)
 129 and maximum entropy (MAXENT) (Phillips *et al.*, 2004). All the models were implemented in the Biomod2 platform
 130 (Thuiller *et al.*, 2013) using R-program (R Development Core Team 2013). The models were fitted using two sets of
 131 predictors:

$$132 \quad \text{Occurrence of species} = \text{GDD3}$$

133 [base model]

$$134 \quad \text{Occurrence of species} = \text{GDD3} + \text{CALC} + \text{RAD} + \text{TWI}$$

135 [full model]

136 Model transferability was assessed using four-fold cross-validation. We applied a commonly used approach
 137 (Zimmermann *et al.*, 2009; le Roux *et al.*, 2013b; Aalto & Luoto, 2014) to compare the predictive power of base (climate-
 138 only) and full (climate-plus-landscape) models by calculating the area under the curve of a receiver operating
 139 characteristic (AUC) plot (Fielding & Bell, 1997), true skill statistics (TSS) and Cohen’s Kappa coefficient (Allouche *et*
 140 *al.*, 2006) based on the four evaluation runs (where the models calibrated with 75% of the data were evaluated with the
 141 withheld 25 % of the data, and repeated for each of the four quarters of the data). A non-parametric Wilcoxon’s test was
 142 employed to examine whether explanatory power and predictive accuracy differed significantly between models.

143 All data were used for projecting species occurrences to current and future climatic conditions.
144 Occurrence probabilities were transformed to binary presence/absence predictions using a TSS cutoff, maximizing model
145 accuracy, defined by BioMod2. In order to account for intermodal variability and to create the final maps of predicted
146 species distributions, we used the consensus approach (Araújo & New, 2007) to construct an ensemble of forecasts that
147 combine the binary predictions from all models. Here we chose a majority's vote of a minimum of three out of five
148 modelling techniques to denote a presence value for a given species inside a given 1 km² grid cell (Figure S1).

149 For each species, variable influence was assessed in Biomod2 by randomizing each variable individually
150 and then projecting the model with the randomized variable while keeping the other variables unchanged. The model
151 predictions containing the randomized variable were then correlated with those of the original models. Finally, the
152 importance of the variable was calculated as one minus the correlation, with higher values indicating higher predictor
153 importance (Thuiller *et al.*, 2009). This was repeated ten times for each modelling method and the resulting variable
154 influence values were averaged.

155 Definition of refugia

156 Areas where the contemporary and future distributions of species overlap may act as refugia (Temunović *et al.*, 2013).
157 By modelling contemporary and projected climates, we can identify these spatiotemporal species distribution overlaps
158 (henceforth referred to as 'resilient cells'). Here, we regard these resilient cells as potential sites for *in situ* refugia where
159 local species may be maintained, i.e. persist even under the assumption of no dispersal (Thomas *et al.*, 2004; Pearson,
160 2006). *In situ* refugia are more likely to function as effective refugia as reaching them requires only local or no range-
161 shifts (Reside *et al.*, 2014) and, by remaining within the present range of a given species, they provide a robust estimate
162 of species persistence (Shoo *et al.*, 2013).

163 To locate *in situ* refugia from SDM predictions we utilized a step-by-step approach as follows (Figure
164 S1): firstly, as refugia are essentially species-specific (Provan & Bennett, 2008; Stewart *et al.*, 2010), we model the species
165 separately to gain insights into their preferred environments. Next, the single-species predictions are projected separately
166 into current climatic conditions and two future climate scenarios to locate cells where species occurrence is maintained
167 both spatially and temporally (Shoo *et al.*, 2013; Temunović *et al.*, 2013). Finally, as favourable environmental conditions
168 supporting refugia may overlap for several species (Hampe & Petit, 2005; Keppel *et al.*, 2012) and to increase prediction
169 robustness, we stack projections of these resilient cells to locate sites where suitable conditions persist for several species.
170 Areas with higher arctic-alpine species diversity are strong refugia candidates as they maximize available information
171 and the probability of this diversity persisting under climate change. Consequently, sites harbouring several refugial

172 species simultaneously are potentially very valuable for future conservation planning. Thus, we set a threshold of a
173 minimum of five persisting species as a prerequisite for a resilient 1 km² grid cell to be classified as a refugia.

174 **Results**

175 The use of topo-edaphic landscape-scale predictors significantly ($p < 0.001$) improves the accuracy of SDMs developed
176 for the studied arctic-alpine species. Based on the cross-validation runs, the mean AUC values improved from 0.77 (base
177 i.e. climate-only model) to 0.86 (full i.e. climate-plus-landscape model). The corresponding TSS values improved from
178 0.51 to 0.62, and Kappa values from 0.39 to 0.46 (Table 2). Lower standard deviations of the evaluation metrics in the
179 full models also indicate greater model accuracy (Table 2).

180 The inclusion of landscape-scale predictors changes contemporary predictions of refugia as well as
181 projections of species distributions and predictions of the availability of suitable environments for arctic-alpine species
182 in a warming climate. Differences arise in both predictions of how many species are predicted to persist in a given refugia,
183 as well as the total count i.e. spatial extent of refugial cells. The full model predicts fewer species range reductions and
184 higher species persistence (Fig 2: b, d; Table S1): out of the 111 species, the full model locates refugia for 77 species in
185 RCP 4.5 (climate-only model: 40 species) and 47 species in RCP 8.5. Considerably less refugia were predicted by the full
186 model into the more extreme RCP 8.5 (207 refugia; compared to 959 in RCP 4.5; Table S1). The base model does not
187 project any refugia into RCP 8.5 (Fig 2: c). The spatial extent of refugial cells predicted by both models into RCP 4.5
188 shows a 79% overlap (Fig 2: a, b). However, over 95% of the refugial cells predicted by the full model for RCP 4.5
189 demonstrate a higher species persistence rate than predictions from the base model (Fig 2: b, d; Table S1).

190 Out of the 111 species included in the analysis, GDD3 was the most influential variable for most of the
191 species ($n=69$); calcareousness for 26 species; TWI for 13 species; and radiation was the most important variable for three
192 species. An example of projected refugia for Alpine Mouse-ear (*Cerastium alpinum*) are shown in Figure S3. This small
193 perennial herb shows a clear preference for high substrate calcareousness and cooler growing conditions offered by the
194 regions refugia. There is a clear spatial prominence of refugial cells around the cooler, more calcareous mountainous
195 regions of the study area (Figs 1 & 2; Table S2; Figure S4: b-d). By including topo-edaphic landscape-scale variables,
196 suitable environments for arctic-alpine species can, however, be projected into cells with a higher GDD3 than by
197 accounting for climate alone. Aside from climatic differences, the refugia derived from full model predictions also exhibit
198 slightly differing landscape conditions, here higher radiation, a higher cover of calcareous substrates and lower TWI
199 (Table S2).

200 Discussion

201 Our results demonstrate more detailed and accurate predictions of refugia following the inclusion of landscape-scale
202 variables reflecting the topo-edaphic conditions to climate-only models. This is due to the limiting effect of using only
203 climatic predictors that overlooks certain fine-resolution spatial and temporal (e.g. extent and persistence of suitable
204 habitat, respectively) aspects of species distributions. In other words, consideration of landscape properties is required to
205 capture the important topo-edaphic conditions shaping refugia distribution patterns in space and time (Austin & Van Niel,
206 2011b), and our study shows that this is true also for high-latitude environments.

207 Refugia defined using both climate and topo-edaphic parameters, compared to refugia defined by climate
208 alone, are more likely to continue to provide suitable conditions for a greater portion of arctic-alpine species growing in
209 a given grid cell (Dobrowski, 2011; Keppel *et al.*, 2012). The inclusion of topo-edaphic variables modifies climate-only
210 models in a way that enables predictions of suitable future conditions into a number of cells with warmer average growing
211 conditions. This significance of accounting for the topo-edaphic setting results partly from how it can modify the climatic
212 conditions experienced by species (Austin & Van Niel, 2011a; Lenoir *et al.*, 2013). As seen for the example species,
213 Alpine Mouse-ear, the climate-only model predicts species persistence in northern refugia, where temperatures are cooler
214 (Figs S2 & S4). The refugia show a more spatially dispersed pattern of persistence in a changed climate, but the forecasts
215 of species persistence are made more optimistic through additions of topographic and edaphic predictors. Moreover,
216 improvements to the models and derived predictions resulting from the incorporation of landscape-scale variables also
217 emphasize an increase in species' tolerance in confronting climatic changes via edaphic conditions (Table S2; Fig S3).
218 The importance of calcareous substrate also shows that adjusting climate models with topographical variables only, e.g.
219 constructing topoclimatic models, may lead to misrepresentative conclusions regarding SDMs and climate change
220 predictions.

221 Possibly owing to the importance of climate as a predictor, the spatial distribution of refugia appears
222 similar between the base and full model predictions for RCP 4.5. This is particularly evident in the northern regions of
223 the study area where GDD3 levels remain favourably low for arctic-alpine species persistence. Areas with the least
224 environmental change are more likely refugia candidates for most species within a grid cell (Reside *et al.*, 2013) and,
225 indeed, considerably less refugia were predicted by the full model into the more extreme RCP 8.5 (207 refugia; compared
226 to 959 in RCP 4.5). The spatial distribution of these 207 refugial cells is more dispersed, suggesting that refugial species
227 in a warmer future would need to rely on more isolated habitats to provide locally favourable landscapes. Though
228 decreased by climatic changes, forecasts of species persistence in refugia are improved by even fairly simple topo-edaphic
229 landscape properties at the mesoscale. Examining such grid cells where species can persist is a robust way of mapping

distributional changes and refugia, though dispersal abilities and possible time lags of species responses to changing landscape configurations are also likely to contribute to the effectiveness of refugia. However, accounting for circumstances of little or no spatial overlap between current and future suitable environments would require information on species dispersal capacities (Reside *et al.*, 2013; Reside *et al.*, 2014) to *ex situ* refugia (Shoo *et al.*, 2013) to enable more realistic estimates of species range-shift abilities to future refugia.

The distributions of a majority of the 111 arctic-alpine species studied here are projected to shrink following increases in temperature with most of the study region appearing as a matrix of unsuitable growing conditions for arctic-alpine species. Heterogeneous mountainous regions, such as the northern areas of our study landscape, are expected to be exceedingly important for refugia and the conservation of species (Loarie *et al.*, 2008; Luoto & Heikkinen, 2008; Scherrer & Körner, 2011), especially for endemic mountain plants at risk from range reduction by upward displacement (Rull & Vegas-Vilarrúbia, 2006). Indeed, our finding that the located refugia are offering conditions of lower soil moisture (as indicated by the variable TWI) is related to the assumption that steeper slopes, such as those found in the Scandes, are less likely to retain water (Beven & Kirkby, 1979). The significance of these mountain refugia, however, is likely to be affected by species' properties such as growth form and dispersal capacity (Engler *et al.*, 2009), and mountain refugia have been disputed to be of little avail to small, scattered populations of species (Birks, 2008). The reduction in available land area at higher elevations may also result in species' local extinctions (Patsiou *et al.*, 2014). Nevertheless, the *in situ* refugial expanse in the mountainous area of the study region is projected to be an important location for providing many arctic-alpine species with suitable conditions for persistence, and its potential to do so deserves further investigation.

More generally, it needs to be acknowledged that evaluating the capability of refugia in promoting species persistence and ecosystem resilience is difficult due to the many uncertainties involved in predicting and modelling climate, changes therein, and species responses to these changes (Wiens *et al.*, 2009; Reside *et al.*, 2013; Shoo *et al.*, 2013). For example, assuming uniform warming across a given landscape, or basing forecasts of changes in local climate on simple temperature-elevation correlations may over-emphasize the importance of relatively cool sites or higher elevations (Ashcroft, 2010). Essential aspects of future refugia research include investigating the dynamics of climatic decoupling between the atmosphere and terrain within arctic-alpine landscapes (Dobrowski, 2011; Scherrer & Körner, 2011; Hylander *et al.*, 2015), as we cannot surmise the effects of topography on climatic stability with the approach used here. A similar modelling framework could also be used to pinpoint individual at-risk species, specific areas or habitat types facing substantial changes in a warmer future (Thuiller *et al.*, 2005; Williams *et al.*, 2005; Loarie *et al.*, 2008). Furthermore, though the landscape-scale predictors used here provide a solid basis for a comprehensive set of ecologically

260 meaningful predictors, enumerating the significance of other variables, such as geomorphological disturbances (Randin
261 *et al.*, 2009; le Roux *et al.*, 2013a; Slaton & Linder, 2015), CO₂ effects (Rickebusch *et al.*, 2008) and biotic interactions
262 (Godsoe *et al.*, 2015; Mod *et al.*, 2015), may also be necessary for more realistic predictions of refugia persistence (sensu
263 Austin and Van Niel 2011b), though their relevance is likely to be further pronounced at finer spatial scales. ur results,
264 highlighting the importance of topo-edaphic landscape-scale factors for modelling refugia as well as demonstrating
265 changes in the extent and connectivity of suitable habitat, show there is a strong case for applying a similar approach at
266 finer or even multiple scales to facilitate a more mechanistic approach to future refugia study.

267 **Conclusions**

268 Climate change projections suggest that returning to historical climatic conditions is highly improbable. Here we show
269 that, despite forecasted future regional warming, several resilient sites found in our study landscape are predicted to
270 maintain suitable local environments due to topo-edaphic conditions. Refugia can be found in places where locally
271 optimal environments remain constant across timescales and in which the effects of landscape preserve environmental
272 conditions more reminiscent of suitable contemporary climates. Predictions of species responses to climate change based
273 on climate alone may be inaccurate in estimating extinction risk as they neglect to consider topographic and edaphic
274 processes affecting species distributions, and may fail to identify refugia where species can persist. Thus, models of
275 species distributions and changes therein need to account for topo-edaphic landscape-scale variables to provide
276 information on relevant spatial and temporal responses to environmental change and should thus be explicitly accounted
277 for in future climate change impact assessment studies.

278 **Acknowledgements**

279 AN was funded by the Lapland Regional Fund (Finnish Cultural Foundation), Societas pro Fauna et Flora Fennica and
280 The Doctoral Program in Geosciences. We acknowledge funding from the Academy of Finland (Project Number
281 1140873). Also, HM was partly founded by the Alfred Kordelin Foundation and The Doctoral Program in Geosciences,
282 and ML was partly funded by the Academy of Finland (Project Number 286950). We would also like to thank the Finnish
283 Museum of Natural History as one of the main contributors of the data on vascular plant species and Juha Aalto for
284 helping with gathering the environmental data and data analysis.

286 *Annina K.J. Niskanen, Department of Geosciences and Geography, P.O. Box 64 (Gustaf Hållströmin katu 2a), FI-00014*
287 *University of Helsinki, Finland*
288 *Email: annina.k.niskanen@helsinki.fi*

290 *Risto K. Heikkinen, Finnish Environment Institute, Natural Environment Centre, P.O. Box 140, FI-00251, Helsinki,*
291 *Finland*
292 *Email: risto.heikkinen@ymparisto.fi*
293
294 *Heidi K. Mod, Department of Geosciences and Geography, P.O. Box 64 (Gustaf Hållströmin katu 2a), FI-00014*
295 *University of Helsinki, Finland*
296 *Email: heidi.mod@helsinki.fi*
297
298 *Miska Luoto, Department of Geosciences and Geography, P.O. Box 64 (Gustaf Hållströmin katu 2a), FI-00014 University*
299 *of Helsinki, Finland*
300 *Email: miska.luoto@helsinki.fi*

301 Aalto, J. & Luoto, M. (2014) Integrating climate and local factors for geomorphological distribution models.
302 *Earth Surface Processes and Landforms*, **39**, 1729-1740.

303 Aalto, J., Le Roux, P.C. & Luoto, M. (2014a) The meso-scale drivers of temperature extremes in high-latitude
304 Fennoscandia. *Climate dynamics*, **42**, 237-252.

305 Aalto, J., Venäläinen, A., Heikkinen, R.K. & Luoto, M. (2014b) Potential for extreme loss in high-latitude
306 Earth surface processes due to climate change. *Geophysical Research Letters*, **41**, 3914-3924.

307 Ackerly, D.D., Loarie, S.R., Cornwell, W.K., Weiss, S.B., Hamilton, H., Branciforte, R. & Kraft, N.J.B. (2010)
308 The geography of climate change: implications for conservation biogeography. *Diversity and*
309 *Distributions*, **16**, 476-487.

310 Allouche, O., Tsoar, A. & Kadmon, R. (2006) Assessing the accuracy of species distribution models:
311 prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, **43**, 1223-1232.

312 Araújo, M.B. & New, M. (2007) Ensemble forecasting of species distributions. *Trends in Ecology & Evolution*,
313 **22**, 42-47.

314 Araújo, M.B. & Peterson, A.T. (2012) Uses and misuses of bioclimatic envelope modeling. *Ecology*, **93**, 1527-
315 1539.

316 Ashcroft, M.B. (2010) Identifying refugia from climate change. *Journal of Biogeography*, **37**, 1407-1413.

317 Austin, M. & Smith, T. (1990) A new model for the continuum concept. *Progress in Theoretical Vegetation*
318 *Science*, pp. 35-47. Springer.

319 Austin, M.P. & Van Niel, K.P. (2011a) Impact of landscape predictors on climate change modelling of species
320 distributions: a case study with *Eucalyptus fastigata* in southern New South Wales, Australia. *Journal*
321 *of Biogeography*, **38**, 9-19.

322 Austin, M.P. & Van Niel, K.P. (2011b) Improving species distribution models for climate change studies:
323 variable selection and scale. *Journal of Biogeography*, **38**, 1-8.

324 Beauregard, F. & de Blois, S. (2014) Beyond a climate-centric view of plant distribution: edaphic variables
325 add value to distribution models. *PloS one*, **9**, e92642.

326 Beven, K.J. & Kirkby, M.J. (1979) A physically based, variable contributing area model of basin hydrology /
327 Un modèle à base physique de zone d'appel variable de l'hydrologie du bassin versant. *Hydrological*
328 *Sciences Bulletin*, **24**, 43-69.

329 Birks, H.H. (2008) The Late-Quaternary history of arctic and alpine plants. *Plant Ecology & Diversity*, **1**, 135-
330 146.

331 Birks, H.J.B. & Willis, K.J. (2008) Alpines, trees, and refugia in Europe. *Plant Ecology & Diversity*, **1**, 147-
332 160.

333 Botkin, D.B., Saxe, H., Araújo, M.B., Betts, R., Bradshaw, R.H.W., Cedhagen, T., Chesson, P., Dawson, T.P.,
334 Ettersson, J.R., Faith, D.P., Ferrier, S., Guisan, A., Hansen, A.S., Hilbert, D.W., Loehle, C., Margules,
335 C., New, M., Sobel, M.J. & Stockwell, D.R.B. (2007) Forecasting the Effects of Global Warming on
336 Biodiversity. *BioScience*, **57**, 227.

337 Breiman, L. (2001) Random forests. *Machine Learning*, **45**, 5-32.

338 Clarke, L., Edmonds, J., Jacoby, H., Pitcher, H., Reilly, J. & Richels, R. (2007) Scenarios of greenhouse gas
339 emissions and atmospheric concentrations. *US Department of Energy Publications*, 6.

340 Dobrowski, S.Z. (2011) A climatic basis for microrefugia: the influence of terrain on climate. *Global Change*
341 *Biology*, **17**, 1022-1035.

342 Dubuis, A., Giovanettina, S., Pellissier, L., Pottier, J., Vittoz, P. & Guisan, A. (2013) Improving the prediction
343 of plant species distribution and community composition by adding edaphic to topo-climatic variables.
344 *Journal of Vegetation Science*, **24**, 593-606.

345 Elith, J., Leathwick, J.R. & Hastie, T. (2008) A working guide to boosted regression trees. *J Anim Ecol*, **77**,
346 802-13.

347 Engler, R., Randin, C.F., Vittoz, P., Czárka, T., Beniston, M., Zimmermann, N.E. & Guisan, A. (2009)
348 Predicting future distributions of mountain plants under climate change: does dispersal capacity
349 matter? *Ecography*, **32**, 34-45.

350 Esri (2013) *ArcScripts*. Available at: (accessed 25 September 2013)

351 Fielding, A.H. & Bell, J.F. (1997) A review of methods for the assessment of prediction errors in conservation
352 presence/absence models. *Environmental conservation*, **24**, 38-49.

353 Fløjgaard, C., Normand, S., Skov, F. & Svenning, J.C. (2009) Ice age distributions of European small
354 mammals: insights from species distribution modelling. *Journal of Biogeography*, **36**, 1152-1163.

Franklin, J. (1995) Predictive vegetation mapping: geographic modelling of biospatial patterns in relation to environmental gradients. *Progress in physical geography*, **19**, 474-499.

Fu, P. & Rich, P.M. (1999) Design and implementation of the Solar Analyst: an ArcView extension for modeling solar radiation at landscape scales. *Proceedings of the Nineteenth Annual ESRI User Conference* (ed by, pp. 1-31.

Godsoe, W., Murray, R. & Plank, M.J. (2015) Information on biotic interactions improves transferability of distribution models. *The American Naturalist*, **185**, 281-290.

Guisan, A. & Thuiller, W. (2005) Predicting species distribution: offering more than simple habitat models. *Ecology letters*, **8**, 993-1009.

Guisan, A., Theurillat, J.P. & Kienast, F. (1998) Predicting the potential distribution of plant species in an alpine environment. *Journal of Vegetation Science*, **9**, 65-74.

Hampe, A. & Petit, R.J. (2005) Conserving biodiversity under climate change: the rear edge matters. *Ecology letters*, **8**, 461-467.

Hastie, T.J. & Tibshirani, R.J. (1990) *Generalized additive models*. CRC Press.

Heikkinen, R.K., Luoto, M., Araújo, M.B., Virkkala, R., Thuiller, W. & Sykes, M.T. (2006) Methods and uncertainties in bioclimatic envelope modelling under climate change. *Progress in Physical Geography*, **30**, 751-777.

Hodd, R.L., Bourke, D. & Skeffington, M.S. (2014) Projected range contractions of European protected oceanic montane plant communities: focus on climate change impacts is essential for their future conservation. *PloS one*, **9**, e95147.

Hugall, A., Moritz, C., Moussalli, A. & Stanisc, J. (2002) Reconciling paleodistribution models and comparative phylogeography in the Wet Tropics rainforest land snail *Gnarosophia bellendenkerensis* (Brazier 1875). *Proceedings of the National Academy of Sciences*, **99**, 6112-6117.

Hylander, K., Ehrlen, J., Luoto, M. & Meineri, E. (2015) Microrefugia: Not for everyone. *Ambio*, **44 Suppl 1**, 60-8.

Keppel, G., Van Niel, K.P., Wardell-Johnson, G.W., Yates, C.J., Byrne, M., Mucina, L., Schut, A.G., Hopper, S.D. & Franklin, S.E. (2012) Refugia: identifying and understanding safe havens for biodiversity under climate change. *Global Ecology and Biogeography*, **21**, 393-404.

Kurtto, A. & Lampinen, R. (1999) Atlas of the distribution of vascular plants in Finland: A digital view of the national floristic database. *Acta Botanica Fennica*, 67-74.

le Roux, P.C., Virtanen, R. & Luoto, M. (2013a) Geomorphological disturbance is necessary for predicting fine-scale species distributions. *Ecography*, **36**, 800-808.

le Roux, P.C., Lenoir, J., Pellissier, L., Wisz, M.S. & Luoto, M. (2013b) Horizontal, but not vertical, biotic interactions affect fine-scale plant distribution patterns in a low-energy system. *Ecology*, **94**, 671-682.

Lenoir, J., Graae, B.J., Aarrestad, P.A., Alsos, I.G., Armbruster, W.S., Austrheim, G., Bergendorff, C., Birks, H.J., Brathen, K.A., Brunet, J., Bruun, H.H., Dahlberg, C.J., Decocq, G., Diekmann, M., Dynesius, M., Ejrnaes, R., Grytnes, J.A., Hylander, K., Klanderud, K., Luoto, M., Milbau, A., Moora, M., Nygaard, B., Odland, A., Ravolainen, V.T., Reinhardt, S., Sandvik, S.M., Schei, F.H., Speed, J.D., Tveraabak, L.U., Vandvik, V., Velle, L.G., Virtanen, R., Zobel, M. & Svenning, J.C. (2013) Local temperatures inferred from plant communities suggest strong spatial buffering of climate warming across Northern Europe. *Global Change Biology*, **19**, 1470-81.

Loarie, S.R., Carter, B.E., Hayhoe, K., McMahon, S., Moe, R., Knight, C.A. & Ackerly, D.D. (2008) Climate change and the future of California's endemic flora. *PLoS One*, **3**, e2502.

Luoto, M. & Heikkinen, R.K. (2008) Disregarding topographical heterogeneity biases species turnover assessments based on bioclimatic models. *Global Change Biology*, **14**, 483-494.

McCullagh, P. & Nelder, J.A. (1989) *Generalized linear models*. CRC press.

McCune, B. & Keon, D. (2002) Equations for potential annual direct incident radiation and heat load. *Journal of Vegetation Science*, **13**, 603-606.

Médail, F. & Diadema, K. (2009) Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *Journal of Biogeography*, **36**, 1333-1345.

Mod, H.K., le Roux, P.C., Guisan, A. & Luoto, M. (2015) Biotic interactions boost spatial models of species richness. *Ecography*, **38**, 913-921.

Moore, I.D., Grayson, R. & Ladson, A. (1991) Digital terrain modelling: a review of hydrological, geomorphological, and biological applications. *Hydrological processes*, **5**, 3-30.

- 409 Moss, R.H., Edmonds, J.A., Hibbard, K.A., Manning, M.R., Rose, S.K., Van Vuuren, D.P., Carter, T.R.,
410 Emori, S., Kainuma, M. & Kram, T. (2010) The next generation of scenarios for climate change
411 research and assessment. *Nature*, **463**, 747-756.
- 412 NASA Land Processes Distributed Active Archive Center (2013) *ASTER DEM*. Available at:
413 https://lpdaac.usgs.gov/data_access (accessed 07. 10. 2013).
- 414 Oksanen, L. & Virtanen, R. (1995) Topographic, altitudinal and regional patterns in continental and suboceanic
415 heath vegetation of northern Fennoscandia. *Acta Botanica Fennica*, **153**, 1-80.
- 416 Patsiou, T.S., Conti, E., Zimmermann, N.E., Theodoridis, S. & Randin, C.F. (2014) Topo-climatic
417 microrefugia explain the persistence of a rare endemic plant in the Alps during the last 21 millennia.
418 *Global Change Biology*, **20**, 2286-300.
- 419 Pearson, R.G. (2006) Climate change and the migration capacity of species. *Trends in Ecology & Evolution*,
420 **21**, 111-113.
- 421 Phillips, S.J., Dudík, M. & Schapire, R.E. (2004) A maximum entropy approach to species distribution
422 modeling. *Proceedings of the twenty-first international conference on Machine learning* (ed by, p.
423 83).
- 424 Pirinen, P., Simola, H., Aalto, J., Kaukoranta, J., Karlsson, P. & Ruuhela, R. (2012) Climatological statistics
425 of Finland 1981–2010. *Finnish Meteorological Institute Reports*, **2012**, 25.
- 426 Provan, J. & Bennett, K.D. (2008) Phylogeographic insights into cryptic glacial refugia. *Trends in Ecology
427 and Evolution*, **23**, 564-71.
- 428 Randin, C.F., Engler, R., Normand, S., Zappa, M., Zimmermann, N.E., Pearman, P.B., Vittoz, P., Thuiller, W.
429 & Guisan, A. (2009) Climate change and plant distribution: local models predict high-elevation
430 persistence. *Global Change Biology*, **15**, 1557-1569.
- 431 Reside, A.E., Welbergen, J.A., Phillips, B.L., Wardell-Johnson, G.W., Keppel, G., Ferrier, S., Williams, S.E.
432 & VanDerWal, J. (2014) Characteristics of climate change refugia for Australian biodiversity. *Austral
433 Ecology*, **39**, 887-897.
- 434 Reside, A.E., VanDerWal, J., Phillips, B.L., Shoo, L.P., Rosauer, D.F., Anderson, B.J., Welbergen, J.A.,
435 Moritz, C., Ferrier, S. & Harwood, T.D. (2013) *Climate change refugia for terrestrial biodiversity*.
436 Available at: <http://www.nccarf.edu.au/publications/climate-change-refugia-terrestrial-biodiversity>
437 (accessed 20. 08. 2013).
- 438 Riahi, K., Grübler, A. & Nakicenovic, N. (2007) Scenarios of long-term socio-economic and environmental
439 development under climate stabilization. *Technological Forecasting and Social Change*, **74**, 887-935.
- 440 Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N. & Rafaj, P.
441 (2011) RCP 8.5—A scenario of comparatively high greenhouse gas emissions. *Climatic Change*, **109**,
442 33-57.
- 443 Rickebusch, S., Thuiller, W., Hickler, T., Araujo, M.B., Sykes, M.T., Schweiger, O. & Lafourcade, B. (2008)
444 Incorporating the effects of changes in vegetation functioning and CO₂ on water availability in plant
445 habitat models. *Biology Letters*, **4**, 556-559.
- 446 Root, T.L., Price, J.T., Hall, K.R., Schneider, S.H., Rosenzweig, C. & Pounds, J.A. (2003) Fingerprints of
447 global warming on wild animals and plants. *Nature*, **421**, 57-60.
- 448 Roux, P.C.I., Virtanen, R., Heikkinen, R.K. & Luoto, M. (2012) Biotic interactions affect the elevational ranges
449 of high-latitude plant species. *Ecography*, **35**, 1048-1056.
- 450 Rull, V. & Vegas-Vilarrúbia, T. (2006) Unexpected biodiversity loss under global warming in the neotropical
451 Guayana Highlands: a preliminary appraisal. *Global Change Biology*, **12**, 1-9.
- 452 Sætersdal, M., Birks, H. & Peglar, S. (1998) Predicting changes in Fennoscandian vascular-plant species
453 richness as a result of future climatic change. *Journal of Biogeography*, **25**, 111-112.
- 454 Scherrer, D. & Körner, C. (2011) Topographically controlled thermal-habitat differentiation buffers alpine
455 plant diversity against climate warming. *Journal of biogeography*, **38**, 406-416.
- 456 Shoo, L.P., Hoffmann, A.A., Garnett, S., Pressey, R.L., Williams, Y.M., Taylor, M., Falconi, L., Yates, C.J.,
457 Scott, J.K. & Alagador, D. (2013) Making decisions to conserve species under climate change.
458 *Climatic Change*, **119**, 239-246.
- 459 Skov, F. & Svenning, J.C. (2004) Potential impact of climatic change on the distribution of forest herbs in
460 Europe. *Ecography*, **27**, 366-380.
- 461 Slaton, M.R. & Linder, P. (2015) The roles of disturbance, topography and climate in determining the leading
462 and rear edges of population range limits. *Journal of Biogeography*, **42**, 255-266.
- 463 Smith, S.J. & Wigley, T. (2006) Multi-gas forcing stabilization with Minicam. *The Energy Journal*, 373-391.

- Sormunen, H., Virtanen, R. & Luoto, M. (2011) Inclusion of local environmental conditions alters high-latitude vegetation change predictions based on bioclimatic models. *Polar Biology*, **34**, 883-897.
- Stewart, J.R., Lister, A.M., Barnes, I. & Dalén, L. (2010) Refugia revisited: individualistic responses of species in space and time. *Proceedings of the Royal Society of London B: Biological Sciences*, **277**, 661-671.
- Svenning, J.C., Normand, S. & Kageyama, M. (2008) Glacial refugia of temperate trees in Europe: insights from species distribution modelling. *Journal of Ecology*, **96**, 1117-1127.
- Taberlet, P. (1998) Biodiversity at the intraspecific level: the comparative phylogeographic approach. *Journal of Biotechnology*, **64**, 91-100.
- Temunović, M., Frascaria-Lacoste, N., Franjić, J., Satovic, Z. & Fernández-Manjarrés, J.F. (2013) Identifying refugia from climate change using coupled ecological and genetic data in a transitional Mediterranean-temperate tree species. *Molecular ecology*, **22**, 2128-2142.
- Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., Erasmus, B.F., De Siqueira, M.F., Grainger, A. & Hannah, L. (2004) Extinction risk from climate change. *Nature*, **427**, 145-148.
- Thuiller, W., Georges, D. & Engler, R. (2013) biomod2: Ensemble platform for species distribution modeling. *R package version*, **2**, r560.
- Thuiller, W., Lafourcade, B., Engler, R. & Araújo, M.B. (2009) BIOMOD - a platform for ensemble forecasting of species distributions. *Ecography*, **32**, 369-373.
- Thuiller, W., Lavorel, S., Araújo, M.B., Sykes, M.T. & Prentice, I.C. (2005) Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 8245-8250.
- Tikkanen, M. (2005) Climate. *The physical geography of Fennoscandia* (ed. by M. Seppälä). Oxford University Press, Oxford
- Waltari, E., Hijmans, R.J., Peterson, A.T., Nyári, Á.S., Perkins, S.L. & Guralnick, R.P. (2007) Locating Pleistocene refugia: comparing phylogeographic and ecological niche model predictions. *PLoS one*, **2**
- Van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram, T., Krey, V. & Lamarque, J.-F. (2011) The representative concentration pathways: an overview. *Climatic change*, **109**, 5-31.
- Wiens, J.A., Stralberg, D., Jongsomjit, D., Howell, C.A. & Snyder, M.A. (2009) Niches, models, and climate change: assessing the assumptions and uncertainties. *Proceedings of the National Academy of Sciences*, **106**, 19729-19736.
- Williams, P., Hannah, L.E.E., Andelman, S., Midgley, G.U.Y., Araújo, M., Hughes, G., Manne, L., Martinez-Meyer, E. & Pearson, R. (2005) Planning for Climate Change: Identifying Minimum-Dispersal Corridors for the Cape Proteaceae. *Conservation Biology*, **19**, 1063-1074.
- Wise, M., Calvin, K., Thomson, A., Clarke, L., Bond-Lamberty, B., Sands, R., Smith, S.J., Janetos, A. & Edmonds, J. (2009) Implications of limiting CO2 concentrations for land use and energy. *Science*, **324**, 1183-1186.
- Zimmermann, N.E., Yoccoz, N.G., Edwards, T.C., Jr., Meier, E.S., Thuiller, W., Guisan, A., Schmatz, D.R. & Pearman, P.B. (2009) Climatic extremes improve predictions of spatial patterns of tree species. *Proceedings of the National Academy of Sciences of the United States of America*, **106 Suppl 2**, 19723-8.

507 **Table 1** Descriptions of the four focal environmental variables with their minimum (Min.), median (Med.), mean and
 508 maximum (Max.) values

Variable	Description	Min.	Med.	Mean	Max.
GDD3 (contemporary)	Growing degree days (annual accumulated	163	810	784	1050
(under RCP 4.5)	daily temperature sum above 3°C; unit °C)	374	1247	1204	1489
(under RCP 8.5)		633	1620	1587	1964
Radiation	Potential annual direct radiation (MJ/cm²/a)	0.28	0.43	0.43	0.62
TWI	Topographic wetness index	5.58	6.93	6.95	8.70
Calcareousness	Cover of calcareous substrates (%)	0	6.7	13.0	90.3

509

510 **Table 2** The model evaluation statistics for the base (climate-only) and full (climate-plus-landscape) models showing the
 511 mean and standard deviation (SD) of the evaluation metrics over four cross-validation runs. Wilcoxon signed rank test p-
 512 values show change in predictive ability (***) = highly significant). AUC improved for 104; TSS for 10; and Kappa for
 513 99 out of 111 species

	AUC		TSS		Kappa	
	Base	Full	Base	Full	Base	Full
Mean ± SD	0.77 ± 0.09	0.86 ± 0.06	0.51 ± 0.14	0.62 ± 0.13	0.39 ± 0.21	0.46 ± 0.16
p-value	< 0.001***		< 0.001***		< 0.001***	

514

515 **Fig 1.** The location and elevation of the study area in northern Fennoscandia. We used data from whole area for training
516 the models, but excluded non-analogue regions from the projections (white dashed line)

517

518 **Fig 2.** Refugia in two Representative Concentration Pathway (RCP) scenarios (RCP 4.5 a-b; RCP 8.5 c-d) of future
519 climate according to base models (climate-only; a, c) and full models (climate-plus-landscape; b, d). The full models
520 predict a higher relative species persistence from contemporary to future climates. No refugia were predicted for scenario
521 RCP 8.5 by the base model. The legend applies throughout

Supporting Information

Figure S1: An outline of the modelling framework used. We combined model predictions for 111 arctic-alpine plant species to estimate current and projected species distribution and identify refugia at a 1 km² resolution. To enable the comparison of species' distributional changes, we ran the predictions for contemporary climate as well as two Representative Concentration Pathways (RCPs 4.5 and RCP 8.5). Two sets of predictors were used: climate-only (growing degree days (GDD3)); and climate-plus-landscape, which included additional landscape-scale predictors: potential solar radiation (RAD); calcareousness (CALC); and topographic wetness index (TWI). Observed species distributions were related to the set of predictors using five statistical modelling techniques: generalized linear modeling (GLM), generalized additive modeling (GAM), generalized boosting methods (GBM), random forest (RF) and maximum entropy (MAXENT). All the models were implemented in the Biomod2 platform under R-program

Table S1: Summary statistics for arctic-alpine species persistence within refugia; the number (n) and proportion (%) of species persisting in refugia according to the two model structures used and two scenarios of future climate. No refugia were predicted for scenario RCP 8.5 by the base model. The total count of cells used in the analysis was 15 622

Figure S2: Predicted contemporary and future distributions of Alpine Mouse-ear (*Cerastium alpinum*) in the study area. The predicted distributions were modelled into two scenarios of future climates (RCP 4.5 and RCP 8.5) according to two sets of predictors: climate-only (base model) and climate-plus-landscape (full model)

Figure S3: Box-and-whisker plot showing the mean variable importance values based on the full model across all species and across the five modelling techniques utilized (boxplot represents median, first and third quartiles; the whiskers show the interquartile range). The variables included in the full model are growing degree days (GDD3), calcareousness (CALC), potential solar radiation (RAD) and topographic wetness index (TWI). See table 1 for a full description of each variable. Outliers are excluded for clarity

Table S2: Summary statistics for refugia located by the base (climate-only) and full (climate-plus-landscape) models under Representative Concentration Pathway (RCP) 4.5 and the full model for RCP 8.5 (the base-model for RCP 8.5 found no refugia). The full set of variables are growing degree days (GDD3), calcareousness, potential solar radiation (RAD) and topographic wetness index (TWI). See table 1 for a full description of each variable and for comparison with contemporary climatic conditions

Figure S4. Spatial characteristics of the predictor variables: growing degree days (GDD3; °C) for the contemporary climate 1981–2010 (a), and GDD3 for the two Representative Concentration Pathway (RCP) scenarios used: RCP 4.5 (b) and RCP 8.5 (c); calcareousness (d; %); potential solar radiation (e; MJ/cm²/a); and topographic wetness index (TWI) (f). See table 1 for a further description of the predictors

Figure 1.

583
584

585
586
587

Figure 2.