<https://helda.helsinki.fi>

ILB 938, a valuable faba bean (*Vicia faba* L) accession

Khazaei, H.

2018-10

Khazaei , H , Link , W , Street , K & Stoddard , F L 2018 , ' ILB 938, a valuable faba bean
(*Vicia faba* L) accession ' , Plant Genetic Resources , vol. 16 , no. 5 , pp. 478-482 . <https://doi.org/10.1017/S1479262118000205>

<http://hdl.handle.net/10138/307483>

<https://doi.org/10.1017/S1479262118000205>

---

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1   **ILB 938, a valuable faba bean (*Vicia faba* L.) accession**

2   H. Khazaei^{1*}, W. Link², K. Street³ and F. L. Stoddard⁴

3   ¹ *Department of Plant Sciences, University of Saskatchewan, Saskatoon, Canada*

4   ² *Department of Crop Sciences, Georg-August-Universität, Göttingen, Germany*

5   ³ *International Center for Agricultural Research in the Dry Areas (ICARDA), Rabat, Morocco*

6   ⁴ *Department of Agricultural Sciences, Viikki Plant Science Centre, and Helsinki Sustainability*
7   *Science Centre, University of Helsinki, Helsinki, Finland*

8

9   *Corresponding author. E-mail: hamid.khazaei@gmail.com or hamid.khazaei@usask.ca

Abstract

Here we review the potential of ILB 938 (IG 12132– doi:10.18730/60FD2), a unique faba bean accession originating from the Andean region of Colombia and Ecuador, maintained at ICARDA - International Center for Agricultural Research in the Dry Areas, with resistance to multiple biotic and abiotic stresses and carrying some useful morphological markers. It has been used as a donor of leaf-related drought adaptation traits and chocolate spot (*Botrytis fabae*) resistance genes in faba bean breeding programs worldwide. From generated populations of recombinant inbred lines, QTLs (quantitative traits loci) associated with these useful traits have been mapped. Other markers, such as a lack of stipule-spot pigmentation and clinging pod wall, show the presence of unusual changes in biochemical pathways that may have economic value in the future.

Keywords: germplasm, faba bean, biotic stress, abiotic stress, mapping population

23   **Introduction**

24   Faba bean (*Vicia faba* L.) seeds are a generous source of plant protein, with a global average
25   protein concentration of 29% on a dry-weight basis (Feedipedia, 2018). It is one of the main
26   sources of affordable protein for human consumption in developing countries (consumed as dry
27   or canned), and for livestock feed in many developed countries. The fresh pods and seeds are
28   widely used as a vegetable crop for fresh seed production. Like other legumes, it symbiotically
29   fixes atmospheric nitrogen, thus improving the soil fertility. As a non-host of many cereal
30   pathogens, faba bean is ideal as a break between grain crops in the rotation (Köpke and
31   Nemecek, 2010). It has a mixed breeding system and is cross-pollinated at frequencies of 4-84%,
32   with the value determined by the interaction between the plant genotype, its environment, and
33   the population of pollinators (Bond and Poulsen, 1983). Its interaction with many species of bee
34   (Stoddard and Bond, 1987) makes it suitable for growing in ecological focus areas (Bues *et al.*,
35   2013). It is widely adapted to cool-temperate agriculture, being grown from Mediterranean
36   climates in southern Australia and Mediterranean basin countries to sub-boreal climates in
37   Finland and Canada. Nevertheless, faba bean cultivation is limited due to its susceptibility to
38   several biotic and abiotic constraints globally (see Stoddard *et al.*, 2006; Torres *et al.*, 2006;
39   Khan *et al.*, 2010). Hence, genetically diverse sources of resistance genes or genes for specific
40   adaptations such as to abiotic stress factors are required in pre-breeding programs worldwide.
41   Some of these germplasm sources, called ‘donors’ may become prominent.

42   Faba bean is represented in germplasm collections by only the cultivated form. Both
43   botanical and molecular data suggest that the wild ancestors of faba bean either have vanished
44   or have not yet been discovered (Maxted, 1993; Duc *et al.*, 2010; Kosterin, 2014; Caracuta *et*
45   *al.*, 2016), which highlights the importance of the accessible diversity within the cultivated

form. The place of origin of faba bean is still unknown. A Near or Middle East centre of origin has been proposed (Cubero, 1974), and the earliest identified remains of faba bean date from 10200 BP in a cave in Israel (Caracuta *et al.*, 2015). Radiation followed in four directions from the proposed centre: Europe, along the North Africa coast to Spain, along the Nile Valley to Ethiopia, and from Mesopotamia to India and China (Lawes *et al.*, 1983). Spanish and European material was taken to South America in the 16th century (Bond, 1976). There are 43,695 faba bean accessions conserved within 37 global genebanks (*ex situ*, FAO, 2010) as well as on-farm conservation (*in situ*, Suso *et al.*, 2005). ICARDA (International Center for Agricultural Research in the Dry Areas) hosts the largest collection of over 9,500 accessions (21% of global collection, FAO, 2010). ICARDA maintains its faba bean germplasm in two classes, ILB (International Legume Bean) accessions from different countries, and BPL (Bean Pure Line) accessions that are derived through selfing from accessions drawn from the ILB collection (Saxena and Varma, 1985).

Accession ILB 938

ILB 938 is the result of mass selection from ILB 438 based on seed size. ILB 438 was brought to ICARDA from the Andean region of Ecuador and Columbia (Robertson, 1984). ICARDA's registered BPL derivatives of ILB 438 and ILB 938 are BPL 710 and BPL 1179, respectively. ILB 938/2 is an inbred line developed at Göttingen for use in genetics and breeding studies. The corresponding "IG" number for ILB 938 in the ICARDA genebank is "IG 12132" (accession doi: 10.18730/60FD2, see <https://www.genesys-pgr.org/10.18730/60FD2>). ILB 438 is registered as IG 11632 in the ICARDA genebank (accession doi: 10.18730/601TB).

Morphological markers

In the wild-type faba bean, the extra-floral nectary on the stipule is coloured black. The presence of stipule spot pigmentation was proposed as an early morphological marker indicating wild-type ‘coloured’ flowers (tannin-containing faba bean), where there is a black spot on each wing petal and dark vein markings on the standard petal (Picard, 1976). The absence of the pigmentation was considered as the corresponding early morphological indicator for the white-flower, zero-tannin trait (Link *et al.*, 2008). ILB 938, however, carries a rare allele (*ssp1*) that decouples pigmentation in flowers from that in stipules, so it has colourless stipules and coloured flowers (Supplementary Figure S1, Khazaei *et al.*, 2014a). An Australian line, AF11212, has the same phenotype and is derived from BPL 710 (Dr. Jeff Paull, The University of Adelaide, Australia; personal communication). Crossing ILB 938/2 with AF11212 (including reciprocal crosses) showed in the F1 and F2 generations uniformly the combination of colourless stipule spots and spotted flowers, confirming that the same gene exists in both accessions (Miller, 2016).

The seed size of ILB 938 is classified as *equina* (horse bean, field bean, flattened seed; 0.6 g / seed) which is expected since it was the medium to large-seeded Mediterranean-adapted faba bean forms that were introduced to Central and South America by immigrants from Spain (Muratova, 1931; Cubero, 1974). The seed coat of ILB 938 is green in colour, which is recessive to the common beige or buff colour (Khazaei *et al.*, 2014b).

A further noticeable morphological character of ILB 938 is the clinging pod wall, where fibres from the inner epidermis of the pod cling to the surface of the seed (Supplementary Figure S2). We have not seen this trait otherwise reported in faba bean germplasm, and while it is of little importance agronomically or economically, it may indicate a difference in cell wall development that has other impacts elsewhere in the plant or in the value chain.

Finally, the funiculus is yellow in ILB 938, in contrast to the common green displayed by other accessions.

Biotic stresses

The resistance of ILB 938 to chocolate spot (CS, caused by *Botrytis fabae* Speg.) has been demonstrated in Egypt (Mohamed *et al.*, 1981 [re-coded NEB 938]; Khalil and Nassib, 1984; Robertson, 1984), Syria (Hanounik, 1982), the United Kingdom (Jellis *et al.*, 1982), Canada (Robertson, 1984), France (Tivoli *et al.*, 1988), and Ethiopia (Beyene *et al.*, 2016). Further, we have noticed its resistance to CS in field conditions of both southern Finland and western Canada.

The resistance of the original source of ILB 938 was confirmed in the Nile Delta after crosses with the local cultivar Giza 3 (ICARDA Caravan, 1998; Zeid *et al.*, 2009). From there it was transferred to locally adapted material that was released as Giza 461 in Egypt (Bond *et al.*, 1994; Dwivedi *et al.*, 2006; El-Komy *et al.*, 2015).

The related bean pure lines BPL 710 and BPL 1179 to ILB 438 and ILB 938, respectively also showed high resistance to CS across environments (Hanounik and Maliha, 1986; Hanounik and Robertson, 1988; Villegas-Fernández *et al.*, 2012). The Australian cultivar Icarus was derived from BPL 710 and released as a cultivar resistant to CS and rust (Dwivedi *et al.*, 2006).

ILB 938 is, furthermore, considered as a consistent source of resistance to rust (*Uromyces viciae-fabae* (Pers.) J. Schrot.) (Rashid and Bernier, 1991; Khalil *et al.*, 1985; Rashid and Bernier, 1986). Both BPL 710 (Australian accession No. AC1269) and BPL 1179 (AC1272) are

registered as rust-resistant accessions in Australia (Ijaz *et al.*, 2018) as well as in ICARDA (1987).

Some studies have suggested that ILB 938 may also carry resistance to crenate broomrape, *Orobanche crenata* Forsk., an achlorophyllous, holoparasitic weed, poses a major constraint to faba bean production in Mediterranean climates (Zeid *et al.*, 2006; 2009).

Abiotic stresses

Drought adaptation is an essential character for faba bean cultivation in arid and semiarid regions. ILB 938 has demonstrated high water use efficiency (WUE) in several studies (e.g., Abdelmula *et al.*, 1999; Link *et al.*, 1999; Stoddard *et al.*, 2006; Khan *et al.*, 2007, 2010; Khazaei *et al.*, 2013; Khazaei *et al.*, 2014b) mainly due to low stomatal conductance, thus minimizing water loss and maintaining yield under drought conditions. Nevertheless, its reduced leaf stomatal conductance was not associated with a highly ramified rooting system (Belachew *et al.*, 2018).

While ILB 938 has relatively low productivity, no yield penalty was observed when it was exposed to drought conditions (Link *et al.*, 1999; Khan *et al.*, 2007; Khazaei *et al.*, 2014b). It maintains a relatively high water status under water deficit conditions, demonstrating high WUE with relatively low yield, because its stomata shut early, reducing potential photosynthesis while limiting water loss.

The response of ILB 938/2 to ultraviolet light differs greatly from that of a contrasting cultivar, Aurora/2 that was developed at low altitudes and high latitudes where incident UV is much weaker than high in the Andes (Yan *et al.*, 2018).

Mapping populations

A population of recombinant inbred lines (RILs) was developed from the cross of Mélodie/2 × ILB 938/2 (along with its reciprocal) at the University of Helsinki (Khazaei *et al.*, 2014a). This population has been mapped for traits related to drought adaptation (Khazaei *et al.*, 2014b), vicine-convicine concentration (v-c, Khazaei *et al.*, 2015), and stipule spot pigmentation (Khazaei *et al.*, 2014a). QTLs (quantitative traits loci) for seed size, seed coat colour, clinging pod wall and yellow funiculus have also been located. ILB 938 and Mélodie differed at two loci affecting stomatal activity at opposite ends of Chromosome II, with each parent contributing a canopy-cooling allele (Khazaei *et al.*, 2014b). The progenies of this population facilitated the development of a reliable molecular marker for v-c in this crop (Khazaei *et al.*, 2017). This population is being phenotyped for salinity response in a collaboration with Egypt and collaborative studies on other traits are in progress. Near-isogenic lines have been derived from heterozygous F5 individuals at Göttingen (Tacke and Link, 2017).

Another RIL population, ILB 938/2 × Disco/2 (Khazaei *et al.*, 2014a), is suitable for CS genetic studies. Disco (low in both tannin and v-c) has been shown to be very sensitive to CS (Villegas-Fernández *et al.*, 2012; Khazaei, Personal observation). A RIL population from ILB 938/2 × Aurora/2 (Khazaei *et al.*, 2014a) will be useful for analyzing the basis of the difference in ultraviolet response of these two lines.

A multi-parent population [(Disco/2 × ILB 938/2) × (IG 114476 × IG 132238)] has been prepared for use in genomic studies (Khazaei *et al.*, 2018). This population is at F4 generation at the time of writing this paper and kept at the University of Reading, UK.

DNA fingerprinting

ILB 938/2 was genotyped using 875 SNP (single nucleotide polymorphism) markers developed by Webb *et al.* (2016). The results showed a high level of homozygosity (99.6%, Webb *et al.*, 2016). The genotyping calls on ILB 938/2 are presented in Supplementary Table S1.

Conclusions

The presence of unusual traits in this material is intriguing, because the crop has been grown in South America for only about 500 of its 10 000 years of domestication. It may be attributable to several causes, including widespread genetic variation introduced by the European settlers, adaptation to extremely varied environments within short distances due to altitude, frequent gene exchanges by pollinators and movement of peoples, and natural selection (Bond *et al.*, 1994), or UV-induced mutation. Recently, several new accessions from Spain, Ecuador, Colombia and Peru with high level of resistance to CS were identified (Maalouf *et al.*, 2016).

Acknowledgments

We thank Dr. Jeff Paull (The University of Adelaide, Australia) for providing the seeds of AF11212. ILB 938 (IG 12132) is also acknowledged for its generous offering. FLS dedicates this paper to the memory of faba bean breeder extraordinaire Dr. David Bond (1929-2017), who was a kind mentor and teacher.

References

- Abdelmula AA, Link W, von Kittlitz E and Stelling D (1999) Heterosis and inheritance of drought tolerance in faba bean, *Vicia faba* L. *Plant Breeding* 118: 458–490.
- Belachew KY, Nagel KA, Fiorani F and Stoddard FL (2018) Diversity in root growth responses to moisture deficit in young faba bean (*Vicia faba* L.) plants. *PeerJ* 6: e4401.
- Beyene AT, John D, Sibiya J and Fikre A (2016) Gene action determining grain yield and chocolate spot (*Botrytis fabae*) resistance in a faba bean. *Euphytica* 207: 293–304.
- Bond DA (1976) Field bean: *Vicia faba* (Leguminosae-Papilionatae). In: Simmonds NW (ed.) *Evolution of Crop Plants*. Longman, London, UK. pp. 179–182.
- Bond DA, Jellis GJ, Rowland GG, Le Guen J, Robertson LD, Khalil SA and Li-Juan L (1994) Present status and future strategy in breeding faba beans (*Vicia faba* L.) for resistance to biotic and abiotic stresses. *Euphytica* 73: 151–166.
- Bond DA and Poulsen MH (1983) Pollination. In: Hebblethwaite PD (ed.) *The Faba Bean (Vicia faba L.)*. Butterworths, London, UK. pp. 77–101.
- Bues A, Preissel S, Reckling M, Zander P, Kuhlman T, Topp K, Watson CA, Lindström K, Stoddard FL and Murphy-Bokern D. (2013). The environmental role of protein crops in the new Common Agricultural Policy. European Parliament, Directorate General for Internal Policies, Policy Department B: Structural and Cohesion Policies, Agricultural and Rural Development IP/B/AGRI/IC/2012-067. 112 pp.
- <http://www.europarl.europa.eu/studies>. ISBN 978-92-823-4521-4, doi: 10.2861/27627.

189 <http://www.europarl.europa.eu/RegData/etudes/etudes/join/2013/495856/IPOL->
190 [AGRI_ET\(2013\)495856_EN.pdf](#)

191 Caracuta V, Barzilai O, Khalaily H, Milevski I, Paz Y, Vardi J, Regev L and Boaretto E (2015)
192 The onset of faba bean farming in the southern Levant. *Scientific Reports* 5: 14370.

193 Caracuta V, Weinstein-Evron M, Kaufman D, Yeshurun R, Silvent J and Boaretto E (2016)
194 14,000-year-old seeds indicate the Levantine origin of the lost progenitor of faba bean.
195 *Scientific Reports* 6: 37399.

196 Cubero J (1974) On the evolution of *Vicia faba* L. *Theoretical and Applied Genetics* 45: 47–51.

197 Duc G, Bao S, Baum M, Redden B, Sadiki M, Suso MJ, Vishniakova M and Zong X (2010)
198 Diversity, maintenance and use of *Vicia faba* L. genetic resources. *Field Crops Research*
199 115: 270–278.

200 Dwivedi SL, Blair MW, Upadhyaya HD, Serraj R, Balaji J, Buhariwalla HK, Ortiz R and Crouch
201 JH (2006) Using genomics to exploit grain biodiversity in crop improvement. *Plant*
202 *Breeding Reviews* 26: 171–310.

203 El-Komy MH, Saleh AA and Molan YY (2015) Resistance/susceptibility of faba bean to *Botrytis*
204 *fabae*: the causal agent of chocolate spot with respect to leaf position. *International*
205 *Journal of Agriculture and Biology* 17: 691–701.

206 FAO (2010) The second report on the state of the world’s plant genetic resources for food and
207 agriculture. Rome, Italy. <http://www.fao.org/docrep/013/i1500e/i1500e00.htm>, accessed
208 01May 2018.

209 Feedipedia (2018) Faba bean (*Vicia faba*). <https://www.feedipedia.org/node/4926>, accessed 01
 210 May 2018.

211 Hanounik SB (1982) Resistance in faba beans to chocolate spot. *FABIS Newsletter* 5: 24–26.

212 Hanounik SB and Maliha N (1986) Horizontal and vertical resistance in *Vicia faba* to chocolate
 213 spot caused by *Botrytis fabae*. *Plant Disease* 70: 770–773.

214 Hanounik SB and Robertson LD (1988) New sources of resistance in *Vicia faba* to chocolate
 215 spot caused by *Botrytis fabae*. *Plant Disease* 72: 696–698.

216 ICARDA (1987) Faba bean pathology progress report 1986–1987. In: Food Legume
 217 Improvement Program, ICARDA, Aleppo, Syria.

218 ICARDA Caravan (1998) Review of agriculture in the dry areas. Issue No. 9.
 219 <https://apps.icarda.org/wsInternet/wsInternet.aspx/DownloadFileToLocal?filePath=Cara>
 220 [van/Caravan9.pdf&fileName=Caravan9.pdf](https://apps.icarda.org/wsInternet/wsInternet.aspx/DownloadFileToLocal?filePath=Cara), accessed 01 May 2018.

221 Ijaz U, Adhikari KN, Stoddard FL and Trethowan RM (2018) Rust resistance in faba bean (*Vicia*
 222 *faba* L.): status and strategies for improvement. *Australasian Plant Pathology* 47: 71–81.

223 Jellis GJ, Bond DA and Old J (1982) Resistance to chocolate spot (*Botrytis fabae*) in ICARDA
 224 accessions of *Vicia faba*. *FABIS Newsletter* 4: 53–54.

225 Khalil SA and Nassib AM (1984) Identification of some sources of resistance to disease in faba
 226 bean. I. Chocolate spot (*Botrytis fabae* Sard.). *FABIS Newsletter* 10: 18–21.

227 Khalil SA, Nassib AM and Mohammed HA (1985) Identification of some sources of resistance
 228 to diseases in faba beans II—Rust (*Uromyces fabae*). *FABIS Newsletter* 11: 18–20.

229 Khan HR, Link W, Hocking TJH and Stoddard FL (2007) Evaluation of physiological traits for
 230 improving drought tolerance in faba bean (*Vicia faba* L.). *Plant and Soil* 292: 205–217.

231 Khan HR, Paull JG, Siddique KHM and Stoddard FL (2010) Faba bean breeding for drought
 232 affected environments: A physiological and agronomic perspective. *Field Crops*
 233 *Research* 115: 279–286.

234 Khazaei H, O’Sullivan DM, Jones H, Pitts N, Sillanpää MJ, Pärssinen P, Manninen O and
 235 Stoddard FL (2015) Flanking SNP markers for vicine-convicine concentration in faba
 236 bean (*Vicia faba* L.). *Molecular Breeding* 35: 38.

237 Khazaei H, O’Sullivan DM, Sillanpää MJ and Stoddard FL (2014a) Genetic analysis reveals a
 238 novel locus in *Vicia faba* decoupling pigmentation in the flower from that in the extra-
 239 floral nectaries. *Molecular Breeding* 34: 1507–1513.

240 Khazaei H, O’Sullivan DM, Sillanpää MJ and Stoddard FL (2014b) Use of synteny to identify
 241 candidate genes underlying QTL controlling stomatal traits in faba bean (*Vicia faba* L.).
 242 *Theoretical and Applied Genetics* 127: 2371–2385.

243 Khazaei H, Purves RW, Song M, Stonehouse R, Bett KE, Stoddard FL and Vandenberg A
 244 (2017) Development and validation of a robust, breeder-friendly molecular marker for the
 245 *vc* locus in faba bean. *Molecular Breeding* 37: 140.

246 Khazaei H, Purves RW, Stoddard FL and Vandenberg A (2018) A multi-parent faba bean (*Vicia*
247 *faba* L.) population for future genomic studies. *Plant Genetic Resources* (in press).

248 Khazaei H, Street K, Bari A, Santanen A and Stoddard FL (2013) Do faba bean (*Vicia faba* L.)
249 accessions from environments with contrasting seasonal moisture availabilities differ in
250 stomatal characteristics and related traits? *Genetic Resources and Crop Evolution* 60:
251 2343–2357.

252 Kosterin OE (2014) The lost ancestors of the broad bean (*Vicia faba* L.) and the origin of plant
253 cultivation in the Near East. *Vavilov Journal of Genetics and Breeding* 18(4/1): 831–840.

254 Köpke U and Nemecek T (2010) Ecological services of faba bean. *Field Crops Research* 115:
255 217–233.

256 Lawes DA, Bond DA and Poulsen MH (1983) Classification, origin, breeding methods and
257 objectives. In: Hebblethwaite PD (ed.) *The Faba Bean* (*Vicia faba* L.). Butterworths,
258 London, UK. pp. 23–76.

259 Link W, Abdelmula AA, Kittlitz E von, Bruns S, Riemer H and Stelling D (1999) Genotypic
260 variation for drought tolerance in *Vicia faba*. *Plant Breeding* 118: 477–483.

261 Link W, Hanafy M, Malenica N, Jacobsen H-J and Jelenić S (2008) Broad bean. In: Kole C and
262 Hall TC (eds) *A Compendium of Transgenic Crop Plants: Transgenic Legume Grains*
263 *and Forages*. Vol. 3. Wiley, New York, pp. 71–88.

264 Maalouf F, Ahmed S, Shaaban K, Bassam B, Nawar F, Singh M and Amri A (2016) New faba
 265 bean germplasm with multiple resistances to *Ascochyta* blight, chocolate spot and rust
 266 diseases. *Euphytica* 211: 157–167.

267 Maxted N (1993) A phenetic investigation of *Vicia* L. subgenus *Vicia* (Leguminosae, Viciaeae).
 268 *Botanical Journal of the Linnean Society* 111: 155–182.

269 Miller M (2016) Genetics of pigmentation in faba bean (*Vicia faba* L.). Undergraduate Thesis.
 270 University of Saskatchewan, Canada.

271 Mohamed HA, Zeid NA and Habib WF (1981) Variation with the fungus *Botrytis fabae* Sard.
 272 *FABIS Newsletter* 3: 49–50.

273 Muratova VS (1931) Common beans (*Vicia faba* L.). *Bulletin of Applied Botany of Genetics and*
 274 *Plant Breeding* 50: 1–298.

275 Picard J (1976) Aperçu sur l'hérédité du caractère absence de tannins dans les graines de
 276 féverole (*Vicia faba* L.). *Annales de l'Amélioration des Plantes* 26: 101–106.

277 Rashid KY and Bernier CC (1986) Selection for slow rusting in faba bean (*Vicia faba* L.) to
 278 *Uromyces viciae-fabae*. *Crop Protection* 5: 218–224.

279 Rashid KY and Bernier CC (1991) The effect of rust on yield of faba bean cultivars and slow
 280 rusting populations. *Canadian Journal of Plant Science* 71: 967–972.

281 Robertson LD (1984) A Note on the I.L.B. Source of *Botrytis fabae* resistance. In: Chapman GP
 282 and Tarawali SA (eds) *Systems for Cytogenetic Analysis in Vicia faba* L. *Advances in*
 283 *Agricultural Biotechnology*. Vol 11. Springer, Dordrecht.

284 Saxena MC and Varma S (1985) Faba beans, Kabuli chickpeas, and lentils in the 1980s. An
 285 international workshop. May 16–20, 1983; proceedings. International Centre for
 286 Agricultural Research in the Dry Areas. Aleppo, Syria.

287 Stoddard FL and Bond DA (1987) The pollination requirements of the faba bean (*Vicia faba* L.).
 288 Bee World 68: 144-152.

289 Stoddard FL, Balko C, Erskine W, Khan HR, Link W and Sarker A (2006) Screening techniques
 290 and sources of resistance to abiotic stresses in cool-season food legumes. *Euphytica* 147:
 291 167–186.

292 Suso MJ, Moreno MT and Cubero JI (1993) Variation among Spanish faba bean cultivars:
 293 taxonomy and evolutionary implications. *Genetic Resources and Crop Evolution* 40:
 294 105–111.

295 Tacke R and Link W (2017) Towards a localization of the “vc-” gene responsible for low vicine
 296 and convicine content in seeds of faba bean (*Vicia faba* L.) and towards a low vicine and
 297 convicine winter faba bean cultivar. In: 8th International Conference on Legume Genetics
 298 and Genomics, September 18–22. Siófok, Hungary. p. 115.

299 Tivoli B, Berthelem P, Leguen J and Onfroy C (1988) A study of the performance of certain faba
 300 bean genotypes in relation to *Botrytis fabae* and *Ascochyta fabae* in France. *FABIS*
 301 *Newsletter* 21: 36–39.

302 Torres AM, Roman B, Avila CM, Satovic Z, Rubiales D, Sillero JC, Cubero JI and Moreno MT
 303 (2006) Faba bean breeding for resistance against biotic stresses: towards applications of
 304 marker technology. *Euphytica* 147: 67–80.

- 305 Villegas-Fernández AM, Sillero JC and Rubiales D (2012) Screening faba bean for chocolate
306 spot resistance: evaluation methods and effects of age of host tissue and temperature.
307 *European Journal of Plant Pathology* 132: 443–453.
- 308 Webb A, Cottage A, Wood T, Khamassi K, Hobbs D, Gostkiewicz K, White M, Khazaei H, Ali
309 M, Street D, Stoddard FL, Maalouf F, Ogbonnaya F, Link W, Thomas J and O'Sullivan
310 DM (2016) A SNP-based consensus genetic map for synteny-based trait targeting in faba
311 bean (*Vicia faba* L.). *Plant Biotechnology Journal* 14: 177–185.
- 312 Yan Y, Neugart S, Stoddard FL and Aphalo PJ (2018) Different responses to solar ultraviolet
313 (UV) and blue radiation in two accessions of *Vicia faba* over two generations. *Aspects of*
314 *Applied Biology* 138, *Advances in Legume Science and Practice*, pp. 63–64.
- 315 Zeid M, Ghazy AI and Link W (2006) Questing for *Orobanche crenata* resistance genes in faba
316 bean. In: *New Life Sciences: Changing Lives*, 26–29 April 2006, BioVision Alexandria,
317 Egypt.
- 318 Zeid M, Mitchell S, Link W, Carter M, Nawar A, Fulton T and Kresovich S (2009) Simple
319 sequence repeats (SSRs) in faba bean: new loci from *Orobanche*-resistant cultivar ‘Giza
320 402’. *Plant Breeding* 128: 149–155.