<https://helda.helsinki.fi>

TILLING, high-resolution melting (HRM), and next-generation sequencing (NGS) techniques in plant mutation breeding

Taheri, Sima

2017-03

Taheri , S , Abdullah , T L , Jain , S M , Sahebi , M & Azizi , P 2017 , ' TILLING, high-resolution melting (HRM), and next-generation sequencing (NGS) techniques in plant mutation breeding ' , Molecular Breeding , vol. 37 , no. 3 , 40 . <https://doi.org/10.1007/s11032-017-0643-7>

<http://hdl.handle.net/10138/307693>

<https://doi.org/10.1007/s11032-017-0643-7>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/316428389>

TILLING, high-resolution melting (HRM), and next-generation sequencing (NGS) techniques in plant mutation breeding

Article in *Molecular Breeding* · February 2017

CITATION

1

READS

199

5 authors, including:

Sima Taheri

University of Malaya

49 PUBLICATIONS 190 CITATIONS

[SEE PROFILE](#)

Thohirah Lee Abdullah

Universiti Putra Malaysia

118 PUBLICATIONS 454 CITATIONS

[SEE PROFILE](#)

Shri Mohan Jain

University of Helsinki

207 PUBLICATIONS 3,913 CITATIONS

[SEE PROFILE](#)

Mahbod Sahebi

Universiti Putra Malaysia

73 PUBLICATIONS 402 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:

Oil Palm [View project](#)

Salt stress effects on rice [View project](#)

REVIEW

TILLING, high-resolution melting (HRM), and next-generation sequencing (NGS) techniques in plant mutation breeding

Sima Taheri · Thohirah Lee Abdullah ·
Shri Mohan Jain · Mahbod Sahebi · Parisa Azizi

Received: 15 June 2016 / Accepted: 19 February 2017
© Springer Science+Business Media Dordrecht 2017

Abstract Induced mutations have been used effectively for plant improvement. Physical and chemical mutagens induce a high frequency of genome variation. Recently, developed screening methods have allowed the detection of single nucleotide polymorphisms (SNPs) and the identification of traits that are difficult to identify at the molecular level by conventional breeding. With the assistance of reverse genetic techniques, sequence variation information can be linked to traits to investigate gene function. Targeting induced local lesions in genomes (TILLING) is a high-throughput technique to

identify single nucleotide mutations in a specific region of a gene of interest with a powerful detection method resulted from chemical-induced mutagenesis. The main advantage of TILLING as a reverse genetics strategy is that it can be applied to any species, regardless of genome size and ploidy level. However, TILLING requires laborious and time-consuming steps, and a lack of complete genome sequence information for many crop species has slowed the development of suitable TILLING targets. Another method, high-resolution melting (HRM), which has assisted TILLING in mutation detection, is faster, simpler and less expensive with non-enzymatic screening system. Currently, the sequencing of crop genomes has completely changed our vision and interpretation of genome organization and evolution. Impressive progress in next-generation sequencing (NGS) technologies has paved the way for the detection and exploitation of genetic variation in a given DNA or RNA molecule. This review discusses the applications of TILLING in combination with HRM and NGS technologies for screening of induced mutations and discovering SNPs in mutation breeding programs.

S. Taheri (✉) · T. L. Abdullah · P. Azizi
Department of Crop Science, Faculty of Agriculture, University
Putra Malaysia, 43400 Serdang, Selangor, Malaysia
e-mail: sima_taheri65@yahoo.com

T. L. Abdullah
e-mail: thohirah@upm.edu.my

P. Azizi
e-mail: bahar3236@yahoo.com

S. M. Jain
Department of Agricultural Sciences, University of Helsinki,
PL-27, Latokartanonkaari 5, Helsinki, Finland
e-mail: mohan.jain@helsinki.fi

M. Sahebi
Laboratory of Plantation Crops, Institute of Tropical Agriculture,
Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia
e-mail: mahbod_sahebi@yahoo.com

M. Sahebi
Department of Land Management, Faculty of Agriculture,
Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

Keywords Mutation breeding · Mutagenesis · EMS ·
TILLING · High resolution melting · Next-generation
sequencing · SNPs

Abbreviations

BSA	Bulk segregant analysis
CEL I	Celery endonuclease
CE	Capillary electrophoresis

CODDLE	Codons optimized to detect deleterious lesions
ComSeq	Compressed sequencing approach
dCARE	Deep candidate resequencing
DGGE	Denaturing gradient gel electrophoresis
dHPLC	Denaturing high performance liquid chromatography
EMS	Denaturing methanesulphonate
HDA	Heteroduplex analysis
HRM	High-resolution melting
IAEA	International Atomic Energy Agency; indels: insertion/deletion
MAB	Mutation-assisted breeding
MAF	Mutant allele frequency
MAS	Marker-assisted selection
mRNA	Messenger RNA
miRNA	microRNA
NGM	Next-generation mapping
NGS	Next-generation sequencing
PCR	Polymerase chain reaction
SNP	Single nucleotide polymorphism
SSCP	Single-strand conformational polymorphism
SSR	Simple sequence repeat
T-DNA	Transfer DNA
TGGE	Temperature gradient gel electrophoresis
TGS	Target genome sequencing
TILLING	Targeting induced local lesions in genomes
TPSeq	Targeted parallel sequencing
WGS	Whole genome sequencing

Introduction

One of the keys to sustainable agriculture is the creation of genetic variation in plant crops (Griggs et al. 2013). Over the past 70 years, mutation breeding has generated thousands of novel crop varieties in hundreds of crop species and billions of dollars in additional revenue, delivering higher yields, increased nutritional value and resilience to the effects of climate change, such as resistance to diseases and tolerance to drought. Mutation refers to sudden and heritable changes to the genetic material in any organism not caused by normal genetic segregation or recombination including spontaneous mutations and induced mutations (Suprasanna et al. 2015; Mba 2013; Austin et al. 2011; Oladosu et al. 2016). Spontaneous mutations occur in nature at a very low rate;

therefore, physical and chemical mutagens are important for mutation induction in crop breeding programs. Mutations theoretically encompass all changes that occur in the DNA sequence and result in changes in the genetic code (van Harten 1998). An induced mutation is an alternative and complementary technique in plant breeding for the introduction of genetic changes and the establishment of new genetic resources. In the 1920s, Lewis Stadler at the University of Missouri discovered the mutagenic effects of X-rays in barley by exposing barley seeds. Since the 1950s, it has been widely used, specifically in crops with low genetic variability and those that are not amenable to improvement through conventional breeding methods. The number of physical and chemical mutagens used in mutation breeding is large and continues to increase (Mba et al. 2010; Mba 2013). According to the International Atomic Energy Agency (IAEA) Mutant Varieties Database (<http://mvgs.iaea.org>), 3233 mutant varieties of 214 different plant species have been officially released worldwide (Joint 2015). Of these, 49.5, 21.9, and 15% belongs to cereals, ornamental and decorative plants and legumes, respectively (Fig. 1). Gamma rays have been most used successfully to develop new mutant varieties of ornamentals with novel leaf and flower colour/shape (Taheri et al. 2014).

In any plant breeding program, genetic variation is necessary for crop improvement. Using mutagenesis to create novel genetic variation is extremely effective to promote natural genetic resources with the induction of random changes throughout the genome, resulting in a single plant with a large number of different mutations.

Chemical and physical mutagenesis has successfully assisted the development of improved and new cultivars among both seed and vegetatively propagated crops (Parry et al. 2009). Seeds of seed-propagated crops are treated with the mutagens while in vegetatively propagated plants stem cuttings, twigs, bulbs, buds, rhizomes and tubers are exposed to the mutagens. Chemical mutagens most often produce point mutations (Kodym and Afza 2003) and chromosome breaks that result in various chromosomal rearrangements that affect plant fertility and cause lethality. Among the different chemical mutagens, ethyl methane sulphonate (EMS) is the most powerful, effective, reliable and frequently used chemical mutagens in plants. Treatment with the mutagen breaks the nuclear DNA, and during the process of DNA repair, new mutations are induced randomly and are heritable. The changes can also occur in cytoplasmic organelles, again resulting in

Fig. 1 Percentage (%) of mutant varieties by crop type reported on the Joint FAO/IAEA Mutant Varieties Database (Joint FAO/IAEA 2015)

chromosomal or genomic mutations that enable plant breeders to select useful mutants such as those with modified flower colour, flower shape, disease resistance and early flowering phenotypes (Jain 2010). Breeders are most interested in point mutations because large-scale changes can have negative effects on chromosome structures. However, large-scale changes in chromosome structure increase the number of recombination events, and the breaking of undesirable linkages is also highly valuable (Parry et al. 2009). EMS mainly produces C/G to T/A transitions with the induction of C to T substitutions, and EMS also generates G/C to C/G or G/C to T/A transversions or A/T to G/C transitions through 7-ethylguanine hydrolysis and through 3-ethyladenine pairing errors at a low frequency (Serrat et al. 2014). More recently, with biotechnology approaches such as in vitro cell and tissue culture and totipotency advantages, i.e. regeneration of the complete plant from individual plant cells, mutation breeding in vegetatively propagated plants has become more effective (Mba 2013). Various important factors affect the results of chemical mutagenesis, including the mutagen concentration, mutagen volume with respect to the sample size, treatment duration, temperature, pre-soaking of seeds, pH, use of catalytic agents and post-treatment handling of the mutants (Mba et al. 2010). Using induced mutations and enhancing mutation frequency rate

and recombination of available desired genes from gene pools and the related plant species by sexual hybridization, plant breeders could successfully develop new cultivars with high yield and resistance to abiotic and biotic stresses. One of the advantages of mutation induction is the chance of obtaining unselected genetic variation when one or few characters of a prominent cultivar are to be modified to improve plants (Jain 2010). However, classical induced mutagenesis has some disadvantage, such as (1) the requirement of a large mutant population generation, (2) occurrence of chimeras, and (3) heterozygosity of the mutated loci. Mutation-assisted breeding (MAB) along with biotechnology approaches can improve these limitations without affecting the requirement of generating large mutant populations. Novel biotechnology approaches have made it feasible to survey specific regions of the genome that control a trait of interest for the introduction of alterations in large mutant populations with desirable genetic backgrounds (Mba 2013). This review briefly discusses the progress in the use of modern technologies such as targeting induced local lesions in genomes (TILLING) in combination with high-resolution melting (HRM) and next-generation sequencing (NGS) techniques for detection of mutations in mutation breeding programs to enhance the efficiency of induced plant mutagenesis in plant breeding era.

Forward and reverse genetic approaches for gene identification

Forward and reverse genetics are two main approaches that connect genotype to phenotype with the aim of determining the function of a gene/genes by screening the phenotype or genotype of individual mutants. In the classical genetics approach, known as forward genetics, mutant phenotypes (spontaneous or induced mutants) are compared with wild type to identify and characterize the causal genes.

In reverse genetics, the function of a gene identifies based on analysing the phenotype resulting from known gene sequence changes, in contrast to the reliance of forward genetics on identification of responsible genes for a specific phenotype (caused by induced mutagenesis or natural variation) (Jankowicz-Cieslak and Till 2015). A mapping population that segregates at the phenotype of interest is then produced, and candidate genes are mapped in a small region of the chromosome. Finally, the gene responsible for the desired phenotype will be identified by applying genetic engineering or reverse genetics methods (Esfeld et al. 2013; Ji 2013).

Despite the many identified responsible genes for the desired phenotype through the use of forward genetics, in most cases, genes of known sequence are not linked to a phenotype. This will happen more in non-model species that forward genetics can be more challenging due to genetic redundancy. In generated large mutant populations, single nucleotide polymorphism (SNP) discovery technologies are very useful for functional genomics, forensic medicine, clinical diagnostics, population genetics, molecular epidemiology and plant and animal breeding (Rigola et al. 2009). Advances in the genetic era, such as the discovery of DNA, the polymerase chain reaction (PCR) and genome sequencing, have led to techniques such as reverse genetic approaches (from genotype to phenotype), in which the phenotypic effect of the manipulation of a particular gene or gene product is assessed.

Classical reverse genetic approaches to identify induced mutations

Conventional and pre-screening mutation scanning methods in large mutant populations for the presence of sequence polymorphisms that require a separation step are discussed in this section. (1) The single-strand conformational polymorphism (SSCP) technique

(Orita et al. 1989) performed under non-denaturing conditions results in the formation of single-stranded DNA sequence conformers with specific characteristic mobilities during polyacrylamide gel electrophoresis. Any changes in sequence results in conformer mobility change. (2) Denaturing gradient gel electrophoresis (DGGE) (Muyzer and Smalla 1998) allows rapid screening for single-base changes in enzymatically amplified DNA. This technique is based on the migration of double-stranded DNA molecules through polyacrylamide gels containing linearly increasing concentrations of a denaturing agent. (3) Heteroduplex analysis (HDA) (Highsmith et al. 1999) is based on the detection of a mutation that results in conformational differences in the duplex DNA produced during polymerase chain reaction (PCR) amplification. The goal of heteroduplex analysis is to differentiate homoduplex DNA from heteroduplex DNA fragments based on their conformations under native conditions. (4) Denaturing high-performance liquid chromatography (dHPLC) (Underhill et al. 1997; Xiao and Oefner 2001) consists of the automatic detection of small insertions and deletions as well as single-base substitutions. In brief, produced heteroduplexes resulting from mixing, denaturing, and re-annealing two or more chromosomes that differ in sequence are retained less than their corresponding homoduplexes are in a unique DNA separation matrix. (5) In Temperature gradient gel electrophoresis (TGGE) (Li et al. 2002) a single chemical denaturant concentration is used to lower the melting temperature of all double-stranded molecules followed by the differential melting of the fragments with a temperature gradient. The sensitivity of a temperature gradient for DNA denaturation can be increased by establishing the gradient perpendicular to the direction of electrophoresis. These methods require a time-consuming step for the separation of PCR products in a gel or other matrix, which increases the risk of contamination in future reactions due to exposure of the PCR products to the environment. “SSCP, as a reverse genetic method, provides a high-throughput strategy for the detection of polymorphisms; however, it has low efficiency for the detection of novel mutations with a size limit of 200 to 300 bp in the target DNA sequence. Moreover, the high cost of performing a microarray and its low detection frequency of less than 50% (Borevitz et al. 2003; Tillib and Mirzabekov 2001) have limited its application.

TILLING: a reverse genetic technique for induced mutation detection

Sixteen years ago, the reverse genetic technique, targeting induced local lesions in genomes (TILLING) was first developed in *Arabidopsis* by McCallum and collaborators (McCallum et al. 2000a, b). TILLING is a high-throughput technique to identify single nucleotide mutations in a specific region of a gene of interest with a powerful detection method that resulted from chemical-induced mutagenesis. In the late 1990s, TILLING was first used by Claire McCallum, a graduate student (in cooperation with the Fred Hutchinson Cancer Research Centre and Howard Hughes Medical Institute), who worked on characterizing the function of two chromomethylase genes in *Arabidopsis* and applied reverse genetic approaches, such as T-DNA (incorporation of a foreign DNA into the genome of interest) lines and antisense RNA, but was unable to successfully apply these approaches to identify mutants for a DNA methyltransferase (chromotransferase) (Barkley and Wang 2008). The successful approach was what is now known as TILLING. Based on previous studies, TILLING can be applied in any species, regardless of its genome size and ploidy level and has been successfully applied in a range of crop plants such as hexaploid and durum wheat, barley, rice, tomato, maize, sorghum, soybean and potato, sunflower, melon, pea, and peanut to identify single base pair changes or small deletions in specific target genes (Uauy et al. 2009; Slade et al. 2005; Wang et al. 2012; Tsai et al. 2013; Singh et al. 2014; Sharp and Dong 2014) (see Table 1 for a list of those). Recently, Gauffier et al. (2016) used a TILLING approach in tomato to develop broad-spectrum resistance to potyviruses is prevented by eIF4E gene redundancy (Gauffier et al. 2016). For TILLING and for polymorphism analysis, the effect of missense mutations on the encoded protein must be evaluated. In this case, codons optimized to detect deleterious lesions (CODDLE) (<http://www.proweb.org/coddle/>) can be used to evaluate whether a missense mutation is likely to have an effect on the encoded protein in a selected specific DNA sequence with a length of 1 kb. CODDLE was developed as a general tool for polymorphism analysis, designing primers for any organism and detecting any mutagen based on the DNA sequence information input provided by users (De-Kai et al. 2006).

High-throughput TILLING comprised of a two-step procedure in which chemical mutagenesis (Koornneeff

et al. 1982) is followed by a sensitive mutation detection instrument. For mutation induction, either biological agents such as transposons and T-DNA (Azpiroz-Leehan and Feldmann 1997; Balcells et al. 1991), or physical agents such as fast neutron, UV, x-ray and gamma-ray radiation (Stadler 1928; Sparrow and Woodwell 1962; Kovacs and Keresztes 2002) or chemical mutagens such as N-methyl-N-nitrosourea (MNU), 1,2:3,4-diepoxybutane (DEB) or ethyl methanesulfonate (EMS) can be used. Among these, T-DNA and transposon mutagenesis generate only destroyer mutants, whereas EMS mutagenesis is the most effective, reliable and powerful which produces different types of mutants within each gene, including nonsense, missense, splicing and cis-regulatory mutants (Krieg 1963; Kim et al. 2006; Kodym and Afza 2003; Greene et al. 2003; Brockman et al. 1984).

In TILLING, a variety of mutation scanning methods have been used for screening the mutants including CEL I endonuclease mutation detection together with electrophoresis conditions (Stemple 2004; Sato et al. 2006), high-resolution melting (HRM) (Botticella et al. 2011; Bovina et al. 2014; Sestili et al. 2015; Acevedo-Garcia et al. 2016) and next-generation sequencing (Tsai et al. 2011). The most favoured mutation detection method for TILLING in plants is the use of mismatch-specific celery nuclease, CEL I, together with the LI-COR gel analyser system (LI-COR Biosciences) (Gottwald et al. 2009). This technique was very popular in early 2000 to detect induced mutations caused by chemical mutagens. With the development of more sensitive and high-throughput techniques, it has been largely superseded.

With HRM, mutations are detected in target genes by using PCR, followed by denaturing and re-annealing of the double-stranded DNA (dsDNA) product which is monitored via a DNA-binding fluorescent dye. On re-annealing of the double-stranded DNA, fluorescence increases and results in high-resolution melt curve (temperature vs fluorescence) (Reed et al. 2007).

Recently, next-generation sequencing technique tremendously facilitated the discovery of mutations in TILLING populations that results in phenotypes, both for focused and for genome-wide discovery. The advent of next-generation sequencing as an important tool for whole-genome sequencing, re-sequencing and de novo sequencing has revolutionized genetic research (Quail et al. 2012). Next-generation sequencing in a single reaction can be used to sequence a large number of

Table 1 Applied mutation detection techniques and affected traits in model and crop plants

Species/ploidy level	Mutagen	Mutation detection technology	M ₂ size	Mutation frequency (1 mutation/kb)	Traits	References
Arabidopsis (2×)	EMS	dHPLC, LI-COR	6912 3072 3712	1/170 1/300 1/89	— — —	Till et al. (2003a, b); Greene et al. (2003); Martin et al. (2009)
Maize (2×)	EMS	LI-COR	750	1/485	Chromomethylase	Till et al. (2004a, b)
Rice (2×)	EMS	LI-COR LI-COR CEL-I, Agarose gel	— 768 6912	1/1000 1/294 1/451	— — —	Wu et al. (2005); Till et al. (2007); Serrat et al. (2014)
Rice (2×)	EMS	TILLING	1860	—	—	Casella et al. (2013)
Rice (2×)	EMS	TILLING by Sequencing	2048	1/293	Phytic acid metabolism	Kim and Tai (2014)
Barley (2×)	EMS	dHPLC	9216	1/1000	Floral organ regulation	Caldwell et al. (2004)
Barley (2×)	EMS	LI-COR	10,279	1/500	Row type morphology and immunity to fungus	Gottwald et al. (2009)
Barley (2×)	NaN ₃	TILLMore CEL-I, Agarose gel	5600	1/374	Starch metabolism	Talamè et al. (2008); Bovina et al. (2011); Sparla et al. (2014)
Wheat (6×)	EMS	TILLING-HRM	2020	1/26	Powdery mildew disease Resistance (TaMlo gene)	Acevedo-Garcia et al. (2016)
Wheat (6×)	EMS	TILLING by sequencing	4500	1/35000	—	King et al. (2015)
Wheat (6×)	EMS	LI-COR PAGE, LI-COR CEL-I, Agarose gel	10,000 1536 2348	1/24 1/38 1/37, 1/23	Starch quality Starch quality Starch quality and grain Hardness	Slade et al. (2005, 2012); Uauy et al. (2009); Dong et al. (2009a, b)
Wheat (6×)	EMS	LI-COR, HRM LI-COR	4500 4244	1/84 —	Starch quality Starch biosynthesis	Botticella et al. (2011); Sestili et al. (2010)
Wheat (6×)	EMS	Agarose gel, PAGE	2610	1/34; 1/47	Spike development	Chen et al. (2012)
Wheat (6×)	EMS	Direct sequencing	630	1/3	Grain hardness	Feiz et al. (2009)
Wheat (4×)	EMS	PAGE-LI-COR	8000 1386	1/40 1/51	Starch quality Starch quality	Slade et al. (2005, 2012); Uauy et al. (2009)
Wheat (4×)	EMS	TILLING-HRM	3992	—	Starch metabolism	Bovina et al. (2014); Sestili et al. (2015)
Wheat (4×)	EMS	CEL I, agarose gel, dHPLC	1140	1/77	Carotenoid metabolism	Colasunno et al. (2016)
Wheat (2×)	EMS	CEL-I	1400	1/1300	Grain quality and lignin Biosynthesis	Rothe (2010)
Wheat (2×)	EMS	CEL-I	1532	1/92	Waxy and lignin	Rawat et al. (2012)
Sorghum (2×)	EMS	LI-COR	1600	1/526	Forage digestibility	Xin et al. (2008)
Sorghum (2×)	EMS	TILLING by sequencing (ComSeq)	1024	<0.1%	—	Nida et al. (2016)

Table 1 (continued)

Species/ploidy level	Mutagen	Mutation detection technology	M ₂ size	Mutation frequency (1 mutation/kb)	Traits	References
Soybean (4×)	EMS	LI-COR	768; 529 40,000	1/550; 1/140 1/1000 to 1/1300	—	Cooper et al. (2008); Anai (2012)
<i>Brassica rapa</i> (2×)	EMS	LI-COR	1344	1/130.8, 1/41.5	—	Wang et al. (2008)
<i>Brassica rapa</i> (2×)	EMS	LI-COR	9216	1/60	DNA methylation	Stephenson et al. (2010)
<i>Brassica oleracea</i> (2×)	EMS	LI-COR	8750	1/447	Wax biosynthesis and dwarf stature	Himelblau et al. (2009)
<i>Brassica napus</i> (canola) (4×)	EMS	LI-COR - NGS	3158	1/109	Mutation identification in genes of interest	Gilchrist et al. (2013)
<i>Lotus japonicus</i> (2×)	EMS	LI-COR, CE	4904	1/502	Nodule development	Perry et al. (2003); Wang and Robson (2014)
<i>Medicago truncatula</i>	EMS	LI-COR	4500, 4350	1/485, 1/242.5	—	(Le Signor et al. 2009)
Sunflower (2×)	EMS	LI-COR	3651	1/475	Fatty acid biosynthetic pathway and downy mildew resistance	Sabetta et al. (2011)
Tomato (2×)	EMS	CE, HRM	8225	1/737	Proline biosynthesis	Gady et al. (2009)
Tomato (2×)	EMS	LI-COR	4741, 1926 4759 3052	1/322, 1/574 1/574 1/1237	Shelf life Virus resistance Shelf life	Minoia et al. (2010); Piron et al. (2010); Gauffier et al. (2016); Okabe et al. (2011, 2012)
Tomato (2×)	EMS	TILLING	3052	—	Ascorbate biosynthesis	Baldet et al. (2013)
Tomato (2×)	EMS	TILLING	—	—	Fruit set mechanisms	Mazzucato et al. (2015)
Tomato (2×)	EMS	LI-COR	—	—	Lycopene synthesis	Silletti et al. (2013)
Peanut (4×)	EMS	LI-COR	3420	1/967	Seed quality	Knoll et al. (2011)
Peanut (4×)	EMS	TILLING by sequencing	768	1/1066	Stress resistance	Guo et al. (2015)
Pea (2×)	EMS	LI-COR	8000	1/669	Gibberellin metabolism	Triques et al. (2007)
Flax	EMS	LI-COR	4894	—	—	Chantreau et al. (2013)
<i>Cucurbita pepo</i>	EMS	LI-COR	1464	—	—	Vicente-Dólera et al. (2014)
<i>Brachypodium distachyon</i> (2×)	NaN ₃	LI-COR	5530	1/396	Lignin biosynthesis	Dalmais et al. (2013)
Tobacco (2×)	EMS	TILLING by sequencing	3072	1/1423	Leaf yield	Reddy et al. (2012)

EMS ethyl methanesulfonate, TILLING targeting induced local lesions in genomes, dHPLC denaturing high performance liquid chromatography, HRM high-resolution melting, NGS next generation sequencing

DNA sequences. The high-quality DNA is sheared into fragments of a specific size and specific adapters are ligated to the 3' and 5' ends (Buermans and Den Dunnen 2014). These DNA templates are immobilized on a flow

cell surface and amplified by sequential addition of nucleotides. Some of these techniques are applicable to many species, and each technique has advantages and limitations (Tables 1 and 2).

Table 2 Comparison of different methods involved in mutation detection in TILLING populations

Method	Advantages	Disadvantages	Reference
TILLING-CEL I	<ol style="list-style-type: none"> 1. High sensitivity 2. High throughput 3. The populations generated for TILLING in any species provide valuable resources for teaching and research 4. Suitable for polyploids with higher mutation frequency 	<ol style="list-style-type: none"> 1. Rely on time and cost consuming enzymatic screening system 2. Needs multi-dimensional pooling 3. Only 5% of the total mutations in the EMS-mutagenized populations are truncations 	Wang et al. (2012); Greene et al. (2003); Bleecker and Kende (2000); Byrne (2006); Eckardt (2007); Gilchrist et al. (2013); Parry et al. (2009); Perry et al. (2003)
TILLING-HRM	<ol style="list-style-type: none"> 1. Non-enzymatic screening system 2. High sensitivity 3. Time and cost saving 	<ol style="list-style-type: none"> 1. Depends strongly on good PCR instruments and dyes. 2. Needs multi-dimensional pooling 3. Small insertions and deletions may be somewhat more difficult to detect than substitutions. (4) Detection sensitivity is limited to amplicons of <450 bp. 	Reed and Wittwer (2004); Lochlainn et al. (2011); Wittwer (2009); Gady et al. (2009); van der Stoep et al. (2009); Simko (2016); Tindall et al. (2009); Chen et al. (2014a)
TILLING-NGS	<ol style="list-style-type: none"> 1. Non-enzymatic screening system 2. High throughput 3. Time saving 4. Identification of mutations through targeted sequencing 5. More efficient in polyploids 6. Mutation detection in pools deeper than eight individuals 	<ol style="list-style-type: none"> 1. Cost is still high 2. Needs multi-dimensional pooling 3. High rate of incorrectly identified DNA bases in the sequence data produced, where billions of base calls translate to millions of errors which are not easily diagnosed or corrected. 4. The processing and storage of massive amounts of sequence data. Data analysis can be time-consuming and may require special knowledge of bioinformatics to garner accurate information from sequence data. 	Zargar et al. (2015); Jünemann et al. (2013); Egan et al. (2012); Meacham et al. (2011); Ganai et al. (2009)

An overview of TILLING methodology

In the basic procedure of TILLING, seeds are mutagenized by treatment with ethyl methanesulphonate (EMS). The alkylating agent (EMS), which has great mutagenic potential, is the mutagen most commonly applied for almost all TILLING populations (Kurowska et al. 2011). Using the EMS mutagen, a high rate of mutation can be achieved that spreads out over the genome randomly without excessive DNA damage (Gilchrist and Haughn 2005). The developed M_1 plants are self-fertilized, and M_2 individuals are used to prepare DNA samples for mutational screening. The DNA samples are pooled and arrayed on microtiter plates followed by PCR amplification of the targeted DNA segment. Amplification products are incubated with a celery endonuclease (CEL I) that cleaves mismatches in heteroduplexes between mutants and wild type by preference. CEL I endonuclease is a member of the S1 nuclease family of single-strand-specific nucleases (Oleykowski et al. 1998) that cleaves on the 3' side of

mismatches and loop outs in heteroduplexes between wild-type and mutant DNA while leaving duplexes intact. Cleavage products generated by endonuclease are separated and electrophoresed using the LI-COR gel analyser system, and gel images are analysed with the aid of a standard commercial image-processing program (Adobe Photoshop; Adobe Systems, Mountain View, CA). Because mutations are detected on complementary strands, amplification products with different amounts of double end-labelling allow rapid visual confirmation and therefore can be easily distinguished from amplification artefacts. Upon mutation detection in a pool, the plant carrying the mutation is identified by screening individual DNA samples. The mutation is discovered by sequencing the target gene segment and identifying the type of nucleotide change (Lee et al. 2014). Figure 2 shows the general TILLING procedure.

Whereas the establishment of the initial screening population and the corresponding ordered DNA samples requires an investment of time and money, a

complementary approach to traditional TILLING, individualized TILLING (iTILLING), has been developed for Arabidopsis and is more cost and time effective (Bush and Krysan 2010). Furthermore, HRM as mutation detection method in TILLING populations (Botticella et al. 2011) with no reliance on enzymatic screening system or ‘TILLING by sequencing’ (Tsai et al. 2011) technologies have improved great potential of TILLING strategy for mutation detection which have been explained in the following sections.

Advantages and disadvantages of TILLING

The generated mutant population for any species using TILLING provides valuable resources for teaching and research. TILLING is an easy technique that does not require complicated manipulations or expensive equipment. It is used to easily screen point mutations to identify the functions of specific genes without the tedious tissue culture procedures that are involved in anti-sense RNA and RNAi. In addition, TILLING provides a high efficiency of mutation detection and high sensitivity attributed to the combination of CEL I, double-end fluorescent dye labelling and the LI-COR system as an alternative to dHPLC, as well as high frequencies of mutagenesis and a high-throughput screening capacity (De-Kai et al. 2006). Among reverse genetic techniques, TILLING can be used to identify genetic variability in either mutagenized or natural populations. In addition, unlike transgenic plants, for which there are some barriers to their creation or marketing, TILLING can be applied in plant research and breeding programs in various species because the variation does not involve the transformation of exogenous genetic material (Gilchrist et al. 2013). Using TILLING in a small and a large number of original research studies for a wide range of plants species (Table 1) supports the power of the TILLING technique for the investigation of genetic alterations in organisms with different genome sizes and ploidy levels. In contrast to the advantages of TILLING, it also has some limitations, such as the high cost and time commitment associated with the development of a mutagenized population for most species. Furthermore, the technique itself is labour-intensive, and the poor cleavage efficiency of the endonuclease CEL I and 59-39 exonuclease activity results in a reduction of signal/noise levels and prevents the performance of pooled sample analyses with more than eight samples per pool (Till et al. 2006). In addition, TILLING cannot provide information about the nature

of the sequence changes and their possible effect on gene function; therefore, once mismatches have been detected, the results must be verified by sequencing, which incurs additional time and cost (Gilchrist and Haughn 2010) (Table 2). There are some bioinformatics tools such as the project aligned related sequences and evaluate SNPs (PARSESNP) program (<http://www.proweb.org/parsesnp/>) (Taylor and Greene 2003) to aid the researcher in deciding which mutations to characterize. Once TILLING process is completed, mutations are automatically analysed by the PARSESNP which provides graphical and tabular information on the location (PSSM difference) and severity (SIFT score) of mutations and provides information on the creation or loss of restriction sites caused by the induced polymorphisms. PSSM score ≥ 10 indicates a mutation that is more likely to have a damage effect on protein function. SIFT scores < 0.05 have been empirically determined to be deleterious (Till et al. 2003a; Xin et al. 2008).

Some critical weak points of TILLING with CEL I endonuclease mutation detection method, as mentioned above, have resulted in the introduction of a diverse set of new screening platforms that do not rely on endonuclease mismatch cleavage, such as high-resolution DNA melting analysis (HRM), in which only simple PCR is performed before melting curve analysis, and next-generation sequencing which has recently been successfully applied in TILLING.

HRM: an alternative screening technique

HRM is a post-polymerase chain reaction (PCR) method that was established as an alternative screening technique for the detection of genetic variations (SNPs, mutations, methylation) in PCR amplicons (Zhou et al. 2005, 2004). Although dsDNA is very stable at room temperature, its two strands begin to separate with increasing temperature. The melting temperature (T_m) is the temperature at which 50% of the DNA is single-stranded. The length and guanine-cytosine (Ling et al. 2013) content of the DNA fragment affect T_m . The three hydrogen bonds of GC base pairs increase their stability compared with adenine-thymine (AT) base pairs linked by only two hydrogen bonds. Therefore, DNA sequences with a high GC content have a higher T_m than do DNA sequences containing a low number of GC base pairs (Druml and Cichna-Markl 2014). Melting of a short double-stranded region occurs at a faster rate if the regions melts in one transition without

Fig. 2 Diagram of the TILLING strategy. TILLING procedure will be started by seed mutagenesis with a chemical mutagen and germinated to produce M₁ plants. M₁ plants are allowed to self-pollinate to generate M₂ plants. M₂ seeds will be collected for banking, and the DNA will be extracted from part of interest of plant for mutation discovery. Eightfold DNA pooling and DNA array in a two-dimensional format on 96-well plates. After PCR amplification of target genes with designed primers, upon heating

and cooling steps, the heteroduplexes are formed and then incubated to digest with a mismatch endonuclease (CELI) for heteroduplex mismatch cleavage. Cut strands are separated by denaturing polyacrylamide/agarose gel electrophoresis and visualized by fluorescence detection using a Li-Cor DNA analyser. After identification of mutations, the polymorphic individuals are sent to sequencing to verify the induced mutation

intermediate states; in contrast, longer PCR products may melt in multiple stages or ‘domains’ (50–500 bps) (Erali and Wittwer 2010). HRM is a new screening

method for DNA polymorphisms that was introduced in 2002 by a collaboration between academics (University of Utah, UT, USA) and industry (Idaho

Technology, UT, USA). Some benefits, such as simplicity, low cost, no need for enzymatic screening system and high sensitivity/specificity, make HRM an important new tool for genotyping, SNP polymorphisms, insertions or deletions (indels) and DNA methylations detection in PCR amplicons (Herrmann et al. 2007; White et al. 2007).

HRM working plan The HRM working plan is based on the changes in fluorescence along with the melting of double-stranded DNA that is measured using a saturated DNA-binding dye and a highly accurate optical detection system without the need for costly labelled probes. Three factors are involved in the HRM process: dye chemistry, instrument resolution and data analysis. HRM involves several main steps: (1) mutagenesis of plant material by EMS (seeds, pollen, leaf, flower, rhizome, root, among others), (2) DNA extraction, (3) PCR amplification with gene-specific primers labelled with a fluorescent dye, (4) high-resolution melting and (5) data analysis. First, PCR is performed in the presence of a dye that binds to double-stranded DNA (dsDNA). This dye emits low levels of fluorescence when unbound but is highly fluorescent in the bound state. Popular dyes for this application include CGreen/LCGreen Plus and SYBR Green I, which have a high sensitivity, stability, reliability and compatibility with many popular HRM instruments (Carén et al. 2006; Herrmann et al. 2006). However, the high cost of these dyes and the difficulties associated with the detection of heteroduplexes have limited their use (Pornprasert et al. 2008; Price et al. 2007; Worm et al. 2001). Recently, EvaGreen and ResoLight HRM dye (He et al. 2014) were found to be a less expensive saturated DNA binding dye with equal binding activity for GC-rich and AT-rich regions, without inhibiting amplification and no sequence preference (Wittwer 2009; Li et al. 2010). Following PCR, as the temperature is gradually increased, the changes in fluorescence in each sample during DNA melting are displayed by the temperature-normalized melting curves. The differences among curves demonstrate the relative difference in fluorescence (Δ Fluorescence, ΔF) of a respective sample in comparison to the non-mutant sample. A ΔF value of 0.05 is considered significant, according to the manufacturer's instruction (Li et al. 2010; Simko 2016; Hofinger et al. 2009). During HRM, even a subtle and small DNA variation that results in allelic differences among PCR amplicons can be detected based on melting curve differences. The

demonstrated advantages of HRM analysis have resulted in the use of this approach to detect DNA polymorphisms in plant species such as apple, barley, grapevine, olive, almond, pepper, sweet cherry and *Solanum lycopersicum* L. (Liew et al. 2004; Wu et al. 2008; Chagné et al. 2008; Lehmensiek et al. 2008; Mackay et al. 2008; Donini et al. 2009; Gady et al. 2009; Ganopoulos et al. 2011; Golding et al. 2010). An exchange between G/C and T/A base pairs results in relatively large changes in T_m of approximately 0.8–1.4 °C (Liew et al. 2004). Duplex melting is generally monitored using intercalating dyes, although fluorescently labelled primers have also been used. HRM detection is based on the principle of melting curve changes during the process of DNA denaturation based on the primary sequence. The operation can be performed immediately after PCR without electrophoresis and in the same reaction tube. In comparison to electrophoresis, the detection time is reduced and the resolution is increased (Yu et al. 2013). Furthermore, HRM in combination with TILLING was used to screen an EMS-mutagenized wheat population to identify SNPs in hexaploid and durum wheat *Sbella* genes (Botticella et al. 2011; Bovina et al. 2014).

Advantages and disadvantages of HRM Among all the available scanning techniques, HRM is simple, rapid and inexpensive, and it is the only method that is performed in a closed tube and can be carried out in the same container used for PCR amplification in almost 15 min without need for any enzymatic screening system. The closed tube makes it possible to send the samples immediately after scanning for genotyping or sequencing (if necessary), without any need for processing or automation. In addition, contamination concerns are decreased when PCR products are not exposed to the environment (Reed and Wittwer 2004). However, the efficiency of HRM depends critically on the DNA concentration and quality, effective PCR, instrument decontamination and resolution and dyes (Herrmann et al. 2007). Some limitations, such as the GC content and length of the amplicon, affect the HRM efficiency, resulting in much shorter typical reads (only covering 150–500 bp) than those obtained with Li-Cor and CE. The usefulness of using HRM in the TILLING approach is highlighted when the target is a specific region with a known impact on protein structure or when the gene of interest contains many short exons and thus a short read length is acceptable. In addition, specialized software is

needed to analyse the different melt curves. In addition, sequencing of the amplicons is required to determine the number and position of mutations within the amplicon (Chateigner-Boutin and Small 2007). Other limitations of HRM are the difficulty associated with the detection of SNPs when the polymorphism occurs too close to either of the primer binding sites, the complex sequence, or the existence of a large number of alleles in the analysed population (Mader et al. 2008), the presence of multi-locus markers, the non-specific PCR amplicons (Distefano et al. 2012) and the presence of stutter bands (Dossett et al. 2010), which prevent reliable interpretation of the melting curve profiles. Nevertheless, HRM can be performed on standard qPCR machines with a simple software upgrade and thus is a suitable platform for initial TILLING screening (Sikora et al. 2012; Simko 2016) (Table 2). Future research should focus on improvements in hardware, software and DNA-binding dyes to increase the accuracy of the melting curves. Additionally, more research should be conducted to improve the HRM capacity to analyse longer amplicons, the reproducibility of the results and the possibility of results that can be transitioned across experiments and laboratories (Simko 2016).

Next-generation sequencing applications for causative EMS-generated mutation identification

Next-generation DNA sequencing technologies provide new opportunities for plant breeding programs. The detection and exploitation of genetic variation have always been an essential part of plant breeding. Next-generation sequencing techniques became commercially available in approximately 2005, providing great opportunities for the life sciences (Egan et al. 2012). Below, we will briefly describe a range of NGS applications in plant mutation breeding. Important applications are as follows.

Mapping by sequencing For many years, to identify genes and mutations in phenotype of interest, forward genetic screens (Page and Grossniklaus 2002; Candela and Hake 2008) has been applied as a powerful method for researchers. However, in classical forward genetic screens, mapping the casual mutation using genetic crosses is a complex and multistep procedure. Traditional mapping or positional cloning of mutations has been replaced by NGS using whole-genome sequencing (Smith et al. 2008; Srivatsan et al. 2008; Blumenstiel

et al. 2009; Irvine et al. 2009; Lister et al. 2009; Schneeberger et al. 2009; Zuryn et al. 2010; Pawełkowicz et al. 2016). Ideally, the sequencing of mutant genomes is a simple way to identify mutations that cause phenotypes of interest. However, due to the numerous unrelated polymorphisms that segregate with the causative mutation in a mutagenized population, a very low signal is obtained. Therefore, some genetic analyses to purify the chromosomal location carrying the causative mutation are needed, even for NGS mapping approaches. Mapping by sequencing approach through NGS accelerated identification of casual mutations at SNP level even in complex genetic backgrounds (Schneeberger 2014; Schneeberger and Weigel 2011). After mutagenesis, mutation screening strategies depend on species, its breeding system, dominant or recessive nature of casual mutation. Nevertheless, mapping by sequencing is not dependent on reference genome sequences, genetic crosses and any kind of linkage information. In NGS mapping approaches, the background noise has been improved first by genetic analysis to refine the genomic region of interest or through bulk analysis of a very large number of mutant lines. In these years, several mapping by sequencing analysis pipelines have been introduced and have been applied in model plants (James et al. 2013), non-model plants (Nordström et al. 2013) and crop species like rice (Abe et al. 2012).

An approach called simultaneous mapping and mutation identification by deep sequencing (SHOREmap) by the principal of mutant allele frequency (MAF) estimation, including in isogenic backcross populations through whole-genome sequencing (WGS), was able to successfully map a causative mutation using Illumina Genome Analyser (GA) sequencing of a population of 500 pooled F₂ lines of *Arabidopsis* (Schneeberger et al. 2009). However, in that study, the strength of SHOREmap was not specified in fewer F₂ lines for mapping mutations with difficult-to-score phenotypes or working with organisms for which it is difficult to propagate large numbers. This straightforward method was applied in non-reference *A. thaliana* accession (Uchida et al. 2011, 2014) and in non-EMS mutations in non-model plants (Guo et al. 2012) to identify mutations that resulted in phenotypes of interest. The employed methodology was bulked segregant analysis (BSA), in which, after crossing a mutant to a different polymorphic accession, whole-genome sequencing data are extracted from pooled F₂ segregants and analysed for single nucleotide polymorphisms (SNPs). The

candidate SNPs are then detected and extracted using linkage analysis of the SNPs in the accessions used to produce the F₁ generation and filtering the linked SNPs with multiple appropriate criteria (Uchida et al. 2014). By contrast, in pools of mutant recombinants, dominant or semi-dominant mutations do not fix commonly.

Later, a DNA-based mapping-by-sequencing approach called next-generation EMS mutation mapping (NGM) by the principle of homozygosity mapping was developed by Austin et al. (2011) in *A. thaliana* using Illumina GA data to reliably and easily map candidate causative mutations even in small F₂ populations. Additionally, no prior mapping information is needed to perform this protocol (Austin et al. 2011). This tool is a Web-based service (<http://bar.utoronto.ca/ngm>) which (1) supports mutation mapping and identification and filtering for their effects and (2) visualizes chromosome-wide SNP densities and mapping scores, including manual refinement of mapping intervals. However, there is an uncertainty in the exact feature(s) of the NGM pipeline for mutant identification in fewer F₂ populations.

Although described approaches are fast and facile, their application is affected by some obstacles including inter-accession crosses necessity that diminishes the success rate of genetic modifier screens and recognition of subtle phenotypic variation of mutants in F₂ populations. Ashelford et al. (2011) applied resequencing of a novel Arabidopsis clock mutant early bird (*ebi-1*) genome that had been backcrossed four times to the parental line to create many candidate mutants. To reduce the cost of whole-genome sequencing of SHOREmap or NGM for large amounts of impractical and uninformative non-target regions, Liu et al. (2012) developed a time and cost-effective and facile targeted parallel sequencing (TPSeq) method without necessity of advanced computational devices and skills and identified three novel nitrate-signalling mutants in Arabidopsis, concurrently.

Furthermore, rice researchers developed another DNA-based mapping-by-sequencing method called MutMap by the principle of MAF estimation in backcross populations to use genome sequencing to identify the genomic position of EMS-induced mutants that are most probably harbouring agronomically important traits such as pale green leaves and semi-dwarfism (Abe et al. 2012). MutMap is based on the crossing of

a M3-M5 mutant of interest directly to the original wild type and followed by selfing of F₁ individuals, allowing clear segregation in second generation (F₂) of subtle phenotypic differences. Then, DNA from 20 F₂ individuals with mutant phenotype will send for Illumina whole-genome sequencing to detect casual SNPs different from the reference genome. Although this approach minimized the number of crosses in crop species and required mutant F₂ progeny, it is not suitable for plants without reference genome sequence and mutants with early development lethality or sterility.

To reduce the number of causal candidate mutations, Hartwig et al. (2012), combined deep candidate resequencing (dCARE) using the new Ion Torrent Personal Genome Machine sequencing platform to identify causative mutations for the suppression of like heterochromatin protein1 (*lhp1*), a gene involved in chromatin-mediated gene repression. Notwithstanding that this method exploits backcross principle and SHORE pipeline for mapping analysis and necessity of targeted deep sequencing of candidates to isolate exact causative SNPs, it is a capable method for mapping-by-sequencing. Thereafter, Allen et al. (2013) screened leaf hyponasty EMS-induced mutants via single backcrossing of the mutant to its parent followed by whole-genome deep sequencing of F₂ population and analysis with NGM approach to clearly identify causal mutation in HASTY gene involved in microRNA biogenesis. Avoiding MutMap obstacle, MutMap+ was developed for the identification of causal SNPs in rice mutants and gene isolation in crops that are recalcitrant to artificial crosses (Fekih et al. 2013).

Further studies on using next-generation sequencing for mutant identification have been done by Mateo-Bonmati et al. (2014). They combined conventional linkage mapping and Illumina whole-genome resequencing to identify the causal mutations in four loss-of-function *angulata* (*anu*) mutants which affect deficits in leaf shape and pigmentation (Mateo-Bonmati et al. 2014). Similarly, in Arabidopsis, next-generation sequencing of backcrossed bulk segregants was applied to identify abscisic acid (ABA)-resistant root elongation (AR) mutants (Thole and Strader 2015; Thole et al. 2014). Recently, Cheng et al. (2015) successfully applied next-generation sequencing in targeted mutation detection in hexaploid crop, *Crambe abyssinica*, to detect causative mutations in *CaFAD2* gene leading to changes in crambe oil composition (Cheng et al. 2015).

SNP discovery through NGS SNP discovery is the most common application of NGS. SNPs have many applications in the construction of linkage mapping, genetic diversity analyses, association mapping and marker-assisted selection in several species (Cortés et al. 2011; Ray and Satya 2014; Thomson 2014; Huq et al. 2016). The discovery of SNPs using NGS has been reported in grapevine (*Vitis*) (Yang et al. 2016), cucumber (*Cucumis sativus* L.) (Pawelkiewicz et al. 2016), pepper (*Capsicum*) (Devran et al. 2015), cabbage (*Brassica oleracea*) (Lee et al. 2015), olive tree (*Olea europaea* L.) (Kaya et al. 2013), sunflower (*Helianthus annuus* L.) (Pegadaraju et al. 2013), soybean (*Glycine max*) (Vidal et al. 2012), pepper (*Capsicum spp.*) (Ashrafi et al. 2012), wheat (Allen et al. 2011; Trebbi et al. 2011; Shavrukov et al. 2014), lupine (*Lupinus angustifolius* L.) (Yang et al. 2012), eggplant (Barchi et al. 2011), rice (Feltus et al. 2004; McNally et al. 2009; Yamamoto et al. 2010; Chen et al. 2014b; Zheng et al. 2016), Arabidopsis (Zhang and Borevitz 2009; Jander et al. 2002), barley (Close et al. 2009; Waugh et al. 2009), sorghum (Nelson et al. 2011), cotton (Byers et al. 2012), common beans (Cortés et al. 2011), soybean (Hyten et al. 2010), potato (Hamilton et al. 2011), flax (FU and Peterson 2012), *Aegilops tauschii* (You et al. 2011), alfalfa (Han et al. 2011), oat (Oliver et al. 2011), maize (Jones et al. 2009) and chickpea (*Cicer arietinum* L.) (Azam et al. 2012; Gaur et al. 2012) to name a few. Small plant genomes with a good reference genome, such as rice and Arabidopsis, are available in which SNP-derived NGS has been performed completely (Yamamoto et al. 2010; Ossowski et al. 2008). The quality of the SNP calling depends on factors such as the presence of repeat elements and incomplete or inaccurate reference genome sequences (Treangen and Salzberg 2012; Kumar et al. 2012).

TILLING by sequencing Although in TILLING a number of platforms has been applied for mutation detection of target genes within genomic DNA samples, most of these techniques rely on PCR amplification of target mutant region and wild type together to detect mismatches by heteroduplex amplicons using CEL I followed by gel electrophoresis or by HRM analysis. Such approaches are labour intensive and challenged by pools exceeding eight individuals, and screening a single target at a time is limited. Moreover, sequencing is required to characterize the mutations. ‘TILLING by sequencing’ technology demonstrates great potential

by applying NGS in the TILLING strategy for mutation detection. An example of the use of Illumina sequencing in TILLING (Tsai et al. 2011; Rigola et al. 2009; Tsai et al. 2015; Granier et al. 2015) is the implementation of a flexible and effective combination of TILLING-NGS methods to detect mutations in targeted loci in rice and wheat. Likely, TILLING by sequencing technology was applied to identify mutations in genes to enhance leaf yield in tobacco (*Nicotiana tabacum*) (Reddy et al. 2012) and to detect mutations in genes related to biotic and abiotic stress resistance in peanut (*Arachis hypogaea*) (Guo et al. 2015). The method is characterized as follows: (1) TILLING-NGS is a high-throughput and reliable mutant identification method that is performed by the application of multidimensional pooling and a probability threshold, (2) it enables the identification of a mutation with an associated base change and effect, (3) it does not rely on labelled fluorescent primers or potentially variable endonuclease digestion, (4) it allows a flexible choice of pooling methods and species and (5) it is scalable in scope and experimental combinations.

In another study, Krothapalli et al. (2013) used next-generation sequencing in EMS-induced mutant of *Sorghum bicolor* defective in hydrogen cyanide release. They identified point mutation that resulted in a premature stop codon in the coding sequence of dhurrinase2, which encode a protein involved in the dhurrin catabolic pathway that is the cause of an acyanogenic sorghum mutant phenotype (Krothapalli et al. 2013).

Recently, Nida et al. (2016) applied a novel method to identify carriers of rare SNPs with frequency of less than 0.1% in TILLING population with compressed sequencing approach (ComSeq) in *S. bicolor*. ComSeq is a combination of NGS with new mathematical field of compressed sensing (CS) (Shental et al. 2010; Erlich et al. 2010). Compared to other TILLING population detection methods (Tsai et al. 2011; Missirian et al. 2011) which are based on ‘multidimensional pooling’ and detect only a single carrier, ComSeq detects many carriers without requiring sophisticated SNP calling methods (Missirian et al. 2011). Furthermore, this approach is a very cost-effective tool for breeders to identify novel alleles.

Mutation scanning by exon capture and next-generation sequencing Although TILLING by sequencing technique has improved traditional TILLING, because of the limit in the number of genes targeted in each run

and normalization of DNA samples at several stages, this method is still labour-intensive. In addition, despite declining costs of next-generation sequencing (Shendure and Ji 2008), it is still costly enough for researchers in eukaryotic species with large genomes. Fortunately, an alternative method was developed to focus on isolating of high-value coding gene sequences (exons) (comprising 2% of eukaryotic genome) and resequencing coding portions genome-wide of large genomes for SNP discovery (Hodges et al. 2007; Cosart et al. 2011). This exon capture method has been applied in durum wheat to identify variations in functionally important regions of the genome (Saintenac et al. 2011) and to identify SNPs within parents of mapping populations of bread wheat to allow the development of high-density maps (Winfield et al. 2012). Furthermore, Mascher et al. (2013) implied and evaluated a whole exome capture platform for cultivated barley and demonstrated its applicability to genome-wide variation discovery in related *Hordeum* species and hexaploid wheat (*Triticum aestivum*) (Mascher et al. 2013). Similarly, Henry et al. (2014) used exome capture and next-generation sequencing of rice TILLING mutants for large-scale mutation discovery. They also extended their study to EMS-induced mutations in tetraploid wheat showing the method is applicable for polyploidy species.

Lately, another study was performed for wheat TILLING evaluation using next-generation sequencing exon capture for mutation scanning (King et al. 2015). The above studies show that mutation detection by TILLING can be a less laborious, more efficient and less expensive method by using exon capture for genome re-sequencing in polyploid wheat, barley and other plant species. However, an improved genomic reference with more complete coverage of homologues is required for accurate mutation calling.

Like other technologies, NGS has some limitations, such as the requirement for costly equipment and reagents and the limited number of identified tags for sample barcoding, despite the large number of nucleotides that can be sequenced in one run. Analysis of the results of NGS sequencing is also challenging because it is necessary to distinguish between false-positive and real SNPs (Kurowska et al. 2011) (Table 2). In the near future, the number of NGS applications will certainly grow, and the available ones will improve. NGS technologies are paving the way toward a new era of scientific discovery and have great potential applications in

plant-breeding programs for the development of superior cultivars of crops and ornamentals. As genome sequencing becomes easier, the additive demand of food will decrease in the coming decades for the next-generation plant breeders (Egan et al. 2012).

Conclusion and prospects

Mutagenesis has played a significant role in plant breeding. However, its effectiveness can be enhanced by using molecular technology and bioinformatics and assist plant breeders with useful induced mutations and develop mutation screening methods to identify novel alleles of target genes for crop improvement. Over the past 16 years, a variety of different techniques have been applied for detecting induced mutations in plant populations such as classical and recent reverse genetic techniques.

The TILLING-CEL I platform is widely applicable, but it involves steps such as endonuclease digestion reactions, cloning and gel electrophoresis runs, which are critical and also time-consuming. In addition, a lack of complete genome sequence information for many crop species has slowed the development of suitable TILLING targets. In contrast, TILLING-HRM is accurate, sensitive, fast and cost-effective, and the only step required is a simple PCR performed prior to the DNA melting curve analysis. However, the number and position of mutations within the amplicon cannot be identified using this method alone. Therefore, amplicon sequencing is needed to identify sequence variants.

With the tremendous improvement of the cost and accuracy of NGS technologies in the past decade, the potential of TILLING-NGS approach for mutation detection has increased.

A comparison of mutation discovery by Illumina sequencing and traditional TILLING was conducted by Gilchrist et al. (2013). The sensitivity level of two approaches was found to be the same in oilseed rape, but sequencing was more expensive. They suggested several ways to decrease the cost of sequencing for mutation detection by barcoding samples during PCR amplifications (instead of post-amplification), using automated platforms for high complexity of PCR amplification and target many regions in multiple lines and applying platforms that provide more sequencing data per lane.

Improvements in target genome sequencing (TGS), TILLING by sequencing, mapping by sequencing and

targeted modification of specific genes will facilitate the discovery of SNPs and indels using NGS technologies to further assist our understanding of plant genetics and genomics.

In the future, NGS-enabled discoveries will continue in the next decade. Bioinformatics and sequence data will simplify the detection of target genes with allelic diversity resulting in the development of agriculture traits to meet the demands of global food insecurity. However, analysing large amounts of generated data after the application of NGS technologies, together with the extraction of bio-information, remains a challenge that needs to be addressed in future research.

Acknowledgements The authors thank University Putra Malaysia (UPM) for supporting this work.

Authors' contributions All authors had substantial contributions to the conception, design and drafting of this work as individual experts in their fields. In particular, authors ST and TLA contributed to writing and organizing the contents of the manuscript and revising it critically. Author SMJ revised the manuscript critically. Authors MS and PA contributed to the writing, figure preparation and formatting of the manuscript.

References

- Abe A, Kosugi S, Yoshida K, Natsume S, Takagi H, Kanzaki H, Matsumura H, Yoshida K, Mitsuoka C, Tamiru M (2012) Genome sequencing reveals agronomically important loci in rice using MutMap. *Nat Biotechnol* 30:174–178
- Acevedo-Garcia J, Spencer D, Thieron H, Reinstädler A, Hammond-Kosack K, Phillips AL, Panstruga R (2016) mlo-based powdery mildew resistance in hexaploid bread wheat generated by a non-transgenic TILLING approach. *Plant Biotechnol J* 2016:1–12
- Allen AM, Barker GL, Berry ST, Coghill JA, Gwilliam R, Kirby S, Robinson P, Brenchley RC, D'Amore R, McKenzie N (2011) Transcript-specific, single-nucleotide polymorphism discovery and linkage analysis in hexaploid bread wheat (*Triticum aestivum* L.). *Plant Biotechnol J* 9:1086–1099
- Allen RS, Nakasugi K, Doran R, Millar TA, Waterhouse P (2013) Facile mutant identification via a single parental backcross method and application of whole genome sequencing based mapping pipelines. *Front Plant Sci* 4:362
- Anai T (2012) Potential of a mutant-based reverse genetic approach for functional genomics and molecular breeding in soybean. *Breed Sci* 61:462–467
- Ashelford K, Eriksson ME, Allen CM, D'Amore R, Johansson M, Gould P, Kay S, Millar AJ, Hall N, Hall A (2011) Full genome re-sequencing reveals a novel circadian clock mutation in *Arabidopsis*. *Genome Biol* 12:1
- Ashrafi H, Hill T, Stoffel K, Kozik A, Yao J, Chin-Wo SR, Van Deynze A (2012) De novo assembly of the pepper transcriptome (*Capsicum annuum*): a benchmark for in silico discovery of SNPs, SSRs and candidate genes. *BMC Genomics* 13:571
- Austin RS, Vidaurre D, Stamatiou G, Breit R, Provart NJ, Bonetta D, Zhang J, Fung P, Gong Y, Wang PW (2011) Next-generation mapping of *Arabidopsis* genes. *Plant J* 67:715–725
- Azam S, Thakur V, Ruperao P, Shah T, Balaji J, Amindala B, Farmer AD, Studholme DJ, May GD, Edwards D (2012) Coverage-based consensus calling (CbCC) of short sequence reads and comparison of CbCC results to identify SNPs in chickpea (*Cicer arietinum*; Fabaceae), a crop species without a reference genome. *Am J Bot* 99:186–192
- Azpiroz-Leehan R, Feldmann KA (1997) T-DNA insertion mutagenesis in *Arabidopsis*: going back and forth. *Trends Genet* 13:152–156
- Balcells L, Swinburne J, Coupland G (1991) Transposons as tools for the isolation of plant genes. *Trends Biotechnol* 9:31–37
- Baldet P, Bres C, Okabe Y, Mauxion J-P, Just D, Boumonville C, Ferrand C, Mori K, Ezura H, Rothan C (2013) Investigating the role of vitamin C in tomato through TILLING identification of ascorbate-deficient tomato mutants. *Plant Biotechnol* 30:309–314
- Barchi L, Lanteri S, Portis E, Acquadro A, Valè G, Toppino L, Rotino GL (2011) Identification of SNP and SSR markers in eggplant using RAD tag sequencing. *BMC Genomics* 12:304
- Barkley N, Wang M (2008) Application of TILLING and EcoTILLING as reverse genetic approaches to elucidate the function of genes in plants and animals. *Curr Genomics* 9: 212
- Bleecker AB, Kende H (2000) Ethylene: a gaseous signal molecule in plants. *Annu Rev Cell Dev Biol* 16:1–18
- Blumenstiel JP, Noll AC, Griffiths JA, Perera AG, Walton KN, Gilliland WD, Hawley RS, Staehling-Hampton K (2009) Identification of EMS-induced mutations in *Drosophila melanogaster* by whole-genome sequencing. *Genetics* 182: 25–32
- Borevitz JO, Liang D, Plouffe D, Chang H-S, Zhu T, Weigel D, Berry CC, Winzler E, Chory J (2003) Large-scale identification of single-feature polymorphisms in complex genomes. *Genome Res* 13:513–523
- Botticella E, Sestili F, Hernandez-Lopez A, Phillips A, Lafiandra D (2011) High resolution melting analysis for the detection of EMS induced mutations in wheat *SbeIIa* genes. *BMC Plant Biol* 11:156
- Bovina R, Talame V, Silvio S, Sanguineti MC, Trost P, Sparla F, Tuberosa R (2011) Starch metabolism mutants in barley: a TILLING approach. *Plant Genet Resour* 9:170–173
- Bovina R, Brunazzi A, Gasparini G, Sestili F, Palombieri S, Botticella E, Lafiandra D, Mantovani P, Massi A (2014) Development of a TILLING resource in durum wheat for reverse-and forward-genetic analyses. *Crop Pasture Sci* 65: 112–124
- Brockman HE, de Serres FJ, Ong T-M, DeMarini DM, Katz AJ, Griffiths AJ, Stafford RS (1984) Mutation tests in *Neurospora crassa*: a report of the US environmental protection agency gene-tox program. *Mutat Res-Rev Mutat* 133: 87–134
- Buermans H, Den Dunnen J (2014) Next generation sequencing technology: advances and applications. *BBA-Mol Basis Dis* 1842:1932–1941

- Bush SM, Krysan PJ (2010) iTILLING: a personalized approach to the identification of induced mutations in Arabidopsis. *Plant Physiol* 154:25–35
- Byers RL, Harker DB, Yourstone SM, Maughan PJ, Udall JA (2012) Development and mapping of SNP assays in allotetraploid cotton. *Theor Appl Genet* 124:1201–1214
- Byrne ME (2006) Shoot meristem function and leaf polarity: the role of class III HD-ZIP genes. *PLoS Genet* 2:e89
- Caldwell DG, McCallum N, Shaw P, Muehlbauer GJ, Marshall DF, Waugh R (2004) A structured mutant population for forward and reverse genetics in barley (*Hordeum vulgare* L.). *Plant J* 40:143–150
- Candela H, Hake S (2008) The art and design of genetic screens: maize. *Nature Rev Genet* 9:192–203
- Carén H, Holmstrand A, Sjöberg R-M, Martinsson T (2006) The two human homologues of yeast UFD2 ubiquitination factor, UBE4A and UBE4B, are located in common neuroblastoma deletion regions and are subject to mutations in tumours. *Eur J Cancer* 42:381–387
- Casella L, Greco R, Bruschi G, Wozniak B, Dreni L, Kater M, Cavigliolo S, Lupotto E, Piffanelli P (2013) TILLING in European rice: hunting mutations for crop improvement. *Crop Sci* 53:2550–2562
- Chagné D, Gasic K, Crowhurst RN, Han Y, Bassett HC, Bowatte DR, Lawrence TJ, Rikkerink EH, Gardiner SE, Korban SS (2008) Development of a set of SNP markers present in expressed genes of the apple. *Genomics* 92:353–358
- Chantreau M, Grec S, Gutierrez L, Dalmais M, Pineau C, Demailly H, Paysant-Leroux C, Tavernier R, Trouvé J-P, Chatterjee M (2013) PT-flax (phenotyping and TILLInG of flax): development of a flax (*Linum usitatissimum* L.) mutant population and TILLInG platform for forward and reverse genetics. *BMC Plant Biol* 13:1
- Chateigner-Boutin A-L, Small I (2007) A rapid high-throughput method for the detection and quantification of RNA editing based on high-resolution melting of amplicons. *Nucleic Acids Res* 35:e114
- Chen L, Huang L, Min D, Phillips A, Wang S, Madgwick PJ, Parry MA, Hu Y-G (2012) Development and characterization of a new TILLING population of common bread wheat (*Triticum aestivum* L.). *PLoS One* 7:e41570
- Chen L, Hao L, Parry MA, Phillips AL, Hu YG (2014a) Progress in TILLING as a tool for functional genomics and improvement of crops. *J Integr Plant Biol* 56:425–443
- Chen H, Xie W, He H, Yu H, Chen W, Li J, Yu R, Yao Y, Zhang W, He Y (2014b) A high-density SNP genotyping array for rice biology and molecular breeding. *Mol Plant* 7:541–553
- Cheng J, Salentijn EM, Huang B, Denneboom C, Qi W, Dechesne AC, Krens FA, Visser RG, Loo EN (2015) Detection of induced mutations in CaFAD2 genes by next-generation sequencing leading to the production of improved oil composition in *Crambe abyssinica*. *Plant Biotechnol J* 13:471–481
- Close TJ, Bhat PR, Lonardi S, Wu Y, Rostoks N, Ramsay L, Druka A, Stein N, Svensson JT, Wanamaker S (2009) Development and implementation of high-throughput SNP genotyping in barley. *BMC Genomics* 10:582
- Colasuonno P, Incerti O, Lozito ML, Simeone R, Gadaleta A, Blanco A (2016) DHPLC technology for high-throughput detection of mutations in a durum wheat TILLING population. *BMC Genet* 17:1
- Cooper JL, Till BJ, Laport RG, Darlow MC, Kleffner JM, Jamaï A, El-Mellouki T, Liu S, Ritchie R, Nielsen N (2008) TILLING to detect induced mutations in soybean. *BMC Plant Biol* 8:9
- Cortés AJ, Chavarro MC, Blair MW (2011) SNP marker diversity in common bean (*Phaseolus vulgaris* L.). *Theor Appl Genet* 123:827–845
- Cosart T, Beja-Pereira A, Chen S, Ng SB, Shendure J, Luikart G (2011) Exome-wide DNA capture and next generation sequencing in domestic and wild species. *BMC Genomics* 12:1
- Dalmaï M, Antelme S, Ho-Yue-Kuang S, Wang Y, Darracq O, d'Yvoire MB, Cézard L, Légée F, Blondet E, Oria N (2013) A TILLING platform for functional genomics in *Brachypodium distachyon*. *PLoS One* 8:e65503
- De-Kai W, Zong-Xiu S, Yue-Zhi T (2006) Application of TILLING in plant improvement. *Acta Genet Sin* 33:957–964
- Devran Z, Kahveci E, Özkaynak E, Studholme DJ, Tör M (2015) Development of molecular markers tightly linked to Pvr4 gene in pepper using next-generation sequencing. *Mol Breed* 35:101
- Distefano G, Caruso M, La Malfa S, Gentile A, Wu SB (2012) High resolution melting analysis is a more sensitive and effective alternative to gel-based platforms in analysis of SSR—an example in citrus. *PLoS One* 7:e44202
- Dong C, Dalton-Morgan J, Vincent K, Sharp P (2009a) A modified TILLING method for wheat breeding. *Plant Genome* 2:39–47
- Dong C, Vincent K, Sharp P (2009b) Simultaneous mutation detection of three homoeologous genes in wheat by high resolution melting analysis and mutation surveyor. *BMC Plant Biol* 9:1
- Donini P, Muleo R, Colao MC, Miano D, Cirilli M, Intrieri MC, Baldoni L, Rugini E (2009) Mutation scanning and genotyping by high-resolution DNA melting analysis in olive germplasm. *Genome* 52:252–260
- Dossett M, Bassil N, Finn C (2010) High resolution melting detects sequence polymorphism in *Rubus occidentalis* monomorphic microsatellite markers. In: XXVIII International Horticultural Congress on Science and Horticulture for People (IHC2010): International Symposium on 926:91–95
- Druml B, Cichna-Markl M (2014) High resolution melting (HRM) analysis of DNA—its role and potential in food analysis. *Food Chem* 158:245–254
- Eckardt NA (2007) Positive and negative feedback coordinate regulation of disease resistance gene expression. *Plant Cell* 19:2700–2702
- Egan AN, Schlueter J, Spooner DM (2012) Applications of next-generation sequencing in plant biology. *Am J Bot* 99:175–185
- Erali M, Wittwer CT (2010) High resolution melting analysis for gene scanning. *Methods* 50:250–261
- Erlach Y, Gordon A, Brand M, Hannon GJ, Mitra PP (2010) Compressed genotyping. *IEEE Trans Inf Theory* 56:706–723
- Esfeld K, Uauy C, Tadele Z (2013) Application of TILLING for orphan crop improvement. In: *Biotechnology of Neglected and Underutilized Crops*. Springer, pp 83–113
- Feiz L, Martin J, Giroux M (2009) Creation and functional analysis of new Puroindoline alleles in *Triticum aestivum*. *Theor Appl Genet* 118:247–257
- Fekih R, Takagi H, Tamiru M, Abe A, Natsume S, Yaegashi H, Sharma S, Sharma S, Kanzaki H, Matsumura H (2013)

- MutMap+: genetic mapping and mutant identification without crossing in rice. *PLoS One* 8:e68529
- Feltus FA, Wan J, Schulze SR, Estill JC, Jiang N, Paterson AH (2004) An SNP resource for rice genetics and breeding based on subspecies indica and japonica genome alignments. *Genome Res* 14:1812–1819
- Fu YB, Peterson GW (2012) Developing genomic resources in two Linum species via 454 pyrosequencing and genomic reduction. *Mol Ecol Resour* 12:492–500
- Gady AL, Hermans FW, Van de Wal MH, van Loo EN, Visser RG, Bachem CW (2009) Implementation of two high through-put techniques in a novel application: detecting point mutations in large EMS mutated plant populations. *Plant Methods* 5:1–14
- Ganal MW, Altmann T, Röder MS (2009) SNP identification in crop plants. *Curr Opin Plant Biol* 12:211–217
- Ganopoulos I, Argiriou A, Tsaftaris A (2011) Microsatellite high resolution melting (SSR-HRM) analysis for authenticity testing of protected designation of origin (PDO) sweet cherry products. *Food Control* 22:532–541
- Gauffer C, Lebaron C, Moretti A, Constant C, Moquet F, Bonnet G, Caranta C, Gallois JL (2016) A TILLING approach to generate broad-spectrum resistance to potyviruses in tomato is hampered by eIF4E gene redundancy. *Plant J* 85:717–729
- Gaur R, Azam S, Jeena G, Khan AW, Choudhary S, Jain M, Yadav G, Tyagi AK, Chattopadhyay D, Bhatia S (2012) High-throughput SNP discovery and genotyping for constructing a saturated linkage map of chickpea (*Cicer arietinum* L.). *DNA Res*: dss018
- Gilchrist EJ, Haughn GW (2005) TILLING without a plough: a new method with applications for reverse genetics. *Curr Opin Plant Biol* 8:211–215
- Gilchrist E, Haughn G (2010) Reverse genetics techniques: engineering loss and gain of gene function in plants. *Brief Funct Genomics* 9:103–110
- Gilchrist EJ, Sidebottom CH, Koh CS, MacInnes T, Sharpe AG, Haughn GW (2013) A mutant *Brassica napus* (canola) population for the identification of new genetic diversity via TILLING and next generation sequencing. *PLoS One* 8:e84303
- Golding B, Jeong H-J, Jo YD, Park S-W, Kang B-C (2010) Identification of capsicum species using SNP markers based on high resolution melting analysis. *Genome* 53:1029–1040
- Gottwald S, Bauer P, Komatsuda T, Lundqvist U, Stein N (2009) TILLING in the two-rowed barley cultivar 'Barke' reveals preferred sites of functional diversity in the gene HvHox1. *BMC Res Notes* 2:258
- Granier F, Lemaire A, Wang Y, LeBris P, Antelme S, Vogel J, Laudencia-Chingcuanco D, Sibout R (2015) Chemical and radiation mutagenesis: induction and detection by whole genome sequencing. In: Vogel JP (ed) *Genetics and genomics of Brachypodium*. Springer, pp 155–170
- Greene EA, Codomo CA, Taylor NE, Henikoff JG, Till BJ, Reynolds SH, Enns LC, Burtner C, Johnson JE, Odden AR (2003) Spectrum of chemically induced mutations from a large-scale reverse-genetic screen in Arabidopsis. *Genetics* 164:731–740
- Griggs D, Stafford-Smith M, Gaffney O, Rockström J, Öhman MC, Shyamsundar P, Steffen W, Glaser G, Kanie N, Noble I (2013) Policy: sustainable development goals for people and planet. *Nature* 495:305–307
- Guo Y-L, Todesco M, Hagmann J, Das S, Weigel D (2012) Independent FLC mutations as causes of flowering-time variation in *Arabidopsis thaliana* and *Capsella rubella*. *Genetics* 192:729–739
- Guo Y, Abernathy B, Zeng Y, Ozias-Akins P (2015) TILLING by sequencing to identify induced mutations in stress resistance genes of peanut (*Arachis hypogaea*). *BMC Genomics* 16:1
- Hamilton JP, Hansey CN, Whitty BR, Stoffel K, Massa AN, Van Deynze A, De Jong WS, Douches DS, Buell CR (2011) Single nucleotide polymorphism discovery in elite north American potato germplasm. *BMC Genomics* 12:1
- Han Y, Kang Y, Torres-Jerez I, Cheung F, Town CD, Zhao PX, Udvardi MK, Monteros MJ (2011) Genome-wide SNP discovery in tetraploid alfalfa using 454 sequencing and high resolution melting analysis. *BMC Genomics* 12:350
- van Harten AM (1998) Mutation breeding: theory and practical applications. Cambridge University Press
- Hartwig B, James GV, Konrad K, Schneeberger K, Turck F (2012) Fast isogenic mapping-by-sequencing of ethyl methanesulfonate-induced mutant bulks. *Plant Physiol* 160: 591–600
- He J, Zhao X, Laroche A, Lu Z-X, Liu H, Li Z (2014) Genotyping-by-sequencing (GBS), an ultimate marker-assisted selection (MAS) tool to accelerate plant breeding. *Front Plant Sci* 5: 484
- Henry IM, Nagalakshmi U, Lieberman MC, Ngo KJ, Krasileva KV, Vasquez-Gross H, Akhunova A, Akhunov E, Dubcovsky J, Tai TH (2014) Efficient genome-wide detection and cataloging of EMS-induced mutations using exome capture and next-generation sequencing. *Plant Cell* 26:1382–1397
- Herrmann MG, Durtschi JD, Bromley LK, Wittwer CT, Voelkerding KV (2006) Amplicon DNA melting analysis for mutation scanning and genotyping: cross-platform comparison of instruments and dyes. *Clin Chem* 52:494–503
- Herrmann MG, Durtschi JD, Wittwer CT, Voelkerding KV (2007) Expanded instrument comparison of amplicon DNA melting analysis for mutation scanning and genotyping. *Clin Chem* 53:1544–1548
- Highsmith WE, Jin Q, Nataraj AJ, O'Connor JM, Burland VD, Baubonis WR, Curtis FP, Kuskawa N, Garner MM (1999) Use of a DNA toolbox for the characterization of mutation scanning methods. I: construction of the toolbox and evaluation of heteroduplex analysis. *Electrophoresis* 20:1186–1194
- Himelblau E, Gilchrist EJ, Buono K, Bizzell C, Mentzer L, Vogelzang R, Osborn T, Amasino RM, Parkin IA, Haughn GW (2009) Forward and reverse genetics of rapid-cycling *Brassica oleracea*. *Theor Appl Genet* 118:953–961
- Hodges E, Xuan Z, Balija V, Kramer M, Molla MN, Smith SW, Middle CM, Rodesch MJ, Albert TJ, Hannon GJ (2007) Genome-wide in situ exon capture for selective resequencing. *Nat Genet* 39:1522–1527
- Hofinger BJ, Jing H-C, Hammond-Kosack KE, Kanyuka K (2009) High-resolution melting analysis of cDNA-derived PCR amplicons for rapid and cost-effective identification of novel alleles in barley. *Theor Appl Genet* 119:851–865
- Huq MA, Akter S, Nou IS, Kim HT, Jung YJ, Kang KK (2016) Identification of functional SNPs in genes and their effects on plant phenotypes. *J Plant Biotechnol* 43:1–11

- Hyten DL, Cannon SB, Song Q, Weeks N, Fickus EW, Shoemaker RC, Specht JE, Farmer AD, May GD, Cregan PB (2010) High-throughput SNP discovery through deep resequencing of a reduced representation library to anchor and orient scaffolds in the soybean whole genome sequence. *BMC Genomics* 11:1
- Irvine DV, Goto DB, Vaughn MW, Nakaseko Y, McCombie WR, Yanagida M, Martienssen R (2009) Mapping epigenetic mutations in fission yeast using whole-genome next-generation sequencing. *Genome Res* 19:1077–1083
- Jain SM (2010) Mutagenesis in crop improvement under the climate change. *ROM BIOTECH LETT* 15:88–106
- James GV, Patel V, Nordström KJ, Klasen JR, Salomé PA, Weigel D, Schneeberger K (2013) User guide for mapping-by-sequencing in *Arabidopsis*. *Genome Biol* 14:1
- Jander G, Norris SR, Rounsley SD, Bush DF, Levin IM, Last RL (2002) *Arabidopsis* map-based cloning in the post-genome era. *Plant Physiol* 129:440–450
- Jankowicz-Cieslak J, Till BJ (2015) Forward and reverse genetics in crop breeding. In: Al-Khayri JM, Jain SM, Johnson DV (eds) *Advances in plant breeding strategies: breeding, biotechnology and molecular tools*. Springer, pp 215–240
- Ji Q (2013) Gene identification: forward genetics. In: Lübberstedt T, Varshney RK (eds) *Diagnostics in plant breeding*. Springer, pp 41–60
- Joint FAO (2015) Plant breeding and genetics newsletter. Joint FAO/IAEA 35: July 2015
- Jones E, Chu W-C, Ayele M, Ho J, Bruggeman E, Yourstone K, Rafalski A, Smith OS, McMullen MD, Bezawada C (2009) Development of single nucleotide polymorphism (SNP) markers for use in commercial maize (*Zea mays* L.) germplasm. *Mol Breed* 24:165–176
- Jünemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U, Kalinowski J, Mellmann A, Goesmann A, von Haeseler A, Stoye J (2013) Updating benchtop sequencing performance comparison. *Nat Biotechnol* 31:294–296
- Kaya HB, Cetin O, Kaya H, Sahin M, Sefer F, Kahraman A, Tanyolac B (2013) SNP discovery by Illumina-based transcriptome sequencing of the olive and the genetic characterization of Turkish olive genotypes revealed by AFLP, SSR and SNP markers. *PLoS One* 8:e73674
- Kim S-I, Tai TH (2014) Identification of novel rice low phytic acid mutations via TILLING by sequencing. *Mol Breed* 34:1717–1729
- Kim Y, Schumaker KS, Zhu J-K (2006) EMS mutagenesis of *Arabidopsis*. In: Salinas J, Sanchez-Serrano JJ (eds) *Arabidopsis Protocols*, vol 2. Springer, pp 101–103
- King R, Bird N, Ramirez-Gonzalez R, Coghill JA, Patil A, Hassani-Pak K, Uauy C, Phillips AL (2015) Mutation scanning in wheat by exon capture and next-generation sequencing. *PLoS One* 10:e0137549
- Knoll JE, Ramos ML, Zeng Y, Holbrook CC, Chow M, Chen S, Maleki S, Bhattacharya A, Ozias-Akins P (2011) TILLING for allergen reduction and improvement of quality traits in peanut (*Arachis hypogaea* L.). *BMC Plant Biol* 11:81
- Kodym A, Afza R (2003) Physical and chemical mutagenesis. In: Grotewold E (ed) *Plant functional genomics*, vol 3. Springer, pp 189–203
- Koornneef M, Dellaert L, Van der Veen J (1982) EMS-and relation-induced mutation frequencies at individual loci in *Arabidopsis thaliana* (L.) Heynh. *Mutat Res-Fund Mol M* 93:109–123
- Kovacs E, Keresztes A (2002) Effect of gamma and UV-B/C radiation on plant cells. *Micron* 33:199–210
- Krieg DR (1963) Ethyl methanesulfonate-induced reversion of bacteriophage T4rII mutants. *Genetics* 48:561
- Krothapalli K, Buescher EM, Li X, Brown E, Chapple C, Dilkes BP, Tuinstra MR (2013) Forward genetics by genome sequencing reveals that rapid cyanide release deters insect herbivory of *Sorghum bicolor*. *Genetics* 195:309–318
- Kumar S, Banks TW, Cloutier S (2012) SNP discovery through next-generation sequencing and its applications. *Int J Plant Genomics* 2012:1–15
- Kurowska M, Daszkowska-Golec A, Gruszka D, Marzec M, Szurman M, Szarejko I, Maluszynski M (2011) TILLING—a shortcut in functional genomics. *J Appl Genet* 52:371–390
- Le Signor C, Savoie V, Aubert G, Verdier J, Nicolas M, Pagny G, Moussy F, Sanchez M, Baker D, Clarke J (2009) Optimizing TILLING populations for reverse genetics in *Medicago truncatula*. *Plant Biotechnol J* 7:430–441
- Lee LS, Till BJ, Hill H, Huynh OA, Jankowicz-Cieslak J (2014) Mutation and mutation screening. In: Henry RJ, Furtado A (eds) *Cereal genomics: methods and protocols*. Springer, pp 77–95
- Lee J, Izzah NK, Jayakodi M, Perumal S, Joh HJ, Lee HJ, Lee S-C, Park JY, Yang K-W, Nou I-S (2015) Genome-wide SNP identification and QTL mapping for black rot resistance in cabbage. *BMC Plant Biol* 15:32
- Lehmensiek A, Sutherland MW, McNamara RB (2008) The use of high resolution melting (HRM) to map single nucleotide polymorphism markers linked to a covered smut resistance gene in barley. *Theor Appl Genet* 117:721–728
- Li Q, Liu Z, Monroe H, Cui CT (2002) Integrated platform for detection of DNA sequence variants using capillary array electrophoresis. *Electrophoresis* 23:1499–1511
- Li YD, Chu ZZ, Liu XG, Jing HC, Liu YG, Hao DY (2010) A cost-effective high-resolution melting approach using the EvaGreen dye for DNA polymorphism detection and genotyping in plants. *J Integr Plant Biol* 52:1036–1042
- Liew M, Pryor R, Palais R, Meadows C, Erali M, Lyon E, Wittwer C (2004) Genotyping of single-nucleotide polymorphisms by high-resolution melting of small amplicons. *Clin Chem* 50:1156–1164
- Ling HQ, Zhao S, Liu D, Wang J, Sun H, Zhang C, Fan H, Li D, Dong L, Tao Y (2013) Draft genome of the wheat A-genome progenitor *Triticum urartu*. *Nature* 496:87–90
- Lister R, Gregory BD, Ecker JR (2009) Next is now: new technologies for sequencing of genomes, transcriptomes, and beyond. *Curr Opin Plant Biol* 12:107–118
- Liu K-h, McCormack M, Sheen J (2012) Targeted parallel sequencing of large genetically-defined genomic regions for identifying mutations in *Arabidopsis*. *Plant Methods* 8:1
- Lochlainn SÓ, Amoah S, Graham NS, Alamer K, Rios JJ, Kurup S, Stoute A, Hammond JP, Østergaard L, King GJ (2011) High resolution melt (HRM) analysis is an efficient tool to genotype EMS mutants in complex crop genomes. *Plant Methods* 7:1–9
- Mackay JF, Wright CD, Bonfiglioli RG (2008) A new approach to varietal identification in plants by microsatellite high resolution melting analysis: application to the verification of grapevine and olive cultivars. *Plant Methods* 4:8

- Mader E, Lukas B, Novak J (2008) A strategy to setup codominant microsatellite analysis for high-resolution-melting-curve-analysis (HRM). *BMC Genet* 9:1
- Martín B, Ramiro M, Martínez-Zapater JM, Alonso-Blanco C (2009) A high-density collection of EMS-induced mutations for TILLING in Landsberg erecta genetic background of *Arabidopsis*. *BMC Plant Biol* 9:1
- Mascher M, Richmond TA, Gerhardt DJ, Himmelbach A, Clissold L, Sampath D, Ayling S, Steuernagel B, Pfeifer M, D'Ascenzo M (2013) Barley whole exome capture: a tool for genomic research in the genus *Hordeum* and beyond. *Plant J* 76:494–505
- Mateo-Bonmati E, Casanova-Sáez R, Candela H, Micol JL (2014) Rapid identification of angulata leaf mutations using next-generation sequencing. *Planta* 240:1113–1122
- Mazzucato A, Cellini F, Bouzayen M, Zouine M, Mila I, Minoia S, Petrozza A, Picarella ME, Ruii F, Carriero F (2015) A TILLING allele of the tomato *Aux/IAA9* gene offers new insights into fruit set mechanisms and perspectives for breeding seedless tomatoes. *Mol Breed* 35:1–15
- Mba C (2013) Induced mutations unleash the potentials of plant genetic resources for food and agriculture. *Agronomy* 3:200–231
- Mba C, Afza R, Bado S, Jain SM (2010) Induced mutagenesis in plants using physical and chemical agents. *Plant Cell Culture: Essential Methods*: 111–130
- McCallum CM, Comai L, Greene EA, Henikoff S (2000a) Targeting induced local lesions in genomes (TILLING) for plant functional genomics. *Plant Physiol* 123:439–442
- McCallum CM, Comai L, Greene EA, Henikoff S (2000b) Targeted screening for induced mutations. *Nat Biotechnol* 18:455–457
- McNally KL, Childs KL, Bohnert R, Davidson RM, Zhao K, Ulat VJ, Zeller G, Clark RM, Hoen DR, Bureau TE (2009) Genomewide SNP variation reveals relationships among landraces and modern varieties of rice. *Proc Natl Acad Sci USA* 106:12273–12278
- Meacham F, Boffelli D, Dhahbi J, Martin DI, Singer M, Pachter L (2011) Identification and correction of systematic error in high-throughput sequence data. *BMC Bioinformatics* 12:451
- Minoia S, Petrozza A, D'Onofrio O, Piron F, Mosca G, Sozio G, Cellini F, Bendahmane A, Carriero F (2010) A new mutant genetic resource for tomato crop improvement by TILLING technology. *BMC Res Notes* 3:69
- Missirlian V, Comai L, Filkov V (2011) Statistical mutation calling from sequenced overlapping DNA pools in TILLING experiments. *BMC Bioinformatics* 12:1
- Muyzer G, Smalla K (1998) Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie Van Leeuwenhoek* 73:127–141
- Nelson JC, Wang S, Wu Y, Li X, Antony G, White FF, Yu J (2011) Single-nucleotide polymorphism discovery by high-throughput sequencing in sorghum. *BMC Genomics* 12:352
- Nida H, Blum S, Zielinski D, Srivastava DA, Elbaum R, Xin Z, Erlich Y, Fridman E, Shental N (2016) Highly efficient de novo mutant identification in a *Sorghum bicolor* TILLING population using the ComSeq approach. *Plant J* 86:349–359
- Nordström KJ, Albani MC, James GV, Gutjahr C, Hartwig B, Turck F, Paszkowski U, Coupland G, Schneeberger K (2013) Mutation identification by direct comparison of whole-genome sequencing data from mutant and wild-type individuals using k-mers. *Nat Biotechnol* 31:325–330
- Okabe Y, Asamizu E, Saito T, Matsukura C, Ariizumi T, Brès C, Rothan C, Mizoguchi T, Ezura H (2011) Tomato TILLING technology: development of a reverse genetics tool for the efficient isolation of mutants from micro-Tom mutant libraries. *Plant Cell Physiol* 52:1994–2005
- Okabe Y, Asamizu E, Ariizumi T, Shirasawa K, Tabata S, Ezura H (2012) Availability of micro-Tom mutant library combined with TILLING in molecular breeding of tomato fruit shelf-life. *Breed Sci* 62:202–208
- Oladosu Y, Rafii MY, Abdullah N, Hussin G, Ramli A, Rahim HA, Miah G, Usman M (2016) Principle and application of plant mutagenesis in crop improvement: a review. *Biotechnol Biotechnol Equip* 30:1–16
- Oleykowski CA, Mullins CRB, Godwin AK, Yeung AT (1998) Mutation detection using a novel plant endonuclease. *Nucleic Acids Res* 26:4597–4602
- Oliver RE, Lazo GR, Lutz JD, Rubenfield MJ, Tinker NA, Anderson JM, Morehead NHW, Adhikary D, Jellen EN, Maughan PJ (2011) Model SNP development for complex genomes based on hexaploid oat using high-throughput 454 sequencing technology. *BMC Genomics* 12:77
- Orita M, Iwahana H, Kanazawa H, Hayashi K, Sekiya T (1989) Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proc Natl Acad Sci USA* 86:2766–2770
- Ossowski S, Schneeberger K, Clark RM, Lanz C, Warthmann N, Weigel D (2008) Sequencing of natural strains of *Arabidopsis thaliana* with short reads. *Genome Res* 18:2024–2033
- Page DR, Grossniklaus U (2002) The art and design of genetic screens: *Arabidopsis thaliana*. *Nature Rev Genet* 3:124–136
- Parry MA, Madgwick PJ, Bayon C, Tearall K, Hernandez-Lopez A, Baudo M, Rakszegi M, Hamada W, Al-Yassin A, Ouabbou H (2009) Mutation discovery for crop improvement. *J Exp Bot* 60:2817–2825
- Pawelkowicz M, Zieliński K, Zielińska D, Pläder W, Yagi K, Wojcieszek M, Siedlecka E, Bartoszewski G, Skarzyńska A, Przybecki Z (2016) Next generation sequencing and omics in cucumber (*Cucumis sativus* L.) breeding directed research. *Plant Sci* 242:77–88
- Pegadaraju V, Nipper R, Hulke B, Qi L, Schultz Q (2013) De novo sequencing of sunflower genome for SNP discovery using RAD (restriction site associated DNA) approach. *BMC Genomics* 14:556
- Perry JA, Wang TL, Welham TJ, Gardner S, Pike JM, Yoshida S, Parniske M (2003) A TILLING reverse genetics tool and a web-accessible collection of mutants of the legume *Lotus Japonicus*. *Plant Physiol* 131:866–871
- Piron F, Nicolai M, Minoia S, Piednoir E, Moretti A, Salgues A, Zamir D, Caranta C, Bendahmane A (2010) An induced mutation in tomato eIF4E leads to immunity to two potyviruses. *PLoS One* 5:e11313
- Pomprasert S, Phusua A, Suanta S, Saetung R, Sanguanserm Sri T (2008) Detection of alpha-thalassemia-1 southeast Asian type using real-time gap-PCR with SYBR Green1 and high resolution melting analysis. *Eur J Haematol* 80:510–514
- Price EP, Smith H, Huygens F, Giffard PM (2007) High-resolution DNA melt curve analysis of the clustered, regularly

- interspaced short-palindromic-repeat locus of campylobacter jejuni. *Appl Environ Microbiol* 73:3431–3436
- Quail MA, Smith M, Coupland P, Otto TD, Harris SR, Connor TR, Bertoni A, Swerdlow HP, Gu Y (2012) A tale of three next generation sequencing platforms: comparison of ion torrent, Pacific biosciences and Illumina MiSeq sequencers. *BMC Genomics* 13:341
- Rawat N, Sehgal SK, Joshi A, Rothe N, Wilson DL, McGraw N, Vadlani PV, Li W, Gill BS (2012) A diploid wheat TILLING resource for wheat functional genomics. *BMC Plant Biol* 12:1
- Ray S, Satya P (2014) Next generation sequencing technologies for next generation plant breeding. *Front Plant Sci* 5:367
- Reddy TV, Dwivedi S, Sharma NK (2012) Development of TILLING by sequencing platform towards enhanced leaf yield in tobacco. *Ind Crop Prod* 40:324–335
- Reed GH, Wittwer CT (2004) Sensitivity and specificity of single-nucleotide polymorphism scanning by high-resolution melting analysis. *Clin Chem* 50:1748–1754
- Reed GH, Kent JO, Wittwer CT (2007) High-resolution DNA melting analysis for simple and efficient molecular diagnostics. *Pharmacogenomics* 8:597–608
- Rigola D, van Oeveren J, Janssen A, Bonné A, Schneiders H, van der Poel HJ, van Orsouw NJ, Hogers RC, de Both MT, van Eijk MJ (2009) High-throughput detection of induced mutations and natural variation using KeyPoint technology. *PLoS One* 4:E4761
- Rothe N (2010) Validation of tilling populations in diploid and hexaploid wheat. Dissertation, Kansas State University, Kansas
- Sabetta W, Alba V, Blanco A, Montemurro C (2011) sunTILL: a TILLING resource for gene function analysis in sunflower. *Plant Methods* 7:20
- Saintenac C, Jiang D, Akhunov ED (2011) Targeted analysis of nucleotide and copy number variation by exon capture in allotetraploid wheat genome. *Genome Biol* 12:1
- Sato Y, Sato K, Nishio T (2006) Interspecific pairs of class II S haplotypes having different recognition specificities between *Brassica oleracea* and *Brassica rapa*. *Plant Cell Physiol* 47:340–345
- Schneeberger K (2014) Using next-generation sequencing to isolate mutant genes from forward genetic screens. *Nature Rev Genet* 15:662–676
- Schneeberger K, Weigel D (2011) Fast-forward genetics enabled by new sequencing technologies. *Trends Plant Sci* 16:282–288
- Schneeberger K, Ossowski S, Lanz C, Juul T, Petersen AH, Nielsen KL, Jørgensen J-E, Weigel D, Andersen SU (2009) SHOREmap: simultaneous mapping and mutation identification by deep sequencing. *Nat Methods* 6:550–551
- Serrat X, Esteban R, Guibourt N, Moysset L, Nogués S, Lalanne E (2014) EMS mutagenesis in mature seed-derived rice calli as a new method for rapidly obtaining TILLING mutant populations. *Plant Methods* 10:5
- Sestili F, Botticella E, Bedo Z, Phillips A, Lafiandra D (2010) Production of novel allelic variation for genes involved in starch biosynthesis through mutagenesis. *Mol Breed* 25:145–154
- Sestili F, Palombieri S, Botticella E, Mantovani P, Bovina R, Lafiandra D (2015) TILLING mutants of durum wheat result in a high amylose phenotype and provide information on alternative splicing mechanisms. *Plant Sci* 233:127–133
- Sharp P, Dong C (2014) TILLING for plant breeding. In: Fleury D, Whitford R (eds) *Crop breeding: methods and protocols*. Springer, pp 155–165
- Shavrukov Y, Suchecki R, Eliby S, Abugalieva A, Kenebayev S, Langridge P (2014) Application of next-generation sequencing technology to study genetic diversity and identify unique SNP markers in bread wheat from Kazakhstan. *BMC Plant Biol* 14:258
- Shendure J, Ji H (2008) Next-generation DNA sequencing. *Nat Biotechnol* 26:1135–1145
- Shental N, Amir A, Zuk O (2010) Identification of rare alleles and their carriers using compressed sequencing. *Nucleic Acids Res* 38:e179–e179
- Sikora P, Chawade A, Larsson M, Olsson J, Olsson O (2012) Mutagenesis as a tool in plant genetics, functional genomics, and breeding. *Int J Plant Genomics* 2011:1–13
- Silletti MF, Petrozza A, Stigliani AL, Giorio G, Cellini F, D'Ambrosio C, Carriero F (2013) An increase of lycopene content in tomato fruit is associated with a novel Cyc-B allele isolated through TILLING technology. *Mol Breed* 31:665–674
- Simko I (2016) High-resolution DNA melting analysis in plant research. *Trends Plant Sci* 21:528–537
- Singh R, Tiwari R, Sharma D, Tiwari V, Sharma I (2014) Mutagenesis for wheat improvement in the genomics era. *J Wheat Res (JWR)* 6:120–125
- Slade AJ, Fuerstenberg SI, Loeffler D, Steine MN, Facciotti D (2005) A reverse genetic, nontransgenic approach to wheat crop improvement by TILLING. *Nat Biotechnol* 23:75–81
- Slade AJ, McGuire C, Loeffler D, Mullenberg J, Skinner W, Fazio G, Holm A, Brandt KM, Steine MN, Goodstal JF (2012) Development of high amylose wheat through TILLING. *BMC Plant Biol* 12:1
- Smith AD, Xuan Z, Zhang MQ (2008) Using quality scores and longer reads improves accuracy of Solexa read mapping. *BMC Bioinformatics* 9:1
- Sparla F, Falini G, Botticella E, Pirone C, Talamè V, Bovina R, Salvi S, Tuberosa R, Sestili F, Trost P (2014) New starch phenotypes produced by TILLING in barley. *PLoS One* 9:e107779
- Sparrow AH, Woodwell GM (1962) Prediction of the sensitivity of plants to chronic gamma irradiation. *Radiat Bot* 2:91N513IN721–512IN620IN1026
- Srivatsan A, Han Y, Peng J, Tehrani AK, Gibbs R, Wang JD, Chen R (2008) High-precision, whole-genome sequencing of laboratory strains facilitates genetic studies. *PLoS Genet* 4:e1000139
- Stadler LJ (1928) Genetic effects of X-rays in maize. *Proc Natl Acad Sci USA* 14:69–75
- Stemple DL (2004) TILLING—a high-throughput harvest for functional genomics. *Nat Rev Genet* 5:145–150
- Stephenson P, Baker D, Girin T, Perez A, Amoah S, King GJ, Østergaard L (2010) A rich TILLING resource for studying gene function in *Brassica rapa*. *BMC Plant Biol* 10:62
- van der Stoep N, van Paridon CD, Janssens T, Krenkova P, Stambergova A, Macek M, Matthijs G, Bakker E (2009) Diagnostic guidelines for high-resolution melting curve (HRM) analysis: an interlaboratory validation of BRCA1

- mutation scanning using the 96-well LightScanner™. *Hum Mutat* 30:899–909
- Suprasanna P, Mirajkar S, Bhagwat S (2015) Induced mutations and crop improvement. In: Bahadur B, Rajam MV, Sahijram L, Krishnamurthy KV (eds) *Plant biology and biotechnology*, vol 1. Springer, pp 593–617
- Taheri S, Abdullah TL, Ahmad Z, Abdullah NAP (2014) Effect of acute gamma irradiation on curcuma alismatifolia varieties and detection of DNA polymorphism through SSR marker. *Biomed Res Int* 2014:1–18
- Talamè V, Bovina R, Sanguineti MC, Tuberosa R, Lundqvist U, Salvi S (2008) TILLMore, a resource for the discovery of chemically induced mutants in barley. *Plant Biotechnol J* 6: 477–485
- Taylor NE, Greene EA (2003) PARSESNP: a tool for the analysis of nucleotide polymorphisms. *Nucleic Acids Res* 31:3808–3811
- Thole J, Strader L (2015) Next-generation sequencing as a tool to quickly identify causative EMS-generated mutations. *Plant Signal Behav* 10:1–4
- Thole JM, Beisner ER, Liu J, Venkova SV, Strader LC (2014) Absciscic acid regulates root elongation through the activities of auxin and ethylene in *Arabidopsis thaliana*. G3 (Bethesda) 4:1259–1274
- Thomson MJ (2014) High-throughput SNP genotyping to accelerate crop improvement. *Plant Breed Biotechnol* 2:195–212
- Till BJ, Reynolds SH, Greene EA, Codomo CA, Enns LC, Johnson JE, Burtner C, Odden AR, Young K, Taylor NE (2003a) Large-scale discovery of induced point mutations with high-throughput TILLING. *Genome Res* 13:524–530
- Till BJ, Colbert T, Tompa R, Enns LC, Codomo CA, Johnson JE, Reynolds SH, Henikoff JG, Greene EA, Steine MN (2003b) High-throughput TILLING for functional genomics. In: Grotewold E (ed) *Plant functional genomics*, vol 3. Springer, pp 205–220
- Till BJ, Reynolds SH, Weil C, Springer N, Burtner C, Young K, Bowers E, Codomo CA, Enns LC, Odden AR (2004a) Discovery of induced point mutations in maize genes by TILLING. *BMC Plant Biol* 4:12
- Till BJ, Burtner C, Comai L, Henikoff S (2004b) Mismatch cleavage by single-strand specific nucleases. *Nucleic Acids Res* 32:2632–2641
- Till BJ, Zerr T, Comai L, Henikoff S (2006) A protocol for TILLING and Ecotilling in plants and animals. *Nat Protoc* 1:2465–2477
- Till BJ, Cooper J, Tai TH, Colowit P, Greene EA, Henikoff S, Comai L (2007) Discovery of chemically induced mutations in rice by TILLING. *BMC Plant Biol* 7:19
- Tillib SV, Mirzabekov AD (2001) Advances in the analysis of DNA sequence variations using oligonucleotide microchip technology. *Curr Opin Biotechnol* 12:53–58
- Tindall EA, Petersen DC, Woodbridge P, Schipany K, Hayes VM (2009) Assessing high-resolution melt curve analysis for accurate detection of gene variants in complex DNA fragments. *Hum Mutat* 30:876–883
- Treangen TJ, Salzberg SL (2012) Repetitive DNA and next-generation sequencing: computational challenges and solutions. *Nat Rev Genet* 13:36–46
- Trebbi D, Maccaferri M, de Heer P, Sørensen A, Giuliani S, Salvi S, Sanguineti MC, Massi A, van der Vossen EAG, Tuberosa R (2011) High-throughput SNP discovery and genotyping in durum wheat (*Triticum durum* Desf.) *Theor Appl Genet* 123: 555–569
- Triques K, Sturbois B, Gallais S, Dalmais M, Chauvin S, Clepet C, Aubourg S, Rameau C, Caboche M, Bendahmane A (2007) Characterization of *Arabidopsis thaliana* mismatch specific endonucleases: application to mutation discovery by TILLING in pea. *Plant J* 51:1116–1125
- Tsai H, Howell T, Nitcher R, Missirian V, Watson B, Ngo KJ, Lieberman M, Fass J, Uauy C, Tran RK (2011) Discovery of rare mutations in populations: TILLING by sequencing. *Plant Physiol* 156:1257–1268
- Tsai H, Missirian V, Ngo KJ, Tran RK, Chan SR, Sundaresan V, Comai L (2013) Production of a high-efficiency TILLING population through polyploidization. *Plant Physiol* 161: 1604–1614
- Tsai H, Ngo K, Lieberman M, Missirian V, Comai L (2015) Tilling by sequencing. In: Alonso JM, Stepanova AN (eds) *Plant functional genomics: methods and protocols*, vol 3. Springer, pp 359–380
- Uauy C, Paraiso F, Colasuonno P, Tran RK, Tsai H, Berardi S, Comai L, Dubcovsky J (2009) A modified TILLING approach to detect induced mutations in tetraploid and hexaploid wheat. *BMC Plant Biol* 9:115
- Uchida N, Sakamoto T, Kurata T, Tasaka M (2011) Identification of EMS-induced causal mutations in a non-reference *Arabidopsis thaliana* accession by whole genome sequencing. *Plant Cell Physiol* 52:716–722
- Uchida N, Sakamoto T, Tasaka M, Kurata T (2014) Identification of EMS-induced causal mutations in *Arabidopsis thaliana* by next-generation sequencing. In: Sanchez-Serrano JJ, Salinas J (eds) *Arabidopsis Protocols*, vol 3. Springer, pp 259–270
- Underhill PA, Jin L, Lin AA, Mehdi SQ, Jenkins T, Vollrath D, Davis RW, Cavalli-Sforza LL, Oefner PJ (1997) Detection of numerous Y chromosome biallelic polymorphisms by denaturing high-performance liquid chromatography. *Genome Res* 7:996–1005
- Vicente-Dólera N, Troadec C, Moya M, del Río-Celestino M, Pomares-Viciano T, Bendahmane A, Picó B, Román B, Gómez P (2014) First TILLING platform in *Cucurbita pepo*: a new mutant resource for gene function and crop improvement. *PLoS One* 9:e112743
- Vidal RO, Nascimento LCD, Maurício Costa Mondego J, Amarante Guimarães Pereira G, Falsarella Carazzolle M (2012) Identification of SNPs in RNA-seq data of two cultivars of *Glycine max* (soybean) differing in drought resistance. *Genet Mol Biol* 35:331–334
- Wang TL, Robson F (2014) TILLING in *Lotus japonicus*. In: Tabata S, Stougaard J (eds) *The Lotus Japonicus genome*, vol IV. Springer, pp 229–243
- Wang N, Wang Y, Tian F, King GJ, Zhang C, Long Y, Shi L, Meng J (2008) A functional genomics resource for *Brassica napus*: development of an EMS mutagenized population and discovery of FAE1 point mutations by TILLING. *New Phytol* 180:751–765
- Wang TL, Uauy C, Robson F, Till B (2012) TILLING in extremis. *Plant Biotechnol J* 10:761–772
- Waugh R, Jannink J-L, Muehlbauer GJ, Ramsay L (2009) The emergence of whole genome association scans in barley. *Curr Opin Plant Biol* 12:218–222
- White HE, Hall VJ, Cross NC (2007) Methylation-sensitive high-resolution melting-curve analysis of the SNRPN gene as a

- diagnostic screen for Prader-Willi and Angelman syndromes. *Clin Chem* 53:1960–1962
- Winfield MO, Wilkinson PA, Allen AM, Barker GL, Coghill JA, Burridge A, Hall A, Brenchley RC, D'Amore R, Hall N (2012) Targeted re-sequencing of the allohexaploid wheat exome. *Plant Biotechnol J* 10:733–742
- Wittwer CT (2009) High-resolution DNA melting analysis: advancements and limitations. *Hum Mutat* 30:857–859
- Worm J, Aggerholm A, GuLDBERG P (2001) In-tube DNA methylation profiling by fluorescence melting curve analysis. *Clin Chem* 47:1183–1189
- Wu J-L, Wu C, Lei C, Baraoidan M, Bordeos A, Madamba M, Suzette R, Ramos-Pamplona M, Mauleon R, Portugal A (2005) Chemical-and irradiation-induced mutants of indica rice IR64 for forward and reverse genetics. *Plant Mol Biol* 59:85–97
- Wu S-B, Wirthensohn MG, Hunt P, Gibson JP, Sedgley M (2008) High resolution melting analysis of almond SNPs derived from ESTs. *Theor Appl Genet* 118:1–14
- Xiao W, Oefner PJ (2001) Denaturing high-performance liquid chromatography: a review. *Hum Mutat* 17:439–474
- Xin Z, Wang ML, Barkley NA, Burow G, Franks C, Pederson G, Burke J (2008) Applying genotyping (TILLING) and phenotyping analyses to elucidate gene function in a chemically induced sorghum mutant population. *BMC Plant Biol* 8:103
- Yamamoto T, Nagasaki H, J-i Y, Ebana K, Nakajima M, Shibaya T, Yano M (2010) Fine definition of the pedigree haplotypes of closely related rice cultivars by means of genome-wide discovery of single-nucleotide polymorphisms. *BMC Genomics* 11:1
- Yang H, Tao Y, Zheng Z, Li C, Sweetingham MW, Howieson JG (2012) Application of next-generation sequencing for rapid marker development in molecular plant breeding: a case study on anthracnose disease resistance in *Lupinus angustifolius* L. *BMC Genomics* 13:318
- Yang S, Fresnedo-Ramírez J, Wang M, Cote L, Schweitzer P, Barba P, Takacs EM, Clark M, Luby J, Manns DC (2016) A next-generation marker genotyping platform (AmpSeq) in heterozygous crops: a case study for marker-assisted selection in grapevine. *Hortic Res* 3:16002
- You FM, Huo N, Deal KR, Gu YQ, Luo M-C, McGuire PE, Dvorak J, Anderson OD (2011) Annotation-based genome-wide SNP discovery in the large and complex *Aegilops tauschii* genome using next-generation sequencing without a reference genome sequence. *BMC Genomics* 12:1
- Yu R, Shan X, Wang S, Li X, Jiang Y, Tan H, Li Y (2013) A screening method for detecting simple sequence repeat (SSR) polymorphism of *Zea mays* using high resolution melting-curve analysis. *Afr J Biotechnol* 10:16443–16447
- Zargar SM, Raatz B, Sonah H, Bhat JA, Dar ZA, Agrawal GK, Rakwal R (2015) Recent advances in molecular marker techniques: insight into QTL mapping, GWAS and genomic selection in plants. *J Crop Sci Biotechnol* 18:293–308
- Zhang X, Borevitz JO (2009) Global analysis of allele-specific expression in *Arabidopsis thaliana*. *Genetics* 182:943–954
- Zheng W, Wang Y, Wang L, Ma Z, Zhao J, Wang P, Zhang L, Liu Z, Lu X (2016) Genetic mapping and molecular marker development for Pi65 (t), a novel broad-spectrum resistance gene to rice blast using next-generation sequencing. *Theor Appl Genet* 129:1035–1044
- Zhou L, Vandersteen J, Wang L, Fuller T, Taylor M, Palais B, Wittwer C (2004) High-resolution DNA melting curve analysis to establish HLA genotypic identity. *Tissue Antigens* 64:156–164
- Zhou L, Wang L, Palais R, Pryor R, Wittwer CT (2005) High-resolution DNA melting analysis for simultaneous mutation scanning and genotyping in solution. *Clin Chem* 51:1770–1777
- Zuryn S, Le Gras S, Jamet K, Jarriault S (2010) A strategy for direct mapping and identification of mutations by whole-genome sequencing. *Genetics* 186:427–430