<https://helda.helsinki.fi>

Factors behind the variability of phosphorus accumulation in Finnish lakes

Tammeorg, Olga

2018-05

Tammeorg , O , Haldna , M , Noges , P , Appleby , P , Mols , T , Niemisto , J , Tammeorg , P & Horppila , J 2018 , ' Factors behind the variability of phosphorus accumulation in Finnish lakes ' , Journal of Soils and Sediments , vol. 18 , no. 5 , pp. 2117-2129 . <https://doi.org/10.1007/s11368-018-1973-8>

<http://hdl.handle.net/10138/307852>

<https://doi.org/10.1007/s11368-018-1973-8>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Factors behind the variability of phosphorus accumulation in Finnish lakes

**Olga Tammeorg^{1,2} • Marina Haldna² • Peeter Nõges² • Peter Appleby³ • Tõnu Möls² • Juha Niemistö¹ •
Priit Tammeorg⁴ • Jukka Horppila¹**

¹Department of Environmental Sciences, University of Helsinki, P.O. Box 65, 00014, Helsinki, Finland

²Centre for Limnology, Estonian University of Life Sciences, 61117 Rannu, Tartumaa, Estonia

³Department of Mathematical Sciences, University of Liverpool, L69 3BX, Liverpool, UK

⁴Department of Agricultural Sciences, University of Helsinki, P.O. Box 27, FI-00014, Helsinki, Finland

✉ Olga Tammeorg

olga.tammeorg@helsinki.fi

Abstract

Purpose. Phosphorus retention (TP_{acc}) is one of the major water quality regulators in lakes. The current study aimed at ascertaining the specific lake characteristics regulating TP_{acc} . Moreover, we were interested whether NAO (North Atlantic Oscillation), a proxy of climatic forcing, can explain variability in TP_{acc} , additionally to that ascribed to lake characteristics.

Materials and methods. Sediment cores were obtained from 21 Finnish lakes, subject to radiometric dating and measurements of TP concentrations. Principal components (PCs) were generated using lake characteristics that are usually included into the modelling of TP_{acc} (e.g., lake area, lake depth, catchment area, P inflow), but also the parameters that the classical models usually missed (e.g., anoxic factor). We used significant principal components (PCs), specific combinations of lake characteristics and monthly NAO values as predictors of TP_{acc} .

Results and discussion. Lake characteristics explained the bulk of TP_{acc} variability. The most influential factors (positive drivers) behind TP_{acc} included PC1 (representing mainly deep lakes), PC2 (small lakes with high levels of anoxia and water column stability), PC3 (productive lakes, with large catchment area and short water residence time), PC4 (lakes with high water column stability, low anoxic factor, and relatively high sediment focusing), and PC5 (lakes with high levels of P inflow, anoxia and long water residence time). Additionally, we found a potential negative effect of NAO in October on the annual TP_{acc} . This NAO was significantly positively related to temperatures in surface and near-bottom water layer (also their difference) in autumn, suggesting the possible implications for the internal P dynamics. Increased mineralization of organic matter is the most likely explanation for the reduced TP_{acc} associated with NAO driven water temperature increase.

Conclusions. The analysis presented here contributes to the knowledge of the factors controlling P retention. Moreover, this spatially and temporally comprehensive sediment data can potentially be a valuable source for modelling climate change implications.

Keywords Lake characteristics • Lakes • NAO • Phosphorus accumulation rate • Phosphorus retention

1 Introduction

Being one of the major regulators of the productivity in waterbodies, the phosphorus (P) retention has been in the focus of aquatic ecosystem modelling for about half of a century. In the mass balance models, the P retention is often determined as the difference between the inflowing and outflowing P. As an alternative that enables to eliminate the need for extensive monitoring programmes, the net P retention can be estimated by multiplying the net sediment accumulation rate with its P content (TP_{acc} ; Dillon and Evans 1993; Boers et al. 1998). There have been many attempts to predict P retention from a number of characteristics (e.g., Dillon and Kirchner 1975; Larsen and Mercier 1976; Vollenweider 1975), whereby the most common predictors of the P retention include phosphorus and hydraulic loading rate, TP particle settling velocity and mean depth. Nevertheless, the large prediction errors were found to be associated with those models, as these do not account for P release (Nürnberg, 1984). Sediments can serve as an important source of P in the years following reduction of external loading until the legacy P pool is reduced or buried in the deeper sediments (Sas 1990; Jeppesen et al. 2005; Søndergaard et al. 2013). Moreover, Benjamin and Brett (2008) demonstrated that the prevailing approach of conceptualization of the P retention overestimates the impact of the parameters usually used. The authors found that the best mass balance model tested could explain 84% of the variability in log-transformed lake TP concentrations, while it explained only 35% of the variability in TP retention and resulted in a large prediction error for individual lakes. The complex coupling of sediment composition, external load, catchment hydrology, lake morphometry, and biogeochemical reactions has been recognized to control P retention (Hupfer and Lewandowski 2008; Søndergaard et al. 2013; Huser et al. 2016). Hence, there is still a need for a P retention model that could better account for the lake specifics.

Climate change can affect P retention via variations in air temperature and precipitation that both influence P transport to the lakes (Jeppesen et al. 2009; Pettersson et al. 2010). Additionally, changes in temperature and wind have considerable implications for the vertical transport of P in lakes (Spears and Jones 2010; Tammeorg et al. 2013; Tammeorg et al. 2016; Woolway et al. 2017). Generally, climate change is associated with increased net P accumulation in lakes due to enhanced external nutrient loading leading also to increased internal P loading (Jeppesen et al. 2011). The North Atlantic Oscillation (NAO) index has performed as a good indicator of climatic forcing in European lakes. NAO index has shown to have a positive correlation with e.g. water temperatures, some lake water chemistry variables (Blenckner et al. 2007), wind speed (Vermaat et al. 2008), particularly in winter and spring, and wave-mixed depths (Spears and Jones 2010). Nevertheless, there is still a lack of knowledge on the relationship of NAO with P retention. As sediment records can reflect climatic variability (Bennion et al. 2006;

Rose et al. 2010; Sánchez-López et al. 2016) connecting TP_{acc} to climatic variation via NAO could be a useful tool to target that knowledge gap.

In the current study, we aimed at ascertaining the specific combinations of lake characteristics, principal components (PCs) that determine TP_{acc} , using data obtained from dated sediment cores collected from 21 Finnish lakes. Principal components (PCs) were generated using lake characteristics that are usually included into the modelling of TP retention (e.g., area and depth of lakes, size of catchment area and P inflow), but also the parameters missed by the classical models (e.g., anoxic factor, Osgood's index). Additionally, we coupled this information and the data on NAO for the time period covered by the sediment cores to elucidate the potential role of climatic variability in TP_{acc} . As climate change is primarily associated with changes in air temperature, which are closely coupled to water temperature, we were particularly interested whether potential NAO effects on TP_{acc} can be attributed to the temperature changes.

2 Methods

2.1 Study area

The 21 lakes of the study were all located in southern Finland, with their areas ranging from 0.25 to 155 km². The mean depth of the lakes varied from 1.3 to 21.0 m (Table 1), and the maximum depth from 3 to 68 m. Most of the lakes had deep areas, which undergo periodic anoxia, generally in winter and during thermal stratification in summer. The values of the anoxic factor (i.e., the product of the duration of anoxia and the percentage of the anaerobic areas) varied from 0 for the nonstratifying lakes (Nürnberg 1984) to 50 d y⁻¹ (Tammeorg et al. 2017). The monitoring data (Finnish Environment Institute) indicated the trophic status ranging from mesotrophic to hypertrophic (Table 1). Mean phosphorus inflow, TP_{in} varied from 104 mg m⁻² y⁻¹ in mesotrophic lakes to 910 mg m⁻² y⁻¹ in (hyper)eutrophic lakes (Tammeorg et al. 2017). The catchments of the eutrophic and hypertrophic lakes have mainly been impacted by agricultural activities (Ekholm and Mitikka 2006). All studied lakes were subject to a variety of restoration methods (including wastewater diversion, biomanipulation, artificial aeration) during past 30 years (Table 1).

99 **2.2 TP accumulation from dated sediment cores**

100 TP accumulation rate (TP_{acc} , $mg\ m^{-2}\ y^{-1}$) was calculated by multiplying the concentration of TP in the sediment
101 layer by the sedimentation rate. For that, sediment cores were collected with HTH gravity corer from the deepest
102 site of the lakes (the Kajaanselkä basin of Lake Vesijärvi was sampled at a site close to the maximally deep due to
103 technical reasons) targeting the accumulation areas (Håkanson and Jansson 1983) in March 2013 and 2014, when
104 the lakes were ice-covered. Sampling at locations that were predominantly stratified and anoxic during summer
105 ensured also minimal core disturbances due to wind activity (sediment resuspension), and bioturbation. Low water
106 temperatures during sampling lowered the risk of temperature-dependent transformations (e.g. P release) at the
107 sediment-water interface. Moreover, as it was identified by visual inspection, sediment surface was oxidized
108 inhibiting the release of P in the lakes studied. Dissolved oxygen concentration in the near-bottom water layer was
109 mainly above $7.0\ mg\ l^{-1}$. Each of the cores was sectioned into 0.5 cm slices to a depth of 20 cm to cover the period
110 for which also TP concentrations in the surface water layer and water temperature data were available, i.e. most
111 recent three decades (1986-2014). All sediment samples (40 samples per lake) were freeze-dried and ground. The
112 TP concentrations from the sediment subsamples were further determined using the methods by Koroleff (1979;
113 Lachat autoanalyzer, QuickChem Series 8000; Lachat instruments, Loveland, USA) after wet digestion with
114 sulphuric acid and hydrogen peroxide (Milestone Ethos 1600 microwave oven; Milestone, Sorisole, Italy).

115 Sedimentation rates were determined by dating cores (40 layers per core) by ^{210}Pb and ^{137}Cs . The analysis was
116 performed at the Liverpool University Environmental Radioactivity Laboratory. Sub-samples from each core were
117 analysed for ^{210}Pb , ^{226}Ra , and ^{137}Cs by direct gamma assay using Ortec HPGe GWL series well-type coaxial low
118 background intrinsic germanium detectors (Appleby et al. 1986). ^{210}Pb was determined via its gamma emissions
119 at 46.5 keV, and ^{226}Ra by the 295 keV and 352 keV γ -rays emitted by its daughter radionuclide ^{214}Pb following 3
120 weeks storage in sealed containers to allow radioactive equilibration. ^{137}Cs was measured by its emissions at 662
121 keV. The absolute efficiencies of the detectors were determined using calibrated sources and sediment samples of
122 known activity. Corrections were made for the effect of self-absorption of low energy γ -rays within the sample
123 (Appleby et al. 1992). ^{210}Pb dates were calculated mainly using CRS model (Appleby and Oldfield 1978). Since
124 in many cases the ^{210}Pb record spanned no more than around three decades, the calculation of reliable dates
125 demanded the use of well-defined ^{137}Cs dates as reference points. The method is described in detail by Appleby
126 (2001).

To quantify the possible impact of sediment focusing at the sampling area, a well-recognized issue (e.g., Eisenreich et al. 1989; Blais and Kalff 1993; Rowan et al. 1995; Lamborg et al. 2002; Heathcote and Downing 2014), focusing factors (FFs), calculated as the ratio of the measured mean ^{210}Pb supply rate (flux) at the core site to the atmospheric flux, were determined for each site. The atmospheric flux (based on fallout data from a number of European sites and records in 36 Finnish cores held in the Liverpool University ERRRC base) was estimated to be $100 \pm 20 \text{ Bq m}^{-2} \text{ y}^{-1}$. Although causes of high ^{210}Pb supply rates at specific sites can also include allochthonous inputs of fallout ^{210}Pb from the catchment (Appleby 2001), this was not thought to be a significant issue in our study, since our cores were collected from sites far from inlet streams at the deepest points of the lake.

Data on water temperature and surface water TP concentrations covering two-three most recent decades in the lakes studied were obtained from Hertta database (Finnish Env. Inst.). Besides air temperature, water temperatures can be affected by wind; thus, we analyzed also wind data. However, the more direct effects of the wind activity (through enhanced sediment resuspension at shallow areas and increased horizontal transport of these sediments to lake deeps, focusing) cannot be ignored. Data on wind speed for the Helsinki-Vantaa airport (8 measurements per day) were obtained from the Finnish Meteorological Institute. Daily NAO values were obtained from: <http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml>. Studied (hydro)meteorological variables were averaged over the months from January to December for the years 1970–2014. Data on TP concentration in the lake water of the surface layer was used to reflect the trophic state history of the lakes.

2.3 Statistical methods

Raw NAO, water temperature, and wind speed data statistics are shown with boxplot diagrams. The trends in rates of sediment accumulation, TP concentrations in the sediments and TP_{acc} , and TP concentration in the surface water layer over the years 1986–2014 for the studied lakes were tested with linear regression analysis.

To ascertain lake characteristics responsible for the TP_{acc} , the Principal Component Analysis (PCA) was carried out. Principal components (PCs) were obtained as weighted linear combinations of the original variables. Original variables included those lake characteristics that were demonstrated to be of paramount importance for controlling lake phosphorus dynamics, i.e. maximum depth (D_{max}), mean depth (D), ratio of D_{max} to D (to represent the potential importance of lateral sediment flux, i.e. sediment focusing), lake area (LA), catchment area (CA), ratio of the CA to LA (used as a proxy of water residence time), inflow of P (TP_{in}), anoxic factor (AF, to represent sediment P release due to anoxia), Osgood's index, or $D \times \text{LA}^{-0.5}$ (OI, to represent water column stability). D_{max}/D

correlated well with the focusing factor ($r = 0.600$, $p = 0.005$, based on the data of 20 sites, in which cores were sampled at the deepest lake location), supporting the use of the parameter to characterize sediment focusing in the lakes. Each characteristic was statistically standardised to have a zero mean and unit standard deviation in the set of all lakes. This approach generates principal components (PCs) as new complex (synthetic) uncorrelated factors that integrate individual characteristics. This approach is justified by the coexistence of different factors (lake characteristics) that correlate with each other. For example, significant positive correlation was found between the AF and $D_{\max}$ ($r = 0.541$, $p = 0.006$), TP_{in} and CA/LA ($r = 0.612$, $p = 0.002$), OI and AF ($r = 0.509$, $p = 0.011$). The effects of the PCs on the TP_{acc} were estimated by using the general multiparametrical linear model (SAS GLM procedure, type III). Initially, all nine PCs were used together as predictors of the TP_{acc} to ascertain *significant PCs*. After that, significant PCs were used singly as the predictors of the TP_{acc} . The significance was adjusted with the Bonferroni's correction. The approach of using PCs as independent variables has proven to be an effective tool in predicting internal P loading and water quality variables (Tammeorg et al. 2017). **Noteworthy, the effect (significant or not) of a PC on TP accumulation does not depend on the proportion of the lake variance that the specific PC describes.**

The correlations between NAO and lake water temperature in the surface and near-bottom water layer, their difference, and wind speed were presented with Pearson correlation coefficient. False discovery rate was applied to multiple testing ($Q=0.25$ was set as the proportion of the rejected null hypotheses which are erroneously rejected; Benjamini and Hochberg 1995). General multiparametrical linear model was used also to elucidate the effect of NAO and water temperature difference between the surface and near-bottom water layer on TP retention that remained after separating the lake specific effects encompassed under the significant PCs. The TP_{acc} values were log-transformed to make data distribution close to normal. Statistical analyses were done with SAS (version 9.2, SAS Institute Inc.).

3 Results

3.1. Variability across sediment cores, and lake water TP concentrations during two-three decades

The majority of the study lakes had well-defined peaks in the ^{137}Cs activity versus depth records that were confidently attributed to the fallout from the 1986 Chernobyl accident. The good resolution of the Chernobyl peaks suggests that sediment mixing has not been significant and that the ^{210}Pb and ^{137}Cs fallout records stored in the

sediments of these lakes, and the sediment accumulation rates (SARs) determined from those records (Tables S1 – S21), are reasonably reliable. Focusing factors (FFs) were less than two at twelve sites, greater than three at three sites, while intermediate values were found for the remaining six locations (Table 1). The mean FF for all 21 cores studied was about two. Six cores had long-term records spanning periods of time ranging from around 60 years (Kajaanselkä, Äimäjärvi) to more than 120 years (Hormajärvi, Punelia, Puujärvi). The remaining 15 had much shorter records, ranging from 46 years (Tuusulanjärvi) to as few as 19 years (Enonselkä). Mean post-1986 sedimentation rates varied widely from $0.021 \text{ g cm}^{-2} \text{ y}^{-1}$ (Lake Punelia) to $0.36 \text{ g cm}^{-2} \text{ y}^{-1}$ (Villikkalanjärvi), being generally higher in the lakes of higher trophic ($R^2 = 0.56, p < 0.0001$; Fig. 1). At twelve sites, the SAR was relatively constant over the last 30 years. Mean SARs at these sites varied by more than an order of magnitude, from $0.021 \text{ g cm}^{-2} \text{ y}^{-1}$ (Punelia) to $0.30 \text{ g cm}^{-2} \text{ y}^{-1}$ (Tiiläänjärvi). Since 1986, there were mainly systematic increases in the SAR at eight sites, and decrease in one site (Pusulanjärvi).

In the lakes studied, the mean post-1986 sediment TP concentrations (TP_{sed}) varied from 1.1 mg g^{-1} (Villikkalanjärvi) to 6.0 mg g^{-1} (Rehtijärvi). TP_{sed} increased significantly ($p < 0.01$) towards the surface of the core (the most recent years) in nine of the lakes studied that were mainly eutrophic (Table 2). There were increases in TP_{sed} in the lakes with increased SARs (Kynäröjärvi, Loppijärvi, Tuusulanjärvi), with constant SARs (Bodominjärvi, Punelia, Puujärvi, Sahajärvi, Karhujärvi), and in Pusulanjärvi that displayed a decrease in SAR. In Rehtijärvi ($R^2 = 0.380, p < 0.0001$), TP_{sed} decreased over the time period of 30 years. In overall, trends in SAR and TP_{sed} coincided at nine sites. Finally, TP_{acc} increased significantly ($p < 0.01$) in 13 of the lakes (Table 2), decreased ($p < 0.05$) in one lake (Rehtijärvi), while showing no changes in the rest of the lakes (mean for the years since 1986 varied from 340 in Punelia to $6038 \text{ mg m}^{-2} \text{ y}^{-1}$ in Pusulanjärvi; Fig. 2). At seven sites, trends observed in TP_{acc} coincided with those of the lake water TP concentration (TP), showing no change (Enäjärvi, Pusulanjärvi, and Pyhäjärvi(O)), and increases (Karhujärvi, Villikkalanjärvi, Loppijärvi, and Hormajärvi). At the rest six sites with increased TP_{acc} , TP either decreased (Kajaanselkä, basin of Lake Vesijärvi, Puujärvi, Tuusulanjärvi) or showed no changes (Bodominjärvi, Punelia, Sahajärvi) over the 30-year period.

3.2 Factors behind the variability in TP accumulation

3.2.1 Lake specifics

The first six and PC8 were found to have significant effect on the TP_{acc} , describing together 60% of the variability of TP_{acc} ($p < 0.0001$). Significant PCs represented about 98% of lake data variability in total (Table 3), whereby

the most of the lakes studied belonged to the types that were characterised by PC1 (39%), PC2 (27%), and PC3 (18%). The effect of PC1 – PC5 on TP_{acc} remained still significant, when these were used as predictors in the simple linear model (Table 4). The PC1 was mainly representative of the deep lakes (Table 3). In PC2, the highest loadings were by OI (0.516), LA (–0.458), and AF (0.420). By importance for PC3, CA (0.627) was followed by TP_{in} (0.504) and CA/LA (0.479). The OI (0.654), AF (–0.569), and D_{max}/D (0.402) were the major constituents of the PC4. The major contributing lake characteristics to PC5 included TP_{in} (0.620), followed by CA/LA (–0.363) and AF (0.357). In PC6, the highest loadings were by LA (0.612) and D_{max}/D (–0.570). The PC8 was primarily determined by CA (–0.607), CA/LA (0.564), and LA (0.516). PC6 and PC8 were not significant drivers of TP_{acc} ($R^2 = 0.025$, $p = 0.070$; $R^2 = 0.011$, $p = 0.679$). In general, TP_{acc} increased gradually with an increase in the values of the PC1 – PC 5 (Table 4; Fig. 3).

3.2.2 Climatic factors

During the years represented in sediment cores, long-term monthly NAO values varied from -1.024 to 1.092 on average (mean values close to zero), whereby somewhat lower values were observed in October (Fig. 4a). Daily mean wind speed was particularly variable during the winter months. Generally, the values decreased towards August, and increased since then (Fig. 4b). The water temperature difference between the surface and bottom layers increased towards July (Fig. 4c), when the highest temperatures reached 20.2 and 13.8 °C in surface water layer and near-bottom water layer, respectively. The temperatures decreased during the following months. As a result, temperature difference between the surface and the near bottom water layer was close to zero in September, October and November (mean values for the corresponding months were 1.2, 0.5 and -0.2 °C).

Monthly NAO correlated significantly with water temperature in the surface and near bottom layer, their difference, and wind speed throughout the year (Table 5). There were many, mainly positive significant correlations of the studied (hydro)meteorological variables with winter-spring NAO and considerably less, mainly negative correlations with the NAO in summer months. Further, a number of (mainly positive) correlations with the NAO increased again in the autumn months. Indeed, reported correlations had the highest significance level mainly in winter-spring, though correlations in winter were as high as in autumn.

Mean NAO value in October showed a potentially significant effect on the TP_{acc} , additional to those ascribed to the significant PCs, increasing predictive ability of the model ($R^2 = 0.613$, $p < 0.0001$). An increase in NAO index in October by one unit decreased TP_{acc} 1.2 times (19%; Fig. 5). Moreover, the mean NAO index in

October correlated significantly positively with the mean surface and bottom water temperatures and their difference in November ($r = 0.300$, $p < 0.01$; $r = 0.296$, $p < 0.01$; $r = 0.281$, $p < 0.01$, respectively; Table 4). The positive effect of the NAO in October on the temperature difference between surface and bottom water layer in November still remained significant (Fig. 5; $p < 0.01$) when the effects associated with the lake specifics were accounted for. At the same time, no significant correlations were found between NAO in October and average wind speed. Average temperature difference in November was -0.2°C (Fig. 4b), being lower in the surface layer than in the near bottom water layer.

4 Discussion

4.1 Variations in TP accumulation and its importance

Our results confirm the high importance of the lake trophic state in regulating rates of net sedimentation, one of the determinants of the P accumulation, reported earlier (e.g., Trolle et al. 2009), as these were considerably higher for eutrophic than for mesotrophic lakes. Thus, in case there are no external loading data with sufficient resolution available for a particular lake, changes in net sedimentation rate could shed light on its trophic state history. Our data showed that in most lakes both the sedimentation rates and water TP concentrations either increased or remained constant on the long-term scale, while a decrease in sediment accumulation rate was very rare among the studied lakes. These observations agree with the water quality monitoring data for twenty years in multiple agricultural Finnish lakes showing no improvement in the lake water quality (based on the chlorophyll *a* concentrations; Ekholm and Mitikka 2006).

Changes in trophic state during recent years can possibly explain an increase in sediment TP concentrations in the lakes that displayed also an increase in net sedimentation rate over the recent 30 years (e.g., Loppijärvi, Kynäjäjärvi). An increase in TP concentrations over the 30-year period (higher concentrations in the topmost sediments) in lakes with constant sedimentation rates (Bodominjärvi, Sahajärvi, Karhujärvi) is most likely due to diagenetic processes (Carignan and Flett 1981; Trolle et al. 2011). In general, we observed such patterns of sediment TP concentrations mainly in eutrophic lakes. Similarly, elevated concentrations in the surficial sediments representing a large pool of recyclable P were associated with lake eutrophic conditions shown by earlier studies (Carey and Rydin 2011). Moreover, it was not a surprise to observe such a phenomenon in mesotrophic lakes (Punelia and Puujärvi), as the sediments in these lakes can have limited P binding capacity (Carey and Rydin 2011;

Dittrich et al. 2013). Although an opposite vertical distribution of TP concentrations with higher levels in deeper sediments would be expected for oligotrophic lakes (due to Al availability; Carey and Rydin 2011), we found such TP distribution in one highly eutrophic lake (Rehtijärvi). It can be due to a combined effect of post-depositional migration and release of P into water column during periods of anoxia, similarly to what was concluded by Dillon and Evans (1993). This agrees with an increase of the lake water TP concentration in this lake on the long-term scale.

Increases in P could, indeed, result from sediment focusing. This could be of particular concern in cases of Pyhäjärvi (S), Ormajärvi, and the Enonselkä basin of Lake Vesijärvi that had FFs greater than three. However, none of those cores displayed an increase in TP_{acc} . Moreover, the mean FF for the study area indicates a rather modest level of bias associated with sediment focusing (Heathcote and Downing 2014). Moreover, the TP_{acc} found in our lakes of mesotrophic and higher trophic state were within the range of the values reported for other lakes of the northern temperate zone (summarized in Tammeorg et al. (2017)), being considerably higher than the values reported for oligotrophic lakes (Dillon and Evans 1993). This increase in TP_{acc} across the trophic gradient provides a support for the accuracy of our estimates. Therefore, the variations in TP_{acc} can be explained to considerable extent by differences in lake trophicity, which is closely coupled to lake morphology (Søndergaard et al. 2003; Hupfer and Lewandowski 2008).

The changes in lake water TP concentration similar to those in TP_{acc} were expected, confirming the high importance of sediments in P budget of lakes (Hupfer and Lewandowski 2008; Søndergaard et al. 2013). There were some lakes in which increased TP_{acc} appeared to sustain or augment eutrophication (e.g., Villikkalanjärvi, Loppijärvi, Karhujärvi). Unchanged lake TP concentration in lakes that showed an increase in TP_{acc} can be also due to possible time lags, a well-known phenomenon (Jeppesen et al. 2005). Moreover, restoration efforts could also have a role. In Tuusulanjärvi, in which increased TP_{acc} co-occurred with decreased lake water TP concentration, food web management applied since 1998 has compensated for the amplified P-cycling, revealed by the decreasing chlorophyll:total P ratio (Horppila et al. 2017).

4.2 Morphometric factors behind variations in TP accumulation

From the multiple external and internal factors controlling TP accumulation on the long-term scale, our model did not consider those that are related to sediment composition. Nevertheless, the simple model based on lake parameters that are usually available could explain a bulk of the variability in TP_{acc} . Similarly, there are numerous studies that have shown the association of P retention with morphometric/ hydraulic characteristics of lakes (e.g.,

Vollenweider 1975; Nürnberg 1984; Dillon and Molot 1996; Brett and Benjamin 2008; Kõiv et al. 2011). Generally, TP_{in} and hydraulic retention time are of key role in regulating TP retention in the classical models. Our model takes into account the additional characteristics that classical models lack, i.e. the anoxic factor reflecting P release and the Osgood's index characterising water column stability (Nürnberg 1984; Nürnberg 2009). Moreover, while P retention is conventionally calculated as a coefficient from mass balance equation (Brett and Benjamin 2008), we connected observed P accumulation rates with lake specific features. Interestingly, despite being quite different, our model gave nearly identical results (similar R^2) to classical ones. Similarly, Benjamin and Brett (2008) concluded that various multiple regression models yield very similar fits to those of Vollenweider type analyses because the terms typically considered in share many variables. While this makes the use of more simple models more preferable, we claim the approach used here to better take into account lake specifics via the use of PCs as independent factors. The relevance of the approach is supported, for example, by the finding of Kõiv et al. (2011) who showed that the retention is much more strongly determined by external P loading and hydrological residence time in large lakes than in smaller lakes.

Although the PCs represent a combination of different lake characteristics, they are somewhat predetermined by the values of some particular drivers (main contributors). Lake depth (D , D_{max}), the main contributor to the PC1, has been generally acknowledged as a factor that favours P accumulation (Håkanson and Jansson 1983), which agrees with the positive effect of PC1 on TP_{acc} in our study. However, the largest proportion of TP_{acc} variability was ascribed to changes in PC2, PC3, and PC5. The P accumulation appeared to be high in small lakes, with high water column stability, and high anoxic factor. The sediment P pool is often small in large lakes because resuspension leads to washout of particulate TP and organic net sedimentation is low, the latter due to high mineralization (Jeppesen et al. 2007). In small lakes, conditions are more favourable for stable stratification, and the relative importance of anaerobic areas is high. Both PC3 and PC5 characterize productive lakes, as TP_{in} is one of the major constituents of those components. In general, high TP_{in} results in the increased deposition of newly-produced P-rich material (Marsden 1989; Carey and Rydin 2011). The productivity is associated with large CA and high CA/LA values in the PC3, while with low CA/LA and relatively high AF in the PC5, suggesting differences in the relative importance of internal and external P loading in lakes. Lower CA/LA values are indicative of longer residence times and higher percentage of P load from internal sources with strong implications of the sediment P sources for productivity and water quality (Huser et al. 2016). High levels of external loading often result in oxygen deficits that sustain the recycling of P to the water column (Gächter and Wehrli 1998; Moosman et al. 2006) through the breakdown of the iron-phosphorus complexes (Einsele 1936;

Mortimer 1941, 1942). On the other hand, PC3 is likely to represent the lakes with productivity determined by external P sources. Allochthonous, mineral-bound particulate matter is more prone to settling, resulting in higher loss of P in lakes with shorter residence time (Brett and Benjamin 2008). Finally, TP accumulation tended to be high in lakes with high water column stability, but low anoxic factor, and relatively high $D_{\max}/D$ characterizing sediment focusing (PC4).

4.3 Climatic variability as a potential factor behind temporal changes in TP accumulation

One of the most interesting findings of the current study is that NAO in October influenced potentially the annual TP_{acc} , explaining its variability in addition to the significant PCs. Previously, the most pronounced implications for lake ecosystems were ascribed to the NAO values in winter-early spring (e.g., Bleckner et al. 2007; Pettersson et al. 2010; Spears and Jones 2010). Similarly, we found numerous significant correlations of NAO with wind speed and water temperatures at this period of time. While interpreting the results, it should be considered that NAO affects simultaneously air temperature, precipitation, wind speed and direction, cloudiness etc., each of them having potential feedbacks to lakes involving different lag periods. In the study region, the mechanisms behind NAO effects on the lakes functioning are mostly related with hydrology and ice regime, as milder temperatures cause more thaw days with increased runoff and shorter duration of ice cover (Nöges et al. 2010; Pettersson et al. 2010). Moisture transported from North-Atlantic causes more precipitation that acts in the same direction increasing the runoff (Hurrell and Van Loon 1997). Also in southern Finland, precipitation in winter was found to strongly associate with NAO (Irannezhad et al. 2014). Increased runoff mostly increases nutrient loading, if available in the catchment (Jeppesen et al. 2009; Trolle et al. 2011), entailing an increase in TP_{acc} . However, also opposite effect can be expected by flushing with meltwater, which is very likely to occur in the studied lakes with small area and depth. Nevertheless, our study revealed a potential importance of the autumnal NAO values for the water temperatures between the surface and near-bottom water layer, and their difference suggesting therefore possible mechanisms behind the changes in TP_{acc} .

The changes in the water temperature difference are most likely linked to NAO via air temperature. This suggestion is supported by the significant positive correlation between NAO and air temperatures in autumn reported for Finland (Irannezhad et al. 2015). Although there are no equivalent data reported for the wind, the results obtained for the nearby areas suggest its relevance as a possible explanatory mechanism from the perspective of both temperature difference and TP retention. Vermaat et al. (2008) showed that a positive NAO leads to increased wind-induced turbulence, and hence to higher resuspension of particle-bound nutrients.

Similarly, Spears and Jones (2010) showed that positive NAO correlated with stronger, more westerly winds, though correlations (including correlation between NAO and wave-mixed depth) were found only for winter and spring. Nevertheless, our data for Finnish lakes did not reveal any significant correlations of autumnal NAO with wind speed during September-November, suggesting the key role of air temperature in regulating autumnal water temperatures and a difference in temperature, which are likely to be linked to TP_{acc} .

Climate warming is generally associated with prolonged stratification in lakes leading to prolonged periods of anoxia, and release of P from sediments (Jeppesen et al. 2009). Similarly, Snorheim et al. (2017) found significant positive correlations between air temperature and anoxic factor for the northern dimictic Lake Mendota, additionally showing that the factor had the greatest potential impact for the stratification conditions (from the other studied factors, as wind speed and humidity). However, our water temperature data showed that stratification can be broken already since September, and water temperature difference in November (potentially linked to decreased TP retention) is negligible. Moreover, the prolonged algal blooms, associated with higher temperatures, are expected to result in higher supply of the organic matter and associated nutrients to the sediment during autumn (Blenckner et al. 2007; Jeppesen et al. 2009; Trolle et al. 2011). Therefore, this scenario suggests an importance of additional drivers to explain reduced TP_{acc} , e.g., flushing. Previously, a significant positive correlation was found between NAO and DOC discharge from the River Oulujoki in autumn (Marttila et al. 2014). However, the most likely mechanism seems to be associated with increased mineralization of the organic material in the sediments due to increased temperatures, as concluded by Gudasz et al. (2010). Over the boreal zone, the increase in organic carbon mineralization in sediments overlain by mixed water due to temperature increase (range of 1.8–4 °C) was predicted to result in a decrease of organic carbon burial of 6–15% in lake sediments (Gudasz et al. 2010). Providing that the sediment P is closely related to organic carbon (Håkanson and Jansson 1983), this would be the most likely explanation for the reduced TP sedimentation potentially associated with NAO driven water temperature increase.

5 Conclusions

The lake characteristics explained bulk of the variability in TP accumulation (TP_{acc}). TP_{acc} tends to be high in the lakes with following features: 1) mainly deep lakes; 2) small lakes with high levels of anoxia and water column stability; 3) lakes with high levels of P inflow, large catchment area and high CA/LA; 4) lakes with high water column stability, low anoxic factor, and relatively high sediment focusing; 5) lakes with high levels of P inflow,

anoxia and low CA/LA. Additionally to the effects attributed to lake specifics, we found a negative effect of NAO in autumn on annual TP_{acc} in Finnish lakes. The temperatures in surface and bottom water layer and their difference in autumn in the lakes studied were potentially related with NAO, suggesting the possible implications for P dynamics. Hence the analysis presented here for an internally consistent dataset (sampled in the same way) seems to better take account of lake specifics than previously. Moreover, this spatially and temporally comprehensive sediment data can potentially be a valuable source for modelling climate change implications.

Acknowledgments. The research was supported by the Academy of Finland (grant nr. 263365), Maa- ja vesitekniikan tuki ry (grant nr. 29159), and the City of Lahti. We thank Heidi Holmroos, Lauri Happonen, Jani Ruohola, and Raija Mastonen for their help in the field and in the lab. J. Weckström is acknowledged for useful discussions.

Funding: This study was funded by the Academy of Finland (grant nr. 263365), Maa- ja vesitekniikan tuki ry (grant nr. 29159), and the City of Lahti.

Conflict of Interest: The authors declare that they have no conflict of interest.

References

- Appleby PG, Richardson N, Nolan PJ (1992) Self-absorption corrections for well-type germanium detectors. *Nucl Instrum Meth B* 71(2):228–233
- Appleby PG, 2001. Chronostratigraphic techniques in recent sediments, in *Tracking Environmental Change Using Lake Sediments Volume 1: Basin Analysis, Coring, and Chronological Techniques*, (eds W M Last & J P Smol), Kluwer Academic, pp 171–203
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Royal Stat Society. Series B (Methodological)* 57(1):289–300
- Bennion H, Carvalho L, Sayer CD, Simpson GL, Wischniewski J (2012) Identifying from recent sediment records the effects of nutrients and climate on diatom dynamics in Loch Leven. *Freshw Biol* 57(10):2015–2029
- Blais JM, Kalff J (1993) Atmospheric loading of Zn, Cu, Ni, Cr, and Pb to lake sediments: The role of catchment, lake morphometry, and physio-chemical properties of the elements. *Biogeochem* 23(1):1–22

420 Blenckner T, Adrian R, Livingstone DM, [Jennings E](#), [Weyhenmeyer GA](#), [George D](#), [Jankowski T](#), [Järvinen M](#),
 421 [Aonghusa CN](#), [Nöges T](#), [Straile D](#) (2007) Large- scale climatic signatures in lakes across Europe: A meta-
 422 analysis. *Global Change Biol* 13(7):1314–1326
 423 Boers PCM, Van Raaphorst W, Van der Molen DT (1998) Phosphorus retention in sediments. *Water Sci*
 424 *Technol* 37(3):31–39
 425 Brett MT, Benjamin MM (2008) A review and reassessment of lake phosphorus retention and the nutrient loading
 426 concept. *Freshw Biol* 53(1):194–211
 427 Carey CC, Rydin E (2011) Lake trophic status can be determined by the depth distribution of sediment
 428 phosphorus. *Limnol Oceanogr* 56(6):2051–2063
 429 Dillon PJ, Evans HE (1993) A comparison of phosphorus retention in lakes determined from mass balance and
 430 sediment core calculations. *Water Res* 27(4):659–668
 431 Dillon PJ, Kirchner WB (1975) Reply to Chapra's Comment. *Water Res Res* 11(6):1035–1036.
 432 Dillon PJ, Molot LA (1996) Long-term phosphorus budgets and an examination of a steady-state mass balance
 433 model for central Ontario lakes. *Water Res* 30(10):2273–2280
 434 Dittrich M, Chesnyuk A, Gudimov A, [McCulloch J](#), [Quazi S](#), [Young J](#), [Winter J](#), [Stainsby E](#), [Arhonditsis G](#) (2013)
 435 Phosphorus retention in a mesotrophic lake under transient loading conditions: Insights from a sediment
 436 phosphorus binding form study. *Water Res* 47(3):1433–1447
 437 Eisenreich SJ, Capel PD, Robbins JA, Bourbonniere R (1989) Accumulation and diagenesis of chlorinated
 438 hydrocarbons in lacustrine sediments. *Environ Sci Technol* 23(9):1989–1126
 439 Ekholm P, Mitikka S (2006) Agricultural lakes in Finland: current water quality and trends. *Environ Monitor*
 440 *Assess* 116(1):111–135
 441 Golterman HL (2001) Phosphate release from anoxic sediments or 'What did Mortimer really
 442 write?' *Hydrobiologia* 450(1):99–106
 443 Gudaszc, Bastviken D, Steger K, Premke K, Sobek S, Tranvik LJ (2010) Temperature-controlled organic carbon
 444 mineralization in lake sediments. *Nature* 466(7305):478–481
 445 Gächter R, Wehrli B (1998) Ten years of artificial mixing and oxygenation: no effect on the internal phosphorus
 446 loading of two eutrophic lakes. *Environ Sci Technol* 32(23):3659–3665
 447 Heathcote AJ, Downing JA (2012) Impacts of eutrophication on carbon burial in freshwater lakes in an intensively
 448 agricultural landscape. *Ecosystems* 15(1):60–70

449 Horppila J, Holmroos H, Niemistö J, [Massa I](#), [Nygrén N](#), [Schönach P](#), [Tapio P](#), [Tammeorg O](#) (2017) Variations of
 450 internal phosphorus loading and water quality in a hypertrophic lake during 40 years of different management
 451 efforts. *Ecol Engineer* 103:264–274
 452 Hupfer M, Lewandowski J (2008) Oxygen Controls the Phosphorus Release from Lake Sediments—a Long-
 453 Lasting Paradigm in Limnology. *International Review of Hydrobiology* 93(4–5):415–432
 454 Huser BJ, Egemose S, Harper H, [Hupfer M](#), [Jensen H](#), [Pilgrim KM](#), [Reitzel K](#), [Rydin E](#), [Futter M](#). (2016)
 455 Longevity and effectiveness of aluminum addition to reduce sediment phosphorus release and restore lake
 456 water quality. *Water Res* 97:122–132
 457 Håkanson L, Jansson M (1983) *Principles of Lake Sedimentology*. Springer–Verlag, Berlin
 458 Irannezhad M, Marttila H, Kløve B (2014) Long- term variations and trends in precipitation in Finland. *Int J*
 459 *Climatol* 34(10):3139–3153
 460 Irannezhad M, Chen D, Kløve B (2015) Interannual variations and trends in surface air temperature in Finland in
 461 relation to atmospheric circulation patterns, 1961–2011. *Int J Climatol* 35(10):3078–3092
 462 Jeppesen E, Kronvang B, Meerhoff M, [Søndergaard M](#), [Hansen KM](#), [Andersen HE](#), [Lauridsen TL](#), [Liboriusen L](#),
 463 [Beklioglu M](#), [Özen A](#), [Olesen JE](#) (2009) Climate change effects on runoff, catchment phosphorus loading
 464 and lake ecological state, and potential adaptations. *J Environ Qual* 38(5):1930–1941
 465 Jeppesen E, Meerhoff M, Jacobsen BA, [Hansen RS](#), [Søndergaard M](#), [Jensen JP](#), [Lauridsen TL](#), [Mazzeo N](#), [Branco](#)
 466 [CW](#) (2007) Restoration of shallow lakes by nutrient control and biomanipulation—the successful strategy
 467 varies with lake size and climate. *Hydrobiologia* 581(1):269–285
 468 Jeppesen E, Søndergaard M, Jensen JP, [Havens KE](#), [Anneville O](#), [Carvalho L](#), [Coveney MF](#), [Deneke R](#), [Dokulil](#)
 469 [MT](#), [Foy BO](#), [Gerdeaux D](#) (2005) Lake responses to reduced nutrient loading—an analysis of contemporary
 470 long- term data from 35 case studies. *Freshw Biol* 50(10):1747–1771
 471 Katsev S, Tsandev I, L'Heureux I, Rancourt DG (2006) Factors controlling long-term phosphorus efflux from lake
 472 sediments: Exploratory reactive-transport modeling. *Chem Geol* 234(1):127–147
 473 Koroleff F (1979) Methods for the chemical analysis for seawater. (In Finnish). *Meri* 7:1–60
 474 Kõiv T, Nõges T, Laas A (2011) Phosphorus retention as a function of external loading, hydraulic turnover time,
 475 area and relative depth in 54 lakes and reservoirs. *Hydrobiologia* 660(1):105–115
 476 Lamborg CH., Fitzgerald WF, Duman AWH, Benoit JM, Balcom PH, Engstrom DR (2002) Modern and historic
 477 atmospheric mercury fluxes in both hemispheres: Global and regional mercury cycling implications. *Glob*
 478 *Biogeochem Cyc* 16(4):1104–1115

479 Larsen DP, Mercier HT (1976) Phosphorus retention capacity of lakes. *J Fish Res Board Can* 33: 1742–1750.
 480 Marttila H, Irannezhad M, Saukkoripi J, Kløve B (2015) Atmospheric circulation patterns influencing variations
 481 in organic carbon fluxes in the River Oulujoki, Finland. *Water Environ J* 29(4):474–481
 482 Moosmann L, Gächter R, Müller B, Wüest, A. (2006). Is phosphorus retention in autochthonous lake sediments
 483 controlled by oxygen or phosphorus? *Limnol. Oceanogr* 51(1):763–771
 484 Moss B, Kosten S, Meerhof M, Battarbee RW, Jeppesen E, Mazzeo N, Havens K, Lacerot G, Liu Z, De Meester
 485 L, Paerl H (2011) Allied attack: climate change and eutrophication. *Inland waters* 1(2):101–105
 486 Noges P, Noges T, Laas A (2010). Climate-related changes of phytoplankton seasonality in large shallow Lake
 487 Võrtsjärv, Estonia. *Aquat Ecosystem Health Manage* 13(2):154–163
 488 Nürnberg GK (1984) The prediction of internal phosphorus load in lakes with anoxic hypolimnia. *Limnol*
 489 *Oceanogr* 29(1):111–124
 490 Nürnberg GK (2009) Assessing internal phosphorus load—problems to be solved. *Lake Reservoir*
 491 *Manage* 25(4):419–432
 492 Pettersson K, George G, Nöges P, Nöges T, Blenckner T (2010) The impact of the changing climate on the supply
 493 and re-cycling of phosphorus. In: *The impact of climate change on European lakes*, Springer, Netherlands,
 494 pp. 121–137
 495 Rose NL, Morley D, Appleby PG, Battarbee RW, Alliksaar T, Guilizzoni P, Jeppesen E, Korhola A, Punning JM
 496 (2011) Sediment accumulation rates in European lakes since AD 1850: trends, reference conditions and
 497 exceedence. *J Paleolimnol* 45(4):447–468
 498 Rowan DJ, Cornett RJ, King K, Risto B (1995) Sediment focusing and ²¹⁰Pb dating: a new approach. *J Paleolimnol*
 499 13:107–118
 500 Sánchez-López G, Hernández A, Pla-Rabes S, Trigo RM, Toro M, Granados I, Sáez A, Masqué P, Pueyo JJ,
 501 Rubio-Inglés MJ, Giral S (2016) Climate reconstruction for the last two millennia in central Iberia: The role
 502 of East Atlantic (EA), North Atlantic Oscillation (NAO) and their interplay over the Iberian
 503 Peninsula. *Quatern Sci Rev* 149:135–150
 504 Snorheim CA, Hanson PC, McMahon KD, Read JS, Carey CC, Dugan HA (2017) Meteorological drivers of
 505 hypolimnetic anoxia in a eutrophic, north temperate lake. *Ecol Model* 343:39–53
 506 Søndergaard M, Jensen JP, Jeppesen E (2003) Role of sediment and internal loading of phosphorus in shallow
 507 lakes. *Hydrobiologia* 506(1):135–145

508 Søndergaard M, Bjerring R, Jeppesen E (2013) Persistent internal phosphorus loading during summer in shallow
 509 eutrophic lakes. *Hydrobiologia* 710(1):95–107
 510 Spears BM, Jones ID (2010) The long-term (1979–2005) effects of the North Atlantic Oscillation on wind-induced
 511 wave mixing in Loch Leven (Scotland). *Hydrobiologia* 646(1):49–59
 512 Tammeorg O, Niemistö J, Möls T, Laugaste R, Panksep K, Kangur K (2013) Wind-induced sediment resuspension
 513 as a potential factor sustaining eutrophication in large and shallow Lake Peipsi. *Aquat Sci* 75(4):559–570
 514 Tammeorg O, Möls T, Niemistö J, Holmroos H, Horppila J (2017) The actual role of oxygen deficit in the linkage
 515 of the water quality and benthic phosphorus release: Potential implications for lake restoration. *Sci Tot*
 516 *Environ* 599:732–738
 517 Trolle D, Hamilton DP, Pilditch CA, Duggan IC, Jeppesen E (2011) Predicting the effects of climate change on
 518 trophic status of three morphologically varying lakes: Implications for lake restoration and
 519 management. *Environ Model Software* 26(4):354–370
 520 Trolle D, Hamilton DP, Hendy C, Pilditch C (2009) Sediment and nutrient accumulation rates in sediments of
 521 twelve New Zealand lakes: influence of lake morphology, catchment characteristics and trophic state. *Marine*
 522 *Freshw Res* 59(12):1067–1078
 523 Vermaat JE, McQuatters-Gollop A, Eleveld MA, Gilbert AJ (2008) Past, present and future nutrient loads of the
 524 North Sea: causes and consequences. *Estuarine, Coastal Shelf Sci* 80(1):53–59
 525 Vollenweider RA (1975) Input-output models with special reference to the phosphorus loading concept in
 526 limnology. *Schweizerische Zeitschrift für Hydrologie* 37(1):53–84
 527 Woolway RI, Meinson P, Nöges P, Jones ID, Laas A (2017) Atmospheric stilling leads to prolonged thermal
 528 stratification in a large shallow polymictic lake. *Climatic Change* 141(4):759–773
 529 Sas H (1990) Lake restoration by reduction of nutrient loading: Expectations, experiences, extrapolations. *Verh*
 530 *Internat Verein Theor Angew Limnol* 24(1):247–251
 531 Tammeorg O, Horppila J, Tammeorg P, Haldna M, Niemistö J (2016) Internal phosphorus loading across a cascade
 532 of three eutrophic basins: A synthesis of short- and long-term studies. *Sci Tot Environ* 572:943–954
 533 Appleby PG, Nolan PJ, Gifford DW, Godfrey MJ, Oldfield FJAN, Anderson NJ, Battarbee RW (1986) ²¹⁰Pb dating
 534 by low background gamma counting. *Hydrobiologia* 143(1):21–27
 535 Carignan R, Flett RJ (1981) Postdepositional mobility of phosphorus in lake sediments. *Limnol Oceanogr* 26(2):
 536 361–366

537 Marsden MW (1989) Lake restoration by reducing external phosphorus loading: the influence of sediment
 538 phosphorus release. *Freshw Biology* 21(2):139–162

539 Einsele W (1936) Über die Beziehungen des Eisenkreislaufs zum Phosphatkreislauf im eutrophen See. *Arch*
 540 *Hydrobiol* 29(6):664–686

541 Mortimer CH (1941) The exchange of dissolved substances between mud and water in lakes. *J Ecol* 29(2):280–
 542 329

543 Mortimer CH (1942) The exchange of dissolved substances between mud and water in lakes. *J Ecol* 30(1):147–
 544 201

545 Hurrell JW, Van Loon H (1997) Decadal Variations in Climate Associated with the North Atlantic Oscillation. In:
 546 Diaz HF, Beniston M, Bradley RS (eds) *Climatic Change at High Elevation Sites*. Springer, Dordrecht

547

Table 1. Basic lake characteristics, including lake trophic status (TI), mean and maximum depth (D, D_{max}), catchment area (CA), lake area (LA), anoxic factor (AF) values, mean phosphorus inflow (TP_{in}), age of the 20-cm sediment cores sampled in 2014, focusing factor (FF), and restoration activities of the study area. For the AF and TP_{in} mean values for the period 1986–2014 are presented.

Lake	Coordinates	TI	D (m)	D _{max} (m)	CA (km ²)	LA (km ²)	AF d y ⁻¹	TP _{in} (mg P m ⁻² y ⁻¹)	Age y	FF	Restoration measures applied
Äimäjärvi	61°03'N 24°10'E	eutr	2.9	9	93	8.5	22.1	228	65	0.7	Wastewater diversion in 1969, biomanipulation since 1997, sedimentation ponds established in 2004
Bodominjärvi	60°15'N 24°40'E	eutr	4.3	12.7	32	4.1	7.2	507	31	2.5	Chemical treatment, in 1980 oxygen-depleted water diversion, aeration from 1970s to 1998
Enäjärvi	60°20'N 24°22'E	eutr	3.2	9.1	34	4.9	1.8	323	89	0.8	Wastewater diversion in 1976, fish removal from 1993, aerated from 1998
Enonselkä	61°00'N 16°36'E	eutr	6.8	33	84	26	24.7	137	19	3.9	1976-1978 sewage diversion, aeration (since 2009), biomanipulation (1989-1993), sedimentation ponds and wetlands
Hormajärvi	60°17'N 24°01'E	meso	7.3	21	16	5.1	36.1	88	121	1.8	Aeration since 2008
Kajaanselkä	61°09'N 25°28'E	meso	6.8	42	138	44	1.2	124	64	1.9	-
Karhujärvi	60°14'N 24°17'E	eutr	2.2	4.9	142	1.9	0.0	253	25	1.9	Dredging of the shore areas in 1997, fish removal since 1996, measures to reduce macrophyte expansion (1992, 1993, 1994-1995)
Katumajärvi	60°59'N 24°31'E	meso	7.1	18.9	51	3.8	29.4	232	38	2.4	Sedimentation ponds established in 2004-2005, biomanipulation 2003-2005
Kyynäröjärvi	61°07'N 24°59'E	eutr	1.3	3	25	0.25	0.0	2263	42	0.6	-
Loppijärvi	60°41'N 24°25'E	eutr	1.8	6.7	82	11.8	0.0	93	34	1.4	Wastewater diversion in 1975, wetlands and sedimentation ponds, fish removal since 1990s
Ormajärvi	61°06'N 24°59'E	meso	9.6	29.4	86	6.6	48.6	128	31	3.3	-
Punelia	60°41'N 24°12'E	meso	3.8	14	102	6.8	12.7	23	126	2.4	-
Pusulanjärvi	60°27'N 23°59'E	eutr	4.9	10.6	226	2.1	49.7	1707	35	1.6	Wastewater load until 1988, intensive fishing, Aeration (first in 1989)
Puujärvi	60°15'N 23°43'E	meso	8.3	21.7	27	6.4	15.0	26	173	1.7	-
Pyhäjärvi (A)	60°43'N 26°00'E	eutr	21	68	459	12.9	32.9	814	34	2.3	-
Pyhäjärvi (S)	61°00'N 22°18'E	meso	5.5	26.2	461	155	4.5	106	27	4.5	Intensive commercial fishing (+ fish removal)
Rehtijärvi	60°51'N 23°29'E	eutr	9.2	30	3	0.4	47.3	628	29	1.0	Wetlands and sedimentation ponds in 1994-1998, biomanipulation
Sahajärvi	60°44'N 25°28'E	eutr	4.3	11	26	1.92	26.2	403	28	2.3	-
Tiiläänjärvi	60°32'N 25°42'E	eutr	4.4	10.3	38	2.1	26.9	1428	25	2.2	-
Tuusulanjärvi	60°25'N 25°04'E	hyper	3.2	10	92	5.9	26.5	960	46	1.1	Wastewater diversion since 1979, aeration since 1970, wetlands and biomanipulation since 1998
Villikkalanjärvi	60°47'N 26°02'E	hyper	3.2	10	413	7.1	18.5	2302	27	1.5	Fish removal

Table 2. Sediment accumulation rate (SAR), sediment phosphorus concentration (TP_{sed}), phosphorus accumulation rate (TP_{acc}), and TP concentration of the lake water (TP) in lakes studied for 1986–2014. Significant trends in SAR, TP_{sed}, TP_{acc}, and TP (increase “+”, decrease “-”) over the 30-year period are shown (T1, T2, T3, and T4, respectively), and “0” indicates no significant changes. Long-term data on water quality were not available for Lake Kyyjärvi (indicated as “na”).

Lake	SAR (g cm ⁻² y ⁻¹)	T 1	TP _{sed} (mg g ⁻¹)	T 2	TP _{acc} (mg P m ⁻² y ⁻¹)	T3	TP (µg l ⁻¹)	T4
Aimajärvi	0.04	+	2.2	0	907	+	42	-
Bodominjärvi	0.16	0	1.5	+	2373	+	32	0
Enäjärvi	0.04	0	2.7	0	1074	0	98	0
Enonselkä	0.18	0	2.7	0	4910	0	34	-
Hormajärvi	0.03	+	5.3	0	1407	+	14	+
Kajaanselkä	0.08	+	1.6	0	1418	+	16	-
Karhujärvi	0.15	0	1.2	+	1827	+	72	+
Katumajärvi	0.06	+	4.6	0	2610	+	19	0
Kyyjärvi	0.16	+	1.6	+	2540	+	50	na
Loppijärvi	0.07	+	2.3	+	1699	+	30	+
Ormajärvi	0.06	0	3.6	0	2073	0	20	-
Punelia	0.02	0	1.8	+	340	+	14	0
Pusulanjärvi	0.19	-	3.2	+	6038	0	49	0
Puujärvi	0.02	0	2.2	+	508	+	12	-
Pyhäjärvi (A)	0.17	0	2.1	0	3502	0	44	0
Pyhäjärvi (S)	0.11	0	2.3	0	2488	0	17	+
Rehtijärvi	0.20	0	6.0	-	12044	-	56	+
Sahajärvi	0.17	0	1.7	+	2906	+	40	0
Tiiläänjärvi	0.30	0	1.5	0	4533	0	80	+
Tuusulanjärvi	0.11	+	1.7	+	1894	+	101	-
Villikkalanjärvi	0.36	+	1.1	-	4097	+	111	+

Table 3. Coefficients for calculating the first six (of eight) principal components (PCs) and corresponding eigenvalues of the correlation matrix calculated for 21 lakes with a full set of all eight contributing characteristics. The characteristics needed include: maximum depth ($D_{\max}$), lake area (LA), catchment area (CA), mean depth (D), ratio of the CA to LA, inflow of P (TP_{in}), anoxic factor (AF, defined as the product of the duration of anoxia and the percentage of the anaerobic areas), Osgood's index, or $D \times LA^{-0.5}$ (OI).

Lake	PC 1	PC 2	PC 3	PC 4	PC 5	PC 6	PC 8
Characteristics							
D	0.441	0.265	0.201	-0.101	-0.334	-0.033	0.076
$D_{\max}$	0.500	0.105	0.177	0.102	-0.174	-0.222	0.130
CA	0.072	-0.334	0.627	-0.097	-0.217	0.164	-0.607
LA	0.318	-0.458	0.113	0.033	0.176	0.613	0.516
AF	0.263	0.420	0.195	-0.569	0.357	-0.056	0.107
CA/LA	-0.374	0.032	0.479	0.177	-0.363	-0.265	0.554
OI	0.157	0.516	-0.002	0.653	-0.040	0.383	-0.150
TP_{in}	-0.270	0.195	0.504	0.163	0.620	0.042	-0.004
$D_{\max}/D$	0.376	-0.339	0.015	0.402	0.367	-0.570	0.044
Eigenvalue	3.480	2.414	1.636	0.493	0.470	0.228	0.079
Proportion of variability	0.387	0.268	0.182	0.057	0.055	0.052	0.009
Cumulative proportion	0.387	0.655	0.837	0.892	0.944	0.969	0.978

Table 4. Most significant predictors of the phosphorus accumulation rate (log-transformed values) according to the linear model. The effects indicate the change of the phosphorus accumulation rate when significant principal component (PC) changes by one unit.

Significant predictors	Effect	R^2	p
PC 1	2.78 ± 0.82	0.044	0.006
PC 2	5.86 ± 0.75	0.197	< 0.0001
PC 3	4.88 ± 0.78	0.137	< 0.0001
PC 4	4.95 ± 0.76	0.046	0.004
PC 5	6.17 ± 0.85	0.141	< 0.0001

Table 5. Significant correlations (indicated by Pearson correlation coefficient) of monthly NAO values with monthly values for the (hydro)meteorological variables including water temperature in the surface layer, near bottom water layer and difference in these water temperatures.

Month		NAO_1	NAO_2	NAO_3	NAO_4	NAO_5	NAO_6	NAO_7	NAO_8	NAO_9	NAO_10	NAO_11	NAO_12
Jan	bot	0.210*											
	wind	0.601****											
Feb	bot	0.197*											
	wind		0.436*										
Mar	surf		0.271****										
	dif		0.142**	0.178***									
	wind			0.337*									
Apr	surf		0.216*	0.485****									
	bot			0.223*									
	dif		0.250*	0.403****									
May	bot		0.163**	0.152*									
	dif		-0.189**										
	wind	-0.347*											
Jun	surf					0.191*							
	dif					0.208**							
Jul	surf			-0.203***									
	bot							0.326*					
	dif			-0.144*									
Aug	surf				-0.227****								
Sep	surf			-0.190*									
	wind		-0.345*										
Oct	bot								-0.180**				
	dif								0.170**				
	wind				-0.310*				-0.427**				
Nov	surf									0.291**	0.300**		
	bot									0.264**	0.256**		
	dif									-0.218*	0.229**	0.281**	
	wind		-0.365*	-0.321*									
Dec	surf								-0.353*				
	bot			0.314*									
	dif			-0.413**				-0.434**					
	wind											0.325*	0.426**

**** - the level of significance $p < 0.0001$; *** - $p < 0.001$; ** - $p < 0.01$; * - $p < 0.05$.

Figure Captions

Fig. 1 Variations of the sediment accumulation rate in mesotrophic (meso), eutrophic (eutr) and hypertrophic (hyper) lakes of southern Finland

Fig. 2. Phosphorus accumulation rate (TP_{acc}) in 21 Finnish lakes over 1986–2014.

Fig. 3. Dependence of the phosphorus accumulation rate (log-transformed values) on the specific combination of lake characteristics represented by PC5 that characterizes mainly productive lakes

Fig. 4. Monthly variations in NAO (a), daily mean wind speed (b), and water temperature differences between surface and near bottom layers of the lakes studied (c) during 1986–2014.

Fig. 5. Dependence of the annual phosphorus accumulation rate (TP_{acc} ; $\log(TP_{acc}) = 7.4 - 0.4 * \text{NAO in October}$), and water temperature difference between the surface and near bottom layer in November (temperature difference in November = $0.4 * \text{NAO in October} - 0.06$) on the NAO in October