<https://helda.helsinki.fi>

Effects of climate change on the distribution of hoverfly species (Diptera : Syrphidae) in Southeast Europe

Milicic, Marija

2018-04

Milicic , M , Vujic , A & Cardoso , P 2018 , ' Effects of climate change on the distribution of hoverfly species (Diptera : Syrphidae) in Southeast Europe ' , Biodiversity and Conservation , vol. 27 , no. 5 , pp. 1173-1187 . <https://doi.org/10.1007/s10531-017-1486-6>

<http://hdl.handle.net/10138/308070>

<https://doi.org/10.1007/s10531-017-1486-6>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Effects of climate change on the distribution of hoverfly species (Diptera: Syrphidae) in Southeast Europe

Marija Miličić^{1,3*}, Ante Vujić², Pedro Cardoso³

¹BioSense Institute - Research Institute for Information Technologies in Biosystems, University of Novi Sad, Trg Dr Zorana Đinđića 1, 21000 Novi Sad, Serbia, ²Department of Biology and Ecology, Faculty of Sciences, University of Novi Sad, Novi Sad, Serbia, ³Finnish Museum of Natural History, Zoology Unit, University of Helsinki, Helsinki, Finland

*Corresponding author: Marija Miličić, BioSense Institute - Research Institute for Information Technologies in Biosystems, University of Novi Sad, Trg Dr Zorana Đinđića 1, Novi Sad, Serbia. Telephone: +381/644849132. E-mail: marija.milicic@biosense.rs

Acknowledgements

We kindly thank John O'Brien for English proofreading and Dr Tamara Jurca for useful comments regarding this paper. This work was supported by the Ministry of Education, Science and Technological Development, Republic of Serbia, under Grant No. 173002 and Grant No. 43002, the Provincial Secretariat for Science and Technological Development under Grant No. 114-457-2173/2011-01 and H2020 project ANTARES under Grant No. 664387.

Abstract: Climate change presents a serious threat to global biodiversity. Loss of pollinators in particular has major implications, with extirpation of these species potentially leading to severe losses in agriculture and, thus, economic losses. In this study, we forecast the effects of climate change on the distribution of hoverflies in Southeast Europe using species distribution modelling and climate change scenarios for two time-periods. For 2041-2060, 19 analysed species were predicted to increase their areas of occupancy, with the other 25 losing some of their ranges. For 2061-2080, 55% of species were predicted to increase their area of occupancy, while 45% were predicted to experience range decline. In general, range size changes for most species were below 20%, indicating a relatively high resilience of hoverflies to climate change when only environmental variables are considered. Additionally, range-restricted species are not predicted to lose more area proportionally to widespread species. Based on our results, two distributional trends can be established: the predicted gain of species in alpine regions, and future loss of species

from lowland areas. Considering that the loss of pollinators from present lowland agricultural areas is predicted and that habitat degradation presents a threat to possible range expansion of hoverflies in the future, developing conservation management strategy for the preservation of these species is crucial. This study represents an important step towards the assessment of the effects of climate changes on hoverflies and can be a valuable asset in creating future conservation plan, thus helping in mitigating potential consequences.

Key words: conservation, global warming, insects, endemism, species distribution modelling

Introduction

Ecosystems across the world are facing severe modifications due to climate change and many species are facing extinction risk as a result. Species tolerance to changing climate is critical from ecological, conservation and evolutionary points of view (Garcia-Robledo et al. 2016). Several studies have shown that climate change influences many species in different ways: they can move their range to find suitable environment (Hickling et al. 2006; Parmesan 2006); alter phenology in order to adapt to new conditions (Visser 2008; Gardner et al. 2011); modify their behaviour, with species opting to change foraging or activity hours, adapt their physiology, or increase metabolism and growth rates (Hughes 2000); shift their preferred habitat; or eventually undergo evolutionary shifts (Bradshaw and Holzapfel 2006; Visser 2008; Williams et al. 2008; Dauterive et al. 2009; Maggini et al. 2011). If none of these is possible or sufficient, extinction is possible (Thuiller et al. 2008; Lurgi et al. 2012). Among range shifts, climate change is expected to force species distributions towards higher elevations and latitudes, leading to extinction of species whose future habitable climate space becomes too small or too isolated from their current geographical ranges (Hill et al. 2002; Midgley et al. 2002; Wilson et al. 2005). Limited dispersal capacity, low reproductive rate and a high degree of habitat specialization are attributes that make species prone to environmental disturbances (Isaac et al. 2009). Species with a limited distribution often possess most of these characteristics. Although widespread species may also be endangered, range-restricted species are particularly vulnerable (Thomas et al. 2004; Wulf et al. 2013). Estimating the effects of climate change on species distributions is an important step in assessing the vulnerability of species to extinction and can provide useful information about the spectrum of possible consequences (Araújo et al. 2005; Gibson et al. 2010; Yates et al. 2010). Species distribution models (SDM; also called environmental niche

models) are often used to predict the effects of climate change and they have been successfully applied in a number of environmental studies (Hannah et al. 2002; Elith et al. 2006; Peterson 2006). SDM assess the relationship between species occurrence at sites and the environmental characteristics of those areas (Franklin 2009) in order to predict the distribution of suitable environmental envelopes for the species in non-sampled areas or time-frames (Elith and Leathwick 2009; Costion et al. 2015). When used in combination with future climate change scenarios, these models can indicate the expected effect of changing climate on species distributions.

Here, we use SDM to assess the potential effects of climate change on Southeast (SE) European hoverflies. Hoverflies are Dipteran insects comprising around 6000 described species (Thompson 2013). They are recognized as an important pollinator group (Fontaine et al. 2005; Petanidou et al. 2011; Jauker et al. 2012; Stenley et al. 2013), and some species are used as biological control agents (White et al. 1995). SE Europe harbours exceptional hoverfly diversity. The Balkan Peninsula, occupying the largest part of SE Europe, is considered a hotspot of European biodiversity (Griffits et al. 2004) owing to its long-term environmental stability (Previšić et al. 2009) and habitat diversity. The great variety of plants and habitat heterogeneity in this region promotes a high diversity of insect fauna. Vujić et al. (2001) revealed that the diversity of hoverflies in the Balkan Peninsula is amongst the highest in Europe. The Aegean islands, a part of our study area, have also been designated as one of the world's hotspots for hoverflies (Vujić et al. 2012, 2016b; Radenković et al. 2011).

Our aims were to: (i) analyse the effects of climate change on the distribution of species by examining predicted changes in range size based on forecasts of current and future potential distribution; (ii) describe and compare species-richness patterns for both present and future scenarios; (iii) verify if owing to their theoretically higher vulnerability, the areas of occupancy of range-restricted species decrease proportionally more than those of widespread species; and (iv) discuss possible consequences to mutualistic networks and implications for conservation of hoverflies.

Material and methods

Occurrence data

Species distribution data for all species in SE Europe were extracted from the database of the Department of Biology and Ecology of the University of Novi Sad, which is the largest database on the region's hoverflies (occurrences of species used in this study are available at:

http://www.db.e.uns.ac.rs/o_departmanu/laboratorije/laboratorija_za_istrazivanje_i_zastitu_biodiverziteta/prilog/milic_et_al_2017_-_species_occurrences_data). This database comprises data from field collecting in the study area from 1950-2015, data obtained from different museum and private collections, and published material referring to this geographic area. Only specimens with precise distributional data were used. If locality coordinates were available, they were checked for accuracy. Records only with locality names were assigned coordinates using Google Earth (Google Inc, 2016). For our analysis, we only used species endemic to SE Europe or whose ranges outside this region do not cover areas with climatic conditions differing from those within the study area (otherwise SDM would reflect only part of the environmental niche of species and, thus, be potentially biased). For reducing sampling bias, we applied the thinning procedure, where we used a threshold of 0.01 of the maximum distance between any two points. The procedure is explained in detail in Miličić et al. (2017). After data processing, all species with less than five occurrence points were dropped (the number of occurrences per species is assessable in occurrence data table, provided on the link above in text).

Selection of predictor variables

We used 19 bioclimatic variables plus elevation data (2.5 arc-minutes resolution, approximately 4.5 km²) taken from the WorldClim dataset (Hijmans et al. 2005) for model building. As future bioclimatic variables, we used climate projections at the same resolution from the global climate models used in the Fifth Assessment report of the Intergovernmental Panel on Climate Change (IPCC 2013). We chose the HadGEM2-ES model with RCP 8.5 (Representative Concentration Pathway), which is a greenhouse gas concentration trajectory that assumes that emissions will continue to rise throughout the 21st century. We deliberately choose the “worst case scenario” because historical and current trends of greenhouse emissions are trailing the RCP 8.5 trajectory (Peters et al. 2013). Modelling was done in two stages. First, we used all variables. Then, using only the stronger predictors for each

species, we built the final models and, in that way, avoiding overfitting the models (see details in Miličić et al. 2017).

Species distribution modelling

For SDM, we used the maxent function of the *dismo* R package (Hijmans et al. 2016). MAXENT is one of the most commonly used algorithms for this purpose (Phillips et al. 2006, 2008; Peterson et al. 2007; Ortega-Huerta and Peterson 2008; Merow et al. 2013). This algorithm shows a generally good performance for presence-only data, even with small sample sizes (Kumar and Stohlgren 2009; Pearson et al. 2007). There are several examples where MAXENT has been used for modelling the potential distributions of range-restricted species. For example, Gibson et al. (2010) used MAXENT to estimate the effect of climate change on a range-restricted marsupial. Costion et al. (2015) and Krause et al. (2015) used it to assess the effect of climate change on endemic species of plants, and Vujić et al. (2016a) used MAXENT to identify favourable habitats for hoverflies of conservation interest in Serbia.

Dataset was split into training and test data. MAXENT default settings were maintained. For each species, maps of current and future potential distributions were created for the year 2050 (average of years 2041-2060) and 2070 (average 2061-2080). These maps were then transformed to binary format (showing suitable/unsuitable areas for species), applying the threshold that maximized the sum of sensitivity and specificity (Liu et al. 2005, 2013).

Binary maps were used to calculate the potential area of occupancy (pAOO) for all species in all time-periods. To assess the predictive performance of the models, we used TSS (True Skill Statistic) as an evaluation measure, which has been shown to be a good measure of accuracy (Allouche et al. 2006; Liu et al. 2013). TSS values range from -1 to +1, with +1 indicating perfect model agreement and values of zero or less indicating a performance no better than random (Allouche et al. 2006).

Calculation of potential species richness

Our second objective was to describe and compare the species richness patterns for both present and future scenarios. Maps for each species under the present scenario were overlaid and summed for species richness. We then did the same for the future scenario. Then, the overall present and future richness maps were subtracted, allowing changes in diversity per cell between time-periods to be determined (see also Ferreira et al. 2016). All maps were created using the software DIVA-GIS (version 7.5).

Our third objective was to test if the ranges of range-restricted species decrease proportionally more than those of widespread species. We calculated the Pearson correlation between present pAOO of all species and the respective predicted relative changes in range size for both time-periods. A significantly negative correlation would indicate that species with smaller ranges would have higher proportional losses of pAOO, confirming our hypothesis.

Range expansion and contraction patterns

In order to test whether range expansion and contraction patterns are related with altitude, Spearman rank correlation among all cells showing difference in species richness (between both future periods and present) and altitude was calculated.

Results

Species distribution models

In total, 44 species of hoverflies were included in our analysis (Tab. 1). TSS values used for evaluation of the models varied between 0.49 and 0.99 (Tab. 1), representing a good fit of the models. The bioclimatic variable contributing to the highest number of models (n=24) was precipitation seasonality (bio15). Other variables contributing to more than 10 final models were mean temperature of the wettest quarter (bio8), mean temperature of the driest quarter (bio9) and precipitation of the driest month (bio14). The list of bioclimatic variables used in each final model is given in Fig. 1.

For 2041-2060, 19 species (43%) were predicted to lose part of their range, while 25 species were predicted to gain in range. However, for 40% of the species, their pAOO changed by less than 20%. For 2061-2080, 20 species (45%) were predicted to reduce their area of occupancy, whereas 24 species (55%) would gain occupancy. Variation in range size for 38% of the species was below 20%. Four different trends can be identified from the overall changes in pAOO: (1) fifteen species (34%) were predicted to lose part of their range for both time-periods; (2) twenty species (45%) would expand their pAOO over both time-periods; (3) four species (9%) were predicted to lose part of their range during the first period and then regain some of it under the second period; and (4) another five species (11%) would first gain range and then lose it.

Species richness

We predicted the species richness hotspots to be similar across time. The Aegean islands and part of the Dinaric mountain range stretching through Bosnia and Herzegovina, Serbia and Montenegro were predicted to have the highest potential number of species in all cases (Fig. 2b, 2c, 2d). The Dinaric mountains, together with the Alpine region in Slovenia, high mountain peaks in central Peloponnese, part of the Carpathian Mountains in Romania and the coastal zone along the Black Sea, spreading into the continental areas of Southwest Bulgaria, are predicted to gain species with time. In contrast, the valleys between the Olympus and Rhodopes mountains, the lowland along the Dinaric mountain range and the peripheral zone of Strandza Mountain in Bulgaria are each predicted to lose between 1 and 3 species in the future (Fig. 2e and 2f). In general, higher loss is predicted for 2070 time period.

Loss of area

Our results indicate that the correlations between present ranges of species and proportional changes in range size for both time-periods were not statistically significant (Tab. 2).

Range expansion and contraction patterns for both future time periods showed slight positive statistically significant correlation with altitude (Tab. 2).

Discussion

In this paper, we forecast the effect of climate change on the distribution of hoverflies in SE Europe using SDM and climate change scenarios for two time-periods. We predict species to be distributed in similar proportions amongst losers and gainers of areas of occupancy, yet individual species distributions change considerably over time leading to divergent patterns for various sub-regions of our study area.

Two recent studies analysed the effects of climate change on the distributions of some species belonging to the two largest hoverfly genera in the region, *Merodon* and *Cheilosia*. However, in both studies, only widespread species (i.e. those not limited to the Balkan Peninsula) and with a large number of occurrences (more than 15 and 30 for *Cheilosia* and *Merodon*, respectively) were included in the analyses. Kaloveloni et al. (2015) predicted *Merodon* species to be relatively equally divided amongst gainers and losers of areas of occupancy, whereas Radenković et al.

(2017) concluded that climate change will have serious consequences for the distributions of almost all studied *Cheilosia* species, causing severe range losses for these species across the entire Balkan Peninsula. Undoubtedly, habitat type and the altitude at which a given species occurs influence species distributions of all hoverflies, regardless of which genus they belong to. Most species included in the analyses of Radenković et al. (2017) are Alpine, while the *Merodon* species analysed by Kaloveloni et al. (2015) are both high mountain and Mediterranean in origin, with our analyses confirming the patterns established in these studies. However, endemic species occurring on some Greek islands, and only included in our analyses, show a mixed response, with some species increasing their area of occupancy and that of others decreasing. Mediterranean and lowland taxa are predicted to expand their ranges, as these species can move their range towards higher altitudes if temperatures increase. Thus, it is not surprising that, in most cases, the regions gaining in terms of species richness are mountainous, such as the Alpine regions of Slovenia, the Dinaric Mountains, or part of the Carpathian Mountains. The bioclimatic variables found to mostly affect hoverfly distribution were related to precipitation seasonality and temperature and precipitation in the driest months, which might be related with these findings. Temperature increases tend to shift species towards areas of higher altitude, which typically have higher levels of precipitation (Beniston 2006). Climate change-induced altitudinal shifts have already been reported in numerous studies for different organisms (Penuales and Boada 2003; Wilson et al. 2005; Hickling et al. 2006; Lenoir et al. 2008). In contrast, lowland areas, such as the valleys between mountains are predicted to lose species. Global warming may render the climatic conditions in such regions too harsh (hot and/or dry) for many hoverflies. It should be noted that this loss of species at low altitude might be compensated by range expansions of species coming from warmer areas in the South and East of Europe, as these species were not modelled here.

Based on our results, three patterns can be established: a) a relatively high resilience of Syrphidae to climate change disturbance; b) future range expansions of some hoverfly species to new locations, mostly mountainous; and c) depletion of syrphid species in lowland areas.

We predict some species (such as *Merodon virgatus* Vujić et Radenković, 2016; see also Tab. 1) to significantly expand their range under a feasible climate change scenario. Thus, it seems that projected climate change will create additional favourable climate space for this and about half the other species we considered here. It is also worth mentioning that, for a considerable number of species, the variation in range size for both time-periods was below

20% (40% and 38% of species for 2050 and 2070, respectively). In addition, we found that range-restricted species are not predicted to decrease their ranges to a greater proportional extent than widespread species. Together, these findings might indicate an overall potential inherent resistance to changing climate amongst hoverflies in SE Europe. If true, hoverflies could become an important alternative leading pollinator group if the number of bees continues to severely decline as a consequence of changing climate, as has been projected (Biesmeijer et al. 2006; Dorman et al. 2008). We note that two factors may decisively influence our conclusions regarding the resilience of species and their future range expansions. These are dispersal capacity and diet specialization. Capacity to disperse to new climatically-suitable regions is a critical factor in species responses to climatic change, as these potential new areas may be out of reach for less vagile species. Considering that the vast majority of larvae of the species we analysed are phytophagous (38 out of 44), contrary to species with saprophagous larvae, the distributions of these hoverfly species are conditioned by the distribution of their hosts. Host plants of stenotopic species may be unavailable in the new locations so that even though the climatic envelope of a new area might be suitable, the habitat perhaps cannot support displaced species. In addition, adult hoverflies are always associated with flowering plants, their food source, which entails that our predictions are necessarily influenced by how the distributions of hoverfly host plants will alter in the future. More detailed knowledge about species biology and ecology, namely functional traits such as dispersal ability and diet, are needed to determine limiting factors for species expansion.

According to our results, loss of hoverfly species from lowlands and their migration to higher altitudinal areas is to be expected. Having in mind that lowlands represent significant areas for agriculture used since ancient times (Turner 1974) because of their higher temperatures and less rugged terrain, these altitudinal shifts might cause a depletion of potential pollinators from agricultural areas, inevitably causing economic losses. Additionally, even if a species remain in lowlands, a changing climate may cause temporal (phenological) and spatial (distributional) mismatches (Hegland et al. 2009), between insect and host plant, leading to partial or complete decoupling of mutualistic partners (Visser and Both 2005). Such decoupling may result in changes to ecosystem dynamics, which again lead to economic losses in agricultural communities (Donnelly et al. 2011). Therefore, assessment of the effects of climate change on mutualistic processes between plants and pollinators is critically needed to fully assess the risk of climate change and the possible consequences on insect communities.

Although predicted future range expansions of some hoverfly species and registered relatively low variations in range size may indicate high resilience of hoverflies to climate change, there are a number of factors that threaten current and predicted future locations for Syrphidae conservation. The areas with the highest predicted species richness for every period, such as the Aegean islands, are dominated by Mediterranean vegetation. Large expanses of this vegetation type are severely affected by land degradation processes, leading to desertification as a result of inadequate land use or because of discordance between economic and conservation priorities (Hill et al. 2008). Many deciduous forests across SE Europe face a similar scenario, harbouring high species richness but are severely endangered due to forestry and land degradation. Jovičić et al. (2017) indicated that land use has a strong influence on the species composition of *Merodon* and *Cheilosia* hoverflies. Changes in habitat availability for species and low tolerance to environmental change increase the risks of severe consequences from climate change. Another factor that can threaten the potential future expansion of hoverflies is intensive agriculture; multiple examples testify to its negative effects on biodiversity (Matson et al. 1997; Sotherton 1998; Tilman et al. 2001; Wickramasinghe et al. 2004). Kremen et al. (2002) found that agricultural intensification has a serious effect on bee populations, causing reductions in both diversity and abundance of species, while Hendrickx et al. (2007) established that total species richness of hoverflies decreases with increasing management intensity in agricultural fields. Agriculture also causes fragmentation of natural habitats, which has a ruinous effect, especially on small and isolated populations (Benton et al. 2003). Tourism also represents serious threat to biodiversity. For example, construction of ski resorts has a strong negative effect on many plant and animal species, including hoverflies, considering that the majority of these species are mountainous. Ristić et al. (2012) addressed the negative effects of the construction of a ski resort on Stara Planina Nature Park in Serbia. As a consequence of the construction of the ski centre, population sizes of several endemic species of birds and plants were significantly reduced or even disappeared from this area rich in hoverflies. Similarly, The Valley of Butterflies on the Greek island of Rhodes, which has been designated as a Natura 2000 site, is predicted to be one of the most species-rich areas for hoverflies under both present and future climate projections in our analysis. However, the numerous tourists visiting this location severely affect its environment, and it is unclear how long the species that this site hosts can resist such anthropogenic pressure (Petanidou et al. 1991). Thus, it might prove crucial to find ways of alleviating the consequences of different threatening factors to preserve imperilled species and biodiversity in general in these regions.

Conclusion

Undoubtedly, climate change will affect species ranges in the future. Hoverflies are in general conjectured to have a relatively high resilience to climate change disturbance, with some species predicted to experience future range expansions to new, mostly mountainous locations, while in lowland areas the depletion of syrphid species is to be expected. Such range shifts (both expansionary and contractionary) are all the more important for species dependent on mutualistic networks and that constitute keystone taxa for several ecosystem services such as pollination. Loss of these species would lead to severe losses in agriculture and, consequently, economic losses. Our study represents an important step towards the assessment of the effects of changing climate on hoverflies and can help in future conservation planning, which could mitigate potential economic loss.

Data availability

All data generated or analysed during this study are included in this published article [or assessable through the link provided in the text].

Compliance with ethical standards

Funding: This work was supported by the Ministry of Education, Science and Technological Development, Republic of Serbia, under Grant No. 173002 and Grant No. 43002, Provincial Secretariat for Science and Technological Development under Grant No. 114–457–2173/2011–01 and H2020 project ANTARES under Grant No. 664387. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of interest: The authors declare that they have no conflict of interest.

References

- Allouche O, Tsoar A, Kadmon R (2006) Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *J Appl Ecol* 43:1223–1232. doi: 10.1111/j.1365–2664.2006.01214.x
- Araújo MB, Whittaker RJ, Ladle RJ, Erhard M (2005) Reducing uncertainty in projections of extinction risk from climate change. *Global Ecol Biogeogr* 14:529–538. doi: 10.1111/j.1466–822X.2005.00182.x

295 Beniston M (2006) Mountain weather and climate: a general overview and a focus on climatic change in the Alps.
 296 *Hydrobiologia* 562:3–16. doi: 10.1007/s10750-005-1802-0

297 Benton TG, Vickery JA, Wilson JD (2003) Farmland biodiversity: is habitat heterogeneity the key? *Trends Ecol*
 298 *Evol* 18:182–188. doi: 10.1016/S0169-5347(03)00011-9

299 Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP, Potts SG, Kleukers R,
 300 Thomas CD, Settele J, Kunin WE (2006) Parallel declines in pollinators and insect-pollinated plants in Britain and
 301 the Netherlands. *Science* 313:351–354. doi: 10.1126/science.1127863

302 Bradshaw WE, Holzapfel CM (2006) Evolutionary response to rapid climate change. *Science* 312:1477–1478. doi:
 303 10.1126/science.1127000

304 Costion CM, Simpson L, Pert PL, Carlsen MM, Kress WJ, Crayn D (2015) Will tropical mountaintop plant species
 305 survive climate change? Identifying key knowledge gaps using species distribution modelling in Australia. *Biol*
 306 *Conserv* 191:322–330. doi: 10.1016/j.biocon.2015.07.022

307 Daufresne M, Lengfellner K, Sommer U (2009) Global warming benefits the small in aquatic ecosystems. *Proc Natl*
 308 *Acad Sci USA*, 106:12788–12793. doi: 10.1073/pnas.0902080106

309 Donnelly A, Caffarra A, O'Neill BF (2011) A review of climate-driven mismatches between interdependent
 310 phenophases in terrestrial and aquatic ecosystems. *Int J Biometeorol* 55:805–817. doi:10.1007/s00484-011-0426-5

311 Dormann CF, Schweiger O, Arens P, Augenstein I, Aviron ST, Bailey D, Baudry J, Billeter R, Bugter R, Bukacek
 312 R, Burel F, Cerny M, De Cock R, De Blust G, DeFilippi R, Diekötter T, Dirksen J, Durka W, Edwards PJ, Frenzel
 313 M, Hamersky R, Hendrickx F, Herzog F, Klotz S, Koolstra B, Lausch A, Le Coeur D, Liira J, Maelfait JP, Opdam
 314 P, Roubalova M, Schermann–Legionnet A, Schermann N, Schmidt T, Smulders MJM, Speelmans M, Simova P,
 315 Verboom J, van Wingerden W, Zobel M, Burel F (2008) Prediction uncertainty of environmental change effects on
 316 temperate European biodiversity. *Ecol Lett* 11:235–244. doi: 10.1111/j.1461-0248.2007.01142.x

317 Elith J, Graham CH, Anderson RP, Dudik M, Ferrier S, Guisan A, Hijmans RJ, Huettmann F, Leathwick JR,
 318 Lehmann A, Li J, Lohmann LG, Loiselle BA, Manion G, Moritz C, Nakamura M, Nakazawa Y, Overton JM,
 319 Peterson AT, Phillips SJ, Richardson K, Scachetti-Pereira R, Schapire RE, Soberon J, Williams S, Wisz MS,

320 Zimmermann NE (2006) Novel methods improve prediction of species' distributions from occurrence data.
 321 Ecography 29:129–151. doi: 10.1111/j.2006.0906–7590.04596.x

322 Elith J, Leathwick JR (2009) Species distribution models: ecological explanation and prediction across space and
 323 time. Annu Rev Ecol Evol Syst 40:677–697 doi: 10.1146/annurev.ecolsys.110308.120159

324 Ferreira MT, Cardoso P, Borges PA, Gabriel R, de Azevedo EB, Reis F, Araújo M, Elias RB (2016) Effects of
 325 climate change on the distribution of indigenous species in oceanic islands (Azores). Climatic Change 138:603–615.
 326 doi: 10.1007/s10584–016–1754–6

327 Fontaine C, Dajoz I, Meriguet J, Loreau M (2005) Functional diversity of plant-pollinator interaction webs enhances
 328 the persistence of plant communities. PLoS Biol 4: e1. doi:10.1371/journal.pbio.0040001

329 Franklin J (2009) Mapping species distributions: spatial inference and prediction. Cambridge University Press,
 330 Cambridge, UK

331 García-Robledo C, Kuprewicz EK, Staines CL, Erwin TL, Kress WJ (2016) Limited tolerance by insects to high
 332 temperatures across tropical elevational gradients and the implications of global warming for extinction. Proc Natl
 333 Acad Sci USA 113:680–685. doi: 10.1073/pnas.1507681113

334 Gardner JL, Peters A, Kearney MR, Joseph L, Heinsohn R (2011) Declining body size: a third universal response to
 335 warming? Trends Ecol Evol 26:285–291. doi: 10.1016/j.tree.2011.03.005

336 Gibson L, McNeill A, de Tores P, Wayne A, Yates C (2010) Will future climate change threaten a range restricted
 337 endemic species, the quokka (*Setonix brachyurus*), in south west Australia? Biol Conserv 143:2453–2461. doi:
 338 10.1016/j.biocon.2010.06.011

339 Google Inc. Google Earth. Mountain View, California, USA. 2013. <https://www.google.com/earth>. Accessed 15
 340 January, 2016

341 Griffiths HI, Kryštufek B, Reed JM (2004) Balkan Biodiversity Pattern and Process in the European Hotspot.
 342 Kluwer Academic Press, Dordrecht

343 Hannah L, Midgley GF, Millar D (2002) Climate change-integrated conservation strategies. Global Ecol Biogeogr
 344 11:485–495. doi: 10.1046/j.1466–822X.2002.00306.x

345 Harrell FE (2001) Regression modelling strategies with applications to linear models, logistic regression and
 346 survival analysis. Springer Series in Statistics, Springer–Verlag, New York

347 Hegland SJ, Nielsen A, Lázaro A, Bjerknes AL, Totland Ø (2009) How does climate warming affect plant-pollinator
 348 interactions? *Ecol Lett* 12:184–195. doi: 10.1111/j.1461-0248.2008.01269.x

349 Hendrickx F, Maelfait JP, Van Wingerden W, Schweiger O, Speelmans M, Aviron S, Augenstein I, Billeter R,
 350 Bailey D, Bukacek R, Burel F, Diekötter T, Dirksen J, Herzog F, Liira J, Roubalova M, Vandomme V, Bugter R
 351 (2007) How landscape structure, land-use intensity and habitat diversity affect components of total arthropod
 352 diversity in agricultural landscapes. *J Appl Ecol* 44:340–351. doi: 10.1111/j.1365-2664.2006.01270.x

353 Hickling R, Roy DB, Hill JK, Fox R, Thomas CD (2006) The distributions of a wide range of taxonomic groups are
 354 expanding polewards. *Glob Change Biol* 12:450–455. doi: 10.1111/j.1365-2486.2006.01116.x

355 Hijmans RJ, Phillips S, Leathwick J, Elith J (2016) dismo: Species Distribution Modeling. R package version 1.1–1.
 356 <https://CRAN.R-project.org/package=dismo>

357 Hijmans RJ, Guarino L, Mathur P (2012) DIVA-GIS, version 7.5. A geographic information system for the analysis
 358 of species distribution data. Manual. <http://www.diva-gis.org>.

359 Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for
 360 global land areas. *Int J Climatol* 25:1965–1978. doi: 10.1002/joc.1276

361 Hill JK, Thomas CD, Fox R, Telfer MG, Willis SG, Asher J, Huntley B (2002) Responses of butterflies to twentieth
 362 century climate warming: implications for future ranges. *Proc Roy Soc Lond B Bio* 269:2163–2171. doi:
 363 10.1098/rspb.2002.2134

364 Hill J, Stellmes M, Udelhoven T, Röder A, Sommer S (2008) Mediterranean desertification and land degradation:
 365 mapping related land use change syndromes based on satellite observations. *Glob Planet Chang* 64:146–157. doi:
 366 10.1016/j.gloplacha.2008.10.005

367 Hughes L (2000) Biological consequences of global warming: is the signal already apparent? *Trends Ecol Evol*
 368 15:56–61. doi: 10.1016/S0169-5347(99)01764-4

369 IPCC (2014) Fifth Assessment Report (AR5). Cambridge University Press

370 Isaac JL, Vanderwal J, Johnson CN, Williams SE (2009) Resistance and resilience: quantifying relative extinction
 371 risk in a diverse assemblage of Australian tropical rainforest vertebrates. *Divers Distrib* 15:280–288. doi:
 372 10.1111/j.1472–4642.2008.00531.x

373 Jauker F, Bondarenko B, Becker HC, Steffan–Dewenter I (2012) Pollination efficiency of wild bees and hoverflies
 374 provided to oilseed rape. *Agric For Entomol* 14:81–87. doi: 10.1111/j.1461–9563.2011.00541.x

375 Jovičić S, Burgio G, Diti I, Krašić D, Markov Z, Radenković S, Vujić A (2017) Influence of landscape structure and
 376 land use on *Merodon* and *Cheilosia* (Diptera: Syrphidae): contrasting responses of two genera. *J Insect Conserv* doi:
 377 10.1007/s10841–016–9951–1

378 Kaloveloni A, Tscheulin T, Vujić A, Radenković S, Petanidou T (2015) Winners and losers of climate change for
 379 the genus *Merodon* (Diptera: Syrphidae) across the Balkan Peninsula. *Ecol Model* 313:201–211. doi:
 380 10.1016/j.ecolmodel.2015.06.032

381 Krause CM, Cobb NS, Pennington DD (2015) Range shifts under future scenarios of climate change: dispersal
 382 ability matters for Colorado Plateau endemic plants. *Nat Areas J* 35:428–438. doi:
 383 <http://dx.doi.org/10.3375/043.035.0306>

384 Kremen C, Williams NM, Thorp RW (2002) Crop pollination from native bees at risk from agricultural
 385 intensification. *Proc Natl Acad Sci USA* 99:16812–16816. doi: 10.1073/pnas.262413599

386 Kumar S, Stohlgren TJ (2009) Maxent modeling for predicting suitable habitat for threatened and endangered tree
 387 *Canacomyrica monticola* in New Caledonia. *J Ecol Nat Environ* 1:094–098.

388 Lenoir J, Gégout JC, Marquet PA, De Ruffray P, Brisse H (2008) A significant upward shift in plant species
 389 optimum elevation during the 20th century. *Science* 320:1768–1771. doi: 10.1126/science.1156831

390 Liu C, Berry PM, Dawson TP, Pearson RG (2005) Selecting thresholds of occurrence in the prediction of species
 391 distributions. *Ecography* 28:385–393. doi: DOI: 10.1111/j.0906–7590.2005.03957.x

392 Liu C, White M, Newell G (2013) Selecting thresholds for the prediction of species occurrence with presence-only
 393 data. *J Biogeogr* 40:778–789. doi: 10.1111/jbi.12058

394 Lurgi M, Lopez B, Montoya J (2012) Climate change impacts on body size and food web structure on mountain
 395 ecosystems. *Philos Trans Roy Soc Lond B Biol Sci* 367:3050–3057. doi: 10.1098/rstb.2012.0239

396 Maggini R, Lehmann A, Kéry M, Schmid H, Beniston M, Jenni L, Zbinden N (2011) Are Swiss birds tracking
 397 climate change? Detecting elevational shifts using response curve shapes. *Ecol Model* 222:21–32. doi:
 398 10.1016/j.ecolmodel.2010.09.010

399 Matson PA, Parton WJ, Power AG, Swift MJ (1997) Agricultural intensification and ecosystem properties. *Science*
 400 277:504–509. doi: 10.1126/science.277.5325.504

401 Merow C, Smith MJ, Silander JA (2013) A practical guide to MaxEnt for modeling species' distributions: what it
 402 does, and why inputs and settings matter. *Ecography* 36:1058–1069. doi: 10.1111/j.1600–0587.2013.07872.x

403 Midgley GF, Hannah L, Millar D, Rutherford MC, Powrie LW (2002) Assessing the vulnerability of species
 404 richness to anthropogenic climate change in a biodiversity hotspot. *Glob Ecol Biogeogr* 11:445–451. doi:
 405 10.1046/j.1466–822X.2002.00307.x

406 Miličić M, Vujić A, Jurca T, Cardoso P (2017) Designating conservation priorities for Southeast European
 407 hoverflies (Diptera: Syrphidae) based on species distribution models and species vulnerability. *Insect Conserv*
 408 *Divers* 10:354–366. doi: 10.1111/icad.12232

409 Ortega-Huerta MA, Peterson AT (2008) Modeling ecological niches and predicting geographic distributions: a test
 410 of six presence-only methods. *Rev Mex Biodiv* 79:205–216

411 Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol Evol Syst*
 412 37:637–669. doi: 10.1146/annurev.ecolsys.37.091305.110100

413 Pearson RG, Raxworthy CJ, Nakamura M, Peterson TA (2007) Predicting species distributions from small numbers
 414 of occurrence records: a test case using cryptic geckos in Madagascar. *J Biogeogr* 34:102–117. doi: 10.1111/j.1365–
 415 2699.2006.01594.x

416 Peñuelas J, Boada M (2003) A global change-induced biome shift in the Montseny Mountains (NE Spain). *Glob*
 417 *Chang Biol* 9:131–140. doi: 10.1046/j.1365–2486.2003.00566.x

418 Petanidou T, Vokou D, Margaritis NS (1991) *Panaxia quadripunctaria* in the highly touristic Valley of Butterflies
 419 (Rhodes, Greece): conservation problems and remedies. *Ambio*, 20:124–128

420 Petanidou T, Vujić A, Ellis WN (2011) Hoverfly diversity (Diptera: Syrphidae) in a Mediterranean scrub
 421 community near Athens, Greece. *Ann Soc Entomol* 47:168–175. doi: dx.doi.org/10.1080/00379271.2011.10697709

422 Peters GP, Andrew RM, Boden T, Canadell JG, Ciais P, Le Quere C, Marland G, Raupach MR, Wilson C (2013)
 423 The challenge to keep global warming below 2 °C. *Nat Clim Chang* 3:4–6. doi:10.1038/nclimate1783
 424 Peterson AT, Papes M, Eaton M (2007) Transferability and model evaluation in ecological niche modeling: a
 425 comparison of GARP and Maxent. *Ecography* 30:550–560. doi: 10.1111/j.0906–7590.2007.05102.x
 426 Peterson AT (2006) Uses and requirements of ecological niche models and related distributional models.
 427 *Biodiversity Informatics* 3:59–72. doi: <https://doi.org/10.17161/bi.v3i0.29>
 428 Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions.
 429 *Ecol Model* 190:231–259. doi: 10.1016/j.ecolmodel.2005.03.026
 430 Phillips SJ, Dudik M (2008) Modeling of species distributions with Maxent: new extensions and a comprehensive
 431 evaluation. *Ecography* 31:161–175. doi: 10.1111/j.0906–7590.2008.5203.x
 432 Previšić A, Walton C, Kučinić M, Mitrikeski PT, Kerovec M (2009) Pleistocene divergence of Dinaric *Drusus*
 433 endemics (Trichoptera, Limnephilidae) in multiple microrefugia within the Balkan Peninsula. *Mol Ecol* 18:634–647.
 434 doi: 10.1111/j.1365–294X.2008.04046.x
 435 Radenković S, Schweiger O, Milić D, Harpke A, Vujić A (2017) Living on the edge: Forecasting the trends in
 436 abundance and distribution of the largest hoverfly genus (Diptera: Syrphidae) on the Balkan Peninsula under future
 437 climate change. *Biol Conserv* 212:216–229. doi: 10.1016/j.biocon.2017.06.026
 438 Radenković S, Vujić A, Stahls G, Perez-Banon C, Petanidou T, Šimić S (2011) Three new cryptic species of the
 439 genus *Merodon* Meigen (Diptera: Syrphidae) from the island of Lesvos (Greece). *Zootaxa* 2735:35–56.
 440 Ristić R, Kašanin-Grubin M, Radić B, Nikić Z, Vasiljević N (2012) Land degradation at the Stara Planina ski resort.
 441 *Environ Manage* 49:580–592. doi: 10.1007/s00267–012–9812–y
 442 Sotherton NW (1998) Land–use changes and the decline of farmland wildlife: an appraisal of the set–aside
 443 approach. *Biol Conserv* 83:259–268. doi: 10.1016/S0006–3207(97)00082–7
 444 Stanley DA, Gunning D, Stout JC (2013) Pollinators and pollination of oilseed rape crops (*Brassica napus* L.) in
 445 Ireland: ecological and economic incentives for pollinator conservation. *J Insect Conserv* 17:1181–1189. doi:
 446 10.1007/s10841–013–9599–z

447 Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, Erasmus BF, De Siqueira MF,
 448 Grainger A, Hannah L, Hughes L (2004) Extinction risk from climate change. *Nature* 427:145–148. doi:
 449 doi:10.1038/nature02121

450 Thompson FC (2013) Family Syrphidae. *Systema Dipterorum*, version 1.5. <http://www.diptera.org>. Accessed 28
 451 September 2016

452 Thuiller W, Albert C, Araújo M, Berry P, Cabeza M, Guisan A, Hickler T, Midgley G, Paterson J, Schurr F, Sykes
 453 M, Zimmermann N (2008) Predicting global change impacts on plant species' distributions: future challenges.
 454 *Perspect Plant Ecol Evol Syst* 9:137–152. doi: 10.1016/j.ppees.2007.09.004

455 Tilman D, Fargione J, Wolff B, D'Antonio C, Dobson A, Howarth R, Schindler D, Schlesinger WH, Simberloff D,
 456 Swackhamer D (2001) Forecasting agriculturally driven global environmental change. *Science* 292:281–284. doi:
 457 10.1126/science.1057544

458 Turner II B L (1974) Prehistoric intensive agriculture in the Mayan lowlands. *Science* 185:118–124.

459 Visser ME (2008) Keeping up with a warming world; assessing the rate of adaptation to climate change. *Proc Roy*
 460 *Soc Lond B Biol Sci* 275:649–659. doi: 10.1098/rspb.2007.0997

461 Visser ME, Both C (2005) Shifts in phenology due to global climate change: the need for a yardstick. *Proc Roy Soc*
 462 *Lond B Biol Sci* 272:2561–2569. doi: 10.1098/rspb.2005.3356

463 Vujić A, Petanidou T, Tscheulin T, Cardoso P, Radenković S, Stahls G, Baturan Ž, Mijatović G, Rojo S, Perez-
 464 Banon C, Devalez J, Andrić A, Jovičić S, Krašić D, Markov Z, Radišić D, Tataris G (2016b) Biogeographical
 465 patterns of the genus *Merodon* Meigen, 1803 (Diptera: Syrphidae) in islands of the eastern Mediterranean and
 466 adjacent mainland. *Insect Conserv Divers* 9:181–191. doi: 10.1111/icad.12156

467 Vujić A, Radenković S, Nikolić T, Radišić D, Trifunov S, Andrić A, Markov Z, Jovičić S, Mudri Stojnić S,
 468 Janković M, Lugonja P (2016a) Prime Hoverfly (Insecta: Diptera: Syrphidae) Areas (PHA) as a conservation tool in
 469 Serbia. *Biol Conserv* 198:22–32. doi: 10.1016/j.biocon.2016.03.032

470 Vujić A, Radenković S, Stahls G, Ačanski J, Stefanović A, Veselić S, Andrić A, Hayat R (2012) Systematics and
 471 taxonomy of the *ruficornis* group of genus *Merodon* Meigen (Diptera: Syrphidae). *Syst Entomol* 37:578–602. doi:
 472 10.1111/j.1365–3113.2012.00631.x

473 Vujić A, Šimić S, Radenković S (2001) Endangered species of hoverflies (Diptera: Syrphidae) on the Balkan
 474 Peninsula. *Acta Entomologica Serbica* 5:93–105.

475 White AJ, Wratten SD, Berry NA, Weigmann U (1995) Habitat manipulation to enhance biological control of
 476 brassica pests by hover flies (Diptera: Syrphidae). *J Econ Entomol* 88:1171–1176. doi:
 477 <https://doi.org/10.1093/jee/88.5.1171>

478 Wickramasinghe LP, Harris S, Jones G, Jennings VN (2004) Abundance and species richness of nocturnal insects on
 479 organic and conventional farms: effects of agricultural intensification on bat foraging. *Conserv Biol* 18:1283–1292.
 480 doi: 10.1111/j.1523–1739.2004.00152.x

481 Williams SE, Shoo LP, Isaac JL, Hoffmann AA, Langham G (2008) Towards an integrated framework for assessing
 482 the vulnerability of species to climate change. *Plos Biol* 6:2621–2626. doi: 10.1371/journal.pbio.0060325

483 Wilson RJ, Gutiérrez D, Gutiérrez J, Martínez D, Agudo R, Monserrat VJ (2005) Changes to the elevational limits
 484 and extent of species ranges associated with climate change. *Ecol Lett* 8:1138–1146. doi: 10.1111/j.1461–
 485 0248.2005.00824.x

486 Wulff AS, Hollingsworth PM, Ahrends A, Jaffré T, Veillon JM, L’Huillier L, Fogliani B (2013) Conservation
 487 priorities in a biodiversity hotspot: analysis of narrow endemic plant species in New Caledonia. *PLoS One*
 488 8:e73371. doi: 10.1371/journal.pone.0073371

489 Yates CJ, McNeill A, Elith J, Midgley GF (2010) Assessing the impacts of climate change and land transformation
 490 on *Banksia* in the South West Australian Floristic Region. *Divers Distrib* 16:187–201. doi: 10.1111/j.1472–
 491 4642.2009.00623.x

Figure legends

Fig. 1 Contribution of bioclimatic variables related with temperature and precipitation in the final species distribution models of 44 analysed species of hoverflies in SE Europe. For detailed information on bioclimatic variables, visit www.worldclim.org

Fig. 2 Geopolitical map of SE Europe with significant localities (a) and projected potential species richness of hoverflies for (b) present, (c) 2050, (d) 2070, and differences between (e) 2050 and present and (f) 2070 and present. Each cell represents the total number of species in defined grid cells

Tables

Tab. 1 TSS values and pAOO values for all time periods, absolute and relative change in pAOO between present and projected future scenarios for 44 species of hoverflies in SE Europe

Tab. 2 Proportional loss of area and connection of range expansion and contraction patterns with altitude