<https://helda.helsinki.fi>

The Ordovician succession adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen

Kroger, Bjorn

2017-09-05

Kroger , B , Finnegan , S , Franeck , F & Hopkins , M J 2017 , ' The Ordovician succession adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen ' , American Museum Novitates , no. 3882 , pp. 1-28 . <https://doi.org/10.1206/3882.1>

<http://hdl.handle.net/10138/308335>

<https://doi.org/10.1206/3882.1>

cc_by

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

The Ordovician succession adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen

Björn Kröger¹, Seth Finnegan², Franziska Franeck³, Melanie J. Hopkins⁴

Abstract: The Ordovician sections along the western shore of the Hinlopen Strait, Ny Friesland, were discovered in the late 1960s and since then prompted numerous paleontological publications; several of them are now classical for the paleontology of Ordovician trilobites, and Ordovician paleogeography and stratigraphy. Our 2016 expedition aimed in a major recollection and reappraisal of the classical sites. Here we provide a first high-resolution lithological description of the Kirtonryggen and Valhallfonna formations (Tremadocian –Darriwilian), which together comprise a thickness of 843 m, a revised bio- and lithostratigraphy, and an interpretation of the depositional sequences. We find that the sedimentary succession is very similar to successions of eastern Laurentia; its Tremadocian and early Floian part is composed of predominantly peritidal dolostones and limestones characterized by ribbon carbonates, intraclastic conglomerates, microbial laminites, and stromatolites, and its late Floian to Darriwilian part is composed of fossil-rich, bioturbated, cherty mud-wackestone, skeletal grainstone and shale, with local siltstone and glauconitic horizons. The succession can be subdivided into five third-order depositional sequences, which are interpreted as representing the SAUK IIIB Supersequence known from elsewhere on the Laurentian platform.

¹ Finnish Museum of Natural History, University of Helsinki, Finland, bjorn.kroger@helsinki.fi

² Department of Integrative Biology, University of California, Berkeley, USA

³ Naturhistorisk Museum, University of Oslo, Norway

⁴ Division of Paleontology, American Museum of Natural History, USA

INTRODUCTION

The Ordovician sections along the western shore of the Hinlopen Strait (Hinlopenstretet), Ny Friesland, were discovered by accident in 1966, when an expedition team from the Cambridge University stopped to collect water from a melt stream and collected several well preserved trilobites (Fortey and Bruton, 2013). This discovery prompted two focused collection trips during the summers of 1967 (Vallance and Fortey, 1968) and 1972 (Fortey and Bruton, 1973) that resulted in a classic series of publications focused on the trilobite successions (e.g. Fortey, 1974a,b, 1975a,b, 1980a) and accompanying fauna (e.g. brachiopods: Hansen and Holmer, 2011; conodonts: Lehnert et al., 2013; graptolites: Fortey, 1971, Cooper and Fortey, 1982; heterostracans: Bockelie and Fortey, 1976; mollusks: Evans and King, 1990; Morris and Fortey, 1976; ostracods: Williams and Siveter, 2008; radiolarians: Maletz and Bruton, 2005). Some of these publications have been influential in shaping approaches to paleogeographical reconstructions: At Hinlopen Strait the interplay between depth-related biofacies (“community types” of Fortey, 1975b), paleogeographical provinces, and sea level fluctuation became evident (e.g., Cocks and Fortey, 1982). Other publications on the Hinlopen Strait succession had a significant impact on discussions about Ordovician stratigraphy. The Valhallan Stage was based on a section of the Hinlopen Strait; it was suggested in Fortey (1980a) to represent a time interval that previously has not been recognised in many North American sections because of a widespread hiatus on the continent. The Valhallan Stage, which is equivalent to the lower Dapingian Stage in modern terms, was never formally accepted, but its recognition was a significant step toward a supraregional stratigraphic correlation of this particular time interval (see Ross et al., 1997).

Despite the scientific importance of the Hinlopen Strait sections very few subsequent expeditions have been made. It took 34 years, after which in short sequence three expeditions made stops in the area, collected Ordovician samples and remeasured the sections: In 2005 a

group of the Polar Marine Geological Research Expedition (PMGRE) reappraised the sedimentary succession in context of a larger mapping project (Kosteva and Teben'kov, 2006); in 2007 a group from the German Bundesanstalt für Geowissenschaften und Rohstoffe (Federal Institute for Geosciences and Natural Resources) measured some sections at Hinlopen Strait and took biostratigraphic (conodont) samples in context of their project Circum-Arctic Structural Events 10 (CASE 10) (Lehnert et al., 2013); and in 2008 a Norwegian-Swedish Group lead by Nils-Martin Hanken of the University of Tromsø visited the area and made some focused paleontological collections, but had bad luck with weather/snow conditions and polar bears (Hansen and Holmer, 2010, 2011). Consequently, no major attempt to resample the sections at Hinlopen Strait has been made since 1972. Further, no attempt has been made by any previous expeditionary team to describe and reconstruct the paleoenvironmental succession in detail.

The goal of our 2016 expedition was to measure and describe in detail the lithology and stratigraphy of the Ordovician rocks at its two main outcrop sites Profilstranda and Olenidsletta (Fig. 1) and to collect paleontological and geochemical samples at high stratigraphic resolution. Herein, we provide the first results of this expedition, a detailed lithological description of the measured sections, a new high resolution stratigraphy, and an interpretation of the depositional sequences and its corresponding relative sea-level changes. The description will be the basis of a number of forthcoming publications with paleontological and geochemical focus.

GEOLOGICAL SETTING

The study area is on the northeastern edge of Ny Friesland on the island of Spitsbergen of the Svalbard archipelago, Norway, adjacent to the Hinlopen Strait (Hinlopenstretet), which divides Spitsbergen from Nordaustlandet (Fig. 1). This region comprises two exposure areas

of lower Paleozoic rocks of the Oslobreen Group, Hinlopenstretet Supergroup: one north (Basissletta) and one south (Olenidsletta) of the Buldrebreen arm of the Valhallfonna glacier at small cliffs along the coast line and in melt stream beds. The exposed sediments are subdivided into the Tokammane, Kirtonryggen, and Valhallfonna formations of the Oslobreen Group (Harland et al., 1966; Fortey and Bruton, 1973; Harland, 1997) and range from the Early Cambrian to the Middle Ordovician. The Kirtonryggen Formation comprises the Spora, Basissletta, and Nordporten members, and the Valhallfonna Formation is subdivided into the Olenidsletta and Profilbekken members (Fortey and Bruton, 1973, see details below). The Kap Sparre Formation in Nordauslandet on the eastern side of Hinlopenstretet is correlative to the Oslobreen Group, but high-resolution comparisons of the two are hampered by a lack of biostratigraphic data in the former (Stouge et al., 2011). The pre-Carboniferous basement of the Svalbard archipelago consists of a number of tectonostratigraphically distinct terranes that were stretched along the margin of Laurentia in pre-Caledonian times (Gee and Page, 1994; Gee and Teben'kov, 2004). The study area is part of the Nordaustlandet terranes of eastern Svalbard which in tectonic reconstructions are placed in close proximity to the Franz Joseph allochthon of North-Eastern Greenland (Smith and Rasmussen, 2008). During the mid-Paleozoic the sediments of the Hinlopenstretet Supergroup underwent minor folding with predominant NNW–SSE strike directions (Harland, 1997). The Hinlopenstretet Supergroup is generally little or not metamorphosed (Gee and Teben'kov, 2004).

In the southern outcrop area a roughly N/S directed Mesozoic dolerite intrusion limits the western expansion of the exposure of the early Paleozoic sediments. The intrusion is part of a complex of Late Jurassic – Early Cretaceous dolerites which are more widespread in the southern part of the Hinlopen Strait (Halvorsen, 1989).

99 The data presented herein are the result of a joint expedition of the authors (BK, FF, MH,
100 SF), and the camp manager Hårvard Kårstad during July and August, 2016. We traveled to
101 Ny Friesland by boat and built a campsite at Profilbekken (79°50'17.9"N / 017°42'19.9"W).
102 The Olenidsletta area was accessible to us from Profilstranda across the Buldrebreven via
103 rubber boat. During our expedition all visited sections were completely ice free.

104 We measured five sections using a Jacob's staff and clinometer, and tape measure (Fig. 1):

105 (1) The Spora River (SR) section and Profilstranda (PS) section form together a nearly
106 continuous outcrop of the Basissletta and Nordporten members. The base of the SR section is
107 within the Spora River channel just west of where the river crosses a small ridge formed by
108 massive pinkish dolostones of the top Tokammane Formation (79°51'51.0"N /
109 017°37'21.2"E). The SR section top is the topmost stromatolite bed at the mouth of the
110 Spora River (79°51'57.4"N / 017°38'33.2"E)

111 (2) The top of the SR section at the mouth of the Spora River is also the base of the PS
112 section. We measured the upper part of the Basissletta Member and the complete Nordporten
113 Member southward along the Profilstranda coastline. The outcrop is nearly continuous in its
114 northern part with a minor fault and exposure gap at 59 m from the base of the section
115 (79°51'41.2"N / 017°39'51.5"E). A number of major exposure gaps occur higher up in the
116 Basissletta Member ca. 50 m - 200 m just north of the small headland that forms the base of
117 the Nordporten Member (79°51'35.1"N / 017°40'36.8"E). There the dolostones are covered
118 by several meter thick layers of modern beach gravel. From the base of the Nordporten
119 Member up to the Olenidsletta Member the outcrop is continuous. The top of the PS section
120 is the uppermost prominent hardground within the uppermost Nordporten Member at
121 79°51'02.9"N / 017°41'24.0"E.

(3) The top of the PS section is the base of the overlying PO section. The PO section is just the southward continuation of the Profilstranda coastline outcrop and ends at a mouth of a small meltstream at 79°50'49''N / 017°42'04''E. The base of the Olenidsletta Member of the Valhallfonna Formation occurs within the first 5 meters of the PO section.

(4) We measured a part of the Profilbekken Member along the Profilbekken river (section PR). The Profilbekken river cuts through the Profilbekken Member beds at a very low angle and the beds form a number of repeating folds and faults which makes a reliable thickness measurement very difficult for a large part of the section. However, it was possible to measure a continuous ca. 45 m thick log from a place ca. 100 m upstream from the river mouth, where massive skeletal grainstone beds form a shallow ridge. This ridge is identical with the "basal algal conglomerate" ridge of Fortey (1980: 17); its top forms the top of the PR section at 79°50'36.1''N / 017°43'04.5''E.

(5) We measured a complete section of the Profilbekken Member at the southern end of the Olenidsletta area. The top of the outcrop and the youngest beds of the Valhallfonna Formation occur at promontory F where the beds form a shallow syncline (79°46'43''N / 017°54'20''E; Fig. 1). We followed the anastomosing meltstream just NW of promontory F in SW direction ca. 300 m upstream, which is stratigraphically downward. The outcrop is partly heavily weathered and in its upper part secondarily dolomitized.

DESCRIPTION OF SECTIONS

TOKAMMANE FORMATION

General features: The uppermost ca. 40 m of the Tokammane Formation consist of a succession of three depositional units. The bases of these units are composed of buff-

weathering, argillaceous, wavy-bedded, bioturbated, dolo-mudstone, flat-pebble and other intraclastic conglomerates, and thin-bedded dolo-siltstone with symmetric fine-scaled ripples. The top of each unit is formed by massive mounds of pinkish dolo-mudstone / dolo-wackestone, which contain thrombolitic and fenestral textures, and local patches of intraclastic conglomerates. The mounds have diameters of several tens of meters and at Spora River are thickest at the top of the formation. Each depositional unit is capped by an unconformity.

Detailed description: The measured section at Spora River (Fig. 2) starts at a ca. 0.5 m-thick massive flat-pebble conglomerate, which has a fining-up tendency of intraclasts and an erosive 3 cm-thick irregular top. The overlying 5 m of buff-weathering, wavy-bedded dolo-mudstone are topped by a ca. 2 m-thick massive, pinkish, fenestral dolo-mudstone. In the top 0.15 m of this pinkish mudstone, domal stromatolites occur that are up to 0.5 m in diameter.

The second unit begins above an erosional base with ca. 10 m of a monotonous, buff-weathering, wavy-bedded, argillaceous dolo-mudstone. In the lowermost beds of this unit parallel tunnels of unknown burrowers with diameters of less than 10 mm are common. At the top of this dolostone, a more massive, pinkish unit above an erosional base marks the base of the third unit. In the lower 0.2 m, this unit is composed of patches of poorly rounded intraclastic conglomerates; above this is a 25 m-thick fenestral dolo-mud and dolo-wackestone, partly with a thrombolitic texture. Around 2 m into this unit (at 19 m in the section), this massive dolostone is interrupted by a 1 m-thick layer of thin-bedded, platy- to wavy-bedded, dolo-siltstone that contains areas with small scale, symmetrical ripples and patches of flat-pebble conglomerates. Above this dolo-siltstone is a covered interval of about 10 m, overlain by the top 13 m of the Tokammane Formation, which are formed by massive, pinkish, thrombolitic mounds (Fig. 3A).

KIRTONRYGGEN FORMATION

SPORA MEMBER

We measured the Spora Member at its type locality at Spora River, which we estimated to have a thickness of ca. 20 m. The base of the member is well defined by a prominent erosional surface with a more than 1 m-high relief, which cuts into the underlying Tokammane Formation. The lower ca. 10 m of the Spora Member consist of massive, dark-grey, burrow-churned, wavy-bedded, fossil-rich dolo-mudstone / dolo-wackestone. At its lower 2 m, black flint nodules and thin flint layers are abundant. At a position of ca. 3 m from the base, a distinctive ca. 0.3 m-thick intraclastic conglomerate with rounded ca. 10-30 mm large pebbles occurs. Fossil abundance peaks at ca. 5-7 m above the base of the member, and the fauna is dominated by small ophiletid gastropods (diameter ca. 20-30 mm) and ellesmerocerid and endocerid cephalopods. Toward the top, the dolostone becomes more light-colored, buff-weathering, partly burrow-mottled (Fig. 3B), and rich in distinct trace fossils that form a network of tunnels with diameters of ca. 10 mm (Fig. 4A). The top of the Spora Member is lithologically transitional toward the lower Basissletta Member.

BASISSLETTA MEMBER

General: The Basissletta Member, exposed at Spora River and Profilstranda, is ca. 289 m thick. We placed the base of the member above the uppermost ca. 1 m-thick burrow-mottled limestone ca. 20 m above the base of the Kirtonryggen Formation. Compared to the Spora Member, the lithology in the lower Basissletta Member is composed of purer, and more light-colored, buff-weathering, platy- to wavy-bedded, burrow-churned to homogenous, sucrose dolo-mud to dolo-siltstone. Burrow-mottled horizons and distinctive trace fossils are nearly absent. The middle part of the member is highly condensed with abundant erosional surfaces,

flat-pebble conglomerates, oolite beds, and horizons with massive dark flint nodules and columnar to dome-shaped stromatolite beds. In its upper part argillaceous beds dominate and interchange with skeletal dolo-wackestone to dolo-packstone that contain abundant gastropods and trilobites. The boundary with the overlying Nordporten Member is transitional.

Details: The lower ca. 58 m of the Basissletta Member consist of buff-weathering, mostly planar-bedded, homogenous to burrow-churned dolo-mudstone to dolo-siltstone. Within this interval two horizons occur that have a burrow-mottled texture and contain rare gastropods, cephalopods and sponge macrofossils (at ca. 15 m, 46 m above member base = at positions 80 m, 111 m in the SR section, respectively). The lowermost stromatolites occur as a compact, up to 0.2 m-thick layer on an erosional surface at ca. 123 m in the section. Columnar and dome-shaped stromatolites, ooid layers, intraclastic and flat-pebble conglomerates, cross-lamination textures, and erosional surfaces are common throughout the following 30 m (Figs 3C, 5). The most prominent erosional surface cuts as a more than 0.5 m thick intraclastic conglomerate with a relief of ca. 0.3 m into a wavy bedded dolostone with low domal stromatolites at ca. 71 m above the base of the member at 136 m in the SR section.

At the mouth of the Spora River, at the top of the SR section, ca. 88 m above the base of the Basissletta Member, the stromatolite-rich interval is succeeded by ca. 40 m of massive yellowish-grey weathering, fine-laminated dolo-siltstone, that is exposed along the coast of Profilstranda (Fig. 6A). This interval contains in some places hummocky cross stratification (Fig. 6D) and is partly rich in flint nodules. Flat-pebble and other intraclastic conglomerates occur at 101 m and 122 m above base of Basissletta Member (at 13 m and 34 m in PS section, respectively, Fig. 7). Toward the top of this interval dark-grey weathering, flint-rich, partly-nodular argillaceous dolostones and ribbon dolostone are more common.

At 135 m and 146 m above the base of the Basissletta Member (at 47 and 58 m in PS section, respectively), two horizons with low domal stromatolites, intraformational and flat-pebble conglomerates, oolites, and erosional surfaces occur. Remarkable are teepee structures in a pyrite-rich dolostone (Fig. 6C), that are underlain by a flat pebble conglomerate and overlain by a dark-weathering, argillaceous, nodular, bioturbated dolostone at ca. 138 m above the base of the Basissletta Member (at PS section 50 m). A ca. 0.5 m-thick oolite bed above a massive intraformational conglomerate and below a bed of stromatolites at 145 m (at PS section 57 m) serves as a good local marker horizon.

The upper part of the Basissletta Member is partly poorly exposed, covered or fault disturbed along the coast along Profilstranda (Fig. 7). The exposed parts can be subdivided into three units. The lower ca. 20 m-thick unit is predominantly a yellowish weathering, wavy-bedded, bioturbated dolo-mudstone rich in large flint nodules. The nodules are up to 0.4 m in diameter and 0.3 m in thickness. The middle ca. 30 m-thick unit is rich in intraclastic and flat-pebble conglomerates, erosional surfaces, channels, cross bedding textures and contains predominantly light yellowish-grey, burrow-mottled, nodular dolostone and ribbon dolostone. Toward the top of this middle unit, hardgrounds locally occur on the top of intraclastic, channel rich dolostone beds. The topmost flat-pebble conglomerate of the Basissletta Member occurs at 198 m above the base of the member (at PS section 110 m). The uppermost unit of the Basissletta Member is a ca. 90 m thick, partly covered alternation of beds of yellowish-grey, argillaceous, wavy-bedded to nodular, burrow-churned dolo-mudstone, more massive stylolitic burrow-churned dolo-wackestone, and tempestitic skeletal - intraclastic dolo-packstone with common trilobite hash and gastropods.

NORDPORTEN MEMBER

243 *General:* The Nordporten Member, exposed at Profilstranda (Fig. 7), is ca. 217 m thick.
244 We placed the base of the member at the basal bed of a succession of massive reddish-grey
245 weathering, burrow-churned dolo-mudstone/dolo-wackestone, rich in gastropods and
246 trilobites, which forms the northernmost point of a small headland at Profilstranda
247 (79°51'35.1''N / 017°40'36.8''E). This bed forms the upper transitional part of an alternation
248 of massive dolo-wackestone / dolo-mudstone and argillaceous mudstone. The transition from
249 the Basissletta Member is gradual into a ca. 50 m-thick more massive succession of burrow-
250 mottled dolostone to ribbon-dolostone with erosional-topped / hardground-based repetitive
251 units. Above this succession another ca. 50 m are principally very similar but contain more
252 commonly hardgrounds, prominent grainstone- packstone lithologies, and scattered
253 intraclastic horizons. The upper ca. 120 m of the Nordporten Member consist predominantly
254 of monotonous massive grey to brownish burrow-churned, cherty dolo-mudstone / dolo-
255 wackestone. Toward the top, the succession gets more massive and hardground horizons are
256 more common.

257 *Details:* The base of the Nordporten Member is drawn at the base of a ca. 5 m-thick
258 massive dolostone within the upper part of the transitional unit of alternating massive
259 dolostone and nodular- to wavy-bedded argillaceous mudstone. At a position of ca. 11 m (PS
260 section 209 m) below a prominent erosional surface / hard ground with an underlying 0.2 m
261 oolite. This discontinuity surface is the lowest of six similar erosional surfaces (at 20 m, 29
262 m, 31 m, 34 m, 44 m above member base = at PS section 219 m, 227 m, 229 m, 232 m, 242
263 m, respectively, Fig. 8), each of them with an erosional relief of less than 0.3 m, that are
264 associated with horizons of intraclastic conglomerates and / or oolite beds. On the top of
265 some of these discontinuity surfaces, characteristic micro-mud mounds or micro-bioherms
266 with diameters of ca. 0.3 m and thicknesses of less than 0.2 m occur that commonly contain
267 trilobites and gastropods. The micro-bioherms are embedded in a matrix of planar-bedded,

greenish-grey, argillaceous mudstone, and form the base of the six repetitive units. In each of the units the basal mudstone grades into wavy-bedded to nodular, greenish-grey mudstone to ribbon-dolostone, and finally into massive, burrow-churned dolo- mudstone/ dolo-wackestone. The more argillaceous intervals commonly contain networks of unbranched burrows of the *Gordia* trace fossil type (Fig. 4B) and hummocky cross-stratification. At ca. 56 m above the base of the Nordporten Member (PS section 254 m), a ca. 0.4 m-thick intraclastic conglomerate bed marks the base of a gradual facies transition towards more massive dolostones with common intraclast horizons, grainstone, and packstone layers. At ca. 65 m above the base of the member (PS section 263 m), a ca. 1.5 m-thick pair of brachiopod-rich (*Hesperonomia* sp.) grain-packstone layers forms a marker horizon, which can be seen as the climax of this intraclast-rich succession. The most pronounced erosional surface cuts with a relief of more than 1 m into an underlying dolo-packstone / dolo-grainstone lithology at 79 m above the base of the member (PS section 277 m, Figs 7, 8B). The uppermost of these discontinuity-capped packstone/grainstone layers occurs at 116 m above the base of the Nordporten (PS section 314 m) within a ca. 50 m-thick succession very rich in chert nodules. The top of a gradual facies change is marked by a prominent hardground at 162 m above the base of the Nordporten Member (PS section 360 m) leading to ca. 60 m of very massive, grey-weathering, nodular, heavily bioturbated mudstone / wackestone lithologies with common discontinuity surfaces and hardgrounds that are most densely concentrated near the top of the member. The top 3 m of the formation are very rich in cephalopods and gradually change toward darker, more argillaceous mudstone lithologies.

VALHALLFONNA FORMATION

OLENIDSLETTA MEMBER

General: The Olenidsletta Member comprises a succession of dark limestone and black mudstone with a transitional lithology at its lower boundary. At Profilstranda the thickness of the Olenidsletta Member is approximately 160 m (Fig. 9). A precise thickness specification is impossible for the Profilstranda section because the uppermost part is locally covered and the boundary interval crops out only a few 100 meters SW in the Profilbekken River section. We placed the base of the Olenidsletta Member 3 m within the cephalopod-rich limestone that marks the transition interval from the Nordporten to the Olenidsletta Member (79°51'02.9''N / 017°41'24.0''E.). This boundary bed marks the top of the underlying massive, hardground-rich limestone unit of the Nordporten Member and is 6 m below the uppermost cephalopod occurrence within the transitional interval. The top of the member is not now exposed at Profilstranda, but at the nearby Profilbekken River, where it can be traced in several places across repeating fold sections (see below). Lithologically, the Olenidsletta Member can be roughly subdivided into four transitional intervals. The lowermost 5 meters are predominantly composed of dark, massive, stylolitic, bioturbated lime-mudstone with cephalopod coquina / cephalopod packstone interlayers. The overlying ca. 87 meters are an alternation of densely laminated, dark lime-mudstone and bituminous black shale with bedding thicknesses varying between 0.05 - 0.3 m, which are intermittently rich in graptolites. This shale-rich interval is overlain by a ca. 35 m-thick interval of a more carbonaceous limestone/shale alternation with abundant hardgrounds and beds of burrow-churned to nodular limestone, rich in trilobites, cephalopods, and inarticulate brachiopods. The upper ca. 30 m of the member are again dominated by black shale intervals.

Details: The lower part of the Olenidsletta Member is characterized by a gradual disappearance of bioturbated limestone horizons, an increase of black shale intervals and a gradual decrease in abundance of cephalopods and trilobites. A collection made at 1.5 m above the base of the member included the trilobites *Tropidopyge alveus* and *Carolinites*

317 *genacinaca nevadensis*, confirming that this part of the section is in the lowermost interval of
 318 the Olenidsletta Member (V1a of Fortey, 1980). At 11 m, trilobites of the overlying V1b
 319 interval (*Balnibarbi ceryx*, *Psilocara patagiatum*) were found. No cephalopods were found in
 320 any beds between 5 m to 97 m above the base of the member. A peak in graptolite abundance
 321 occurs ca. 70-90 m above the base of the Olenidsletta Member. The presence of *Balnibarbi*
 322 *pulvurea* and *Balnibarbi erugata* at 73 m and *B. pulvurea* at 90 m above the base of the
 323 member indicates that this part of the section is in the V1c interval of Fortey (1980). A
 324 conspicuous flint layer that serves as a local marker horizon occurs at 87 m above the base of
 325 the member. A flat conspicuous hardground at 94.4 m above the base of the member caps a
 326 ca. 1.5 m-thick bioturbated limestone interval with thickening up tendency. This hardground
 327 marks the onset of a succession of more than a dozen very similar repetitive units. Many of
 328 these units start with bituminous black shale rich in large inarticulate brachiopods, that grade
 329 into massive, nodular, bioturbated lime-mudstone lithologies containing abundant large
 330 trilobites and orthoconic cephalopods. At 99 m above the member base, the hard ground
 331 marking the top of one of the repetitive units is overlain by a thin intraclastic conglomerate.
 332 At 119 m, and 126-129 m, the limestone contains abundant flint nodules. The bituminous
 333 lower sections of the repetitive units are especially rich in large inarticulate brachiopods
 334 (mostly *Ectenoglossa*) in the interval 95-110 m, and 123 m above the base of the member.
 335 This interval (V2a of Fortey, 1980) is also characterized by the large asaphid trilobite *Gog*
 336 *catillus*, the lowest occurrence of which was found at 103 m above the base of the member.
 337 The top layers of the repetitive units at 119 m, 121 m, and 128 m are exceptionally rich in
 338 cephalopods and trilobites and contain horizons with trilobite hash / cephalopod shell
 339 packstone. The orthoconic cephalopod shells are current aligned (Fig. 10A). We place the
 340 boundary between V2a and V2b of Fortey (1980) around 123 m above the base of the
 341 member (Fig. 9) based on the occurrence of late-form *Bienvillia stikta*, late-form

Symphysurus arcticus, and *Hypermecaspis sp* in the interval 123-125 m; this is supported by the occurrence of *Ampyx porcus* and *Lyrapyge ebriosus* (both V2a) at 120.3 m above the base of the member. The uppermost shale-rich part of the member begins above the hard ground at the top of one of the repetitive units at 128 m. The boundary between V2 and V3 is transitional (Fortey 1980), but we place it no higher than 129 m above the base of the member. In this part of the member, the bituminous black shale/limestone is especially rich in large olenid trilobites. The upper part of the member consists of repetitive units with thickness of up to 3 m, each topped by a massive decimeter-thick limestone bed, which is partly bioturbated and capped by a flat hardground. Small orthoconic cephalopods (less than 3 mm in diameter), trilobite fragments, and 3D preserved graptolites are common within the top limestone layers of each of the units (Fig. 10C).

Although not exposed along Profilstranda, the base of the Profilbekken Member is marked by an inarticulate brachiopod-rich phosphatic, skeletal packstone which represents a thin horizon covering the topmost of these repetitive units. This boundary layer is exposed in Profilbekken and near promontory F in the Olenidsletta area south of Buldrebreen (Fig. 1).

PROFILBEKKEN MEMBER

General: The thickness and the lithology of the Profilbekken Member varies across the outcrop area. The complete section of the Profilbekken Member exposed along the F-promontory melt stream (section FP, Fig. 11) is ca.157 m thick. The member is also well exposed at Profilbekken River, but difficult to measure, because of faulting and folding of the beds. A part of the middle Profilbekken Member could be measured at Profilbekken River, but the correlation is problematic because of the lack of common distinct marker beds (Fig. 12). At the FP section the Profilbekken Member largely consists of an alternation of well-

366 bedded, slightly silicified, yellowish-grey, banded lime-mudstone (Fig. 13A) and wavy-
367 bedded, bioturbated to nodular grey lime-mudstone, with a few prominent hardground
368 horizons and more argillaceous greenish-grey intervals. In contrast, the section at
369 Profilbekken River is divided into distinct silty-glaucinitic beds and horizons dominated by
370 intraclastic-skeletal grainstone and packstone lithologies. The upper boundary of the member
371 is only exposed in the southern outcrop area and at section FP it is drawn at the base of the
372 gray-green siltstone-shale unit that caps the massive limestone of the top Profilbekken
373 Member. The upper ca. 10 m of the Profilbekken Member are composed of a massive, light
374 grey weathering, bioturbated lime-mudstone / lime-wackestone with abundant hardgrounds
375 and omission surfaces. The top of the limestone succession is formed by a prominent
376 hardground and / or erosional surface with a ca. 0.1 m relief. A greenish bed of siltstone -
377 marl, rich in orchid brachiopods and pelmatozoan debris overlies this top hardground with an
378 exposed thickness of 4 meters and forms the youngest of the beds of the Valhallfonna
379 Formation.

380 *Details:* The base of the member is a thin (few centimeters) skeletal packstone layer, rich
381 in inarticulate brachiopod (obolids) shell hash and trilobite cuticle fragments (Fortey and
382 Bruton, 1973). The horizon has been found at section FP and in several places along the
383 Profilbekken River, where it was impossible to reconstruct a coherent profile. At
384 Profilbekken River it is evident that similar more or less phosphate-rich, thin, skeletal
385 packstone beds exist in the top few meters of the Olenidsletta Member. These beds contain
386 trilobite hash, brachiopod shell hash and small orthoconic cephalopods in a varying amount
387 and are representative of the termination of individual meter scale repetitive units with
388 thickening-up limestones at the top of the Olenidsletta Member (Fig. 10C). At section FP
389 (Figure section FP, Fig. 11), no such phosphate rich horizon exists above the base of the
390 Profilbekken Member but the dark, partly bituminous argillaceous-silty, laminated lime-

391 mudstone, characteristic of the upper Olenidsletta Member continues for ca. 3 m until it
392 grades into a 21 m-thick succession of well-bedded, grey lime-mudstone which is partly
393 silicified and during weathering forms characteristic yellowish-grey bands. Hardgrounds and
394 discontinuity surfaces occur in some places in this banded limestone. A prominent
395 hardground 11 m above the base of the member contains abundant large orthoconic endocerid
396 cephalopods and trilobite hash. The particular limestone below the hardground is well
397 bioturbated and irregularly bedded and directly above the hardground a thin intraclastic
398 conglomerate occurs. A similar prominent hardground horizon occurs at 33 m above the
399 member base. The monotonous alternation of banded silicified limestone and wavy bedded
400 bioturbated limestone at section FP is only interrupted by two more-argillaceous, greenish
401 weathering, partly flint, nodule-rich beds with transitional boundaries at 52-53 m and at 78-
402 79 m from the base. It is questionable if and how these two argillaceous limestone beds relate
403 to the distinct glauconite intervals that occur at Profilbekken River.

404 At Profilbekken River the lithology is much more variable in the middle part of the
405 member with two prominent glauconite intervals that serve as local marker horizons (Fig.
406 12). The upper glauconite interval consists of three distinct up to 1.5 m-thick glauconite-
407 siltstone beds at 24 m, 29 m, and 31 m (Fig.13B). Each of the glauconite beds caps a
408 prominent bored hardground on top of the respective underlying bioturbated massive
409 limestone. The lower glauconite interval at 4 m is comprised of a succession of 4-5
410 glauconite silt beds, each with a thickness of ca. 0.1 m. The hardgrounds of the lower
411 glauconite interval differ from those in the upper interval in being strongly iron stained (Fig.
412 13B). Generally the lower glauconite interval appears to be very rich in dispersed pyrite. The
413 lower glauconite interval also differs from the upper interval in containing abundant large
414 orthoconic endocerid cephalopods, trilobites, gastropods (Fig. 10D). Large monaxon sponge
415 spicules and orchid brachiopods are common in the limestone beds between the two

glaucinite intervals. Crinoid ossicles are abundant throughout the measured Profilbekken section, but clearly increase in abundance toward the top, where several up to 2 m-thick algal-pelmatozoan grainstone beds form the top of individual hardground-capped repetitive units.

STRATIGRAPHICAL SUBDIVISION AND CORRELATION

PROBLEMS IN LITHOSTRATIGRAPHIC SUBDIVISION

The lithostratigraphic subdivision of the sections described herein is based on units established by Gobbet and Wilson (1960), Harland et al. (1966), and Vallance and Fortey (1968). The formations and members are adopted without changes from Fortey and Bruton (1973). This is common practice in all publications about the Hinlopen Strait sections since the 1970s. Nonetheless, the thickness estimates given in the some of these publications vary dramatically. The most extreme differences exist between measurements of Kosteva and Teben'kov (2006) and Lehnert et al. (2013) with total thicknesses of the Kirtonryggen and Valhallfonna formations of 722 m and 1055 m, respectively. Our own measurement with 843 m is nearly 100 m more than the thickness given in Fortey and Bruton (1973).

The reason for these differences must not be sought for in dissenting practices of placing boundaries of the two formations because these two boundaries are unmistakably defined by Fortey and Bruton (1973). Instead we assume that most of the discrepancies result from differences in the combination of sections from either the northern Profilbekken or the southern Olenidsletta outcrop area and/or from differences in ice conditions. During the 1971 expedition David Bruton recorded much ice in the sections and ice was also present in the sections during the 1967 expedition (David Bruton, pers. comm). This is in contrast to our 2016 expedition with complete ice and snow free conditions. Our combined section of the Profilbekken and Olenidsletta area is based on the combination of one more or less

441 continuous outcrop in the northern area which starts at the Spora River in the north and
442 continues south along the shore of the Profilbekken and a second one along the F-promontory
443 melt stream in the southernmost part of the outcrop area. We did not find any major fault
444 related repetition or gap in these outcrops. But an additional short section of the Profilbekken
445 Member along the Profilbekken River illustrates the existence of significant lateral facies
446 changes between the northern and the southern outcrop area in the Profilbekken Member with
447 greatly reduced thickness in the northern outcrop area (Fig. 14). The opposite trend seems to
448 exist within the Olenidsletta Member: we measured a thickness of ca. 160 m in the northern
449 area whereas Fortey and Bruton (1973) report a total thickness of 145 m based on
450 measurements in the type area in the southern outcrop. We also found slightly different
451 thicknesses than Fortey and Bruton (1973). Our data of the Olenidsletta Member are in
452 general agreement with the biostratigraphy summarized in a pull out chart at the end of the
453 Fortey (1980) volume with all trilobite species listed. The Olenidsletta Member trilobite
454 range zones V1, and V2 are reported to have a thickness of 77 m, 29 m, respectively in
455 Fortey (1980), measured at the Olenidsletta type section. These correspond to our thicknesses
456 of 95 m, 34 m measured at Profilstranda (Fig. 9).

457 It continues to be impossible to create a combined section exclusively from either the
458 northern or southern outcrop area. Hence, any combined section of the Hinlopen Strait
459 Ordovician reflects an individual synthesis across a relatively large area with significant
460 facies and thickness changes.

461 Additionally it can be expected that discrepancies in the individual attempts to correlate
462 between the sections led to differences in the composite log; e.g., the largest difference
463 between our measurement and that of Fortey and Bruton (1973) is within the Basissletta
464 Member. We measured a total thickness of 289 m for the Basissletta Member whilst Fortey
465 and Bruton (1973) measured 250 m at the same section. This difference is best explained by

improved outcrop conditions since the 1970s due to less ice and snow coverage. In our section the Basissletta Member contains two intervals with stromatolites; one crops out at ca. 60–90 m above the base of the member within the Spora River and another at ca. 140–150 m above the base along the Profilbekken shoreline. The Fortey and Bruton (1973) section contains only our lower stromatolite interval. But the Profilbekken shoreline, which was easily accessible to us, was difficult to access 44 years ago (Fortey and Bruton, 1973: 2232). We did not find any signs of a major tectonic repetition of the section and are confident that the Basissletta Member contains two stromatolite-rich intervals. In our opinion the missing ca. 40 m of section between two measurements are a result of a missing middle part of the Basissletta Member in the Fortey and Bruton (1973) section.

A third source of discrepancies is the sometimes ambiguous placing of the member boundaries in section intervals with gradual lithology changes. The original description in Fortey and Bruton (1973) is explicit, but did not prevent quite permissive reinterpretation by some authors. The base of the Nordporten Member, e.g., is defined as “a 10 m-thick, massive, fossiliferous gray limestone forming a small headland 2 km to the north of the Profilbekken” (Fortey and Bruton, 1973: 2232). But it appears that Lehnert et al. (2013) placed their Nordporten Member base at a position ca. 80 m down in the section at a place some hundred meters toward the north of this small headland.

It is important to have these discrepancies in mind when comparing individual fossil occurrences or lithologies from the different published composite sections. These discrepancies and differing practices in logging also set important constraints for high resolution biostratigraphic and chronostratigraphic correlations.

PROBLEMS IN BIOSTRATIGRAPHY AND CORRELATION WITH CHRONOSTRATIGRAPHY

The biostratigraphy of the Kirtonryggen and Valhallfonna formations is primarily based on trilobites, and was first established in great detail within the Valhallfonna Formation with nine successive trilobite faunal (or assemblage) zones (Fortey, 1980). The list of trilobite faunal zones was completed for the Kirtonryggen Formation by Fortey and Bruton (2013). The conodont stratigraphy is based on Fortey and Barnes (1977), Lehnert et al. (2013), and few samples discussed in Fortey and Bruton (2013). A graptolite based stratigraphy was published only for the Olenidsletta Member (Archer and Fortey, 1974; Cooper and Fortey, 1982).

Conflicting biostratigraphic data occur partly within the Kirtonryggen Formation. The Spora Member contains trilobites which have been correlated with the Laurentian *Leiostegium*–*Tesselecauda* trilobite zones of Ross et al. (1997), Stairsian Regional Stage, by Fortey and Bruton (2013). In contrast, the only productive conodont sample from the Spora Member, comes from the very top of the member and contains a fauna of the Laurentian *Rossodus manitouensis* conodont zone, Skullrockian-Stairsian regional stages (Lehnert et al., 2013). Currently no additional fossil data are available to further constrain the age of the Spora Member.

The biostratigraphy of the Basissletta Member is problematic, because the fossils are rare and endemic. Trilobites collected at the base of the lower stromatolite interval (ca. 60 m above the base of the Basissletta Member) and in fossil-poor dolo-mudstones ca. 104 m above the base of the Basissletta Member are interpreted to represent an interval that is not deposited elsewhere on Laurentia (Fortey and Bruton, 2013). Conodonts recovered from the top of the lowermost stromatolite interval $\pm$ 90 m above the base of the member contain the index taxon of the *Macerodus diana*e conodont zone (sample BS122 of Lehnert et al., 2013). A sample from the topmost Basissletta Member contains conodonts indicative of the Laurentian *Oepikodus communis* conodont zone (Fortey and Bruton, 2013: 9-10) and

trilobites in the uppermost Basissletta–lowermost Nordporten Member can be correlated with the Laurentian *Benthamaspis rochmotis*–*Petigurus cullisoni* trilobite zones interval (Fortey and Bruton, 2013: 14), which in turn correlates with the lower part of the *Oepikodus communis* conodont zone. Hence, the base of the Floian Stage can be expected within the upper part of, but not at the top of, the Basissletta Member.

The biostratigraphy of the Nordporten to lower Profilbekken members is well resolved and the trilobite, graptolite, and conodont data are consistent. But few data are available for the Profilbekken Member. Conodonts collected from a bed in between the two main glauconite horizons at Profilbekken River indicate a stratigraphic position within the *Paroistodus originalis* - *Baltoniodus norrlandicus* conodont zones and within the *Isograptus victoriae maximodivergens* - *Levisograptus* graptolite zones (sample BS66 of Lehnert et al., 2013) near the Dapingian - Darriwilian stage boundary. The trilobites above a level of ca. 30 m in the Profilbekken Member section of Fortey and Bruton (1973) are considered to be correlative with the Whiterockian *Orthidiella* trilobite zone of Laurentia, but this level could not be located with confidence in our sections, and it is questionable if it is below BS66 of Lehnert et al. (2013) at the position near the lowermost glauconitic horizons.

This short overview reveals that large intervals of the Kirtonryggen and Valhallfonna formations need a better biostratigraphic resolution, but with the current data the possibilities are limited. A higher biostratigraphic resolution of the sections is also crucial for a reliable interregional correlation of the depositional sequences of the Hinlopen sections.

SEQUENCE STRATIGRAPHY

The succession of the Kirtonryggen and Valhallfonna formations is in several aspects very similar to time equivalent carbonate successions of the eastern paleomargin of Laurentia, and

539 can be directly compared with the Early to early Middle Ordovician carbonates of the central
540 Appalachian Basin (compare Pope and Read, 1997; Brezinski et al., 2012) and of western
541 Newfoundland (compare Pratt and James, 1986, Knight and James, 1987; Knight et al.,
542 2007). The Kirtonryggen Formation is mainly composed of fossiliferous bioturbated mud-
543 wackestone, ribbon carbonate, intraclastic (predominately flat-pebble) conglomerate,
544 grainstone and oolite, and microbial laminite. These lithologies form units of shallowing-up
545 meter- to tens-of-meter-scale parasequences.

546 In comparison, the Lower Ordovician part of the Knox Group of the central Appalachians
547 consists of a number of third-order depositional sequences that are composed of dozens of
548 meter-scale shallowing upward peritidal parasequences with common flat-pebble
549 conglomerates and oolites at their base, low domal stromatolites in their middle parts, and
550 cryptal laminites at their top. The deeper depositional settings of this part of the Knox Group
551 consists of subtidal ribbon carbonates and more massive microbial bioherms (Pope and Read,
552 1998). On western Newfoundland the Lower Ordovician part of the St. George Group
553 consists of peritidal carbonates that range from supratidal cryptalgal laminates, intertidal
554 ribbon carbonates, and stromatolites to deeper subtidal thrombolite mounds and fossiliferous
555 wackestone (Pratt and James, 1986).

556 This general similarity of the carbonate successions of eastern Laurentia, including those
557 of Spitsbergen, is even more compelling when considering their general change in
558 sedimentation style toward the Middle and Late Ordovician. The uppermost parts of the
559 Nordporten Member and Olenidsletta Member record a substantial rise in sea level
560 accompanied by a change of the lithology toward dark, bituminous mud-wackestone and
561 shale lithologies. At the top of the Olenidsletta Member and within the Profilbekken Member
562 the sea level returned to levels comparable to that of the Nordporten Member, but the
563 lithology does not return to a Kirtonryggen facies. Instead the Profilbekken lithologies are

564 dominated by massive mud-wackestone, partly rich in dispersed silica and/or chert, skeletal
565 pack-grainstone (rich in pelmatozoan ossicles and algal fragments), shale-siltstone, and
566 glauconite-rich beds. In the Profilbekken Member, the limestone forms parasequences with
567 thicknesses of a few to tens of meters with shale-siltstone beds at their bases, mud-
568 wackestone beds as main parts, and partly skeletal grain-packstone layers at their top. The
569 parasequences are capped by prominent hardgrounds, which are heavily bored on, and may
570 be phosphate or iron encrusted. With the exception of the glauconite beds, this facies is very
571 similar to the High Bridge and St. Paul Group of Kentucky and Virginia (Pope and Read,
572 1998). This Early to Late Ordovician facies change across eastern Laurentia has been
573 interpreted as reflecting major climatic change during the Ordovician (Pope and Read, 1998;
574 Pope and Steffen, 2003).

575 Despite this general similarity in facies pattern it is difficult to correlate individual
576 depositional sequences of the Kirtonryggen and Valhallfonna formations with that other
577 regions of Laurentia. Five third-order depositional sequences can be distinguished within the
578 Kirtonryggen and Valhallfonna formations (Fig. 14). Generally, the five sequences can be
579 interpreted as part of the Laurentian SAUK IIIB Supersequence (Morgan, 2012), which is
580 sandwiched above the prominent basal Tremadocian Stonehenge transgression (Taylor et al.
581 1992) and below the Darriwilian base of the Tippecanoe Supersequence. The exact
582 stratigraphic range of the hiatus at the base of the Kirtonryggen Formation is not known, and
583 the onset of sedimentation of the Spora Member was either during the latest Skullrockian or
584 during the earliest Stairsian regional stages (see above). But it is clear that the fossil-rich
585 bioturbated dolostone of the Spora Member represents a deep setting that is not represented
586 again until the upper part of the Basissletta Member. The interval between the Spora Member
587 and the upper Basissletta Member contains two major shallowing events; both are Stairsian in
588 age. Therefore, the Spora Member most likely represents the late part of the Stonehenge

transgression, and the two successive lowstand intervals within the Basissletta Member can be interpreted as equivalents of the widespread Laurentian early Stairsian unconformity (e.g.: top Stonehenge Formation, central Appalachians; base Boat Harbor Formation, western Newfoundland; base Rochdale Formation, New York; Tule Valley lowstand, Utah; see Morgan, 2012), and late Stairsian unconformity (e.g.: Rochdale/Fort Cassin formations unconformity, New York; Boat Harbor unconformity, western Newfoundland; see Morgan, 2012), respectively. Consequently, our sequence I would be roughly equivalent with the upper part of the Lower Boat Harbour Formation in western Newfoundland, and our sequence II would be partly equivalent with the middle Boat Harbor Formation in western Newfoundland, and the Rochdale Formation in New York (Fig. 15). This correlation is consistent with the biostratigraphic data of the Basissletta Formation (Fortey and Bruton, 2013; Lehnert et al., 2013). A detailed correlation of the overlying sequence III is not possible with the available data, but the massive deepening at the topmost Nordporten Member / basal Olenidsletta Member can be confined with some confidence to a prominent early Floian Laurentian transgression around the *O. communis* / *O. evae conodont* zone boundary (e.g. Lagnet Point Member, western Newfoundland; Fort Cassin Formation, New York; *evae*-transgression, northeast Greenland; see Morgan, 2012). The massive sea level drop at the lower Profilbekken Member is in accordance with the “basal widespread Whiterockian regression” and a succession of unconformities at the Floian/Dapingian boundary ultimately mark the top of the Laurentian SAUK supersequence (Morgan, 2012).

CONCLUSIONS

The Kirtonryggen and Valhallfonna formations comprise 843 m of mostly carbonaceous Early to early Middle Ordovician sediments. The sedimentary succession is in several aspects very similar to other successions of eastern Laurentia; its Tremadocian and early Floian part

is composed of predominantly peritidal dolostones and limestones characterized by ribbon carbonates, intraclastic conglomerates, microbial laminites, and stromatolites, and its late Floian to Darriwilian part is composed of fossil-rich, bioturbated, cherty mud-wackestone, skeletal grainstone and shale, with local siltstone and glauconitic horizons. The succession would be consistent with a general trend of Early to Middle Ordovician climate cooling.

Lateral facies differentiation complicates the local correlation of the upper Valhallfonna Formation, especially in the absence of a high-resolution biostratigraphy on this part of the succession. The biostratigraphic resolution of the Kirtonryggen and Valhallfonna formations greatly varies, depending on the position within the succession. The more restricted, shallow peritidal carbonates of the Basissletta Member contain a sparse and endemic assemblage, which cannot be directly correlated with other Laurentian areas. A comparatively high-resolved biostratigraphy is possible within the graptolite rich shales and mud-limestones of the Olenidsletta Member.

Within the Kirtonryggen and Valhallfonna formations, five third-order depositional sequences can be subdivided, and which are interpreted as representing the SAUK IIIB Supersequence known from elsewhere on the Laurentian platform. A detailed correlation of the individual third-order sequences is currently difficult because of limited biostratigraphic control, but the available data suggest that especially the Stairsian and Middle Ordovician lowstand intervals are much more complete than in other Laurentian sections.

ACKNOWLEDGEMENTS

We are grateful for the general support during preparation and realization of the project by David Bruton (NHM Oslo, Norway), Ebony (Longyearbyen, Svalbard, Norway), Ethan (Profilbekken, Svalbard, Norway), Richard Fortey (NHM London), Nils Martin Hanken

638 (Tromsø / Oslo, Norway), Håvard Hansen (Norsk Polarinstitut, Svalbard, Norway), Stig
 639 Hennigsen and family (Longyearbyen, Svalbard, Norway), Jim Johansen (Norsk Polarinstitut,
 640 Svalbard, Norway), Håvard Kårstad (Ørsta, Norway), Henriika Kivilä (Helsinki, Finland),
 641 Rupert Krapp (Norsk Polarinstitut, Svalbard, Norway), Lee Hsiang Liow (NHM Oslo,
 642 Norway), Ivar Midtkandal (Oslo, Norway), Hans Arne Nakrem (NHM Oslo, Norway), Svend
 643 Stouge (Københavns Universitet, Denmark), Mariah Slovacek (AMNH, USA), and thank
 644 also the Syssemlannen på Svalbard for Permissions of Fieldwork. Fieldwork was conducted
 645 under permit 2016/00110-2. The project was financially supported by the Niarchos
 646 Foundation (MJH), the National Geographic Society (FF), and the Societas Scientiarum
 647 Fennica (BK). This work is part of Research in Svalbard (RIS) ID 10467.

648

649 REFERENCES

- 650 Archer, J.B., and R.A. Fortey. 1974. Ordovician graptolites from the Valhallfonna Formation,
 651 northern Spitsbergen. *Special Papers in Palaeontology* 13: 87–98.
- 652 Bergström, S.M., X. Chen, J.C. Gutiérrez-Marco, and A. Dronov. 2009. The new
 653 chronostratigraphic classification of the Ordovician System and its relation to major
 654 regional series and stages and to $\delta^{13}\text{C}$ chemostratigraphy. *Lethaia* 42: 97–107.
- 655 Bockelie, T., and R.A. Fortey. 1976. An early Ordovician vertebrate. *Nature* 260: 36–38.
- 656 Boyce, W.D., L.M.E. McCobb, and I. Knight. 2011. Stratigraphic studies of the Watts Bight
 657 Formation (St. George Group), Port au Port Peninsula, western Newfoundland. *Current*
 658 *Research*. Government of Newfoundland and Labrador, Department of Natural
 659 Resources, Mines Branch, Report, 11–11.
- 660 Brezinski, D.K., J.F. Taylor, and J.E. Repetski. 2012. Sequential development of platform to
 661 off-platform facies of the great American carbonate bank in the central Appalachians. *In*

662 J.R. Derby, R.D. Fritz, S.A. Longacre, W.A. Morgan, and C.A. Sternbach, (editors), The
 663 Great American Carbonate Bank: The geology and economic resources of the Cambrian-
 664 Ordovician Sauk megasequence of Laurentia: 383–420. American Association of
 665 Petroleum Geologists.

666 Cocks, L.R.M., and R.A. Fortey. 1982. Faunal evidence for oceanic separations in the
 667 Palaeozoic of Britain. *Journal of the Geological Society* 139: 465–478.

668 Cooper, R.A., and R.A. Fortey. 1982. The Ordovician graptolites of Spitsbergen. *Bulletin of*
 669 *the British Museum (Natural History)*, *Geology* 36: 157–302.

670 Cooper, R.A., P.M. Sadler, O. Hammer, and F.M. Gradstein. 2012. Chapter 20 - The
 671 Ordovician Period. *In* Gradstein, F.M., J.G.O.D. Schmitz, and G.M. Ogg, (editors), *The*
 672 *Geologic Time Scale 2012*: 489–523. Boston: Elsevier.

673 Evans, D.H., and A.H. King. 1990. The affinities of early oncocerid nautiloids from the
 674 Lower Ordovician of Spitsbergen and Sweden. *Palaeontology* 33: 623–630.

675 Fortey, R.A. 1974. The Ordovician Trilobites of Spitsbergen: 1. Olenidae. *Skrifter Norsk*
 676 *Polarinstitut* 160: 1–81.

677 Fortey, R.A. 1975a. Early Ordovician trilobite communities. *Fossil and Strata* 4: 331–352.

678 Fortey, R.A. 1975b. The Ordovician Trilobites of Spitsbergen: 2. Asaphidae, Nileidae,
 679 Raphiophoridae and Telephinidae of the Valhallfonna Formation. *Skrifter Norsk*
 680 *Polarinstitut* 162: 1–125.

681 Fortey, R.A., and D.L. Bruton. 2013. Lower Ordovician trilobites of the Kirtonryggen
 682 Formation, Spitsbergen. *Fossils and Strata* 59: 1–116.

683 Fortey, R.A. 1971. *Tristichograptus*, a triserial graptolite from the Lower Ordovician
 684 of Spitsbergen. *Palaeontology* 14: 188–199.

- 685 Fortey, R.A. 1980. The Ordovician Trilobites of Spitsbergen: 3. Remaining trilobites of the
686 Valhalla Formation. *Skrifter Norsk Polarinstitut* 171: 1–113.
- 687 Fortey, R.A., and D.L. Bruton. 1973. Cambrian-Ordovician rocks adjacent to Hinlopenstretet,
688 North Ny Friesland, Spitsbergen. *Bulletin of the Geological Society of America* 84:
689 2227–2242.
- 690 Fortey, R.A. 1974. A new pelagic trilobite from the Ordovician of Spitsbergen, Ireland and
691 Utah. *Palaeontology* 17: 111–124.
- 692 Gee, D.G., and L.M. Page. 1994. Caledonian terrane assembly on Svalbard: new evidence
693 from $^{40}\text{Ar}/^{39}\text{Ar}$ dating in Ny Friesland. *American Journal of Science* 294: 1166–1186.
- 694 Gee, D.G., and A.M. Teben'kov. 2004. Svalbard: a fragment of the Laurentian margin.
695 Geological Society, London, *Memoirs* 30: 191–206.
- 696 Gobbett, D.J., and C.B. Wilson. 1960. The Oslobreen Series, Upper Hecla Hoek of Ny
697 Friesland, Spitsbergen. *Geological Magazine* 97: 447–460.
- 698 Halvorsen, E. 1989. A paleomagnetic pole position of Late Jurassic/Early Cretaceous
699 dolerites from Hinlopenstretet, Svalbard, and its tectonic implications. *Earth and*
700 *Planetary Science Letters* 94: 398–408.
- 701 Hansen, J., and L.E. Holmer. 2010. Diversity fluctuations and biogeography of the
702 Ordovician brachiopod fauna in Northeastern Spitsbergen. *Bulletin of Geoscience* 85:
703 497–504.
- 704 Hansen, J., and L.E. Homer. 2011. Taxonomy and biostratigraphy of Ordovician brachiopods
705 from northeastern Ny Friesland, Spitsbergen. *Zootaxa* 3076: 1–122.

706 Harland, T.L., R.H. Wallis, and R.A. Gayer. 1966. A revision of the Lower Hecla Hoek
 707 succession in central north Spitsbergen and correlation elsewhere. *Geological Magazine*
 708 103: 70–97.

709 Harland, W.B. 1997. Chapter 7 Northeastern Spitsbergen. Geological Society, London,
 710 *Memoirs* 17: 110–131.

711 Knight, I., K. Azmy, M.G. Greene, and D. Lavoie. 2007. Lithostratigraphic setting of
 712 diagenetic, isotopic, and geochemistry studies of Ibexian and Whiterockian carbonate
 713 rocks of the St. George and Table Head groups, western Newfoundland. *Current*
 714 *Research, Newfoundland and Labrador Department of Natural Resources Geological*
 715 *Survey Report* 07-1: 55-84.

716 Kosteva, N.N., and A.M. Teben'kov. 2006. *Litologicheskaya kharakteristika kembrijsko-*
 717 *ordoviksikh otloshenij zapadnogo pobereshya proliva khinlopen, arhipelag spitsbergen.*
 718 *Complex investigations of Spitsbergen Nature* 6: 109–119.

719 Landing, E., J.A. Adrain, S.R. Westrop, and B. Kröger. 2011. Tribes Hill–Rochdale
 720 formations in east Laurentia: proxies for Early Ordovician (Tremadocian) eustasy on a
 721 tropical passive margin (New York and west Vermont). *Geological Magazine* 149: 93–
 722 123.

723 Lavoie, D., A. Desrochers, G. Dix, I. Knight, and O. Salad Hersi. 2012. The great American
 724 carbonate bank in Eastern Canada: An Overview. In Derby, J.R., R.D. Fritz, S.A.
 725 Longacre, W.A. Morgan, and C.A. Sternbach, (editors), *The Great American Carbonate*
 726 *Bank: The geology and economic reseources of the Cambrian-Ordovician Sauk*
 727 *megasequence of Laurentia: 499–523. American Association of Petroleum Geologists*
 728 *Memoir* 98.

729 Lehnert, O., S. Stouge, and P.A. Brandl. 2013. Conodont biostratigraphy in the Early to
730 Middle Ordovician strata of the Oslobreen Group in Ny Friesland, Svalbard. *Zeitschrift*
731 *der Deutschen Gesellschaft für Geowissenschaften* 164: 149–172.

732 Maletz, J., and D.L. Bruton. 2008. The Middle Ordovician *Provenocitum procerulum*
733 radiolarian assemblage of Spitsbergen and its biostratigraphic correlation. *Palaeontology*
734 51: 1181–1200.

735 Morgan, W.A. 2012. Sequence stratigraphy of the Great American Carbonate Bank. *In*
736 Derby, J.R., R.D. Fritz, S.A. Longacre, W.A. Morgan, and C.A. Sternbach, (editors), *The*
737 *Great American Carbonate Bank: The geology and economic resources of the Cambrian-*
738 *Ordovician Sauk megasequence of Laurentia: 37–79. American Association of Petroleum*
739 *Geologists Memoir* 98.

740 Morris, N.J., and R.A. Fortey. 1976. *Tironucola* gen. nov. and its significance in bivalve
741 evolution. *Journal of Paleontology* 50: 701–709.

742 Pope, M.C., and J.B. Steffen. 2003. Widespread, prolonged late Middle to Late Ordovician
743 upwelling in North America: A proxy record of glaciation? *Geology* 31: 63–66.

744 Pope, M., and J.F. Read. 1998. Ordovician metre-scale cycles: implications for climate and
745 eustatic fluctuations in the central Appalachians during a global greenhouse, non-glacial
746 to glacial transition. *Palaeogeography, Palaeoclimatology, Palaeoecology* 138: 27–42.

747 Pratt, B.R., James, N. P. 1986. The St. George Group (Lower Ordovician) of western
748 Newfoundland: tidal flat island model for carbonate sedimentation in shallow epeiric
749 seas. *Sedimentology* 33: 313–343.

750 Ross, R.J.J., L.F. Hintze, R.L. Ethington, J.F. Miller, M.E. Taylor, and J.E. Repetski. 1997.
751 *The Ibexian, Lowermost Series in the North American Ordovician. U.S. Geological*
752 *Survey Professional Paper* 1579-A: 1–49.

753 Smith, M.P., and J.A. Rasmussen. 2008. Cambrian–Silurian development of the Laurentian
754 margin of the Iapetus Ocean in Greenland and related areas. *Geological Society of*
755 *America Memoirs* 202: 137–167.

756 Stouge, S., Christiansen, J.L., and Holmer, L.E. 2011. Lower Palaeozoic stratigraphy of
757 Murchinsonfjorden and Sparreneset, Nordauslandet, Svalbard. *Geografiska Annaler*,
758 *Series A, Physical Geography* 93: 209–226.

759 Taylor, J.F., J.E. Repetski, and R.C. Orndorff. 1992. The Stonehenge transgression: A rapid
760 submergence of the central Appalachian platform in the Early Ordovician. In Webby,
761 B.D., and J.R. Laurie, (editors), *Global perspectives on Ordovician geology*: 409–418.
762 Balkema.

763 Vallance, G., and R.A. Fortey. 1968. Ordovician succession in North Spitsbergen.
764 *Proceedings of the Geological Society of London* 1648: 91–97.

765 Williams, M., and D.J. Siveter. 2008. The earliest leperdicope arthropod: A new genus from
766 the Ordovician of Spitsbergen. *Journal of Micropalaeontology* 27: 97–101.

767

768

769

FIGURE CAPTIONS

770

771 Figure 1. Map of the outcrop area adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen
772 (Basissletta area in the north and the Olenidsletta in the south). Measured sections in circles:
773 FP, F-Promontory section; PO, Profilstranda-Olenidsletta Member section; PR, Profilbekken
774 River section; PS, Profilstranda section; SR, Spora River section.

775

778 Figure 2. Spora River (SR) section. Abbreviations: Fm, Formation; Mbr, Member.

779 Explanation of symbols: 1, wavy bedded dolostone; 2, planar bedded limestone; 3, wavy

780 bedded dolostone, burrow churned dolostone, ribbon dolostone; 4, laminated, planar bedded

781 dolostone; 5, wavy narrowly bedded limestone; 6, laminated, planar bedded limestone; 7,

782 argillaceous-shaly; 8, silty; 9, glauconitic; 10, thrombolitic, fenestral; 11, stromatolites; 12,

783 oolites; 13, flint nodules; 14, flat pebble conglomerate; 15, general intraclastic conglomerate;

784 16, dispersed silica; 17, erosional surface; 18, hardground; 19, not exposed; 20, trilobites; 21,

785 gastropods; 22, cephalopods; 23, sponges; 24, echinoderms; 25, articulate brachiopods; 26,
786 inarticulate brachiopods.

787

Figure 3. Field photographs of details of the Tokammane and Kirtonryggen formations, Furongian?, Cambrian – Tremadocian, Ordovician, Spora River (SR) section, Ny Friesland, Spitsbergen. A, pinkish thrombolitic - fenestral dolostone of the uppermost Tokammane Formation, at SR section c. 40 m; B, burrow-mottled dolostone of the uppermost Spora Member, at SR section 63 m; C, partly silicified domal stromatolite with underlying flat pebble conglomerate, at SR section 148 m.

798

799 Figure 4. Field photographs of bedding surfaces with trace fossils in originally soft
800 sediments of the Kirtonryggen and Valhallfonna formations, Tremadocian–Darriwilian,
801 Ordovician, adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen. A, Upper Spora Member,
802 at Spora River section 62 m; B. Nordporten Member, at Profilstranda (PS) section 234 m; C,
803 upper Basissletta Member, at PS section 107 m; D, uppermost Profilbekken Member, at F-
804 promontory section 5 m.

805

806

Figure 5. Field photographs of intraclastic horizons of the middle Basissletta Member, Kirtonryggen Formation, Tremadocian, Ordovician, Spora River (SR) section, Ny Friesland, Spitsbergen. A, intraclastic conglomerate over erosional surface, at SR section 142 m; B, oolitic ripple horizon within laminated dolo-mudstone at SR section 143 m; C, flat pebble conglomerate, at SR section 147 m; D, flat pebble conglomerate with gutter cast and vertically oriented clasts, at SR section 150 m.

Figure 6. Field photographs of bedding features of the Basissletta Member, Kirtonryggen Formation, Tremadocian, Ordovician, Ny Friesland, Spitsbergen. A, laminated dolomudstone, at Profilstranda (PS) section 5 m; B, argillaceous interval above flat pebble conglomerate bed, at PS section 87-88 m; C, teepee structures in laminated dolomudstone, at PS section 50 m; D, hummocky cross stratification, at PS section 4 m.

823

Profilstranda - Section (PS)

824

825 Figure 7. Profilstranda (PS) section, Basissletta and Nordporten members, Kirtonryggen
826 Formation, Tremadocian–Floian, Ordovician, adjacent to Hinlopenstretet, Ny Friesland,
827 Spitsbergen. For explanation of symbols see Figure 2.

828

829

Figure 8. Field photographs of discontinuity surfaces within the Nordporten Member, Kirtonryggen Formation, Floian, Ordovician, adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen. A, at Profilstranda (PS) section 219 m; B, at PS section 277 m.

Profilstranda -
Section (PO)

835

836

837 Figure 9. Profilstranda - Olenidsletta Member (PO) section, Olenidsletta Member,
 838 Valhallfonna Formation, Floian–Dapingian, Ordovician, adjacent to Hinlopenstretet, Ny
 839 Friesland, Spitsbergen. For explanation of symbols see Fig. 2.

840

Figure 10. Field photographs of bedding surfaces of the Valhallfonna Formation, Floian–Darriwilian, Ordovician, Basissletta area, adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen. A, surface with masses of fragments of orthoconic cephalopods, current aligned, upper Olenidsletta Member, Profilbekken River (PR) section, bed correlates with bed at 128 m at Profilstranda - Olenidsletta Member (PO) section; B, flat hardground with small borings and ophiletid gastropod, Olenidsletta Member, at PO section 90 m; C, bedding surfaces with masses of minute orthoconic cephalopods and trilobite hash, uppermost Olenidsletta Member, PR section; D, endocerid cephalopod with microbial overgrowth on iron (limonitic) stained hardground in Profilbekken Member, at PR section 0 m.

852

853 Figure 11. F-promontory (FP) section, Profilbekken Member, Valhallfonna Formation,
854 Dapingian–Darriwilian, Ordovician, adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen.
855 Note that the thickness measure in meters counts downward. For explanation of symbols see
856 Figure 2.

857

Figure 12. Profilbekken River (PR) section, Profilbekken Member, Valhallfonna Formation, Dapingian–Darriwilian, Ordovician, adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen. For explanation of symbols see Figure 2.

863

864

865 Figure 13. Field photographs of features of the Profilbekken Member, Valhallfonna
866 Formation, Dapingian–Darriwilian, Ordovician, adjacent to Hinlopenstretet, Ny Friesland,
867 Spitsbergen. A, typical banded, silicified, lime-mudstone of the Profilbekken Member, at PR
868 section c. 10 m; B, iron (limonitic) stained hardground with prominent vertical borings, at PR
869 section 0 m; C, upper glauconite bed, at PR section 32 m.

870

871

872

873 Figure 14. Stratigraphic scheme of the Ordovician adjacent to Hinlopenstretet, Ny
874 Friesland, Spitsbergen.1, Fortey (1980), Fortey and Bruton (2013); 2, Lehnert et al. (2013); 3,
875 Cooper and Fortey (1982).

Figure 15. Correlation of the Ordovician adjacent to Hinlopenstretet, Ny Friesland, Spitsbergen with selected successions of eastern Laurentia (eastern US and Canada). Based on Cooper et al. (2012); Kröger and Landing (2011), Lavoie et al. (2012), ° stage slices of Bergström et al. (2009), * range according to Boyce et al. (2011).