<https://helda.helsinki.fi>

Palaeogeographic distribution and diversity of cephalopods during the Cambrian-Ordovician transition

Fang, Xiang

2019

Fang , X , Kröger , B , Zhang , Y-D , Zhang , Y-B & Chen , T-E 2019 , ' Palaeogeographic distribution and diversity of cephalopods during the Cambrian-Ordovician transition ' , Palaeoworld , vol. 28 , no. 1-2 , pp. 51-57 . <https://doi.org/10.1016/j.palwor.2018.08.007>

<http://hdl.handle.net/10138/308817>

<https://doi.org/10.1016/j.palwor.2018.08.007>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Accepted Manuscript

Title: Palaeogeographic distribution and diversity of cephalopods during the Cambrian–Ordovician transition

Authors: Xiang Fang, Björn Kröger, Yuan-Dong Zhang, Yun-Bai Zhang, Ting-En Chen

PII: S1871-174X(18)30046-5
DOI: <https://doi.org/10.1016/j.palwor.2018.08.007>
Reference: PALWOR 465

To appear in: *Palaeoworld*

Received date: 30-3-2018
Revised date: 23-7-2018
Accepted date: 23-8-2018

Please cite this article as: Fang, Xiang, Kröger, Björn, Zhang, Yuan-Dong, Zhang, Yun-Bai, Chen, Ting-En, Palaeogeographic distribution and diversity of cephalopods during the Cambrian–Ordovician transition. *Palaeoworld* <https://doi.org/10.1016/j.palwor.2018.08.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Palaeogeographic distribution and diversity of cephalopods during the Cambrian–Ordovician transition

Xiang Fang ^{a *}, Björn Kröger ^b, Yuan-Dong Zhang ^{c, d}, Yun-Bai Zhang ^a, Ting-En Chen

^a

^a Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Palaeoenvironment, Chinese Academy of Sciences, Nanjing, China

^b Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland

^c CAS Key Laboratory of Economic Stratigraphy and Palaeogeography, Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Palaeoenvironment, Chinese Academy of Sciences, Nanjing, China

^d University of Chinese Academy of Sciences, Beijing, China

* Corresponding author. *E-mail address*: xfang@nigpas.ac.cn

Abstract

Since the early 20th Century, when the first cephalopods from late Cambrian strata were discovered in North China, more than 160 species belonging to 39 genera in nine families and five orders, have been described from both North and South China, together with North America, Siberia and Kazakhstan. We compiled and analysed all published Cambrian cephalopod occurrences in these regions: the results show that the oldest undisputed cephalopods are from the Jiangshanian Stage of North China. After their origination, cephalopods reached their first diversity peak in the late Cambrian *Acaroceras*–*Sinoeremoceras* Biozone (early Stage 10). This initial diversity peak was followed by the “late Trempealeauan Eclipse”, which eradicated nearly 95% of late Cambrian genera. The extinction event coincides with similar extinctions of trilobites and some other groups of marine life. The rapid subsequent diversification of cephalopods during the Tremadocian (Early Ordovician) was paralleled by a

diversification of graptoloids and radiolarians.

Keywords: Cephalopoda; palaeobiogeography; GOBE; diversity; extinction; late Cambrian

1. Introduction

There is dispute with the timing of the earliest appearance of cephalopods. The problematic middle Cambrian animal *Nectocaris pteryx* from Burgess Shale, along with the problematic lower Cambrian taxa *Petalilium* and (probably) *Vetustovermis*, were interpreted as primitive, non-mineralized cephalopods (Smith and Caron, 2010). However, this interpretation was not supported by Kröger et al. (2011), who proposed that nectocarids do not show homologies with coleoids and rather seems to be an independent branch in the bilaterian Tree of Life, which is the view accepted here.

The oldest undoubted cephalopod fossil is probably *Plectronoceras cambria* (Walcott, 1905), from the lower part of the Fengshan Formation (upper Jiangshanian, Furongian, Cambrian) in Shandong Province, North China (see also Smith and Caron, 2010; Kröger et al., 2011; Mazurek and Zatoń, 2011; Runnegar, 2011; Smith, 2013; Vinther, 2015). In slightly younger Cambrian strata, cephalopods have been commonly recorded worldwide (e.g., Flower, 1964; Teichert, 1967; Yochelson et al., 1973; Chen and Teichert, 1983a, 1983b; Landing and Kröger, 2009), showing a rapid diversification in the latest Cambrian. So far, more than 160 species, belonging to 39 genera, in nine families and five orders, have been described from the late Cambrian interval in North and South China, North America and possibly Siberia and Kazakhstan (Fig. 1). However, this initial pulse of cephalopod evolution has not been addressed since the seminal work of Chen and Teichert (1983a, 1983b). Here, for the first time, we provide a complete review of these earliest cephalopods from the existing literature and conduct an analysis of their palaeogeographic distribution and diversity patterns.

Fig. 1 here

2. Materials and methods

To quantitatively investigate the palaeogeographic distribution of early cephalopods, we compiled all the published cephalopod occurrences in the late Cambrian into a binary data set (presence or absence). The main data sources included Flower (1964), Yochelson et al. (1973), Chen and Qi (1982), Peng and Chen (1983), Chen and Teichert (1983a, 1983b) and Li (1984). Our diversity calculations were based on the number of generic occurrences compiled from these sources and the Tremadocian occurrences published by Kröger and Zhang (2009). We estimated the diversity as the “Hill numbers” following the approach of Chao et al. (2014) and used the function “estimateD()” of the package iNext version 2.0.12 (Hsieh et al., 2016) of the R statistical software version 3.50 (R Core Team, 2014). Hill numbers are a mathematically unified family of diversity indices that incorporate relative abundance and species richness and overcome many shortcomings (Chao et al., 2014). The quantitative palaeobiogeographic analysis, including hierarchical cluster analysis (HCA) with the UPGMA linkage and non-metric multidimensional scaling (NMDS) (Hammer and Harper, 2006) was carried out using the program PAST version 3.15 (Hammer et al., 2001), and adopting the Ochiai (Ochiai, 1957) and Jaccard (Jaccard, 1901a, 1901b) similarity coefficients, which are commonly used in palaeobiogeographic studies (e.g., Shi, 1993; Rong et al., 1995; Shen and Shi, 2000, 2004).

3. Early evolution of cephalopods

The oldest cephalopods occur in the *Tsinania–Ptychaspis* Biozone (Jiangshanian Stage) of North China and include four species of *Plectronoceras* (Chen et al., 1979a; Chen and Qi, 1982). *Plectronoceras* is typified by its small and cryptoconic external shell with simple septa and a ventral siphuncle, short and straight septal necks, and expanded segments, without any cameral and endosiphuncular deposits (Chen et al., 1979a).

In the slightly younger *Changia* Biozone of North China and the *Lotagnostus americanus–Hedinaspis regalis* Biozone of South China (Fig. 2; Peng, 2009), which

fall within the Stage 10 of the Cambrian System, 13 species assigned to nine genera, five families and three orders have been recorded so far. Most diverse in these assemblages are elements of the orders Ellesmerocerida and Plectronocerida.

Fig. 2 here

In the succeeding *Acaroceras*–*Sinoeremoceras* Biozone (upper part of Stage 10), cephalopods reached their first Cambrian diversity peak. Within this time interval, two families of the Order Plectronocerida, i.e., Family Plectronoceratidae and Balkoceratidae, and four families of the Order Ellesmerocerida, i.e., the Family Ellesmeroceratidae, Acaroceratidae, Huaiheceratidae and Xiaoshanoceratidae, are recorded. Moreover, the orders Yanhecerida and Endocerida are distinctive components of this horizon (Chen and Teichert, 1983a; Li, 1984).

However, this cephalopod diversity peak was shortly succeeded by a sharp diversity decline in the latest Cambrian *Mictosaukia* Biozone, which led to the “late Trempealeauan Eclipse” (Stage 10, Furongian) of the cephalopods (see Chen and Teichert, 1983a). This diversity drop indicates a possible extinction event characterized by the abrupt and cryptic disappearances of nearly 95% of the existing cephalopod genera. Only a few forms, including *Ectenolites* and *Clarkoceras* of Ellesmeroceratidae and possibly some elements of Order Endocerida, i.e., *Eocameroceras cambria* Li (see Li, 1984), survived into the earliest Ordovician.

The early Tremadocian (Early Ordovician) cephalopod faunas were of low diversity and dominated by few representatives of orders Ellesmerocerida and Endocerida, which were likely descendants of Furongian ancestors. During the mid Tremadocian, the first species of the Order Tarphycerida appeared in Laurentia (Kröger and Landing, 2008), and the first species of the Order Orthocerida were recorded in the Montagne Noir of France, on marginal Gondwanan continent and close to palaeo south pole (Kröger and Evans, 2011). It was also in this time interval that the orders Endocerida and Ellesmerocerida become the main cephalopod faunas in both South and North China (Lai, 1982). The earliest representatives of the Order Actinocerida

originated in the late Floian, although their diversity was low at that time (Flower, 1976).

4. Palaeobiogeography of early cephalopods

During the *Acaroceras*–*Sinoeremoceras* Biozone of Stage 10 (Cambrian), highly diverse cephalopod faunas appeared in several regions of the world, i.e., North and South China, North America, and probably also Siberia and Kazakhstan, indicating that cephalopod habitats were largely restricted to tropical regions (Kröger, 2013; Fig. 3). Both the results of Ochiai and Jaccard measures of HCA and NMDS indicate that the cephalopod faunas of North China have strong affinities with those of South China (Fig. 4), which probably resulted from the close palaeogeographic position of these two regions during the late Cambrian. In addition, the cephalopod fauna of Kazakhstan and that of Hunan, South China, cluster because of the presence of shared common elements, i.e., *Chabactoceras* and *Huaiheceras*. The clustering of these two regions indicates there were possible channels connecting them.

Fig. 3 here

However, as noted by Flower (1964) and Landing and Kröger (2009), the age of the cephalopod faunas from Kazakhstan and Siberia is dubious. On the basis of the associated organisms, e.g., trilobites, the age could be earliest Ordovician. In addition, the affinity of these cephalopods is questionable due to their poor and fragmentary preservation (Landing and Kröger, 2009).

Fig. 4 here

5. Discussion

The palaeoecology of the earliest cephalopods has been poorly known, largely due to poor preservation and their remarkably sporadic occurrences within the time interval of a few million years. A recent study of the internal microstructure of plectronocerids and ellesmerocerids demonstrated that the oldest cephalopods already possessed two

distinct types of connecting rings, which contain an advanced and calcified ring in plectronocerids and a *Nautilus*-type ring in ellesmerocerids (Mutvei, 2015). It was concluded that the earliest cephalopods possessed low capacity jet-power locomotion and probably had a demersal lifestyle (Mutvei et al., 2007). After their origination during the early Furongian, cephalopods diversified rapidly and reached a diversity peak in a stepwise pattern. It is noteworthy that the rapid diversification of cephalopods during the late Cambrian approximately coincides with the origination and diversification of several other groups (e.g., graptolites and radiolarians, Fig. 2; Zhang et al., 2010; Servais et al., 2016). However, in Cambrian Stage 10, cephalopods disappear suddenly, with only two genera surviving into the earliest Ordovician. This high turnover rate suggests that there may indeed have been a cephalopod extinction event (the “late Trempealeauan Eclipse”) during the Cambrian–Ordovician transition, both in taxonomic diversity and in abundance. In addition, the stepwise diversification of cephalopods during the latest Cambrian is similar to that of the acritarchs, which was halted in the upper part of Stage 10 by the extinction event (Nowak et al., 2015). On the other hand, the occurrence data could be affected by sampling biases, which is usually difficult to avoid. In this case, it is important that, the nearly all of the published cephalopod occurrences during the late Cambrian have been collected and analysed. To reduce any potential sampling biases, we also used “Hill numbers” and the function “estimatedD()” to assess the real cephalopod diversity.

There was also a mass extinction of trilobites in the *Mictosaukia* Biozone (CTI40) during the late Fengshanian (Stage 10, Furongian) marked by the high extinction and low origination rates of species (Zhen and Zhou, 2008, p. 316). A large number of existing non-agnostoid trilobites characterizing the late Cambrian faunas, e.g., Kaolishaniidae, Kingstoniidae, Pterocephalidae, Ptychaspididae, Ptychopariidae and Saukiidae, made their last appearance during this time interval; in addition, only one-third of agnostids survived this extinction (Zhen and Zhou, 2008). This event, called “Late Fengshanian (or Dolgellian) Event”, coincides with the Trempealeauan/Ibex Event of North America (Brasier, 1996; Zhen and Zhou, 2008). The global diversity of lingulids, acrotretids and other brachiopod taxa also experienced a sharp decline during

the latest Cambrian (Sepkoski, 1995). Both the trilobite and brachiopod extinction events correspond to the “late Trempealeauan Eclipse” of cephalopods, and their contemporary timing in turn coincides with a major late Cambrian sea-level regression (Brasier, 1996; Brasier and Lindsay, 2001).

Furthermore, our analysis shows a high cephalopod endemism during the latest Cambrian. More than 80% of all genera occurred on a single palaeo-plate, and only *Ectenolites* achieved a widespread distribution on four palaeo-plates (Fig. 5). As *Ectenolites* is one of the two genera that survived the end-Cambrian, its survival may have been due to the effect of its widespread distribution. Similar effects have been shown by evidence from some other widespread fossil groups (Boucot, 1975; Erwin, 1998; Rong et al., 2004). This may indicate that widespread taxa strategically limit their geographic ranges during an extinction interval, and re-expand their habitats after the extinction.

Fig. 5 here

6. Conclusions

Cephalopods first appeared during the late Cambrian in North China and rapidly dispersed to South China and North America, and probably also to Siberia and Kazakhstan. Shortly afterwards, nearly all of the existing early cephalopods disappeared abruptly in the latest Cambrian, with only two genera surviving into the earliest Ordovician. The surviving cephalopod taxa are characterized by their widespread distribution, in contrast to the extinct taxa that had remarkable restricted distributions. This suggests that those more widespread cephalopod taxa had more chances to survive whereas those more endemic groups were prone to extinction.

Acknowledgements

We are indebted to Lucy Muir and David Harper for polishing the English language. We are grateful to Christian Klug and Claude Monnet for their constructive suggestions of the manuscript. Financial support from the National Natural Science Foundation of

China (Grant No. 41521061), the Chinese Academy of Sciences (Grant No. XDB10010100), State Key Laboratory of Palaeobiology and Stratigraphy (Grant No. 20172111) and the Ministry of Science and Technology of China (Grant Nos. 2013FY111000, 2017ZX05036-001-004) is acknowledged. This is a contribution to IGCP Project 653 ('The Onset of the Great Ordovician Biodiversification Event').

References

- Boucot, A.J., 1975. *Evolution and Extinction Rate Controls*. Elsevier Scientific Publishing Company, Amsterdam, Oxford, New York, 427 pp.
- Brasier, M.D., 1996. The basal Cambrian transition and Cambrian bio-events (from terminal Proterozoic extinctions to Cambrian biomes). In: Walliser, O.H. (Ed.), *Global Events and Event Stratigraphy in the Phanerozoic*. Springer, Berlin, pp. 113–138.
- Brasier, M.D., Lindsay, J.F., 2001. Did supercontinental amalgamation trigger the “Cambrian Explosion”? In: Zhuravlev, A.Y., Riding, R. (Eds.), *The Ecology of the Cambrian Radiation*. Columbia University Press, New York, pp. 69–89.
- Chao, A., Gotelli, N.J., Hsieh, T.C., Sander, E.L., Ma, K.H., Colwell, R.K., Ellison, A.M., 2014. Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. *Ecological Monographs* 84, 45–67.
- Chen, J.Y., Qi, D.L., 1982. Upper Cambrian cephalopods from western Zhejiang. *Geological Society of America Special Papers* 187, 137–142.
- Chen, J.Y., Teichert, C., 1983a. Cambrian cephalopods. *Geology* 11, 647–650.
- Chen, J.Y., Teichert, C., 1983b. Cambrian cephalopods from China. *Palaeontographica Abt. A* 181, 1–102.
- Chen, J.Y., Zou, X.P., Chen, T.E., Qi, D.L., 1979a. Late Cambrian Ellesmerocerida (Cephalopoda) of North China. *Acta Palaeontologica Sinica* 18 (2), 103–122 (in Chinese, with English abstract).
- Chen, J.Y., Zou, X.P., Chen, T.E., Qi, D.L., 1979b. Late Cambrian cephalopods of North China — Plectronocerida, Protactinocerida (Ord. nov.) and Yanhecerida (Ord. nov.). *Acta Palaeontologica Sinica* 18 (1), 1–24 (in Chinese, with English abstract).
- Erwin, D.H., 1998. After the end: recovery from extinction. *Science* 279, 1324–1325.
- Flower, R.H., 1964. The nautiloid Order Ellesmeroceratida (Cephalopoda). State Bureau of Mines and Mineral Resources, New Mexico Institute of Mining and Technology, Memoir 12, 1–234.
- Flower, R.H., 1976. Ordovician cephalopod faunas and their role in correlation. In:

- Bassett, M.G. (Ed.), The Ordovician System. University of Wales Press, Cardiff, pp. 523–552.
- Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M., 2012. The Geologic Time Scale 2012. Elsevier, Boston, 1144 pp.
- Hammer, Ø., Harper, D.A.T., 2006. Paleontological Data Analysis. Blackwell Publishing, Oxford, 351 pp.
- Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4, 1–9.
- Hsieh, T.C., Ma, K.H., Chao, A., McInerney, G., 2016. iNEXT: An R package for interpolation and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* 7, 1451–1456.
- Jaccard, P., 1901a. Distribution de la flore alpine dans le Bassin des Dranses et dans quelques régions voisines. *Bulletin de la Société Vaudoise des Sciences Naturelles* 37, 241–272.
- Jaccard, P., 1901b. Étude comparative de la distribution florale dans une portion des Alpes et du Jura. *Bulletin de la Société Vaudoise des Sciences Naturelles* 37, 547–579.
- Kröger, B., 2013. Cambrian–Ordovician cephalopod palaeogeography and diversity. In: Harper, D.A.T., Servais, T. (Eds.), *Early Palaeozoic Biogeography and Palaeogeography*. Geological Society, London, Memoirs 38, 429–448.
- Kröger, B., Evans, D., 2011. Review and palaeoecological analysis of the late Tremadocian–early Floian (Early Ordovician) cephalopod fauna of the Montagne Noire, France. *Fossil Record* 14, 5–34.
- Kröger, B., Landing, E., 2008. Onset of the Ordovician cephalopod radiation — evidence from the Rochdale Formation (middle Early Ordovician, Stairsian) in eastern New York. *Geological Magazine* 145, 490–250.
- Kröger, B., Zhang, Y.B., 2009. Pulsed cephalopod diversification during the Ordovician. *Palaeogeography, Palaeoclimatology, Palaeoecology* 273, 174–183.
- Kröger, B., Vinther, J., Fuchs, D., 2011. Cephalopod origins and evolution: A congruent

- picture emerging from fossils, development and molecules. *Bioessays* 33, 602–613.
- Lai, C.G., 1982. *Stratigraphy of China 5. Ordovician of China*. Geological Publishing House, Beijing, 297 pp. (in Chinese).
- Landing, E., Kröger, B., 2009. The oldest cephalopods from east Laurentia. *Journal of Paleontology* 83 (1), 123–127.
- Li, L.Z., 1984. Cephalopods from the upper Cambrian Siyangshan Formation of western Zhejiang. In: Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences (Ed.), *Stratigraphy and Palaeontology of Systemic Boundaries in China Cambrian — Ordovician Boundary* (1). Anhui Science and Technology Publishing House, Hefei, pp. 187–265.
- Mazurek, D., Zatoń, M., 2011. Is *Nectocaris pteryx* a cephalopod? *Lethaia* 44, 2–4.
- Mutvei, H., 2015. Characterization of two new superorders Nautilosiphonata and Calciosiphonata and a new order Cyrtocerinida of the subclass Nautiloidea; siphuncular structure in the Ordovician nautiloid *Bathmoceras* (Cephalopoda). *GFF* 137, 164–174.
- Mutvei, H., Zhang, Y.B., Dunca, E., 2007. Late Cambrian plectronocerid nautiloids and their role in cephalopod evolution. *Palaeontology* 50, 1327–1333.
- Nowak, H., Servais, T., Monnet, C., Molyneux, S.G., Vandenbroucke, T.R.A., 2015. Phytoplankton dynamics from the Cambrian Explosion to the onset of the Great Ordovician Biodiversification Event: A review of Cambrian acritarch diversity. *Earth-Science Reviews* 151, 117–131.
- Ochiai, A., 1957. Zoogeographic studies on the soleoid fishes found in Japan and its neighbouring regions. *Nippon Suisan Gakkaishi* 22, 526–530.
- Peng, S.C., 2009. The newly-developed Cambrian biostratigraphic succession and chronostratigraphic scheme for South China. *Chinese Science Bulletin* 54, 4161–4170.
- Peng, S.C., Chen, J.Y., 1983. Discovery of late Cambrian nautiloids from Cili. *Geology of Hunan* 2 (2), 13–17 (in Chinese).
- R Core Team, 2014. R: A language and environment for statistical computing. R

- Foundation for Statistical Computing, Vienna.
- Rong, J.Y., Li, R.Y., Kul'kov, N.P., 1995. Biogeographic analysis of Llandovery brachiopods from Asia with a recommendation of use of affinity indices. *Acta Palaeontologica Sinica* 34, 428–453 (in Chinese, with English abstract).
- Rong, J.Y., Chen, X., Zhou, Z.Y., Chen, J.Q., 2004. Response of major organism group to global environmental perturbations through the Ordovician–Silurian transition in South China. In: Rong, J.Y., Fang, Z.J., (Eds.), *Mass Extinction and Recovery — Evidences from the Palaeozoic and Triassic of South China*. University of Science and Technology of China Press, Hefei, pp. 235–256 (in Chinese, with English abstract).
- Runnegar, B., 2011. Once again: is *Nectocaris pteryx* a stem-group cephalopod? *Lethaia* 44, 373–373.
- Sepkoski, J.J., 1995. The Ordovician radiations: Diversification and extinction shown by global genus-level taxonomic data. In: Cooper, J.D., Droser, M.L., Finney, S.C. (Eds.), *Ordovician Odyssey: Short Papers for the Seventh International Symposium on the Ordovician System*. Pacific Section, Society for Sedimentary Geology (SEPM), California, pp. 393–369.
- Servais, T., Perrier, V., Danelian, T., Klug, C., Martin, R., Munnecke, A., Nowak, H., Nützel, A., Vandenbroucke, T.R.A., Williams, M., Rasmussen, C.M.Ø., 2016. The onset of the ‘Ordovician Plankton Revolution’ in the late Cambrian. *Palaeogeography, Palaeoclimatology, Palaeoecology* 458, 12–28.
- Shen, S.Z., Shi, G.R., 2000. Wuchiapingian (early Lopingian, Permian) global brachiopod palaeobiogeography: a quantitative approach. *Palaeogeography, Palaeoclimatology, Palaeoecology* 162, 299–318.
- Shen, S.Z., Shi, G.R., 2004. Capitanian (Late Guadalupian, Permian) global brachiopod palaeobiogeography and latitudinal diversity pattern. *Palaeogeography, Palaeoclimatology, Palaeoecology* 208, 235–262.
- Shi, G.R., 1993. Multivariate data analysis in palaeoecology and palaeobiogeography — a review. *Palaeogeography, Palaeoclimatology, Palaeoecology* 105, 199–234.
- Smith, M.R., 2013. Nectocaridid ecology, diversity, and affinity: early origin of a

- cephalopod-like body plan. *Paleobiology* 39, 297–321.
- Smith, M.R., Caron, J., 2010. Primitive soft-bodied cephalopods from the Cambrian. *Nature* 465, 469–472.
- Teichert, C., 1967. Major features of cephalopod evolution. In: Teichert, C., Yochelson, E.L. (Eds.), *Essays in Paleontology and Stratigraphy*. University of Kansas Department of Geology Special Publication 2, 162–210.
- Vinther, J., 2015. The origins of molluscs. *Palaeontology* 58, 19–34.
- Walcott, C.D., 1905. Cambrian faunas of China. *Proceedings of United States National Museum* 29, 1–106.
- Yochelson, E.L., Flower, R.H., Webers, G.F., 1973. The bearing of the new Late Cambrian monoplacophoran genus *Knightoconus* upon the origin of the Cephalopoda. *Lethaia* 6, 275–310.
- Zhang, Y.D., Zhan, R.B., Fan, J.X., Cheng, J.F., Liu, X., 2010. Principal aspects of the Ordovician biotic radiation. *Science China: Earth Sciences* 53 (3), 382–394.
- Zhen, Y.Y., Zhou, Z.Y., 2008. Chapter 11. History of trilobite biodiversity: a Chinese perspective. In: Zhou, Z.Y., Zhen, Y.Y. (Eds.), *Trilobite Record of China*. Science Press, Beijing, pp. 301–330.

Figure captions

Figure 1. Global distribution of Cambrian cephalopods. Data sources: Flower (1964), Yochelson et al. (1973), Chen and Qi (1982), Peng and Chen (1983), Chen and Teichert (1983a, 1983b) and Li (1984).

Taxonomy			North China					South China		Kazakhstan	Siberia	Texas
Order	Family	Genus	Liaoning	Shandong	Anhui	Shanxi	Inner Mongolia	Zhejiang	Hunan			
Plectronocerida	Plectronoceratidae	<i>Plectronoceras</i>	✓	✓	✓							
		<i>Paraplectronoceras</i>			✓							
		<i>Lunanoceras</i>		✓	✓							
		<i>Eodiaphragmoceras</i>		✓								
		<i>Jiagouceras</i>			✓							
		<i>Rectseptoceras</i>			✓							
		<i>Palaeoceras</i>										✓
		<i>Parapalaeoceras</i>						✓				
		<i>Ruthenoceras</i>									✓	
	Balkoceratidae	<i>Theskeloceras</i>	✓									
		<i>Balkoceras</i>										✓
Ellesmerocerida	Ellesmeroceratidae	<i>Hunyuanoceras</i>				✓						
		<i>Eburoceras</i>		✓	✓							
		<i>Anhuiceras</i>			✓		✓	✓				
		<i>Chabactoceras</i>			✓			✓	✓	✓		
		<i>Dongshanoceras</i>			✓			✓				
		<i>Eoclarkoceras</i>	✓	✓	✓			✓				
		<i>Eoectenolites</i>			✓							
		<i>Pseudendoceras</i>			✓							
		<i>Qilushougouceras</i>	✓									
		<i>Sinolebetoceras</i>			✓							
		<i>Tanycameroceras</i>	✓		✓							
		<i>Ectenolites</i>	✓		✓			✓		✓		✓
		<i>Clarkoceras</i>			✓							
		<i>Walcottoceras</i>						✓				
	Acaroceratidae	<i>Acaroceras</i>	✓	✓	✓	✓	✓	✓		✓		
		<i>Weishanhuceras</i>		✓								
		<i>Bambusoceras</i>						✓				
	Huaiheceratidae	<i>Huaihecera</i>	✓	✓	✓			✓	✓	✓		
		<i>Huaihecerina</i>	✓									
		<i>Antacaroceras</i>						✓				
		<i>Cilioceras</i>							✓			
		<i>Zhuibianoceras</i>						✓				
	Xiaoshanoceridae	<i>Xiaoshanoceras</i>						✓				
Protactinocerida	Protactinoceratidae	<i>Wanwanoceras</i>	✓	✓	✓							
		<i>Sinoeremoceras</i>	✓	✓	✓		✓					
		<i>Protactinoceras</i>	✓	✓								
		<i>Benxioceras</i>	✓									
		<i>Mastoceras</i>	✓									
		<i>Physalactinoceras</i>	✓	✓								
Yanhecerida	Yanheceratidae	<i>Yanhecera</i>			✓			✓				
		<i>Aetheloxoceras</i>			✓							
		<i>Archendoceras</i>		✓				✓				
		<i>Oonendoceras</i>			✓							
Endocerida	Proterocameroceratidae	<i>Eocameroceras</i>						✓				

Figure 2. Range chart of early cephalopod faunas through the Cambrian and Ordovician transition and the diversity of some other major marine organisms during the onset of the Great Ordovician Biodiversification Event (GOBE). Biozone information is modified from Chen and Teichert (1983a, 1983b) and Peng (2009). Data sources: cephalopods, solid lines indicate the number of genera reported, dashed lines indicate the estimated diversity with the 95% confidence levels for the diversity; acritarchs, Servais et al. (2016); graptolites, Zhang et al. (2010); trilobites (China), Zhen and Zhou (2008); $\delta^{13}\text{C}$ curve, Gradstein et al. (2012) and Servais et al. (2016). SPICE, Steptoean Positive Carbon isotope Excursion; TOCE, Top of Cambrian isotope Excursion. Family Plectronoceratidae and Balkoceratidae belong to Order Plectronocerida. Family Protactinoceratidae belongs to Order Protactinocerida. Family Huaiheceratidae, Acaroceratidae, Ellesmeroceratidae and Xiaoshanoceridae belong to Order Ellesmerocerida. Family Yanheceratidae belongs to Order Yanhecerida. Fossil illustrations: 1. *Plectronoceras* cf. *cambria* (Walcott) (Chen et al., 1979a; Shandong, China); 2. *Acaroceras endogastrum* Chen et al. (Chen et al., 1979b; Anhui, China); 3. *Ectenolites petilus* Chen and Qi (Chen and Qi, 1982; Anhui, China); 4. *Zhuibianoceras zhejiangense* Li (Li, 1984; Zhejiang, China); 5. *Xiaoshanoceras jini* Chen and Teichert (Chen and Teichert, 1983b; Zhejiang, China); 6. *Yanheceras endogastrum* Chen (Chen et al., 1979a; Guizhou, China).

Figure 3. Palaeobiogeographic distribution of cephalopods during the latest Cambrian. 1. Hunan, South China; 2. Zhejiang, South China; 3. Liaoning, North China; 4. Anhui, North China; 5. Shandong, North China; 6. Inner Mongolia, North China; 7. Karatau Mountains, Kazakhstan; 8. Angara Uplift, Siberia; 9. Texas, North America; 10. Nevada, North America; 11. New York, North America. Solid circles mark unequivocal evidence; unfilled circles mark possible occurrences (see Kröger, 2013). The base maps of palaeogeographic reconstructions are produced by PaleoGIS 4.0 for ArcGIS (2011), The Rothwell Group, L.P. 545 (www.paleogis.com).

Late Cambrian

Figure 4. HCA with UPGMA and NMDS of the late Cambrian (*Acaroceras*–*Sinoeremoceras* Biozone) cephalopods based on the Ochiai and Jaccard similarity

measures. NC, North China; SC, South China.

Figure 5. Frequency analysis of cephalopod distribution in the latest Cambrian indicates most cephalopod genera were yielded in a single palaeo-plate (82% palaeo-plates involved in the analysis include the North and South China, Kazakhstan, Siberia and Laurentia).