<https://helda.helsinki.fi>

Preoperative Risk Stratification of Endometrial Carcinoma : L1CAM as a Biomarker

Pasanen, Annukka

2017-09

Pasanen , A , Loukovaara , M , Tuomi , T & Butzow , R 2017 , ' Preoperative Risk Stratification of Endometrial Carcinoma : L1CAM as a Biomarker ' , International Journal of Gynecological Cancer , vol. 27 , no. 7 , pp. 1318-1324 . <https://doi.org/10.1097/IGC.0000000000001043>

<http://hdl.handle.net/10138/311349>

<https://doi.org/10.1097/IGC.0000000000001043>

cc_by_nc

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Preoperative risk stratification of endometrial carcinoma: L1 cell adhesion molecule as a biomarker

Objective: Pre- or intraoperative risk assessment models are used to stratify patients with endometrial carcinoma to lymphadenectomy. Our aim was to determine whether preoperative analysis of L1 cell adhesion molecule (L1CAM) can improve risk assessment.

Methods: Immunohistochemical L1CAM staining was performed on endometrial biopsies of 241 patients and paired hysterectomy samples of 75 patients. Risk assessment models based on preoperative histological type and grade, myometrial invasion and/or tumor diameter and alternative models incorporating preoperative L1CAM were compared with regard to their capability of predicting lymph nodal or distant metastasis. Soluble L1 levels were measured by ELISA in serum samples of 40 patients with endometrial carcinoma.

Results: The concordance rate between L1CAM staining results of preoperative and hysterectomy samples was moderate (kappa 0.586, $P < 0.0001$). Preoperative L1CAM expression was associated with non-endometrioid histology, lymph node involvement, advanced stage and positive peritoneal cytology. Receiver operating characteristic (ROC) analyses showed that L1CAM did not significantly improve risk stratification algorithms based on traditional risk factors. Intraoperative tumor diameter was an effective surrogate for myometrial invasion. There was no statistical difference between L1 serum levels of patients with a L1CAM-positive or L1CAM-negative endometrial carcinoma ($P = 0.786$).

Conclusions: L1CAM expression in endometrial biopsy correlates with high risk features of endometrial carcinoma but does not significantly improve risk stratification algorithms based on traditional factors. Soluble L1 detected in the serum of patients with endometrial carcinoma does not correlate with tumoral L1CAM expression.

Keywords endometrial cancer, risk stratification, lymphadenectomy, L1CAM, soluble L1CAM

31 **Introduction**

32 The standard primary treatment of endometrial carcinoma consists of surgery with total
33 hysterectomy and bilateral salpingo-oophorectomy, complemented with pelvic and para-aortic
34 lymphadenectomy in selected cases [1]. There is no compelling evidence that lymphadenectomy as
35 such is therapeutically beneficial and its main motivation is to more accurately stage and stratify
36 patients to postoperative adjuvant therapy. However, extensive surgery may cause morbidity and
37 the rate of unnecessary treatment should be minimized [2]. At most institutions the decision on
38 lymphadenectomy is made pre- or intraoperatively based on features of the primary tumor,
39 including histologic type, grade of differentiation, and depth of myometrial invasion, as evaluated
40 by preoperative histology, frozen section analysis and imaging.

41 The most validated algorithm (so called Mayo criteria, [3]) defines low risk endometrial carcinoma
42 as endometrioid G1-2 carcinoma with tumor diameter ≤ 2 cm and myometrial invasion $\leq 50\%$.
43 When this algorithm is applied, approximately 30% of the patients fall into the category of low risk
44 for lymphatic dissemination and may avoid lymphadenectomy [4,5]. The remaining 70% of the
45 patients should undergo lymphadenectomy, yet only 19-22% of them present with lymph node
46 metastases [4,5]. More recently a risk stratification model was presented, according to which over
47 40% of the patients could be spared lymphadenectomy with a false positive rate of 57.2% and false
48 negative rate of 0% [6]. This model is based on tumor grade (G1-2 vs G3), diameter (50-mm cut-
49 off) and depth of myometrial invasion (MI, three-tiered). A major difficulty related to these
50 algorithms is the often inaccurate assessment of myometrial invasion by preoperative imaging or
51 gross visualization [7-9]. Frozen section diagnosis is not readily available in many institutions [10]
52 and various investigators question its accuracy [11-13]. Intraoperatively assessed tumor diameter
53 has been proposed as a surrogate for myometrial invasion [6,14,15]

54 Current risk assessment algorithms are burdened with a high frequency of presumably unnecessary
55 lymphadenectomies. To overcome this problem, attention has been paid to the potential value of
56 molecular markers (such as ER/PR status in predicting lymph node involvement [16]).
57 Nevertheless, molecular markers do not have an established role in this setting nor have they been
58 integrated in randomized clinical trials of surgical therapies.

59 A promising prognostic marker, L1 cell adhesion molecule (L1CAM, CD171) predicts disease
60 progression and poor prognosis in many types of cancer including endometrial carcinoma [17-23].
61 The association between L1CAM expression and lymph node involvement of endometrial
62 carcinoma suggests that L1CAM could be a useful biomarker for stratifying patients to
63 lymphadenectomy [21-23]. Also, a soluble form of L1CAM (sL1) exists and has been detected in
64 the serum and ascites of patients carrying a tumor expressing this antigen [17,24-26].

65 Based on the association of L1CAM expression with lymphatic dissemination in endometrial
66 carcinoma, we wanted to evaluate the power of L1CAM in algorithms aimed at stratifying patients
67 to lymph node dissection. To further clarify the potentiality of L1CAM as a biomarker, we
68 compared serum L1CAM concentrations in patients with negative and positive L1CAM expression
69 in tumor sections.

70

71

72 **Material and methods**

73 Patients who underwent primary surgical treatment for endometrial carcinoma at the Department of
74 Obstetrics and Gynecology, Helsinki University Hospital, between January 1, 2007 and December
75 31, 2009 were identified. Patients with a preoperative endometrial sample available for L1CAM
76 analysis were included in the study (n = 241). Approvals of the Institutional Review Board and the
77 National Authority for Medicolegal Affairs of Finland were obtained. During 2007-09, according to

78 the treatment guidelines of our hospital, bilateral pelvic lymphadenectomy was performed in patients
79 with grade 1-2 endometrioid carcinoma with <50% myometrial invasion, the depth of invasion being
80 assessed by vaginal ultrasound and gross visual inspection. In other patients, both pelvic and para-
81 aortic lymphadenectomies were performed. There was some variation in practice patterns because the
82 decision to perform lymphadenectomy and the extent of the procedure depended on patient age and
83 surgical risks. Total rate of lymphadenectomy was 79.7 %. Pertinent patient characteristics and
84 surgical data are shown in Table 1.

85 Factors selected for statistical analyses were: preoperative L1CAM expression, FIGO
86 2009 stage [27], lymph node involvement, histologic type (endometrioid/non-endometrioid), grade
87 of differentiation, depth of myometrial invasion, tumor diameter, cervical stromal invasion, peritoneal
88 cytology status, patient age at surgery, and body mass index (BMI). The cut off values for the numeric
89 variables ($\geq 50\%$ and $>33\%$ for myometrial invasion, 2cm and 5 cm for tumor diameter, 65 years for
90 age, 30 kg/m² for BMI) were based on earlier reports [6,28-30]. Tumor size was measured
91 intraoperatively by the surgeon or after formaldehyde fixation by the pathologist. Primary tumor
92 diameter was defined as the largest dimension of the tumor. If more than 1 lesion was present, the
93 lesion with the largest diameter was considered. Primary tumor diameter was unknown in 14 patients.
94 The presence of cervical stromal invasion was unknown in 2 patients. Peritoneal cytology was
95 considered positive if adenocarcinoma cells were detected in the peritoneal washing, regardless of
96 the number of cancer cells. One case that was positive due to a concomitant borderline serous ovarian
97 tumor was considered negative for endometrial cancer. Peritoneal cytology status was unknown in 4
98 patients.

99 Preoperative L1CAM staining was assessed in tissue samples obtained by uterine aspiration biopsy
100 or curettage. Uterine biopsy was the primary (>90%) sampling method. Uterine curettage was
101 performed when biopsy was insufficient for diagnosis or failed due to cervical stenosis. For
102 immunohistochemical stainings, slides were stained with Ventana Benchmark XT automated slide

103 preparation system (Ventana Medical Systems, Inc., USA) or with Autostainer LV1 (Lab Vision
104 Corporation, USA). Briefly, slides were deparaffinized and heat-induced epitope retrieval was
105 performed following standard protocol. Tissue sections were incubated with primary monoclonal
106 antibodies against L1CAM (CD171; clone 14.10, catalog number SIG-3911-1000, Covance Inc., NJ,
107 USA). The antibody binding site was visualized using a DAB Detection Kit. Sections were
108 counterstained with Mayer's hematoxylin, dehydrated, cleared in xylene, and mounted. L1CAM
109 positivity was defined as >10% of the carcinoma cells staining in one representative slide evaluated
110 by a pathologist (Supplementary figure 1). Neural cells of an appendix slide served as an external
111 positive control and myometrial nerves as an internal positive control (for whole sections). For
112 concordance studies we stained the corresponding hysterectomy sections of all the patients with a
113 positive (n = 50) and of 25 patients with a negative preoperative sample.

114 Starting from November 2014, we have obtained a preoperative blood sample from voluntary
115 patients with endometrial carcinoma treated at the Department of Obstetrics and Gynecology,
116 Helsinki University hospital. Blood fractionation was carried out by centrifugation for 10 min at
117 2000×g and the samples were stored at -70°C. The serum samples of all the patients with an
118 immunohistochemically verified L1CAM-positive (n = 17) and 23 patients with an L1CAM-
119 negative endometrial carcinoma were retrieved. To determine the serum level of L1CAM we used a
120 commercial enzyme linked immunosorbent assay (ELISA) kit (LifeSpan Biosciences Inc., WA,
121 USA, Catalog No. LS-F24209). Standards, controls and samples were processed for sandwich
122 ELISA according to the manufacturer's instructions and duplicate wells were ran for each sample.
123 Final serum dilution (1:2000) was chosen after running test reactions on serial dilutions. The
124 absorbance at 450 nm was measured by an automatic ELISA reader (Multiskan EX, Thermo Fisher
125 Scientific, USA). Results were expressed in ng/ml according to the established standard curve. The
126 limit of detection was 93.75–6000 pg/ml.

Continuous variables (sL1) were compared using the Mann-Whitney U test. Pearson χ^2 analyses were used to compute odds ratios (OR) along with 95% confidence intervals (CI) for the associations between preoperative L1CAM staining and various risk parameters in the cohort. Multiple regression analysis was used to estimate the independent effect of selected risk parameters on either preoperative L1CAM staining or lymph node/distant metastasis (stage IIIC-IV disease). Cohen's kappa statistics were calculated to measure the agreement of preoperative L1CAM staining and tumor histology with corresponding postoperative findings. Based on kappa references outlined by Landis and Koch [31], the strength of agreement was considered moderate for kappa values between 0.41 and 0.60 and substantial for kappa values between 0.61 and 0.80.

Multivariable models were created to test the capability of preoperative L1CAM to predict lymph node and distant metastasis in conjunction with other risk parameters. The estimated weight of each parameter included in a risk model, was determined by rounding statistically significant odds ratios in the multivariable models to the nearest integer. These risk points of each factor were summed to generate a risk score potentially predicting the probability of advanced disease. The risk scores were used to test the discriminating abilities of the risk models with the 2-tailed receiver operating characteristic (ROC) curve area comparison test. Alternative models were created by eliminating selected variables from the models. Statistical significance was set at $P < 0.05$. Data were analyzed using IBM SPSS version 22 software (IBM Corp., Armonk, NY, USA).

Results

Of the 241 preoperative endometrial samples, 64 (26.6%) were L1CAM positive. L1CAM expression was observed in 22.3% (43/193) of grade 1-2 endometrioid carcinomas, 27.6% (8/29) of grade 3 endometrioid carcinomas, and 68.4% (13/19) of non-endometrioid carcinomas ($P < 0.0001$). According to kappa statistics in 75 sample pairs, preoperative L1CAM staining showed moderate

152 agreement with findings in the whole section (κ 0.586, $P < 0.0001$). By comparison, in the whole
153 study population of 241 patients, κ value was 0.551 ($P < 0.0001$) for the agreement of
154 preoperative histology with final histology in detecting high risk cases (grade 3 or non-endometrioid
155 carcinoma). We did not observe any special L1CAM staining pattern, such as preferential positivity
156 at the myoinvasive front, in the whole sections of hysterectomy specimens.

157 Preoperative L1CAM positivity was associated with disease spread beyond the uterine
158 corpus, lymph node involvement, non-endometrioid histology, positive peritoneal cytology, and high
159 age (Table 2). Logistic regression analysis indicated that non-endometrioid histology was
160 independently associated with L1CAM positivity, whereas the effect of disease spread beyond uterine
161 corpus, positive peritoneal cytology or high age was not significant (Table 3). Preoperative high risk
162 histology (grade 3 or non-endometrioid carcinoma), myometrial invasion ($>33\%$ or $\geq 50\%$), tumor
163 diameter (≥ 2 cm or ≥ 5 cm) and preoperative L1CAM positivity were included in logistic regression
164 models, with lymph node and distant metastasis as the dependent variable. Patients with available
165 data for all the variables were included in each model ($n \geq 225$). Tumor size ≥ 2 cm was the only
166 variable that failed to display a significant independent effect on the dependent variable (Table 4).
167 Addition of L1CAM in the models did not significantly improve the AUCs of the risk stratification
168 algorithms ($P > 0.28$, Table 5). Elimination of myometrial invasion from Cox Bauer's model (TD ≥ 5
169 cm, MI $> 33\%$), did not significantly diminish the AUC of the score ($P = 0.429$).

170 There was no statistically significant difference between the concentrations of soluble L1 (s-L1) in
171 the serum samples of patients with L1CAM positive or negative tumors ($P = 0.786$). The mean ($\pm$
172 SD) soluble L1 concentration was 3235.49 ± 808.60 ng/ml for L1CAM positive cases and $3163.27 \pm$
173 765.90 ng/ml for L1CAM negative cases. Median (25th and 75th percentiles) soluble L1 values were
174 3033.90 (2680.60 and 3637.60 respectively) ng/ml in the patients with L1CAM positive tumor and
175 2992.20 (2649.75 and 3467.10 respectively) ng/ml in the L1CAM negative controls.

176

177 **Discussion**

178 Modern management of endometrial cancer is based on personalized surgical and adjuvant treatment.
179 Reliable pre- or intraoperative risk stratification plays a key role in tailoring optimal surgical
180 treatment. Currently used risk assessment methods suffer from inaccuracy and definite indications
181 for lymphadenectomy are yet to be established.

182 L1CAM is a promising prognostic marker that independently predicts poor outcome and lymph nodal
183 involvement in endometrial carcinoma [21-23]. Since L1CAM expression pattern is heterogeneous
184 in endometrial carcinoma (10-100% of the carcinoma cells staining in a positive
185 immunohistochemical assay) and endometrial aspiration biopsy represents only a small portion of the
186 tumor, the true value of L1CAM as a preoperative marker has to be studied on preoperative diagnostic
187 samples. Despite the heterogeneous staining pattern of L1CAM, we observed a moderate
188 concordance rate (kappa 0.586, $P < 0.0001$) between L1CAM staining in preoperative and
189 hysterectomy samples. It is noteworthy, that the concordance between pre- and postoperative L1CAM
190 staining was superior compared to the concordance of pre- and postoperative histology (low vs high
191 grade). L1CAM expression was associated with disease spread beyond uterine corpus (OR 2.5, $P =$
192 0.003), but its significant effect was lost once other factors were taken into account. Further, L1CAM
193 did not significantly improve the performance of risk assessment algorithms based on traditional risk
194 factors. These results imply, that L1CAM is not a useful tool for preoperative treatment planning of
195 endometrial carcinoma.

196 Considering the common difficulties in assessing the depth of myometrial invasion preoperatively,
197 we wanted to test a model without MI as a parameter. Intraoperative tumor diameter is a more feasible
198 measure since it can be reliably evaluated by gross inspection (by the surgeon) even when frozen
199 section analysis is not available. In our study cohort the risk assessment model presented by Cox
200 Bauer et al. performed equally well independently of the presence of MI as a parameter (AUC > 0.8 ;
201 $P = 0.429$), suggesting that intraoperative tumor diameter could be used as an alternative to MI to

202 identify high risk disease, as indicated by previous studies [14,15]. The ideal cut off value for tumor
203 diameter that determines high risk disease needs to be established by further studies.

204 No serum markers have any established role in the treatment of endometrial carcinoma. Few studies
205 have addressed the potential clinical usefulness of serum L1CAM. Fogel et al. detected sL1 in the
206 blood of patients with an advanced L1CAM-positive ovarian or uterine carcinoma, but not in
207 healthy subjects or patients with other types of tumors, suggesting that sL1 could be used in
208 diagnostics or follow up of ovarian and uterine carcinoma [17]. Using a commercial ELISA-kit
209 optimized for serum samples, we were not able to confirm the results of the earlier report. Based on
210 our results, soluble L1CAM is not a useful marker of L1CAM positivity of endometrial carcinoma.

211 A strength of our study was its unselected cohort of patients with endometrial carcinoma treated at a
212 single tertiary care center with well-defined diagnostic and operative standards and systematic
213 follow-up procedures. The relatively high lymphadenectomy rate (192/241, 79.7 %) in the study
214 cohort improved the diagnostics of occult nodal disease permitting more accurate staging. In our
215 institution frozen section is not used to determine the depth of MI and data on MI had to be
216 extrapolated from final pathological reports.

217 In summary, we found a moderate concordance for L1CAM status between endometrial biopsies and
218 corresponding hysterectomy specimens. Preoperative L1CAM expression was associated with lymph
219 nodal and distant metastasis, but L1CAM did not significantly improve risk stratification algorithms
220 based on preoperative histology, tumor diameter and/or myometrial invasion. Interestingly, the
221 performance of risk stratification models did not depend on the presence of myometrial invasion as a
222 variable, suggesting that the more feasible tumor diameter could be used as a surrogate variable.
223 Based on our results, preoperative L1CAM cannot be recommended as a tool for stratifying patients
224 with endometrial carcinoma to lymphadenectomy.

225

226

227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265

Figure legend

Supplementary figure 1.

Figure 1. L1CAM immunohistochemical staining patterns in biopsies containing endometrial carcinoma. a) diffuse; b,c) heterogeneous (b and c from the same biopsy)

[1] Burke WM, Orr J, Leitaio M, Salom E, Gehrig P, Olawaiye AB, et al. Endometrial cancer: A review and current management strategies: Part I. *Gynecol.Oncol.* 2014 8;134:385-392.

[2] May K, Bryant A, Dickinson HO, Kehoe S, Morrison J. Lymphadenectomy for the management of endometrial cancer. *Cochrane Database Syst.Rev.* 2010 Jan 20;(1):CD007585. doi:CD007585.

[3] Mariani A, Webb MJ, Keeney GL, Haddock MG, Calori G, Podratz KC. Low-risk corpus cancer: Is lymphadenectomy or radiotherapy necessary? *Obstet.Gynecol.* 2000 6;182:1506-1519.

[4] Mariani A, Dowdy SC, Cliby WA, Gostout BS, Jones MB, Wilson TO, et al. Prospective assessment of lymphatic dissemination in endometrial cancer: a paradigm shift in surgical staging. *Gynecol.Oncol.* 2008 Apr;109:11-18.

[5] Kumar S, Podratz KC, Bakkum-Gamez JN, Dowdy SC, Weaver AL, McGree ME, et al. Prospective assessment of the prevalence of pelvic, paraaortic and high paraaortic lymph node metastasis in endometrial cancer. *Gynecol.Oncol.* 2014 1;132:38-43.

[6] Cox Bauer CM, Greer DM, Kram JJ, Kamelle SA. Tumor diameter as a predictor of lymphatic dissemination in endometrioid endometrial cancer *Gynecol.Oncol.* 2016 May;141:199-205.

[7] Mavromatis ID, Antonopoulos CN, Matsoukis IL, Frangos CC, Skalkidou A, Creatsas G, et al. Validity of intraoperative gross examination of myometrial invasion in patients with endometrial cancer: a meta-analysis *Acta Obstet.Gynecol.Scand.* 2012 Jul;91:779-793.

[8] Antonsen SL, Jensen LN, Loft A, Berthelsen AK, Costa J, Tabor A, et al. MRI, PET/CT and ultrasound in the preoperative staging of endometrial cancer - a multicenter prospective comparative study. *Gynecol.Oncol.* 2013 Feb;128:300-308.

[9] Alcazar JL, Dominguez-Piriz J, Juez L, Caparros M, Jurado M. Intraoperative Gross Examination and Intraoperative Frozen Section in Patients With Endometrial Cancer for Detecting Deep Myometrial Invasion: A Systematic Review and Meta-analysis *Int.J.Gynecol.Cancer* 2016 Feb;26:407-415.

266 [10] Soliman PT, Frumovitz M, Spannuth W, Greer MJ, Sharma S, Schmeler KM, et al.
267 Lymphadenectomy during endometrial cancer staging: Practice patterns among gynecologic
268 oncologists. *Gynecol.Oncol.* 2010 11;119:291-294.

269 [11] Frumovitz M, Slomovitz BM, Singh DK, Broaddus RR, Abrams J, Sun CC, et al. Frozen
270 section analyses as predictors of lymphatic spread in patients with early-stage uterine cancer
271 *J.Am.Coll.Surg.* 2004 Sep;199:388-393.

272 [12] Case AS, Rocconi RP, Straughn Jr. JM, Conner M, Novak L, Wang W, et al. A
273 prospective blinded evaluation of the accuracy of frozen section for the surgical management
274 of endometrial cancer. *Obstet.Gynecol.* 2006;108:1375-1379.

275 [13] Kumar S, Bandyopadhyay S, Semaan A, Shah JP, Mahdi H, Morris R, et al. The role of
276 frozen section in surgical staging of low risk endometrial cancer. *PLoS One* 2011;6:e21912.

277 [14] AlHilli MM, Podratz KC, Dowdy SC, Bakkum-Gamez JN, Weaver AL, McGree ME, et
278 al. Preoperative biopsy and intraoperative tumor diameter predict lymph node dissemination
279 in endometrial cancer. *Gynecol.Oncol.* 2013 Feb;128:294-299.

280 [15] Tuomi T, Pasanen A, Luomaranta A, Leminen A, Butzow R, Loukovaara M. Risk-
281 stratification of endometrial carcinomas revisited: A combined preoperative and
282 intraoperative scoring system for a reliable prediction of an advanced disease. *Gynecol.Oncol.*
283 2015 Apr;137:23-27.

284 [16] Trovik J, Wik E, Werner HM, Krakstad C, Helland H, Vandenput I, et al. Hormone
285 receptor loss in endometrial carcinoma curettage predicts lymph node metastasis and poor
286 outcome in prospective multicentre trial *Eur.J.Cancer* 2013 Nov;49:3431-3441.

287 [17] Fogel M, Gutwein P, Mechtersheimer S, Riedle S, Stoeck A, Smirnov A, et al. L1
288 expression as a predictor of progression and survival in patients with uterine and ovarian
289 carcinomas. *The Lancet* 2003 9/13;362:869-875.

290 [18] Zeimet AG, Reimer D, Huszar M, Winterhoff B, Puistola U, Azim SA, et al. L1CAM in
291 early-stage type I endometrial cancer: results of a large multicenter evaluation. *J.Natl.Cancer*
292 *Inst.* 2013 Aug 7;105:1142-1150.

293 [19] Bosse T, Nout RA, Stelloo E, Dreef E, Nijman HW, Jürgenliemk-Schulz IM, et al. L1 cell
294 adhesion molecule is a strong predictor for distant recurrence and overall survival in early
295 stage endometrial cancer: Pooled PORTEC trial results. *Eur.J.Cancer* 2014 10;50:2602-2610.

296 [20] Altevoigt P, Doberstein K, Fogel M. L1CAM in human cancer *Int.J.Cancer* 2015 Jun 25.

297 [21] Dellinger TH, Smith DD, Ouyang C, Warden CD, Williams JC, Han ES. L1CAM is an
298 independent predictor of poor survival in endometrial cancer - An analysis of The Cancer
299 Genome Atlas (TCGA) *Gynecol.Oncol.* 2016 May;141:336-340.

300 [22] van der Putten LJ, Visser NC, van de Vijver K, Santacana M, Bronsert P, Bulten J, et al.
301 L1CAM expression in endometrial carcinomas: an ENITEC collaboration study *Br.J.Cancer*
302 2016 Aug 9.

303 [23] Pasanen A, Tuomi T, Isola J, Staff S, Butzow R, Loukovaara M. L1 Cell Adhesion
304 Molecule as a Predictor of Disease-Specific Survival and Patterns of Relapse in Endometrial
305 Cancer *Int.J.Gynecol.Cancer* 2016 Oct;26:1465-1471.

306 [24] Gutwein P, Stoeck A, Riedle S, Gast D, Runz S, Condon TP, et al. Cleavage of L1 in
307 exosomes and apoptotic membrane vesicles released from ovarian carcinoma cells
308 *Clin.Cancer Res.* 2005 Apr 1;11:2492-2501.

309 [25] Zander H, Rawnaq T, von Wedemeyer M, Tachezy M, Kunkel M, Wolters G, et al.
310 Circulating levels of cell adhesion molecule L1 as a prognostic marker in gastrointestinal
311 stromal tumor patients. *BMC Cancer* 2011 May 22;11:189:1-7.

312 [26] Aktas B, Kasimir-Bauer S, Wimberger P, Kimmig R, Heubner M. Utility of mesothelin,
313 L1CAM and Afamin as biomarkers in primary ovarian cancer *Anticancer Res.* 2013
314 Jan;33:329-336.

315 [27] Pecorelli S. Revised FIGO staging for carcinoma of the vulva, cervix, and endometrium.
316 Int.J.Gynaecol.Obstet. 2009 May;105:103-104.
317 [28] Milam MR, Java J, Walker JL, Metzinger DS, Parker LP, Coleman RL, et al. Nodal
318 metastasis risk in endometrioid endometrial cancer. Obstet.Gynecol. 2012 Feb;119:286-292.
319 [29] Benedetti Panici P, Basile S, Salerno MG, Di Donato V, Marchetti C, Perniola G, et al.
320 Secondary analyses from a randomized clinical trial: age as the key prognostic factor in
321 endometrial carcinoma. Am.J.Obstet.Gynecol. 2014 Apr;210:363.e1-363.e10.
322 [30] Secord AA, Hasselblad V, Von Gruenigen VE, Gehrig PA, Modesitt SC, Bae-Jump V, et
323 al. Body mass index and mortality in endometrial cancer: A systematic review and meta-
324 analysis. Gynecol.Oncol. 2016 Jan;140:184-190.
325 [31] Landis JR, Koch GG. The measurement of observer agreement for categorical data.
326 Biometrics 1977 Mar;33:159-174.
327 [
328

Figure 1. L1CAM immunohistochemical staining patterns in biopsies containing endometrial carcinoma.
a) diffuse, b,c) heterogeneous (b and c from the same biopsy)

Table 1. Clinicopathologic data (n = 241).

Age (years) (mean $\pm$ SD)	67.3 $\pm$ 10.4
Body mass index (kg/m ²) (mean $\pm$ SD)	28.5 $\pm$ 6.5
Pelvic lymphadenectomy (number of cases, percent)	162 (67.2%)
Pelvic-aortic lymphadenectomy (number of cases, percent)	30 (12.4%)
Adjuvant therapy (number of cases, percent)	
Vaginal brachytherapy	133 (55.2%)
Whole pelvic radiotherapy	28 (11.6%)
Chemotherapy	6 (2.5%)
Chemotherapy and vaginal brachytherapy	13 (5.4%)
Chemotherapy and whole pelvic radiotherapy	33 (13.7%)
Histology (number of cases, percent)	
Endometrioid carcinoma	222 (92.1%)
Clear cell carcinoma	7 (2.9%)
Serous carcinoma	4 (1.7%)
Undifferentiated carcinoma	2 (0.8%)
Carcinosarcoma	5 (2.1%)
Neuroendocrine carcinoma	1 (0.4%)
Grade (number of cases, percent) (For endometrioid only, n = 222)	
Grade 1	128 (57.7%)
Grade 2	65 (29.3%)
Grade 3	29 (13.1%)
FIGO 2009 stage (number of cases, percent)	
IA	130 (53.9%)
IB	54 (22.4%)
II	13 (5.4%)
IIIA	12 (5.0%)
IIIB	1 (0.4%)
IIIC1	18 (7.5%)
IIIC2	7 (2.9%)
IVA	0 (0%)
IVB	6 (2.5%)

Table 2. Clinicopathologic characteristics according to L1CAM expression in preoperative endometrial samples, univariate analysis

Variable	Negative L1CAM	Positive L1CAM	OR (95% CI)	P
Stage II-IV	34/177 (19.2%)	24/64 (37.5%)	2.5 (1.3-4.7)	0.003
Positive pelvic and/or para-aortic lymph nodes ^a	12/174 (6.9%)	13/61 (21.3%)	3.7 (1.6-8.5)	0.002
Non-endometrioid carcinoma	6/177 (3.4%)	13/64 (20.3%)	7.3 (2.6-20)	<0.0001
Grade 3 (endometrioid only)	21/171 (12.3%)	8/51 (15.7%)	1.3 (0.55-3.2)	0.526
Myometrial invasion $\geq 50\%$	71/177 (40.1%)	27/64 (42.2%)	1.1 (0.61-1.9)	0.772
Tumor size ≥ 2 cm	47/166 (28.3%)	16/61 (26.2%)	0.90 (0.46-1.7)	0.756
Cervical stromal invasion	24/176 (13.6%)	9/63 (14.3%)	1.1 (0.46-2.4)	0.898
Positive peritoneal cytology	7/174 (4.0%)	11/63 (17.5%)	5.0 (1.9-14)	0.001
Age >65 years	94/177 (53.1%)	44/64 (68.8%)	1.9 (1.1-3.6)	0.030
Body mass index ≥ 30 kg/m ²	65/177 (36.7%)	17/64 (26.6%)	0.62 (0.33-1.2)	0.141

^a Stage IV cancers excluded

Table 3. Clinicopathological characteristics associated with L1CAM expression in preoperative endometrial samples, multivariate analysis

Variable	OR (95% CI)	P
Stage II-IV	1.4 (0.66-2.9)	0.389
Non-endometrioid carcinoma	4.4 (1.4-14)	0.010
Positive peritoneal cytology	2.6 (0.84-8.2)	0.097
Age >65 years	1.8 (0.92-3.3)	0.086

Table 4. Risk factors associated with advanced (stage IIIC-IV) endometrial carcinoma, analysis by multivariate risk models. Patients with available data for all risk factors were included.

	Models with L1CAM		Models without L1CAM	
	OR (95% CI)	P	OR (95% CI)	P
<i>Model HR-TD5cm-MI33%</i>	<i>1 (n = 225)</i>		<i>2 (n = 225)</i>	
Preoperative L1CAM	4.1 (1.5-11)	0.007		
Preoperative histology	3.7 (1.3-10)	0.012	5.2 (2.0-14)	0.001
Tumor size ≥ 5 cm	3.3 (1.2-8.7)	0.017	3.1 (1.2-7.9)	0.019
Myometrial invasion (MI)				
MI $\leq 33\%$	1		1	
33% < MI $\leq 66\%$	3.3 (0.72-15)	0.125	3.0 (0.67-14)	0.151
MI > 66%	11 (2.7-45)	0.001	8.2 (2.2-32)	0.002
<i>Model HR-TD5cm</i>	<i>3 (n = 228)</i>		<i>4 (n = 228)</i>	
Preoperative L1CAM	3.0 (1.2-7.7)	0.021		
Preoperative histology	4.8 (1.8-13)	0.002	6.6 (2.7-17)	<0.0001
Tumor size ≥ 5 cm	5.4 (2.2-14)	<0.0001	5.0 (2.1-12)	<0.0001
<i>Model HR-TD2cm-MI50%</i>	<i>5 (n = 227)</i>		<i>6 (n = 227)</i>	
Preoperative L1CAM	3.3 (1.3-8.3)	0.010		
Preoperative histology	4.3 (1.7-11)	0.003	5.8 (2.3-15)	<0.0001
Tumor size ≥ 2 cm	1.3 (0.33-5.3)	0.687	1.3 (0.33-5.2)	0.701
Myometrial invasion $\geq 50\%$	4.9 (1.7-14)	0.003	4.3 (1.5-12)	0.007
<i>Model HR- MI50%</i>	<i>7 (n = 241)</i>		<i>8 (n = 241)</i>	
Preoperative L1CAM	3.0 (1.2-7.4)	0.015		
Preoperative histology	4.0 (1.6-10)	0.003	5.4 (2.2-13)	<0.0001
Myometrial invasion $\geq 50\%$	6.1 (2.3-17)	<0.0001	5.3 (2.0-14)	0.001

HR = high risk histology (G3 or non-endometrioid); TD = tumor diameter; MI = myometrial invasion

Table 5. Areas under curve (AUC) for risk models predicting stage IIIC-IV endometrial carcinoma.

Risk assessment model	AUC (95% CI)	P (2-tailed)
1. HR-TD5cm-MI33%-L1CAM	0.879 (0.828-0.930)	
2. HR-TD5cm-MI33%	0.870 (0.813-0.928)	P = 0.882 vs. Model 1
3. HR-TD5cm-L1CAM	0.852 (0.778-0.925)	
4. HR-TD5cm	0.818 (0.730-0.906)	P = 0.613 vs. Model 3
5. HR-TD2cm-MI50%-L1CAM	0.841 (0.777-0.905)	
6. HR-TD2cm-MI50%	0.805 (0.717-0.894)	P = 0.602 vs. Model 5
7. HR-MI50%- L1CAM	0.833 (0.770-0.896)	
8. HR-MI50%	0.759 (0.659-0.859)	P = 0.289 vs. Model 7

HR = high risk histology (G3 or non-endometrioid); TD = tumor diameter; MI = myometrial invasion