<https://helda.helsinki.fi>

Sugar sensing by ChREBP/Mondo-Mlx - new insight into downstream regulatory networks and integration of nutrient-derived signals

Havula, Essi

2018-04

Havula , E & Hietakangas , V 2018 , ' Sugar sensing by ChREBP/Mondo-Mlx - new insight into downstream regulatory networks and integration of nutrient-derived signals ' , Current Opinion in Cell Biology , vol. 51 , pp. 89-96 . <https://doi.org/10.1016/j.ceb.2017.12.007>

<http://hdl.handle.net/10138/311367>

<https://doi.org/10.1016/j.ceb.2017.12.007>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

**Sugar sensing by ChREBP/Mondo-Mlx – new insight into downstream
regulatory networks and integration of nutrient-derived signals**

Essi Havula¹ & Ville Hietakangas^{2,3,*}

1. Charles Perkins Centre, School of Life and Environmental Sciences, The
University of Sydney, Sydney, NSW 2006, Australia

2. Faculty of Biological and Environmental Sciences, University of Helsinki, Finland.

3. Institute of Biotechnology, University of Helsinki, Finland.

* Correspondence: ville.hietakangas@helsinki.fi

Short title: ChREBP/Mondo-Mlx in sugar sensing

Body text word count: 3090

Highlights

- ChREBP/Mondo-Mlx is an intracellular sugar sensor of animals with conserved target genes and physiological functions
- ChREBP/Mondo-Mlx integrates multiple inputs through direct sensing of sugar and fatty acid-derived metabolites and cross-talk with other nutrient sensing pathways
- ChREBP/Mondo-Mlx controls several second-tier regulators, including transcription factors and hormones

Abstract

Animals regulate their physiology with respect to nutrient status, which requires nutrient sensing pathways. Simple carbohydrates, sugars, are sensed by the basic-helix-loop-helix leucine zipper transcription factors ChREBP/Mondo, together with their heterodimerization partner Mlx, which are well-established activators of sugar-induced lipogenesis. Loss of ChREBP/Mondo-Mlx in mouse and *Drosophila* leads to sugar intolerance, *i.e.* inability to survive on sugar containing diet. Recent evidence has revealed that ChREBP/Mondo-Mlx responds to sugar and fatty acid-derived metabolites through several mechanisms and cross-connects with other nutrient sensing pathways. ChREBP/Mondo-Mlx controls several downstream transcription factors and hormones, which mediate not only readjustment of metabolic pathways, but also control feeding behavior, intestinal digestion, and circadian rhythm.

Preface

Nutrient intake of animals displays a high degree of variation, which requires fine-tuned control of metabolic pathways. Animals regulate their metabolic homeostasis by continuously reading and converging signals originating from both within the organism and the outside environment. Macronutrients (sugars, amino acids and lipids) are sensed by nutrient sensing pathways that detect different dietary inputs and orchestrate an integrated adaptive response that is appropriate for that particular set of macronutrients. Simple carbohydrates, sugars, are sensed by several mechanisms, of which the so-called intracellular sugar sensing by ChREBP/Mondo-Mlx transcription factors is among the most conserved ones. Studies from mouse and *Drosophila* have shown the importance of this system in providing sugar tolerance; animals lacking ChREBP/Mondo-Mlx die rapidly on a diet containing sugars at a level, which is normally physiologically

tolerated [1,2]. We highlight here recent work, which has revealed several novel aspects of the ChREBP/Mondo-Mlx physiological function and regulation by nutrient-derived cues.

ChREBP/Mondo-Mlx – the conserved regulator of metabolic homeostasis

Carbohydrate response element binding protein (ChREBP, also known as MondoB and MLXIPL) and its paralog MondoA (MLXIP) are intracellular sugar sensors that are activated by glucose-6-phosphate (G6P) and other phosphorylated hexoses [3]. ChREBP and MondoA act together with their common heterodimerization partner Mlx, a necessary step for transcriptional activation. Although both are ubiquitously expressed, ChREBP is more prominently expressed in the liver and adipose tissue whereas MondoA is most highly expressed in the skeletal muscle. Muscle breaks down glucose via the glycolytic pathway to release energy that is used for muscle contraction and in fact many of the MondoA-Mlx target genes are involved in the glycolytic pathway [4,5]. Liver is the main site for *de novo* lipogenesis (DNL), where dietary sugars are converted into triacylglycerols (TAGs) and further transported to adipose tissue for storage. ChREBP target genes include lipogenic genes, such as *fatty acid synthase* and *acetyl-CoA-carboxylase*, both of which possess well-established ChREBP-Mlx binding sites, known as the carbohydrate response element (ChoRE), in their promoters [6,7]. ChREBP-Mlx is also a key regulator of glycolytic genes, such as *glucokinase regulatory protein*, *glyceraldehyde-3-phosphate dehydrogenase* and *L-type pyruvate kinase*, in the liver and adipose tissue [6-10]. Moreover, ChREBP-Mlx regulates the glucose-induced expression of *glucose-6-phosphate dehydrogenase* (*G6PDH*), the rate-limiting enzyme of the pentose phosphate pathway (PPP) [8].

The physiological function and key metabolic target genes of ChREBP/Mondo-Mlx are well conserved in animals and studies in *Drosophila* have revealed new insight into the physiological roles and tissue specificity of intracellular sugar sensing. Whereas the vertebrate genome encodes for two Mondo proteins, the fruit fly, *Drosophila*, has only one Mondo protein, displaying high expression in the fat body (corresponding to liver and adipose tissue), midgut (corresponding to small intestine) and Malpighian tubules (corresponding to kidney) [2]. Mondo-Mlx mediates a substantial proportion of *Drosophila* sugar-induced transcription and it activates conserved genes involved in glycolysis, lipogenesis, and the PPP, which is the most strongly over-represented

Mondo-Mlx target pathway in *Drosophila* [11]. The PPP has a key role in producing NADPH reductive power, which is needed for the biosynthesis of lipids. Indeed, inhibition of the Mondo-Mlx induced activation of glucose-6-phosphate dehydrogenase leads to a dramatic reduction of TAG stores in *Drosophila* larvae [11]. The PPP is also essential for sugar tolerance. The reductive power of NADPH generated by the PPP prevents oxidative damage upon high sugar feeding through the glutathione system, which contributes to the animal's ability to tolerate a high sugar diet [11,12].

In addition to conserved metabolic pathways, studies in *Drosophila* have revealed novel tissue-specific functions for sugar sensing. Mondo-Mlx is essential for the sugar-induced regulation of digestive enzymes in the midgut. Sugar feeding represses the expression of Amylases and Maltases, which are needed for digestion of complex carbohydrates [13,14]. This mechanism might have a role in regulating glucose intake, in order to prevent glucose overload [15]. Mondo-Mlx also regulates Malpighian tubule-specific glucose transporters, possibly in order to control excretion of excess glucose [11]. Furthermore, Mondo-Mlx is essential for normal flight muscle function and structure. Muscle-specific knockdown leads to accumulation of glycogen stores, possibly reflecting reduced glucose utilization [16]. In conclusion, ChREBP/Mondo-Mlx is a conserved sugar sensor within animals, with essential roles in sugar-induced lipogenesis and coordinating whole body glucose homeostasis by controlling carbohydrate digestion and transport.

ChREBP/Mondo-Mlx controls a gene regulatory network

In addition to direct regulation of key metabolic genes Mondo-Mlx is emerging as a master regulator of other regulatory genes, including transcription factors (TFs) (Figure 1). Mondo-Mlx directly regulates Krüppel-like factor 10 (KLF10, also known as TIEG1) and its *Drosophila* ortholog Cabut. In *Drosophila*, the *cabut* promoter has two high affinity ChoREs. *cabut* is the most strongly downregulated gene in *mlx* mutants and depletion of Cabut by RNAi induces sugar intolerance in *Drosophila* larvae [2]. Similarly, in rat primary hepatocytes KLF10 expression is activated by glucose stimulation and hampered by dominant negative Mlx [17]. Elevated KLF10 expression is also observed *in vivo* in mouse following feeding on a high fat/high sugar diet [18]. Adenoviral overexpression of KLF10 in primary hepatocytes suppressed the glucose-induced activation of ChREBP and several of its target genes, suggesting that KLF10

can act as a negative feedback regulator of ChREBP [17]. A different outcome was observed in loss-of-function experiments in *Drosophila*, where Cabut mediated downregulation of a gene set in response to sugar feeding [19]. Among the most strongly repressed Cabut target genes was *phosphoenolpyruvate carboxykinase* (*pepck*), which is a direct Cabut target and also a target of KLF10 in mouse [19,20]. PEPCCK is the rate-limiting enzyme of cataplerotic flux from the TCA cycle to mediate channeling of carbon towards gluco- and glyceroneogenesis. Consistent with high expression of *pepck*, *Drosophila mlx* mutants display high levels of glycerol and trehalose (a circulating form of glucose in insects), which could be suppressed by simultaneous loss of *pepck* [19]. Interestingly, Cabut also links sugar sensing to regulation of circadian rhythm. Deregulation of Cabut has a dramatic impact on the cycling of circadian metabolic gene output, while having no impact on the cycling of core clock genes [19]. Notably, in rat hepatocytes ChREBP-Mlx also directly regulates BHLHB2/DEC1, which has a well-established role in circadian regulation [21,22]. In conclusion, through direct regulation of downstream transcription factors, ChREBP/Mondo-Mlx-mediated sugar sensing appears to be closely coupled to the circadian clock. Considering the observations on re-setting of the peripheral circadian clock through timed feeding [23], it will be interesting to further explore the physiological importance of the observed interplay between these TFs.

ChREBP/Mondo-Mlx also controls the expression of TFs involved in lipid homeostasis. The Gli-similar (GLIS) transcription factor Sugarbabe has been long known as the earliest and strongest sugar-responsive gene in *Drosophila* [14]. Recently, *sugarbabe* was found to be a direct target of Mondo-Mlx. Loss- and gain of function analyses revealed that Sugarbabe contributes to the lipid homeostasis *in vivo*. It regulates fatty acid synthesis downstream of Mondo-Mlx by inducing the expression of lipogenic genes *ACC*, *FAS*, *AcCoAS* and *ATPCL* in response to sugar feeding [11]. Gli-similar transcription factor 2, the closest human homologue of Sugarbabe, was originally identified as a bifunctional transcription factor, regulating both the activation and repression of its target genes involved in kidney development [24]. Interestingly, variants of another *sugarbabe* homolog, GLIS3, are now established as one of the strongest known genetic risk factors for both type 1 and type 2 diabetes [25]. In addition to direct regulation of other TFs, ChREBP-Mlx also regulates downstream effectors indirectly. One such example is the fatty acid responsive nuclear receptor PPAR γ [26].

ChREBP expression is elevated during adipocyte differentiation and ChREBP activity is needed to maintain maximal transcriptional output of adipogenic PPAR γ . Consequently, ectopic activation of ChREBP promotes adipocyte differentiation. ChREBP likely affects PPAR γ indirectly through lipid biosynthesis as the activating effects of ChREBP depend on functional Fatty acid synthase [26]. In conclusion, ChREBP/Mondo-Mlx contributes to lipid homeostasis through its direct targets as well as through secondary TFs.

New hormonal targets of ChREBP/Mondo-Mlx

In addition to downstream transcription factors, ChREBP/Mondo-Mlx contributes to systemic regulation through hormonal signals, including FGF21. In cultured hepatocytes ChREBP-Mlx regulates the expression of *FGF21* in a glucose-dependent manner [8,27]. Levels of circulating FGF21 are rapidly increased in response to fructose in humans as well as mice [28,29]. Interestingly, both baseline and fructose-stimulated levels of FGF21 are significantly elevated in subjects with metabolic syndrome [28]. This is consistent with findings showing elevated activation of hepatic ChREBP in obese individuals [30]. Loss of FGF21 in turn leads to impaired *de novo* lipogenesis in mouse liver, which is accompanied by liver fibrosis following prolonged fructose exposure [29]. In addition to metabolic regulation, FGF21 also regulates sugar feeding. Mice lacking FGF21 consume elevated levels of sucrose, whereas ectopic FGF21 suppresses the intake of sugar in mice as well as cynomolgus monkeys [31,32]. The liver-derived FGF21 suppresses sugar feeding in mice by acting through the FGF21 receptor complex in the paraventricular nucleus of the hypothalamus [31]. A similar regulatory system likely exists in humans, since variants of *FGF21* are strongly associated with sweet preference, as judged by the consumption of candy [33].

Research in *Drosophila* has revealed new connections between Mondo-Mlx and hormonal metabolic regulation via TGF- β /Activin signaling. TGF- β /Activin ligand Dawdle is highly expressed in the *Drosophila* fat body, from where it is secreted [15]. Sugar feeding strongly elevates *dawdle* expression, while loss of *mlx* leads to impaired *dawdle* activation [11,15]. Moreover, Mondo-Mlx directly binds to a ChoRE at the *dawdle* promoter [11]. Dawdle is essential in metabolic regulation as mutants of *dawdle* show elevated circulating trehalose and glucose as well as high levels of glycogen and

triacylglycerol [34]. Dawdle acts through multiple target tissues, for example it acts on *Drosophila* insulin producing cells to promote insulin-like peptide secretion [34]. Moreover, Dawdle activates its receptor Baboon in intestinal enterocytes and suppresses the expression of genes encoding carbohydrate digestive enzymes, Amylases and Maltases, thus inhibiting carbohydrate digestion in response to excess systemic glucose [15]. Interestingly, Dawdle functionally cooperates with the Mondo-Mlx target TF Sugarbabe, which also contributes to the sugar-induced repression of amylase expression [11]. In conclusion, studies in mammals and *Drosophila* have opened new insight into the role of ChREBP/Mondo-Mlx on the systemic regulation of metabolism via hormones secreted by the liver and fat body, respectively.

Direct regulation of ChREBP/Mondo-Mlx by nutrient-derived cues

ChREBP/Mondo TFs respond to sugars via their N-terminal Glucose sensing module (GSM), which can be functionally divided into low-glucose inhibitory domain (LID) and glucose response activation conserved element (GRACE) [35]. Although direct structural evidence about ChREBP/Mondo activation mechanism is still missing, ample indirect evidence suggests that direct binding of G6P, and possibly other phosphorylated hexoses, lead to loss of intramolecular inhibition of GRACE by LID [36,37]. Subsequently activated ChREBP/Mondo-Mlx dissociates from 14-3-3, translocates to the nucleus, and binds to the ChoRE to activate its target genes (Figure 2A) [38-40]. Furthermore, glucose-responsive dephosphorylation as well as acetylation have also been shown to contribute to ChREBP activation [41,42]. A detailed representation of ChREBP/Mondo-Mlx activation has been presented elsewhere [3]. This core activation mechanism allows ChREBP/Mondo-Mlx to control gene expression in response to sugar availability. However, additional regulatory mechanisms and functional interplay with other nutrient-responsive TFs to integrate other nutrient-derived signals are beginning to emerge. One such signaling mechanism is O-GlcNacylation. The synthesis of the substrate for O-GlcNacylation, UDP-GlcNAc, through the hexosamine biosynthesis pathway (HBP), depends on availability of nutrients, such as glucose and glutamine [43]. Thereby, O-GlcNacylation is considered as a nutrient sensing mechanism. ChREBP is O-GlcNAc modified [44-46] and recent mass-spectrometric analysis have revealed multiple target sites [47]. Consistent with the glucose sensitivity of the HBP, the levels of ChREBP O-GlcNAc modification are increased upon exposure to high glucose [44,45]. O-GlcNAc modification increases

ChREBP protein stability and increases the expression of ChREBP target genes [44]. Pancreatic β -cells have an inherent glucose sensing capacity and ChREBP activation seems to involve a distinct mechanism in this setting. ChREBP forms a cytoplasmic complex with a Calcium binding protein Sorcin [48]. Calcium influx in response to high glucose releases ChREBP allowing its nuclear localization. In conclusion, ChREBP activity is regulated by sugars through multiple parallel mechanisms.

In addition to posttranslational modifications, ChREBP/Mondo-Mlx is also regulated at the transcriptional level. In mammals, the *ChREBP* gene encodes two isoforms, *ChREBP- α* and *ChREBP- β* , which are transcribed from different promoters [49]. ChREBP- α contains the N-terminal GSM and is thus directly activated by G6P. ChREBP- α stimulates the expression of the shorter ChREBP- β , which lacks inherent glucose sensing ability, but has a 20-fold higher transcriptional activity than ChREBP- α . The promoter of *ChREBP* contains a binding site for Liver X receptor (LXR), and it has been shown that LXR regulates the expression of *ChREBP* in the liver [50,51]. However, the effect of LXR on *ChREBP* expression appears to be independent of nutrient as the expression of *ChREBP* is not altered in high-sugar fed LXR mutant mice [52]. Hepatocyte nuclear factor 4 alpha (HNF-4 α) in turn has been shown to activate the expression of both *ChREBP* isoforms in response to glucose [53]. Moreover, HNF-4 α has been also shown to physically interact with ChREBP, and this interaction is promoted by glucose [53,54].

While ChREBP is activated in the liver of animals on a high sugar diet, a high lipid diet has an inhibitory effect. Furthermore, administration of fatty acids inhibits nuclear translocation and activation of ChREBP (Figure 2B) [55,56]. Fatty acids are oxidized into ketone bodies, such as β -hydroxybutyrate and acetoacetate. By using an *in vitro* binding assay with purified ChREBP and 14-3-3 as well as protein-free hepatocyte extracts, Nagakawa and coworkers showed that β -hydroxybutyrate and acetoacetate increase the binding between ChREBP and 14-3-3 [57]. This increased binding anchors ChREBP to the cytoplasm, preventing its target gene activation. Fatty acid administration also significantly elevates the levels of intracellular AMP [55]. AMP activates the AMP-activated protein kinase (AMPK), which phosphorylates ChREBP on Ser568 and inhibits its transcriptional activity, while having no impact on the nuclear

localization [55,58]. Interestingly, a recent study suggests that AMP can also inhibit ChREBP directly, through an allosteric mechanism. AMP binds to ChREBP and increases its affinity to 14-3-3, which prevents ChREBP nuclear translocation [58]. In conclusion, recent studies suggest that ChREBP acts as a direct sensor for several fatty acid-derived metabolites, thus integrating multiple nutrient-derived cues.

Cross-talk between ChREBP and other nutrient sensing pathways

In addition to nutrient sensing through direct allosteric regulation and posttranslational mechanisms, ChREBP/Mondo acts in close synergy with other nutrient sensing systems, including insulin signaling. A well-established example is SREBP, a target of insulin signaling, which controls synergistically lipogenic gene expression with ChREBP [59]. In *Drosophila*, Mondo was shown to act synergistically with Salt-inducible kinase 3 (SIK3), which is also a target of the insulin signaling pathway [12,60]. Mondo and SIK3 act cooperatively to activate the PPP upon sugar feeding [12]. Similarly to Mondo, SIK3 is essential for sugar tolerance and animals with reduced expression of both Mondo and SIK3 display extremely low tolerance towards dietary sugars [12]. Moreover, ChREBP O-GlcNacylation is inhibited by Foxo1, which is an inhibitory target of insulin signaling, thus providing additional cross-talk between insulin-mediated systemic glucose homeostasis and intracellular glucose sensing [46]. ChREBP also likely modulates insulin signaling, since genome-wide analysis of ChREBP chromatin binding in mouse liver and adipose tissue revealed significant enrichment in ChREBP binding near genes encoding components of the insulin signaling pathway [61].

Evidence about cross-talk between sugar sensing and amino acid sensing is also emerging (Figure 2C). Glutamine is a critical metabolic intermediate, acting as a carbon and ammonium carrier. Glutaminolysis is critical in replenishing the TCA cycle, when TCA cycle carbon is channeled to biosynthesis. Interestingly, glutamine modifies glucose sensing by inhibiting the transcriptional output of MondoA [62]. Consequently, glutamine inhibits the glucose-induced activation of *TXNIP*, a negative regulator of glucose transporter GLUT1 [63], thus increasing cellular glucose uptake [62]. Glutamine has no impact on MondoA nuclear localization or DNA binding, but it likely impacts MondoA cofactors [62]. MondoA can also control the uptake of glutamine. In endothelial cells infected with Kaposi's Sarcoma-associated Herpesvirus, glutamine

transporter SLC1A5 is activated in a MondoA-Mlx-dependent manner to facilitate glutaminolysis [64]. In conclusion, MondoA coordinates the intracellular balance of glucose and glutamine catabolism.

The mechanistic target of rapamycin (mTOR) is a conserved intracellular amino acid sensing system, which also has been shown to modulate the output of ChREBP/Mondo-Mlx. mTOR inhibitors increase the expression of MondoA-Mlx target gene *TXNIP* [65]. mTOR physically interacts with MondoA and sequesters it from forming an active complex with Mlx [65]. Similarly to MondoA, ChREBP is inhibited by mTOR through protein-protein interaction [66]. mTOR inhibition leads to increased expression of *TXNIP*, which causes death of pancreatic β -cells [66]. Interestingly, the levels of *TXNIP* are strongly elevated in diabetic islets, further suggesting that retaining the interaction between cellular amino acid and glucose sensing is essential for normal β -cell homeostasis. Recent evidence from *C. elegans* implies that Mondo-Mlx has a capacity to modulate mTOR signaling, since loss of *C. elegans* homolog of Mondo and Myc, MLL-1, leads to mTOR activation [67]. MLL-1 represses the expression of leucyl-tRNA synthetase *lars-1*, whose activity promotes mTOR activation via RAG-GTPases [67,68]. Through the regulation of mTOR signaling, MLL-1 is essential for promoting longevity [67]. In sum, mTOR and ChREBP/Mondo pathways appear to be closely interconnected, which potentially provides cross-coordination between phenotypic outputs of dietary sugars and amino acids.

Concluding remarks

To conclude, recent work in the field has revealed that ChREBP/Mondo-Mlx regulates a large regulatory network, including transcription factors and hormones, which controls not only metabolic pathways, but also feeding activity and intestinal digestion. In addition to intracellular sugar sensing, ChREBP/Mondo-Mlx integrates multiple nutrient-responsive inputs, placing it at the very heart of an interlinked regulatory system mediating physiological re-adjustment in response to nutrient intake. Such integrated control allows animals to flourish in highly variable nutrient landscapes.

However, several questions remain to be addressed. These include elucidating the structural basis of the function of the Glucose sensing module. This would allow better

understanding of the “nutrient sensor” mechanism of ChREBP/Mondo and possibly to enable design of new pharmaceuticals to modulate its activity. Given the availability of animal models, understanding of the physiological roles of ChREBP/Mondo in different tissues and cell types will certainly arise. Moreover, a wider perspective is needed to understand the natural variation in dietary sugar intake, and the variation between species and individuals with respect to their healthy nutrient landscape. As ChREBP/Mondo-Mlx is a key determinant in sugar tolerance in mice and *Drosophila*, it will be interesting to analyze its function in animals with differential sugar intake - and perhaps differential inherent sugar tolerance. Understanding the genetic factors predisposing humans to disease on a high sugar diet would allow disease prevention through personalized nutrition.

Acknowledgements

We thank Ryan Giblin and Mari Teesalu for feedback. Our research is supported by Academy of Finland (grant no. 286767 to V.H.), Sigrid Juselius Foundation (to VH), Novo Nordisk Foundation (NNF16OC0021460 to V.H.), Helsinki Institute of Life Science (to VH), and the Finnish Diabetes Research Foundation (to V.H.).

Figures

Figure 1. ChREBP/Mondo-Mlx is a master regulator of a sugar-induced gene regulatory network. In response to sugars, ChREBP/Mondo-Mlx activates the expression of a number of metabolic genes, but also other regulatory genes involved in the control of carbohydrate and lipid metabolism, sugar feeding and circadian rhythm. The figure summarizes key targets of mammalian ChREBP and MondoA as well as *Drosophila* Mondo. Abbreviations: ACC: acetyl-CoA carboxylase, Cbt: Cabut, DEC1: deleted in esophageal cancer 1, FAS: fatty acid synthase, FGF21: fibroblast growth factor 21, G6PDH: glucose-6-phosphate dehydrogenase, KLF10: krüppel-like factor 10, L-PK: L-type pyruvate kinase, PEPCK: phosphoenolpyruvate carboxykinase, PPAR γ : peroxisome proliferator-activated receptor gamma, TXNIP: thioredoxin-interacting protein.

Figure 2. ChREBP/Mondo-Mlx activity is regulated by many nutrient-derived cues. ChREBP/Mondo-Mlx is activated by glucose-6-phosphate (G6P) (A) and inhibited both by ketone bodies and AMP, whose levels depend on fatty acids (B). Direct binding of TOR to MondoA and ChREBP inhibits their function and glutamine inhibits the transcriptional activity of ChREBP-Mlx (C).

Selected references

Of special interest *

11. Mattila J, Havula E, Suominen E, Teesalu M, Surakka I, Hynynen R, Kilpinen H, Vaananen J, Hovatta I, Kakela R, et al.: **Mondo-Mlx Mediates Organismal Sugar Sensing through the Gli-Similar Transcription Factor Sugarbabe.** *Cell Rep* 2015, **13**:350-364.

Genome-wide analysis of Mondo-Mlx regulated genes in *Drosophila* reveals Gli-similar TF Sugarbabe and Activin ligand Dawdle as direct Mondo-Mlx targets.

26. Witte N, Muenzner M, Rietscher J, Knauer M, Heidenreich S, Nuotio-Antar AM, Graef FA, Fedders R, Tolkachov A, Goehring I, et al.: **The Glucose Sensor ChREBP Links De Novo Lipogenesis to PPARgamma Activity and Adipocyte Differentiation.** *Endocrinology* 2015, **156**:4008-4019.

Reveals the role of ChREBP in adipocyte differentiation through indirect control of PPARgamma via lipogenesis.

32. Talukdar S, Owen BM, Song P, Hernandez G, Zhang Y, Zhou Y, Scott WT, Paratala B, Turner T, Smith A, et al.: **FGF21 Regulates Sweet and Alcohol Preference.** *Cell Metab* 2016, **23**:344-349.

Shows that FGF21 inhibits sugar feeding in primates, in addition to mice.

61. Pongvarin N, Chang B, Imamura M, Chen J, Moolsuwan K, Sae-Lee C, Li W, Chan L: **Genome-Wide Analysis of ChREBP Binding Sites on Male Mouse Liver and White Adipose Chromatin.** *Endocrinology* 2015, **156**:1982-1994.

Genomewide targets of ChREBP in liver and adipose tissues suggests ChREBP-mediated control of the insulin signalling pathway. A valuable resource for future studies.

*Of outstanding interest ***

31. von Holstein-Rathlou S, BonDurant LD, Peltekian L, Naber MC, Yin TC, Claflin KE, Urizar AI, Madsen AN, Ratner C, Holst B, et al.: **FGF21 Mediates Endocrine Control of Simple Sugar Intake and Sweet Taste Preference by the Liver.** *Cell Metab* 2016, **23**:335-343.

Uncovers that FGF21, whose expression is activated in the liver by ChREBP, inhibits feeding of sugars.

58. Sato S, Jung H, Nakagawa T, Pawlosky R, Takeshima T, Lee WR, Sakiyama H, Laxman S, Wynn RM, Tu BP, et al.: **Metabolite Regulation of Nuclear Localization of Carbohydrate-response Element-binding Protein (ChREBP): ROLE OF AMP AS AN ALLOSTERIC INHIBITOR.** *J Biol Chem* 2016, **291**:10515-10527.

Shows biochemical and structural evidence for AMP as a novel ligand to inhibit ChREBP nuclear localization by enhancing the interaction between ChREBP and 14-3-3.

65. Kaadige MR, Yang J, Wilde BR, Ayer DE: **MondoA-Mlx transcriptional activity is limited by mTOR-MondoA interaction.** *Mol Cell Biol* 2015, **35**:101-110.

Demonstrates direct interaction between mTOR and MondoA to inhibit MondoA activation.

67. Nakamura S, Karalay O, Jager PS, Horikawa M, Klein C, Nakamura K, Latza C, Templer SE, Dieterich C, Antebi A: **Mondo complexes regulate TFEB via TOR inhibition to promote longevity in response to gonadal signals.** *Nat Commun* 2016, **7**:10944.

Demonstrates Mondo-mediated regulation of TOR activity and its impact for longevity in *C. elegans*.

References

1. Iizuka K, Bruick RK, Liang G, Horton JD, Uyeda K: **Deficiency of carbohydrate response element-binding protein (ChREBP) reduces lipogenesis as well as glycolysis.** *Proc Natl Acad Sci U S A* 2004, **101**:7281-7286.
2. Havula E, Teesalu M, Hyotylainen T, Seppala H, Hasygar K, Auvinen P, Oresic M, Sandmann T, Hietakangas V: **Mondo/ChREBP-Mlx-Regulated Transcriptional Network Is Essential for Dietary Sugar Tolerance in Drosophila.** *PLoS Genet* 2013, **9**:e1003438.
3. Havula E, Hietakangas V: **Glucose sensing by ChREBP/MondoA-Mlx transcription factors.** *Semin Cell Dev Biol* 2012.
4. Sans CL, Satterwhite DJ, Stoltzman CA, Breen KT, Ayer DE: **MondoA-Mlx heterodimers are candidate sensors of cellular energy status: mitochondrial localization and direct regulation of glycolysis.** *Mol Cell Biol* 2006, **26**:4863-4871.
5. Billin AN, Eilers AL, Coulter KL, Logan JS, Ayer DE: **MondoA, a novel basic helix-loop-helix-leucine zipper transcriptional activator that constitutes a positive branch of a max-like network.** *Mol Cell Biol* 2000, **20**:8845-8854.
6. Ishii S, Iizuka K, Miller BC, Uyeda K: **Carbohydrate response element binding protein directly promotes lipogenic enzyme gene transcription.** *Proc Natl Acad Sci U S A* 2004, **101**:15597-15602.
7. Jeong YS, Kim D, Lee YS, Kim HJ, Han JY, Im SS, Chong HK, Kwon JK, Cho YH, Kim WK, et al.: **Integrated expression profiling and genome-wide analysis of ChREBP targets reveals the dual role for ChREBP in glucose-regulated gene expression.** *PLoS One* 2011, **6**:e22544.
8. Ma L, Robinson LN, Towle HC: **ChREBP**Mlx* is the principal mediator of glucose-induced gene expression in the liver.** *J Biol Chem* 2006, **281**:28721-28730.
9. Ma L, Tsatsos NG, Towle HC: **Direct role of ChREBP.*Mlx* in regulating hepatic glucose-responsive genes.** *J Biol Chem* 2005, **280**:12019-12027.
10. Yamashita H, Takenoshita M, Sakurai M, Bruick RK, Henzel WJ, Shillinglaw W, Arnot D, Uyeda K: **A glucose-responsive transcription factor that regulates carbohydrate metabolism in the liver.** *Proc Natl Acad Sci U S A* 2001, **98**:9116-9121.
11. Mattila J, Havula E, Suominen E, Teesalu M, Surakka I, Hynynen R, Kilpinen H, Vaananen J, Hovatta I, Kakela R, et al.: **Mondo-Mlx Mediates Organismal Sugar Sensing through the Gli-Similar Transcription Factor Sugarbabe.** *Cell Rep* 2015, **13**:350-364.
12. Teesalu M, Rovenko BM, Hietakangas V: **Salt-Inducible Kinase 3 Provides Sugar Tolerance by Regulating NADPH/NADP⁺ Redox Balance.** *Curr Biol* 2017, **27**:458-464.

13. Benkel BF, Hickey DA: **Glucose Repression of Amylase Gene Expression in DROSOPHILA MELANOGASTER.** *Genetics* 1986, **114**:137-144.
14. Zinke I, Schutz CS, Katzenberger JD, Bauer M, Pankratz MJ: **Nutrient control of gene expression in Drosophila: microarray analysis of starvation and sugar-dependent response.** *EMBO J* 2002, **21**:6162-6173.
15. Chng WB, Bou Sleiman MS, Schupfer F, Lemaitre B: **Transforming growth factor beta/activin signaling functions as a sugar-sensing feedback loop to regulate digestive enzyme expression.** *Cell Rep* 2014, **9**:336-348.
16. Polak GL, Pasqualino A, Docherty JE, Beck SJ, DiAngelo JR: **The Regulation of Muscle Structure and Metabolism by Mio/dChREBP in Drosophila.** *PLoS One* 2015, **10**:e0136504.
17. Iizuka K, Takeda J, Horikawa Y: **Kruppel-like factor-10 is directly regulated by carbohydrate response element-binding protein in rat primary hepatocytes.** *Biochem Biophys Res Commun* 2011, **412**:638-643.
18. Kim JK, Lee KS, Chang HY, Lee WK, Lee JI: **Progression of diet induced nonalcoholic steatohepatitis is accompanied by increased expression of Kruppel-like-factor 10 in mice.** *J Transl Med* 2014, **12**:186.
19. Bartok O, Teesalu M, Ashwall-Fluss R, Pandey V, Hanan M, Rovenko BM, Poukkula M, Havula E, Moussaieff A, Vodala S, et al.: **The transcription factor Cabut coordinates energy metabolism and the circadian clock in response to sugar sensing.** *EMBO J* 2015, **34**:1538-1553.
20. Guillaumond F, Grechez-Cassiau A, Subramaniam M, Brangolo S, Peteri-Brunback B, Staels B, Fievet C, Spelsberg TC, Delaunay F, Teboul M: **Kruppel-like factor KLF10 is a link between the circadian clock and metabolism in liver.** *Mol Cell Biol* 2010, **30**:3059-3070.
21. Iizuka K, Horikawa Y: **Regulation of lipogenesis via BHLHB2/DEC1 and ChREBP feedback looping.** *Biochem Biophys Res Commun* 2008, **374**:95-100.
22. Honma S, Kawamoto T, Takagi Y, Fujimoto K, Sato F, Noshiro M, Kato Y, Honma K: **Dec1 and Dec2 are regulators of the mammalian molecular clock.** *Nature* 2002, **419**:841-844.
23. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U: **Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus.** *Genes Dev* 2000, **14**:2950-2961.
24. Zhang F, Nakanishi G, Kurebayashi S, Yoshino K, Perantoni A, Kim YS, Jetten AM: **Characterization of Glis2, a novel gene encoding a Gli-related, Kruppel-like transcription factor with transactivation and repressor functions. Roles in kidney development and neurogenesis.** *J Biol Chem* 2002, **277**:10139-10149.
25. Yang Y, Chan L: **Monogenic Diabetes: What It Teaches Us on the Common Forms of Type 1 and Type 2 Diabetes.** *Endocr Rev* 2016, **37**:190-222.
26. Witte N, Muenzner M, Rietscher J, Knauer M, Heidenreich S, Nuotio-Antar AM, Graef FA, Fedders R, Tolkachov A, Goehring I, et al.: **The Glucose Sensor ChREBP Links De Novo Lipogenesis to PPARgamma Activity and Adipocyte Differentiation.** *Endocrinology* 2015, **156**:4008-4019.
27. Iizuka K, Takeda J, Horikawa Y: **Glucose induces FGF21 mRNA expression through ChREBP activation in rat hepatocytes.** *FEBS Lett* 2009, **583**:2882-2886.

28. Dushay JR, Toschi E, Mitten EK, Fisher FM, Herman MA, Maratos-Flier E: **Fructose ingestion acutely stimulates circulating FGF21 levels in humans.** *Mol Metab* 2015, **4**:51-57.
29. Fisher FM, Kim M, Doridot L, Cunniff JC, Parker TS, Levine DM, Hellerstein MK, Hudgins LC, Maratos-Flier E, Herman MA: **A critical role for ChREBP-mediated FGF21 secretion in hepatic fructose metabolism.** *Mol Metab* 2017, **6**:14-21.
30. Eissing L, Scherer T, Todter K, Knippschild U, Greve JW, Buurman WA, Pinnschmidt HO, Rensen SS, Wolf AM, Bartelt A, et al.: **De novo lipogenesis in human fat and liver is linked to ChREBP-beta and metabolic health.** *Nat Commun* 2013, **4**:1528.
31. von Holstein-Rathlou S, BonDurant LD, Peltekian L, Naber MC, Yin TC, Claflin KE, Urizar AI, Madsen AN, Ratner C, Holst B, et al.: **FGF21 Mediates Endocrine Control of Simple Sugar Intake and Sweet Taste Preference by the Liver.** *Cell Metab* 2016, **23**:335-343.
32. Talukdar S, Owen BM, Song P, Hernandez G, Zhang Y, Zhou Y, Scott WT, Paratala B, Turner T, Smith A, et al.: **FGF21 Regulates Sweet and Alcohol Preference.** *Cell Metab* 2016, **23**:344-349.
33. Soberg S, Sandholt CH, Jespersen NZ, Toft U, Madsen AL, von Holstein-Rathlou S, Grevengoed TJ, Christensen KB, Bredie WLP, Potthoff MJ, et al.: **FGF21 Is a Sugar-Induced Hormone Associated with Sweet Intake and Preference in Humans.** *Cell Metab* 2017, **25**:1045-1053 e1046.
34. Ghosh AC, O'Connor MB: **Systemic Activin signaling independently regulates sugar homeostasis, cellular metabolism, and pH balance in Drosophila melanogaster.** *Proc Natl Acad Sci U S A* 2014, **111**:5729-5734.
35. Li MV, Chang B, Imamura M, Pongvarin N, Chan L: **Glucose-dependent transcriptional regulation by an evolutionarily conserved glucose-sensing module.** *Diabetes* 2006, **55**:1179-1189.
36. Davies MN, O'Callaghan BL, Towle HC: **Activation and repression of glucose-stimulated ChREBP requires the concerted action of multiple domains within the MondoA conserved region.** *Am J Physiol Endocrinol Metab* 2010, **299**:E665-674.
37. McFerrin LG, Atchley WR: **Evolution of the Max and Mlx networks in animals.** *Genome Biol Evol* 2011, **3**:915-937.
38. Sakiyama H, Wynn RM, Lee WR, Fukasawa M, Mizuguchi H, Gardner KH, Repa JJ, Uyeda K: **Regulation of nuclear import/export of carbohydrate response element-binding protein (ChREBP): interaction of an alpha-helix of ChREBP with the 14-3-3 proteins and regulation by phosphorylation.** *J Biol Chem* 2008, **283**:24899-24908.
39. Eilers AL, Sundwall E, Lin M, Sullivan AA, Ayer DE: **A novel heterodimerization domain, CRM1, and 14-3-3 control subcellular localization of the MondoA-Mlx heterocomplex.** *Mol Cell Biol* 2002, **22**:8514-8526.
40. Merla G, Howald C, Antonarakis SE, Reymond A: **The subcellular localization of the ChoRE-binding protein, encoded by the Williams-Beuren syndrome critical region gene 14, is regulated by 14-3-3.** *Hum Mol Genet* 2004, **13**:1505-1514.
41. Kabashima T, Kawaguchi T, Wadzinski BE, Uyeda K: **Xylulose 5-phosphate mediates glucose-induced lipogenesis by xylulose 5-phosphate-activated protein phosphatase in rat liver.** *Proc Natl Acad Sci U S A* 2003, **100**:5107-5112.

42. Bricambert J, Miranda J, Benhamed F, Girard J, Postic C, Dentin R: **Salt-inducible kinase 2 links transcriptional coactivator p300 phosphorylation to the prevention of ChREBP-dependent hepatic steatosis in mice.** *J Clin Invest* 2010, **120**:4316-4331.
43. Hanover JA, Krause MW, Love DC: **The hexosamine signaling pathway: O-GlcNAc cycling in feast or famine.** *Biochim Biophys Acta* 2010, **1800**:80-95.
44. Guinez C, Filhoulaud G, Rayah-Benhamed F, Marmier S, Dubuquoy C, Dentin R, Moldes M, Burnol AF, Yang X, Lefebvre T, et al.: **O-GlcNAcylation increases ChREBP protein content and transcriptional activity in the liver.** *Diabetes* 2011, **60**:1399-1413.
45. Park MJ, Kim DI, Lim SK, Choi JH, Han HJ, Yoon KC, Park SH: **High glucose-induced O-GlcNAcylated carbohydrate response element-binding protein (ChREBP) mediates mesangial cell lipogenesis and fibrosis: the possible role in the development of diabetic nephropathy.** *J Biol Chem* 2014, **289**:13519-13530.
46. Ido-Kitamura Y, Sasaki T, Kobayashi M, Kim HJ, Lee YS, Kikuchi O, Yokota-Hashimoto H, Iizuka K, Accili D, Kitamura T: **Hepatic FoxO1 integrates glucose utilization and lipid synthesis through regulation of ChREBP O-glycosylation.** *PLoS One* 2012, **7**:e47231.
47. Yang AQ, Li D, Chi L, Ye XS: **Validation, Identification, and Biological Consequences of the Site-specific O-GlcNAcylation Dynamics of Carbohydrate-responsive Element-binding Protein (ChREBP).** *Mol Cell Proteomics* 2017, **16**:1233-1243.
48. Noordeen NA, Meur G, Rutter GA, Leclerc I: **Glucose-induced nuclear shuttling of ChREBP is mediated by sorcin and Ca(2+) ions in pancreatic beta-cells.** *Diabetes* 2012, **61**:574-585.
49. Herman MA, Peroni OD, Villoria J, Schon MR, Abumrad NA, Bluher M, Klein S, Kahn BB: **A novel ChREBP isoform in adipose tissue regulates systemic glucose metabolism.** *Nature* 2012, **484**:333-338.
50. Cha JY, Repa JJ: **The liver X receptor (LXR) and hepatic lipogenesis. The carbohydrate-response element-binding protein is a target gene of LXR.** *J Biol Chem* 2007, **282**:743-751.
51. Gauthier K, Billon C, Bissler M, Beylot M, Lobaccaro JM, Vanacker JM, Samarut J: **Thyroid hormone receptor beta (TRbeta) and liver X receptor (LXR) regulate carbohydrate-response element-binding protein (ChREBP) expression in a tissue-selective manner.** *J Biol Chem* 2010, **285**:28156-28163.
52. Denechaud PD, Bossard P, Lobaccaro JM, Millatt L, Staels B, Girard J, Postic C: **ChREBP, but not LXRs, is required for the induction of glucose-regulated genes in mouse liver.** *J Clin Invest* 2008, **118**:956-964.
53. Meng J, Feng M, Dong W, Zhu Y, Li Y, Zhang P, Wu L, Li M, Lu Y, Chen H, et al.: **Identification of HNF-4alpha as a key transcription factor to promote ChREBP expression in response to glucose.** *Sci Rep* 2016, **6**:23944.
54. Adamson AW, Suchankova G, Rufo C, Nakamura MT, Teran-Garcia M, Clarke SD, Gettys TW: **Hepatocyte nuclear factor-4alpha contributes to carbohydrate-induced transcriptional activation of hepatic fatty acid synthase.** *Biochem J* 2006, **399**:285-295.
55. Kawaguchi T, Osatomi K, Yamashita H, Kabashima T, Uyeda K: **Mechanism for fatty acid "sparing" effect on glucose-induced transcription: regulation of carbohydrate-responsive element-binding protein by AMP-activated protein kinase.** *J Biol Chem* 2002, **277**:3829-3835.

56. Dentin R, Benhamed F, Pegorier JP, Fougelle F, Viollet B, Vaulont S, Girard J, Postic C: **Polyunsaturated fatty acids suppress glycolytic and lipogenic genes through the inhibition of ChREBP nuclear protein translocation.** *J Clin Invest* 2005, **115**:2843-2854.
57. Nakagawa T, Ge Q, Pawlosky R, Wynn RM, Veech RL, Uyeda K: **Metabolite regulation of nucleo-cytosolic trafficking of carbohydrate response element-binding protein (ChREBP): role of ketone bodies.** *J Biol Chem* 2013, **288**:28358-28367.
58. Sato S, Jung H, Nakagawa T, Pawlosky R, Takeshima T, Lee WR, Sakiyama H, Laxman S, Wynn RM, Tu BP, et al.: **Metabolite Regulation of Nuclear Localization of Carbohydrate-response Element-binding Protein (ChREBP): ROLE OF AMP AS AN ALLOSTERIC INHIBITOR.** *J Biol Chem* 2016, **291**:10515-10527.
59. Dentin R, Girard J, Postic C: **Carbohydrate responsive element binding protein (ChREBP) and sterol regulatory element binding protein-1c (SREBP-1c): two key regulators of glucose metabolism and lipid synthesis in liver.** *Biochimie* 2005, **87**:81-86.
60. Wang B, Moya N, Niessen S, Hoover H, Mihaylova MM, Shaw RJ, Yates JR, 3rd, Fischer WH, Thomas JB, Montminy M: **A hormone-dependent module regulating energy balance.** *Cell* 2011, **145**:596-606.
61. Pongvarin N, Chang B, Imamura M, Chen J, Moolsuwan K, Sae-Lee C, Li W, Chan L: **Genome-Wide Analysis of ChREBP Binding Sites on Male Mouse Liver and White Adipose Chromatin.** *Endocrinology* 2015, **156**:1982-1994.
62. Kaadige MR, Looper RE, Kamalanaadhan S, Ayer DE: **Glutamine-dependent anapleurosis dictates glucose uptake and cell growth by regulating MondoA transcriptional activity.** *Proc Natl Acad Sci U S A* 2009, **106**:14878-14883.
63. Wu N, Zheng B, Shaywitz A, Dagon Y, Tower C, Bellinger G, Shen CH, Wen J, Asara J, McGraw TE, et al.: **AMPK-dependent degradation of TXNIP upon energy stress leads to enhanced glucose uptake via GLUT1.** *Mol Cell* 2013, **49**:1167-1175.
64. Sanchez EL, Carroll PA, Thalhofer AB, Lagunoff M: **Latent KSHV Infected Endothelial Cells Are Glutamine Addicted and Require Glutaminolysis for Survival.** *PLoS Pathog* 2015, **11**:e1005052.
65. Kaadige MR, Yang J, Wilde BR, Ayer DE: **MondoA-Mlx transcriptional activity is limited by mTOR-MondoA interaction.** *Mol Cell Biol* 2015, **35**:101-110.
66. Chau GC, Im DU, Kang TM, Bae JM, Kim W, Pyo S, Moon EY, Um SH: **mTOR controls ChREBP transcriptional activity and pancreatic beta cell survival under diabetic stress.** *J Cell Biol* 2017, **216**:2091-2105.
67. Nakamura S, Karalay O, Jager PS, Horikawa M, Klein C, Nakamura K, Latza C, Templer SE, Dieterich C, Antebi A: **Mondo complexes regulate TFEB via TOR inhibition to promote longevity in response to gonadal signals.** *Nat Commun* 2016, **7**:10944.
68. Han JM, Jeong SJ, Park MC, Kim G, Kwon NH, Kim HK, Ha SH, Ryu SH, Kim S: **Leucyl-tRNA synthetase is an intracellular leucine sensor for the mTORC1-signaling pathway.** *Cell* 2012, **149**:410-424.