<https://helda.helsinki.fi>

Decreasing Trend in the Use and Long-Term Use of Benzodiazepines Among Young Adults

Tähkäpää, Sanna-Mari

2018-03-14

Tähkäpää , S-M , Saastamoinen , L , Airaksinen , M , Tuulio-Henriksson , A , Aalto-Setälä , T & Kurko , T 2018 , ' Decreasing Trend in the Use and Long-Term Use of Benzodiazepines Among Young Adults ' , Journal of Child and Adolescent Psychopharmacology , vol. 28 , no. 4 , pp. 279-284 . <https://doi.org/10.1089/cap.2017.0140>

<http://hdl.handle.net/10138/311658>

<https://doi.org/10.1089/cap.2017.0140>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Sanna-Mari Tähkää¹, Leena Saastamoinen^{2*}, Terhi Kurko², Annamari Tuulio-Henriksson^{2,3}, Terhi Aalto-Setälä^{4,5}, Marja Airaksinen⁶

Decreasing trend in the use and long-term use of benzodiazepines among young adults

1 Tikkurilan Uusi Apteekki, Talvikkitie 37, 01300 Vantaa; Finland

2 Research unit, The Social Insurance Institution, Po. Box 450, 00056 Kela, Finland

3Department of Psychology and Logopedics, University of Helsinki, Helsinki, Finland.

4 The Social Insurance Institution, Po. Box 450, 00056 Kela, Finland

5 Family Counselling Center, Helsinki, Finland.

6 Clinical Pharmacy Group, Division of Pharmacology and Pharmacotherapy, Faculty of Pharmacy, University of Helsinki. P.O. Box 56, 00014 University of Helsinki, Finland

*Corresponding author: Terhi Kurko, terhi.kurko@kela.fi, +358-50-5219905

29

30 **Objective** Patterns of benzodiazepine (BZD) use and long-term use among young adults are not well
31 known. Our aim was to study trends in BZD use and long-term use among 18–25-year-old young
32 adults by gender and active substance in a nationwide retrospective longitudinal register-based
33 setting.

34 **Methods** All Finns aged 18–25 years with reimbursed purchases of BZDs in 2006–2014 recorded to
35 the Finnish Prescription Register were included. Annual prevalence rates of BZD use and long-term
36 use among young adults were reported overall, and according to gender, drug group (anxiolytic or
37 hypnotic) and active substance. Long-term use of BZDs was defined as purchasing ≥ 180 Defined
38 Daily Doses (DDD) in at least two drug purchases during a calendar year.

39 **Results** Overall prevalence of BZD use among young adults decreased from 24.0 to 18.8 per 1000
40 inhabitants in 2006–2014. Prevalence of long-term use decreased from 5.5 to 3.3 per 1000 inhabitants.
41 Overall BZD use was higher among females, whereas long-term use was more common among males.
42 Use of anxiolytics was more common than use of hypnotics. Oxazepam, alprazolam, zopiclone and
43 zolpidem were the most used BZDs, whereas alprazolam and clonazepam were the substances with
44 most long-term use. The use and long-term use of BZDs have decreased annually since 2008 among
45 Finnish young adults. Further research is needed to investigate the reasons behind the decline.

46

47

48 **Keywords:** Benzodiazepines, young adults, anxiolytics, hypnotics

49

50

51

52

53

54

55

56

57 Introduction

58 Young adulthood is a developmentally critical phase associated with several challenges, such as
59 achievement of independence, identity formation and important decision making for the future (Patel
60 et al. 2007). Many mental disorders become prevalent during adolescence and young adulthood, and
61 approximately three fourth of lifetime psychiatric disorders have emerged by the age of 24 (Kessler
62 et al.2005). Studies have shown that psychotropic drug use has increased among adolescents and
63 young adults in recent decades (Autti-Rämö et al 2009; Hsia et al 2009, Hartz et al. 2016). However,
64 most of the pharmacoepidemiological research assessing the use and long-term use of BZDs has
65 focused on adult population or the aged (Kurko et al. 2016) who consume BZDs more commonly
66 than younger age groups and are at an increased risk for adverse effects associated with regular BZD
67 use (Barker et al. 2004, Glass et al. 2005)

68
69 Previous studies on BZD use among young adults have concerned specific populations, such as
70 patients with substance use disorders, or focus on abuse of BZDs (McCabe et al 2005, Kornor et al.
71 2010, Riska et al. 2014). In studies assessing BZD use in general populations, young adults have been
72 included as one of the age-groups (Lagnaoui et al 2004, Cunningham et al. 2010, Hollingworth et al.
73 2010, Nordfjaern et al 2012, Olfson et al. 2015). In United States, the prevalence of BZD use was 2.6
74 % and 0.4 % for long-term use among young adults (aged 18–35) (Olfson et al. 2015). In a Norwegian
75 study cohort, 4 % of young adults aged 18–25 years had received a prescription for BZDs, 1 % used
76 BZDs at least 180 DDD per year, and 0.5 % were defined as chronic users (Nordfjaern et al. 2012).
77 Among Taiwanese incident BZD users (aged 16–19), 5 % were long-term users (Yeh et al. 2011)

78
79 There is limited population-based knowledge on the use of BZDs among young people due to the fact
80 that most research has been conducted either among adults in general or among older adults (Kurko
81 et al 2016). The objective of this study was to assess nationwide trends in the prevalence of BZD use
82 and long-term use among young adults 18–25 years of age over a nine-year period of 2006–2014
83 overall and according to gender and active substance.

84 Materials and Methods

86 Data sources

87
88 This nationwide, retrospective register-based study is based on data from the Finnish Prescription
89 Register of the Social Insurance Institution of Finland. The Finnish National Health Insurance covers

all Finnish residents. The Prescription Register contains individual level information on all reimbursed drug purchases dispensed to Finnish residents in outpatient care. In the database, all drugs are classified according to the World Health Organization's, (WHO 2017) Anatomical Therapeutic Chemical (ATC) classification system. The data compiled for the present study contains information on each patient's personal identifier based on person's personal identification number in encrypted form, age at the end of the year, gender, ATC code of the dispensed drug, dispensing date, number of dispensed packages and number of defined daily doses (DDDs) dispensed. According to Finnish legislation, ethical approval was not required to conduct this register-based study with anonymized patient data. Permission to use data was given by the register holder, the Social Insurance Institution of Finland.

The BZDs included in this study are presented in supplementary table 1. These were anti-epileptic clonazepam (ATC-code N03AE01), traditional BZD anxiolytics (N05BA), traditional BZD hypnotics (N05CD), BZD related drugs, also called z-hypnotics (N05CF) and combination of chlordiazepoxide and amitriptyline (N06CA01). Purchases of orally administered dosage forms were included in this study with the exception of oral suspensions.

The study population consisted of all 18–25-year-old young adults with at least one reimbursed purchase of BZD between 1 January 2006 and 31 December 2014. Young adults were selected as a study population because the majority (94.3%, n=68 167) of 0–25-year-old BZD users (n=72 266) were 18–25-year-old young adults (Figure 1), and the definition for DDD by WHO (2016) concerns only adults. Person's age was calculated at the end of each calendar year. Numbers of BZD users, dispensed prescriptions and persons aged 0–25 covered by National Health Insurance in 2014 are presented in Figure 1.

Study variable construction

Anti-epileptic clonazepam was included in this study when used for other than epileptic indications, i.e. when purchased without special reimbursement for epilepsy. In the Finnish system, patients' entitlement for special reimbursement is based on doctor's certificate fulfilling certain diagnosis criteria. These purchases with special reimbursements are distinguished in a register with special

coding. In the non-epileptic indications, the dosing is lower compared to dosing for epilepsy (8mg). The dosage of 1mg was used as the defined daily dose for clonazepam use based on the literature and clinical experience (White 2009, Drug and Alcohol Services 2012, Ashton 2013).

127

BZDs in ATC-classes N03AE01, N05BA, N06CA01 were classified as anxiolytics, and drugs in ATC-classes N05CD and N05CF as hypnotics.

130

131

Reimbursed drug purchase of BZDs was used as a proxy for BZD use (Haukka et al 2007). Long-term use of BZDs was defined as at least 180 DDD purchases of BZDs and two or more drug purchases during a calendar year corresponding to six months' use. This definition of six months' use was based on the WHO's recommendation as the definition for long-term use and a systematic review concerning long-term use BZD use (World Health Organization 1996; Kurko et al 2015). Drug consumption of BZDs per patient per year was calculated by using DDDs recommended by WHO (2016). DDDs of each patient's drug purchases were combined according to the patient's personal identifier.

140

One-year prevalence of BZD use and long-term BZD use were assessed overall and according to gender, drug group (anxiolytic or hypnotic, based on the ATC-classification) and active substance during 2006–2014. Prevalence rates (number of users per 1000 inhabitants) were calculated according to the number of the same age Finnish population.

145

146 Statistical analysis

While this study concerns a whole population, data were analyzed with descriptive statistical methods. Binomial distribution was used to compare changes in prevalence rates between 2006 and 2014. For these prevalence rates, the 95% confidence intervals were calculated. The overall yearly trends of each studied drug were assessed by comparing these confidence intervals. If the confidence intervals were not overlapping, the trend during the assessed period was considered to be statistically significantly changed. Descriptive data analyses were performed using statistical software IBM SPSS Statistics version 22.0 and binomial distribution was tested by R version 3.1.3.2015.

154

155

156 Results

157 Overall prevalence of BZD use among young adults decreased statistically significantly, by 21.5%
158 from 24.0 to 18.8 / 1000 inhabitants over a nine-year period (Figure 2). The annual prevalence of
159 BZD use increased between 2006 and 2008, the highest prevalence (28.6 / 1000 inhabitants) being
160 observed in 2008. The use of BZDs was more common in females than males during the study period.
161 In 2014, prevalence of BZD use among females was 22.8 / 1000 inhabitants and among males 15.1 /
162 1000 inhabitants.

163
164 Overall prevalence of long-term use of BZDs among 18–25-year-old Finns decreased statistically
165 significantly, by 39.8% from 5.5 to 3.3 per 1000 inhabitants during 2006–2014. In 2014, the last
166 studied year, 17.5% of all 18–25-year-old BZD users were defined as long-term users. The prevalence
167 of long-term use was higher among males than females during the entire study period.

168
169 Use of anxiolytic BZDs was more common than use of hypnotics among Finnish young adults (Figure
170 3). The overall prevalence of anxiolytic use decreased 20.6%, from 15.2 to 12.1 / 1000 inhabitants
171 during the study period. The use of hypnotics decreased from 12.5 to 8.5 / 1000 inhabitants, 31.8%
172 between 2006 and 2014. Long-term use of anxiolytics was more common than long-term use of
173 hypnotics among young adults during the study period. In 2014, the prevalence of long-term use of
174 anxiolytics was 2.6 / 1000 inhabitants and hypnotics 0.7 / 1000 inhabitants. Both long-term use of
175 anxiolytics and long-term use of hypnotics decreased during 2006–2014.

176
177 Oxazepam and alprazolam were the most commonly used anxiolytics. The most commonly used
178 hypnotics were z-drugs zopiclone and zolpidem. Over the nine-year study period, there was a decline
179 in the use of almost every studied substance. Alprazolam use decreased the most (53.5%) between
180 2006 and 2014. Only the prevalence of oxazepam use increased by 17.4%, from 5.3 in 2006 to 6.2
181 per 1000 inhabitants in 2014. Use of diazepam and clonazepam was lower than that of oxazepam and
182 alprazolam. During the study period, prevalence of diazepam use ranged between 2.0 and 3.4 / 1000
183 inhabitants, while clonazepam use ranged between 1.6 and 2.3 / 1000 inhabitants, respectively.

184
185 Long-term use of BZDs was relatively low among young adults during the study period. The highest
186 prevalence of long-term use was observed in alprazolam users during 2006–2009 and in clonazepam
187 users during 2010–2014. The substance-level prevalence of long-term use decreased in users of all
188 BZD substances between 2006 and 2014. The decline of long-term use was most significant in
189 diazepam (58.7 %) and alprazolam (57.9 %) users. Compared to other BZDs, long-term use was more

190 prevalent among subjects using diazepam and clonazepam. The relative proportions of long-term use
191 among diazepam and clonazepam users were higher compared to other BZDs. More than half (59.9–
192 64.6 %) of clonazepam users were long-term users during 2006–2014. Corresponding proportions
193 were 24.9–37.5% among diazepam users and 26.8–32.3% among alprazolam users during the study
194 period.

195

196

197

198 Discussion

199 To our best knowledge, the present study is the first nationwide register-based study focusing on use
200 and long-term use of BZDs among young adults. Therefore, the present study offers clinically
201 significant new information on BZD use and long-term use of this less studied user group. We found
202 a decreasing trend of both BZD use and long-term use among Finnish young adults during the
203 investigated nine-year study period. The overall prevalence of BZD use decreased by one fifth and
204 long-term use of BZDs decreased by over one third between 2006 and 2014. Overall BZD use was
205 more common among females, whereas long-term use was more common among males. The four
206 most commonly used BZDs among young adults were oxazepam, alprazolam, zopiclone and
207 zolpidem. Oxazepam was the only substance with increased use between 2006 and 2014. Alprazolam
208 and clonazepam were the substances mostly used for long-term.

209

210 This study was based on data from the Finnish Prescription Register covering all reimbursed BZD
211 purchases in Finland among whole population during the investigated period. Key strengths of this
212 study are the nationwide data and a long nine-year study period. Completeness and accuracy of
213 records in The Finnish Prescription Register is high.

214

215 Drug purchase was used as a proxy for BZD use, because there is concordance between prescription
216 database information and self-reported BZD use (Haukka et al 2007). However, this study may
217 underestimate the total BZD use among young adults in Finland. The Finnish Prescription Register
218 does not record non-reimbursed drug purchases, inpatients' drug use or illegal drug use (Furu et al
219 2010). Moreover, some diazepam and oxazepam purchases in small package sizes are not recorded
220 in the Prescription Register, as they are not usually reimbursable. The coverage of the register was

221 76–90% annually compared to wholesale data of BZD provided by the Finnish Medical Agency
222 (2017). Information on patient’s diagnosis or severity of psychiatric condition is not included in the
223 Prescription Register.

224

225 Previous studies on BZD use and long-term use have varied considerably in terms of definition for
226 long-term use, time-period of the study and age group making it difficult to compare the prevalence
227 rates between studies. There may also be differences in treatment practices between countries.
228 However, prevalence of BZD use (1.9 %) among Finnish young adults in 2014 was similar to that
229 reported in France (1.8 %) but lower than reported in studies in the U.S. (2.6 %), Norway (4.0 %) and
230 Taiwan (6.7 %) (Lagnaoui et al 2004, Yeh et al. 2011, Nordfjaern et al. 2012, Olfson et al. 2015).
231 Long-term use of BZDs (0.3 %) by Finnish young adults was slightly less common than that reported
232 among U.S. young adults (0.4 %) in 2008 (Olfson et al. 2015), and among a sample of French young
233 adults (0.6 %) in 2001 (Lagnaoui et al. 2004). Further, the prevalence of long-term use in Finland
234 was lower than that observed in a Norwegian study cohort (1 %) during 2004–2008 (Nordfjaern
235 2012), and among Taiwanese incident BZD users (5 %) in 2001–2005 (Yeh et al 2011).

236

237 It was not possible to include assessment of indications of BZD prescriptions in the present study.
238 However, results showed that BZDs classified as anxiolytics were more commonly used than
239 hypnotics, which may suggest that BZDs are more often used in anxiety than sleep disorders among
240 young adults. At a substance-level, decreasing trend was observed among almost all BZDs except
241 oxazepam, which was the only substance with increased use. In the last studied year, oxazepam was
242 the most commonly used BZD among Finnish young adults. Increased use of oxazepam may be due
243 to its lower abuse liability (Griffiths et al 1984). This study found that long-term use among oxazepam
244 users was relatively uncommon. However, a recent study showed that oxazepam users were more
245 prone to dose escalation compared with diazepam users (Tvete et al 2016). Despite the decreasing
246 trend of long-term use overall and at a substance-level, long-term use was common among users of
247 high or medium potency BZDs including clonazepam, diazepam and alprazolam. In this study, a
248 majority of all clonazepam users used the substance for long-term. Similarly, a recent study found
249 that clonazepam use is often associated with dose escalation (Alessi-Severini et al. 2016). On the
250 other hand, in the study of Cloos et al. (2015), clonazepam was associated with a lower risk for high-
251 dose use compared to hypnotics and anxiolytics alprazolam and prazepam, which were associated
252 with the highest risk for high-dose use.

253

254 The observed gender difference in overall BZD use is consistent with other studies showing that use
255 of BZDs is more common among females than males (Petitjean et al. 2007, Lagnaoui et al 2004,
256 Cunningham et al. 2010, Olfson et al 2015). The higher prevalence of anxiety and sleep disorders in
257 females during young adulthood partly explains this finding (Wittchen et al. 1998, Suvisaari et al.
258 2009). Yet, in our study, long-term use of BZDs was more prevalent in males than females. This
259 could be partly explained by the previous finding reporting non-medical, e.g. self-reported possible
260 abuse or misuse, use of anxiolytics and hypnotics in adolescence to be significantly more common in
261 males than females in Finland compared to many other European countries (Kokkevi et al 2008).
262 However, our study did not assess the possible abuse of BZDs.

263

264 Decreasing trend in BZD use found in this study suggests that prescribing patterns of BZDs have
265 changed during recent years. Similar decreasing trend of BZD utilization has also been observed in
266 the entire Finnish population (Saastamoinen et al. 2016). National guidelines on rational BZD
267 prescribing recommend to restricting the length of BZD treatment to some weeks (National
268 Supervisory Authority for Health and Welfare 2015, Working Group set up by the Finnish Medical
269 Society Duodecim and the Finnish Sleep Research Society 2017). However, we found that nearly
270 one out of five young adult BZD users could be defined as a long-term user. Decreasing trend of long-
271 term BZD use among young adults raises questions about the factors associated with decreased use.
272 It is probable that a part of BZD use has been replaced by other pharmaceutical use or psychosocial
273 treatments. During recent years, mental health services among adolescents and young adults have
274 increased, which reflects the enhanced recognition of the treatment need, but may also suggest that
275 stigma associated with mental health service use has decreased (Pylkkänen and Laukkanen 2011).
276 Availability of mental health services may have improved, and low threshold online therapies are
277 available, too (Stenberg et al 2016).

278

279 Based on the findings of this study we suggest that patterns of BZD long-term use among young
280 people need more research. Long-term treatment with BZDs can be considered medically justified
281 for some patients. Therefore, it would be of importance to identify specific patient groups, which
282 would benefit from long-term BZD treatment. However, equally important would be to recognize
283 those patients who may experience harm of BZD use.

284

285 In conclusion, both overall use and long-term use of BZDs have decreased annually since the year
286 2008 among Finnish young adults. Decreasing use may be the consequence of increased awareness
287 of problems related to long-term BZD use and recommendations introduced to rationalize the use of

288 BZDs. Further research is needed to identify factors associated with the decline in BZD use among
289 young adults.

290

291

292 Clinical Significance

293 In this first nationwide study on BZD use and long-term use focusing on young adults, we found a
294 declining trend in use and long-term BZD use. Despite this decline, still a nearly one out of five
295 young adult BZD users could be defined as a long-term user and part of this long-term use may also
296 be inappropriate. Our findings urge physicians to pay more attention to BZD prescribing and
297 monitoring for young men who are especially prone to long-term use. In addition, special concern
298 regarding the prescription practices of high potency BZDs alprazolam and clonazepam is needed, as
299 these substances were most commonly used for long-term.

300

301 Conflicts of interest

302 Terhi Aalto-Setälä, Marja Airaksinen, Leena Saastamoinen, Annamari Tuulio-Henriksson, Sanna
303 Tähkäpää: Have no conflicts of interests

304 Terhi Kurko has owned a few stocks of pharmaceutical company Orion Pharma in 2014-2015 and
305 has received an expert fee from Pfizer in 2015.

306

307 References

308 Alessi-Severini S, Bolton JM, Enns MW, Dahl ME, Chateau D, Collins DM, Sareen J. Sustained Use of
309 Benzodiazepines and Escalation to High Doses in a Canadian Population. *Psychiatr Serv* 67:1012–1018,
310 2016.

311 Ashton H. Benzodiazepines: How they work and how to withdraw. 2013 Available at:
312 <http://www.benzo.org.uk/manual/bzcha01.htm>. Accessed Jun 14, 2017.

313 Autti-Rämö I, Seppänen J, Raitasalo R, Martikainen JE, Sourander A. [The use of psychotropics by
314 adolescents and young adults has increased during 2000's]. *Suomen lääkirilehti - Finlands läkartidning*
315 64:477–482, 2009.

316 Barker MJ, Greenwood KM, Jackson M, Crowe SF. Persistence of cognitive effects after withdrawal from
 317 long-term benzodiazepine use: a meta-analysis. *Archives of Clinical Neuropsychology* 4;19(3):437–454,
 318 2004.

319 Cloos J, Bocquet V, Rolland-Portal I, Koch P, Chouinard G. Hypnotics and Triazolobenzodiazepines - Best
 320 Predictors of High-Dose Benzodiazepine Use: Results from the Luxembourg National Health Insurance
 321 Registry. *Psychother Psychosom* 84:273–283, 2015.

322 Cunningham CM, Hanley GE, Morgan S. Patterns in the use of benzodiazepines in British Columbia:
 323 Examining the impact of increasing research and guideline cautions against long-term use. *Health Policy*
 324 97:122–129, 2010

325 Drug and Alcohol Services South Australia, Government of South Australia. Benzodiazepine conversion
 326 chart. Adelaide, Australia, 2012.

327 Finnish Medicines Agency Fimea. Drug consumption in years 2012-2015. Available at:
 328 <http://raportit.nam.fi/raportit/kulutus/laakekulutus.pdf>. Accessed Jun 14, 2017.

329 Furu K, Wettermark B, Andersen M, Martikainen JE, Almarsdottir AB, Sørensen HT. The Nordic Countries as
 330 a Cohort for Pharmacoepidemiological Research. *Basic Clin Pharmacol Toxicol* 106(2):86-94, 2010.

331 Glass J, Lanctôt KL, Herrmann N, Sproule BA, Busto UE. Sedative hypnotics in older people with insomnia:
 332 meta-analysis of risks and benefits. *BMJ* 331(7526), 2005.

333 Griffiths RR, McLeod DR, Bigelow GE, Liebson IA, Roache JD, Nowowieski P. Comparison of diazepam and
 334 oxazepam: preference, liking and extent of abuse. *J Pharmacol Exp Ther* 229:501–508, 1984.

335 Hartz I, Skurtveit S, Steffenak AKM, Karlstad Ø, Handal M. Psychotropic drug use among 0–17 year olds
 336 during 2004-2014: A nationwide prescription database study. *BMC Psychiatry* 16:12, 2016.

337 Haukka J, Suvisaari J, Tuulio-Henriksson A, Lönngqvist J. High concordance between self-reported medication
 338 and official prescription database information. *Eur J Clin Pharmacol* 63:1069–1074, 2007.

339 Hollingworth SA, Siskind DJ. Anxiolytic, hypnotic and sedative medication use in Australia.
 340 *Pharmacoepidemiol Drug Saf* 19:280–288, 2010.

341 Hsia Y, MacLennan K. Rise in psychotropic drug prescribing in children and adolescents during 1992–2001:
 342 A population-based study in the UK. *Eur J Epidemiol* 24:211–216, 2009

343 Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE..Lifetime prevalence and age-of-onset
 344 distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*
 345 62:593–602, 2005.

346 Kokkevi A, Fotiou A, Arapaki A, Richardson C. Prevalence, Patterns, and Correlates of Tranquilizer and
 347 Sedative Use among European Adolescents. *J Adolesc Health* 43:584–592, 2008.

348 Kornor H, Pedersen W, von Soest T, Rossow I, Bramness JG. [Use of benzodiazepines and cannabis in young
 349 adults]. *Tidsskr Nor Laegeforen* 130:928–931, 2010.

350 Kurko TAT, Saastamoinen LK, Tähkää S, Tuulio-Henriksson A, Taiminen T, Tiihonen J, Airaksinen MS,
 351 Hietala J. Long-term use of benzodiazepines: Definitions, prevalence and usage patterns - A systematic
 352 review of register-based studies. *Eur Psychiatry* 30:1037–1047, 2016.

353 Lagnaoui R, Depont F, Fourrier A, Abouelfath A, Bégaud B, Verdoux H, Moore N. Patterns and correlates of
 354 benzodiazepine use in the French general population. *Eur J Clin Pharmacol* 60:523–529, 2004.

355 McCabe SE. Correlates of nonmedical use of prescription benzodiazepine anxiolytics: Results from a
 356 national survey of U.S. college students. *Drug Alcohol Depend* 79:53–62, 2005.

357 National Supervisory Authority for Welfare and Health. Prescribing Benzodiazepines. 2015 Available at:
 358 [http://www.valvira.fi/terveydenhuolto/hyva-](http://www.valvira.fi/terveydenhuolto/hyva-ammatinharjoittaminen/laakehoito/bentsodiatsepiinien_maaraaminen_2)
 359 [ammatinharjoittaminen/laakehoito/bentsodiatsepiinien_maaraaminen_2](http://www.valvira.fi/terveydenhuolto/hyva-ammatinharjoittaminen/laakehoito/bentsodiatsepiinien_maaraaminen_2). Accessed Jun 14, 2017.

360 Nordfjærn T A population-based cohort study of anxiety, depression, sleep and alcohol outcomes among
 361 benzodiazepine and z-hypnotic users. *Addict Behav* 37:1151–1157, 2012.

362 Olfson M, King M, Schoenbaum M. Benzodiazepine use in the United States. *JAMA Psychiatry* 72:136–142,
 363 2015.

364 Patel V, Flisher AJ, Hetrick S P, McGorry P. Mental health of young people: a global public-health challenge.
 365 *Lancet*; 369:1302–1313, 2007.

366 Pylkkänen K, Laukkanen E (2011) [Adolescents do not fear the use of mental health services] *Suomen*
 367 *lääkärilehti - Finlands läkartidning* 66:2316–2317.

368 Riska BS, Skurtveit S, Furu K, Engeland A, Handal M. Dispensing of benzodiazepines and benzodiazepine-
 369 related drugs to pregnant women: A population-based cohort study. *Eur J Clin Pharmacol* 70:1367–1374,
 370 2014

371 Saastamoinen LK, Kalliokoski A, Martikainen J, Voipio T. Medicines in 2015. In: Finnish Statistics on
 372 Medicines 2015. Finnish Medicines Agency Fimea and The Social Insurance Institution, 2016. Available at:
 373 [https://www.fimea.fi/documents/160140/1188389/Suomen_l%C3%A4%C3%A4ketilasto_2015.pdf/a813feac-](https://www.fimea.fi/documents/160140/1188389/Suomen_l%C3%A4%C3%A4ketilasto_2015.pdf/a813feac-1560-4cbf-80e1-44049449e0bf)
 374 [c-1560-4cbf-80e1-44049449e0bf](https://www.fimea.fi/documents/160140/1188389/Suomen_l%C3%A4%C3%A4ketilasto_2015.pdf/a813feac-1560-4cbf-80e1-44049449e0bf) Accessed Jun 13, 2017.

375 Stenberg J, Sequeiros SB, Holm M, Kampman O, Kieseppä T, Korkeila J, Mäki P, Wahlbeck K, Joffe G, Häll P,
 376 Joutsenniemi K. [Mental health through ePsychiatry?] *Suomen lääkarilehti - Finlands läkartidning* 71:2106-
 377 2111a, 2016.

378 Suvisaari J, Aalto-Setälä T, Tuulio-Henriksson A, Härkänen T, Saarni SI, Perälä J, Schreck M, Castaneda A,
 379 Hintikka J, Kestilä L, Lähteenmäki S, Latvala A, Koskinen S, Marttunen M, Aro H, Lönngqvist J. Mental
 380 disorders in young adulthood. *Psychol Med* 39:287–299, 2009.

381 Tvette IF, Bjørner T, Skomedal T. A 5-year follow-up study of users of benzodiazepine: starting with
 382 diazepam versus oxazepam. *Br J Gen Pract* 66:e241–247, 2016.

383 White MP. Medication dosing in anxiety disorders: What the evidence shows. *Prim Psychiatry* 16:21–28,
 384 2009.

385 Wittchen HU, Nelson CB, Lachner G. Prevalence of mental disorders and psychosocial impairments in
386 adolescents and young adults. *Psychol Med* 28:109–126, 1998.

387 Working group set up by the Finnish Medical Society Duodecim and the Finnish Sleep Research Society.
388 Insomnia. Current Care Guideline, 2017. Available at:
389 <http://www.kaypahoito.fi/web/kh/suosituksset/suositus?id=hoi50067>. Accessed Oct 18, 2017.

390 World Health Organization Rational use of benzodiazepines. 1996

391 World Health Organization. ATC/DDD Index 2017. Available at: http://www.whocc.no/atc_ddd_index/.
392 Accessed Jun 14, 2017.

393 World Health Organization: DDD. Definition and general considerations. 2016 Available at:
394 http://www.whocc.no/ddd/definition_and_general_considera/. Accessed Jun 14, 2017.

395 Yeh HH, Chen CY, Fang SY, Chang IS, Wu EC, Lin KM. Five-year trajectories of long-term benzodiazepine use
396 by adolescents: patient, provider, and medication factors. *Psychiatr Serv* 62:900–907, 2011.