<https://helda.helsinki.fi>

Soil biota in boreal urban greenspace : Responses to plant type and age

Francini, G.

2018-03

Francini , G , Hui , N , Jumpponen , A , Kotze , D J , Romantschuk , M , Allen , J A & Setälä , H 2018 , ' Soil biota in boreal urban greenspace : Responses to plant type and age ' , Soil Biology & Biochemistry , vol. 118 , pp. 145-155 . <https://doi.org/10.1016/j.soilbio.2017.11.019>

<http://hdl.handle.net/10138/312553>

<https://doi.org/10.1016/j.soilbio.2017.11.019>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1Soil biota in boreal urban greenspace: responses 2to plant type and age

3

4G. Francini¹, A. Jumpponen², D.J. Kotze¹, N. Hui¹, M. Romantschuk¹, J.A.

5Allen¹, H. Setälä¹

6

71) Department of Environmental Sciences, University of Helsinki, Niemenkatu
873, 15140, Lahti, FINLAND

9

102) Division of Biology, Kansas State University, Manhattan, KS, USA

11

12**Corresponding author:** gaia.francini@gmail.com, nan.hui@helsinki.fi

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29**Key words:** Microbial biomass (PLFA), Urban greenspace, Nematodes,

30Earthworms

31

32Abstract

33

34Plant functional type influences the abundance and distribution of soil biota.
 35With time, as root systems develop, such effects become more apparent. The
 36relationship of plant type and time with the structure and abundance of soil
 37microbial and invertebrate communities has been widely investigated in a
 38variety of systems. However, much less is known about long-term soil
 39community dynamics within the context of urban environments. In this study,
 40we investigated how soil microbes, nematodes and earthworms respond to
 41different plant functional types (lawns only and lawns with deciduous or
 42evergreen trees) and park age in 41 urban parks in southern Finland. As non-
 43urban controls we included deciduous and evergreen trees in 5 forest sites. We
 44expected that microbial biomass and the relative abundance of fungi over
 45bacteria would increase with time. We also expected major differences in soil
 46microbial and nematode communities depending on vegetation: we
 47hypothesized that i) the presence of trees, and evergreens in particular, would
 48support a greater abundance of fungi and fungal-feeding nematodes over
 49bacteria and bacterial-feeding nematodes and ii) the fungi to bacteria ratio
 50would be lowest in lawns, with deciduous trees showing intermediate values.
 51In contrast to our predictions, we showed that old deciduous trees, rather than
 52evergreens, supported the highest fungal abundances and fungal-feeding
 53nematodes in the soil. Consistent with our predictions, microbial biomass in
 54urban park soils tended to increase with time, whereas – in contrast to our
 55hypotheses – fungal-feeding nematode abundance declined. Even in the oldest
 56parks included in the current study, microbial biomass estimates never
 57approximated those in the minimally managed natural forests, where biomass
 58estimates were three times higher. Anecic earthworm abundance also
 59increased with time in urban parks, whereas abundances of fungal-feeding,
 60plant-feeding and omnivorous nematodes, as well as those of epigeic and
 61endogeic earthworms remained constant with time and without any distinct
 62differences between urban parks and the control forests. Our findings highlight
 63that although urban park soils harbor diverse soil communities and

64considerable microbial biomass, they are distinct from adjacent natural sites in
65community composition and biomass.
66

67**Highlights**

68

- 69• In cities, soil communities under different plant types diverge
- 70 increasingly with time
- 71• Microbial biomass associated with deciduous trees increases with park
- 72 age
- 73• Earthworm biomass increases with park age
- 74• Adjacent non-urban sites maintain a greater microbial biomass than
- 75 urban parks

761. Introduction

77

78Urbanized areas are rapidly expanding at the expense of natural habitats.
 79Urban green spaces, including public parks, assume a pivotal role as surrogates
 80for these lost natural environments. These green spaces provide a vast array of
 81ecosystem services (Costanza et al., 1997; Bolund and Hunhammar, 1999)
 82including carbon and nitrogen sequestration (Raciti et al., 2011; Setälä et al.,
 832016), storm water interception and purification (Valtanen et al., 2015),
 84biodiversity and climate regulation (Bolund and Hunhammar, 1999). These
 85ecosystem services depend strictly on the soil, and in turn on the soil biota
 86hosted therein.

87 Soil microbes and invertebrates are directly linked to biogeochemical
 88processes that take place in the soil and thus promote a variety of soil-derived
 89ecosystem services (Lavelle et al. 2006; Balser and Firestone, 2005; Blouin et
 90al., 2013 Ledin, 2000; Haritash and Kaushik, 2009). Yet, the abundance and
 91distribution of soil biota are often directly linked with the distribution of plant
 92species, and indirectly to soil properties that are modified by the plants
 93(Wardle et al., 2004) over time (Bardgett et al., 2005). Although the
 94successional trajectories of soil communities in relation to plant type and soil
 95characteristics over time have been widely investigated in, e.g., primary
 96(Ohtonen et al., 1999; Doblas-Miranda et al., 2008; Brown and Jumpponen
 972014) and secondary succession (Pižl, 1992; Maharning et al., 2009), much less
 98is known about these dynamics in urban park soils.

99 The linkage between plant functional types and soil microbial and
 100invertebrate communities has received substantial interest in contemporary
 101soil- and ecosystem ecology (Orwin et al., 2010; Thomson et al., 2010). There is
 102also a growing body of literature describing soil biota in urban parks
 103(microbes: Baxter et al., 1999; Xu et al., 2014; Ramirez et al., 2014; Hui et al.
 1042017a, b; nematodes: Pavao-Zuckerman and Coleman 2007; Amossé et al.,
 1052016; earthworms: Steinberg et al., 1997; Smetak et al., 2007; Amossé et al.,
 1062016). In a nutshell, these studies collectively provide evidence that
 107urbanization can substantially change in soil microbial and faunal
 108communities. Yet, there is a paucity of studies that simultaneously account for

different trophic groups in large and well-replicated experimental designs within the context of urban ecosystems.

Plant species identity is linked to the quantity and quality of inputs provided to the soil, either via litter deposition or through root exudates, which in turn are largely responsible for the composition of soil microbial (Grayston et al., 1998; Bardgett and McAlister, 1999; Marschner et al., 2004), nematode (Ilieva-Makulec et al., 2006) and earthworm communities (Curry, 2004). Whilst plants that produce labile, nitrogen-rich litter, such as grasses, herbs and deciduous trees often support bacterial-dominated soil microflora, plants producing more recalcitrant litter, such as evergreens, more commonly support fungal-based soil food webs (Wardle et al. 2004). Evergreen trees are adapted to low nutrient availability and thus have leaves with low nutrient contents (Kattge et al. 2011). In contrast, deciduous broadleaf trees have higher foliar nutrient content (Kattge et al. 2011) with *Tilia cordata* and *Acer platanoides* particularly in particular producing high quality litter (Aerts and Chapin 2000, Hobbie et al. 2014). Moreover, soil acidification promoted by evergreen trees (Setälä et al. 2016) is also associated with the prevalence of fungi over bacteria (Bååth and Anderson 2003). Finally, the quantity of root exudates, in boreal systems, tends to be higher in evergreen trees than deciduous trees (Gower et.al 2001), suggesting that evergreen trees may allocate a higher percentage of net primary productivity to, i.e. ectomycorrhizal fungi than do deciduous trees. Such plant-associated distinctions in soil microbial communities are reflected in the relative proportions of bacterial- and fungal-feeding fauna (Trofymow and Coleman, 1982; de Vries et al., 2013). This is particularly interesting as fungal-based food webs are characterized by slow nutrient cycling and a high capacity to retain nutrients, whereas bacterial dominated food webs are characterized by high nutrient turnover and nutrient leaching (de Vries et al., 2006; de Vries et al., 2012). Consequently, the ratio between fungi and bacteria in the soil is crucial when considering fundamental ecological processes such as the rate of organic matter decomposition (Coleman et al., 1983; Wardle, 2002; Moore et al., 2005; Paterson et al., 2008) and thus carbon and nutrient sequestration.

141 The type and availability of plant-derived resources can also change
 142 during plant community succession (Berendse, 1990; Knops and Tilman,
 143 2000). As soil organic matter (OM), C and N accumulate over time, so too do the
 144 biomasses of soil microbes (Zak et al., 1990; Ohtonen et al., 1999), nematodes
 145 (Háněl, 2010) and earthworms (Pižl, 1992). Changes in microbial abundances
 146 are not only quantitative, but also qualitative. For instance, the relative
 147 abundance of fungi over bacteria tends to increase with time (Ohtonen et al.,
 148 1999; Zeller et al., 2001; Bardgett and Walker, 2004). Furthermore, temporal
 149 changes in the fungal to bacterial ratio during succession are often also
 150 mirrored by changes in the ratio of fungal feeding to bacterial feeding
 151 nematodes (Brzeski, 1995; Ferris and Matute, 2003; Háněl, 2010). Only a few
 152 studies have investigated such successional trajectories of soil biota in urban
 153 parks (Smetak et al., 2007; Amossé et al., 2016; Hui et al., 2017a, b).

154 This study is part of a larger project that aims to shed light on the
 155 influence of divergent plant types on the physico-chemical and biological soil
 156 characteristics in urban parks of diverging ages. Our previous work has shown
 157 that plant type and park age are strong determinants of soil characteristics
 158 (Setälä et al., 2016; Setälä et al. 2017) and soil microbial community
 159 composition (Hui et al., 2017a, b). However, the response of microbial biomass,
 160 a measure that strongly relates to the functional activity and capacity of soils,
 161 to plant type and park age remains unresolved. Moreover, in this study, we
 162 incorporate the microbial consumer responses – key actors in providing soil-
 163 based ecosystem services and contributing to nutrient turnover (Wardle et al.,
 164 2004). We aim to investigate how park age and plant functional type affect the
 165 biomass of soil microbes and two important functional groups of soil fauna:
 166 nematodes and earthworms.

167 Given the clear effects that plant functional type and park age have on
 168 soil properties (see Setälä et al. 2016; Setälä et al. 2017) we tested the
 169 following hypotheses: plant functional type affects the soil food web so that i)
 170 evergreen trees producing recalcitrant litter promote an increase of fungal
 171 biomass over bacterial biomass and ii) deciduous trees, the lawn in particular,
 172 producing more labile litter, promote the establishment of bacterial biomass.
 173 Then we test iii) if changes in the soil microbial community also cascade up to

higher trophic levels. We expect that higher densities of fungal feeding nematodes (compared to bacterial feeders) associate with evergreen trees and higher densities of earthworms associate with lawns and deciduous trees. Also, we test iv) whether time since park construction promotes changes in soil microbial and invertebrate community structure and abundance. We hypothesize that soil biota in old parks resemble natural communities more than in young parks. This is because the capacity of plants (especially trees) to modify soils is park-age dependent (Setälä et al. 2016, 2017) with young parks not having had the time to develop plant-soil interactions that are typical of natural forests.

184

185

1862. Materials and Methods

187

1882.1. Study area

189

The study was conducted in two municipalities in the boreal forest zone in southern Finland; Helsinki metropolitan area (60° 10' 15'' N 24° 56' 15'' E, with a population of ca. 1.4 million people) and Lahti (60° 58' 57'' N 25° 39' 41'' E, population ca. 110 000). Winters are cold and wet, while rainfall is moderate all year round. The annual mean temperature is 5.3°C in Helsinki and 4.5°C in Lahti; the annual mean precipitation is 628 mm in Helsinki and 636 mm in Lahti. Summer lasts for approximately 110-120 days, winter 135-145 days and the temperature can span from -35 to +35°C (Finnish Meteorological Institute). The Helsinki and Lahti regions are classified by NRCS (National Resource Conservation Service) as having soils of primarily the Spodosol suborder. However, all urban parks in the two cities are constructed and none showed detectable signs of podzolisation at the time of sampling.

We selected 41 parks in the two cities and five additional control forests (see Setälä et al., 2016 and Hui et al., 2017a for details). Parks of three ages were selected: young parks (between 7 and 15 years old), intermediate parks (ca. 50 years old) and old parks (> 100 years). Control forests, situated in the outskirts of the city of Lahti, are typified as unmanaged, conifer and linden

dominated forests (> 80 years of age). Park size varied from one to several hectares. The selected parks were subjected to routine maintenance, including mowing (mowing residues not removed) and raking of tree leaves in the fall. However, the parks were not irrigated or commonly fertilized. Until the early 1990s, some of the older parks in the city of Lahti were occasionally fertilized - commonly with saltpeter (N, P, K, S), while some of the park lawns in Helsinki have received and still receive light refurbishment fertilization.

In each park we selected, where possible, three vegetation-types: deciduous (represented by *Tilia x vulgaris* 93% and *Acer platanoides* 7%) and evergreen trees (spruce, *Picea sp.* 43.3%, *Abies sp.* 20%, *Pseudotsuga menziesii* 13.3%, *Pinus sylvestris* 13.3%, *Larix sp.* 10%) and a non-treed lawn with grass (including herbs such as *Trifolium pratense*, *Plantago major*). Lawn cover extended also under tree canopies. In some cases we selected parks including only deciduous trees and lawn or only evergreen trees and lawn. The control sites never had lawns, but deciduous (forest linden, *Tilia cordata*) and evergreen trees (Norway spruce, *Picea abies*) were always present at each site. In the parks, distance between the two tree types was always greater than the height of the nearest tree. Plant age was considered as coinciding with park age, except for young parks, where ca. 10 year old saplings were planted at the time of park construction. In order to have at least 10 replicates per park age and plant functional type, we selected 41 urban parks, resulting in 91 sampling locations as described by Setälä et al. (2016) plus five control forests with 10 additional sampling locations, totaling 101 sampling locations.

230

2312.2. Soil sampling and measurements for edaphic responses

232

In urban parks, soils were sampled at the edge of the canopy projection of each tree and in the middle of the lawn area (when present) in October 2014. In the control forests, soils were sampled under the canopy projection of each tree in May 2015. Samples were obtained using a metal corer that was sterilized in ethanol between samples. At each sampling point, 3 subsample soil cores were collected and then pooled. The soil from each sample was homogenized and larger stones, roots and fresh or recognizable plant material removed. Samples

for microbial analysis were stored in resalable plastic bags on ice in the field and frozen at -20 °C in the laboratory. Soil pH was measured in a 1/5 (vol/vol) soil/distilled water suspension. Soil was weighed and analyzed for percent moisture by drying for 48 hr at 105 °C. Total carbon (hereafter C) and nitrogen (N) were obtained by dry combustion at 1350 °C using a LECO CNS2000 Elemental Analyzer (0.07% C and 0.09% N detection limits) and reported as percentage dry mass.

247

2.3. Soil microbial analysis, PLFA

249

Before analysis, the frozen samples were thawed at room temperature and sieved through a 2 mm mesh to remove leaf litter, rocks, large particles and roots. Then, part of the soil was freeze-dried and an aliquot of 4 g for urban park soil and 2 g for forest soil was taken for phospholipid fatty acids (PLFA) analysis. Fatty acids (FA) were extracted following a modified procedure as described in Macnaughton et al. (1997), Bligh and Dyer (1959), Frostegård et al. (1991) and White et al. (1979). The extraction was performed using a Dionex ASE 350 machine (Thermo Scientific). Briefly, the samples were mixed with diatomaceous earth and placed in the extraction vessels between two cellulose filters prewashed in chloroform. FA were extracted with Bligh and Dyer (methanol/chloroform/citrate buffer, 2/1/0.8 vol/vol/vol) at 80 °C and the samples underwent two static cycles of 15 min each. Once FA acids were extracted, chloroform and citrate buffer were added, for a final volume of chloroform/methanol/citrate buffer of 1/1/0.9, vol/vol/vol. The extracts were left overnight to separate the aqueous phase from the lipidic phase. The supernatant was discharged and the lower phase collected, which was then evaporated under a flux of N₂ at 40 °C. The resulting pellet was suspended in chloroform and applied to silica columns (Bond Elut LRC, Agilent). Chloroform, acetone and methanol were applied in sequence to the column to separate neutral lipids, glycolipids and phospholipids. The extracts were then dried under N₂. The phospholipid fraction so obtained was methylated and the fatty acid methyl esters extracted with hexane/chloroform (4/1, vol/vol), dried under N₂ and redissolved in 1 ml hexane. We used 19:0 (methyl

nonadecanoate) as an internal standard. All the solvents used were of GC grade and glassware was baked in an oven at 450 °C for 4 h prior to use.

The samples were analyzed in a Shimadzu GCMSQP2010 Ultra with an Agilent J&WDB23 column. The column oven temperature was set to 60 °C and the injection temperature was 250 °C in splitless mode. MS ion source temperature was 200 °C and the interface temperature was 250 °C. As reference library we used the bacterial acid methyl ester (BAME) mix (SigmaAldrich), while the Supelco 37 Component FAME mix (SigmaAldrich) was used for calibration for PLFA biomass calculation. Peaks were cross-checked against the NIST spectral library. A total of 29 PLFAs were identified. Of the identified peaks, the following were considered to be of bacterial origin: i15:0, 15:0, 17:0, 17:0cy (eubacteria); a15:0, i16:0, i17:0 (gram-positive bacteria); 16:1 ω 9, 18:1 ω 7 (gram-negative bacteria). The PLFA 18:2 ω 6 was considered to be mainly of fungal origin (Frostegård and Bååth, 1996). PLFAs were expressed as $\mu\text{g g}^{-1}$ soil dry weight. The totality of identified peaks was used to investigate coarse microbial community changes, as explained in section 2.5 below. The microbial to fungal (F/B) ratio was calculated as the ratio of PLFA 18:2 ω 6 to bacterial PLFAs.

291

292 2.4. Soil Fauna

293

294 2.4.1. Nematodes

295

Nematodes were collected in early June 2016 in Lahti, and in July-August 2016 in Helsinki. Similarly to the soil sampling, nematodes were collected at the edge of the canopy projection of each tree and in the middle of the lawn (when present) area. In the control forests, soils were sampled under the canopy projection of each tree type. Each sample consisted of six pooled 5 cm diameter soil cores of the uppermost 10 cm. The samples were pooled to make one composite sample and immediately placed in plastic bags and stored in a cooler. In the laboratory, the soil was gently mixed and 10 g (fresh weight) of soil was used for nematode extraction. Nematodes were extracted for 48 h without lights/heating following the wet funnel method described in Sohlenius

306(1979). Nematode feeding guilds were assigned following Yeates et al. (1993).
 307We calculated the nematode channel ratio (NCR) as described in Yeates (2003)
 308and the maturity index (MI) as in Bongers (1990). NCR is calculated as the ratio
 309of bacterial feeding (BF) to bacterial-feeding plus fungal-feeding (FF)
 310nematodes: $BF/(BF+FF)$, and is constrained to have values between 1 (totally
 311bacterial-mediated) and 0 (totally fungal-mediated) (Yeates, 2003). MI is the
 312sum of the relative abundance of each taxon multiplied by their colonizer-
 313persister (c.p.) value. The c.p. value of each family was assigned following
 314Bongers (1990, 1999). MI provides information on the degree of
 315environmental disturbance, with higher values indicating a less disturbed
 316environment.

317

3182.4.2. Earthworms

319

320Earthworms were collected in August 2016 close to where soil microbes and
 321nematodes were sampled. We followed a modified version of the hot mustard
 322liquid method described by Gunn (1992). Each sample consisted of two (in
 323public parks) or four (in control sites) 25 × 25 cm frames irrigated with a hot
 324mustard liquid. The hot mustard slurry was prepared by mixing 15 g of hot
 325mustard powder (Colman's powder) in 100 ml of water, and then allowing it to
 326sit for at least 4 h. Immediately prior to sampling, the mustard slurry was
 327added to approximately 3.5 L of tap water in a watering can. This 3.5 L of
 328mustard water was applied to each frame over a span of 15–20 min. The
 329collected worms were immediately placed in tap water for some minutes, then
 330placed into plastic bags containing moist tissue paper, and stored in a cooler. In
 331the laboratory the earthworms were placed in a refrigerator and stored for 48
 332h to empty their guts. The worms were then identified, weighed and stored in
 3334% formaldehyde solution.

334

3352.5. Data analysis

336

337Generalized linear mixed models (GLMM) were used to investigate the
 338relationship between our focal response variables (see below) and park age (a

factor with three levels; young, intermediate, old), plant functional type (a factor with three levels; evergreen, deciduous, lawn), soil pH, % soil C, % soil N, C/N ratio, soil OM, and % soil moisture. Our response variables were: total microbial biomass (PLFA), PLFA fungal/bacterial ratio, PLFA fungal marker (18:2 ω 6), PLFA gram-/gram+ ratio; nematode NCR ratio, BF nematode, FF nematode, PF (plant-feeding) nematode, PO (predators and omnivores) nematode, anecic earthworm *Lumbricus terrestris* abundance and endogeic and epigeic earthworms (*Lumbricus rubellus*, epigeic; *Allolopophora caliginosa*, endogeic; *Octolasion cyaneum*, endogeic - collected in one sample only; *Allolopophora chlorotica*, endogeic - collected in one sample only; *Dendrobaena octaedra*, epigeic). These GLMMs did not include the control forest samples in Lahti. Response variables were log or square-root transformed, when necessary, to satisfy assumptions of normality. In the GLMM analysis, city was considered a random effect, with park identity nested within city. We performed a stepwise model selection procedure by removing insignificant predictor variables, one at a time, if their p-values were greater than 0.05 and if the AIC subsequently decreased. However, park age and vegetation functional type were always retained in the final model, irrespective of significance. Then, we compared the control forest with old parks in Lahti, using ANOVA, with park age and vegetation functional type (and their interaction) as factors. Lawns were excluded from this analysis. Finally we evaluated whether FF nematodes, BF nematodes and NCR were correlated, respectively, with PLFA 18:2 ω 6, bacterial PLFA and the PLFA fungal/bacterial ratio, using Pearson correlation.

Microbial (based on the relative abundance of PLFA biomasses) and nematode community structures were assessed using nonmetric multidimensional scaling analysis (NMDS). The earthworm community was dominated by one species, resulting in low species richness and deemed unsuitable for NMDS. For the NMDS analysis we used Euclidean distances for PLFAs and Bray-Curtis distances for nematodes. The “envfit” function was used to assess the significance of the relationship between these communities and the environment (factors: city, park age, plant functional type; abiotic parameters: pH, % soil C, % soil N, C/N ratio, soil OM, % soil moisture). To test

our specific hypotheses, NMDS analyses were performed on three different datasets; i) we included all park samples to evaluate the effects of plant functional type and park age on microbial and nematode communities; ii) we included only trees belonging to intermediate and old parks to focus on differences mediated by tree type; iii) we included only old deciduous and evergreen trees in Lahti parks and control forests in order to specifically explore differences between urban parks and natural forest. R scripts and data are provided as supplementary material (Supplementary material, S1-S25).

All statistical analyses were performed in R version 3.2.1. (R Core Team, 2015), using the packages *vegan* (Oksanen et al. 2007), *lme4* (Bates et al. 2014), *nmle* (Pinheiro et al. 2007) and *car* (Fox et al. 2011).

383

384

3853. Results

386

3873.1. Microbial biomass and community composition

388

The GLMM models provided little evidence for systematic responses in PLFA-based total microbial, bacterial and fungal biomasses (18:206) to plant functional type or park age (Table 1, Fig. 1). Rather, these effects highlighted interactions between park age and plant type: PLFA biomasses (total, bacterial, fungal) were lowest under deciduous trees in young parks, yet highest under deciduous trees in old parks (Fig. 1 a, b). Total microbial and bacterial biomasses were mainly negatively correlated with the C/N-ratio (Table 1, Fig. 2a) and positively correlated with % soil C (Fig. 2b). The fungal marker responded positively to % soil N (Table 1, Fig. 2c).

398

In old Lahti parks, total microbial and bacterial biomass (Supplementary material, Table S26) of evergreen and deciduous tree soils was about a third lower than in control forests (total microbial PLFA, age: $F_{1,16} = 15.18$, $p < 0.01$; bacterial PLFA, age: $F_{1,16} = 76.74$, $p < 0.01$). Within forests and old parks, total microbial PLFA biomass did not differ under evergreen and deciduous trees, but bacterial biomass was higher under deciduous ($F_{1,16} = 5.13$, $p = 0.03$) than evergreen trees. Fungal biomass (Supplementary material, Table

405S26) showed an interaction between land use type (control forest vs. urban
 406park) and tree type ($F_{1,16} = 5.13$, $p = 0.03$): fungal biomass was similar among
 407deciduous trees in forest and parks, whereas under evergreens it was
 408substantially higher in forests than in parks.

409 The PLFA-based fungi/bacteria (F/B) ratios suggest that, regardless of
 410park age and plant type, all soils were bacteria dominated (Fig. 1c). However,
 411the relative proportion of fungi over bacteria was higher under deciduous trees
 412than evergreen trees and lawns, regardless of park age (Table 1, Fig. 1c). None
 413of the measured soil parameters (Supplementary material, Table S26)
 414correlated with the F/B-ratio, and the F/B ratio among old parks and forests
 415did not differ (Table 1).

416 The ratio between gram-/gram+ bacteria did not respond to park age or
 417to vegetation type, and did not differ between control forests and old Lahti
 418parks. However, the gram-/gram+ ratio correlated positively with soil pH
 419(Table 1, Fig. 2d).

420 Relative abundances of PLFAs were used in NMDS analyses to
 421investigate changes in the soil microbial communities in parks and control
 422forest. Although microbial biomasses responded, we observed distinct changes
 423in community composition only when the following datasets were analyzed; i)
 424all plant types and park ages and ii) control forest and old deciduous and
 425evergreen vegetation types in Lahti. When all plant types and park ages were
 426included in the NMDSs, plant functional type had no effect on the soil microbial
 427community composition in urban parks (Fig. 3, panel 1). However, park age
 428influenced the composition of these communities ($R^2 = 0.06$, $p = 0.02$) (Fig. 3,
 429panel 2). The soil microbial communities also differed ($R^2 = 0.11$, $p = 0.01$)
 430between the two cities. Environmental variables – primarily % soil C ($R^2 = 0.14$,
 431 $p < 0.01$), % soil N ($R^2 = 0.12$, $p < 0.01$), the C/N-ratio ($R^2 = 0.08$, $p = 0.02$) and
 432soil OM ($R^2 = 0.07$, $p = 0.02$) – correlated with the soil microbial community
 433composition in the parks.

434 Comparing natural forest to deciduous and evergreen trees in old parks
 435in Lahti showed that plant functional type clearly influenced ($R^2 = 0.14$, $p =$
 4360.04) the soil microbial communities (Fig. 3, panel 5). However, the soil

microbial community did not differ among old parks and control forests (Fig. 3, panel 6).

439

3.2. Nematodes

441

Bacterial and fungal feeding nematodes were more abundant under evergreen trees than under lawns (Table 1). However, while the abundance of bacterial feeders declined with park age, fungal feeders did not differ across park age classes. The abundance of bacterial feeders declined with park age and positively correlated with soil % soil C (Table 1). Fungal feeders negatively correlated with soil OM (Table 1). Plant feeders and omnivorous nematodes, although not responding to plant type or park age, positively correlated with soil OM and the C/N ratio (Table 1). When forests were compared with old parks in Lahti, bacterial feeding nematodes were more abundant ($F_{1,16} = 7.06$, $p = 0.01$) under evergreen trees than under deciduous trees in old parks and control forests, while the other groups did not differ (see supplementary material, Table S26).

Plant type affected the nematode channel ratio (NCR) ($F = 4.89$, $p = 0.01$). However, lawn and evergreen or deciduous and evergreen did not differ (Table 1, Fig. 4a), but rather, the differences can be attributed to those between deciduous trees and lawn. NCR negatively correlated with soil OM in the soil (Table 1). The NCR did not differ between old parks and control forest in Lahti. The maturity index (MI) showed a significant interaction between plant type and park age: when considering only deciduous trees, MI had the lowest values in young and old parks and highest values in intermediate parks (Table 1, Fig. 4b). MI also positively correlated with the soil C/N-ratio (Table 1). MI was higher ($F_{1,16} = 7.92$, $p = 0.01$) in old Lahti parks than forest controls (see supplementary material, Table S26).

We observed no relationships (Pearson correlations) between (i) the fungal marker and fungal feeding nematodes, (ii) bacterial biomass and bacterial feeding nematodes, and (iii) the PLFA fungal/bacterial-ratio and NCR.

The NMDSs revealed that nematode communities depended neither on plant functional type nor park age (Fig. 3, panels 7, 8). Park age did, however,

have an effect when deciduous and evergreen trees in old and intermediate parks were compared (Fig. 3, panel 10) ($R^2 = 0.63$, $p < 0.01$). Forests had distinct ($R^2 = 0.48$, $p = 0.01$) nematode communities compared to old Lahti parks (Fig. 3, panel 12). When all park samples were included in the analysis, % soil C was an important determinant of nematode community structure ($R^2 = 0.07$, $p = 0.04$). However, when only evergreen and deciduous trees in intermediate and old urban parks were investigated, only soil moisture correlated with nematode community structure ($R^2 = 0.15$, $p = 0.04$). Soil characteristics seemed to strongly control soil nematode communities (Fig. 1 panels 11, 12) when forests were compared to old parks, with only deciduous and evergreen trees included in the analysis (moisture: $R^2 = 0.61$, $p < 0.01$; pH: $R^2 = 0.56$, $p < 0.01$; OM: $R^2 = 0.39$, $p = 0.01$; % soil C: $R^2 = 0.39$, $p = 0.01$; % soil N: $R^2 = 0.33$, $p = 0.02$; C/N-ratio: $R^2 = 0.57$, $p < 0.01$). Nematode community compositions differed among the two cities when the comparison was made between all parks ($R^2 = 0.13$, $p < 0.01$) and when deciduous and evergreen trees in old and intermediate parks in Helsinki and Lahti were included in the analysis ($R^2 = 0.19$, $p = 0.04$).

487

488 3.3. Earthworms

489

GLMM results of the anecic earthworm *Lumbricus terrestris* showed that its biomass positively correlated with park age (Table 1), with the highest biomasses in intermediate and old parks (Supplementary material, Table S2). Plant type was not associated with *L. terrestris* biomass. *L. terrestris* correlated negatively with soil OM content (Table 1). The *L. terrestris* biomass did not differ between old parks and control forests in Lahti.

The epigeic and endogeic earthworm biomasses responded to plant type (Table 1), with highest abundances in lawns (Supplementary material, Table S2). Here, soil pH negatively correlated with the biomass of earthworms (Table 1). The biomass of these earthworm taxa did not differ between old parks and control forests, nor between evergreen and deciduous trees in the two habitat types.

502

503

5044. **Discussion**

505

5064.1. Effects of plant functional type on the soil biota

507

508We hypothesized that different plant functional types – due to their
 509documented divergent effects on soil characteristics and plant derived
 510resources – will lead to changes in soil communities and the abundances of
 511species and functional groups. Although overall communities were
 512compositionally largely invariable, the biomasses and relative abundances of
 513different microbial and nematode functional groups did respond to plant
 514functional type.

515 Importantly, the PLFA-based F/B ratio was lowest in lawns, suggesting
 516that bacteria dominated in lawn soils. Yet, to our surprise, soils under
 517deciduous trees, and not evergreen trees, had the highest F/B and lowest NCR.
 518NCR represents the relative abundance of bacterial feeders over fungal feeders,
 519and thus mirrors the relative amount of bacteria over fungi in the soil (Yeates
 5202003). In general, the relative amount of fungi over bacteria tends to be higher
 521under deciduous and especially under evergreen trees than lawn. Thus, our
 522data contrast those reported (e.g. Wardle et al., 2004). The reasons for this
 523remain unclear, and given our data, we can only speculate. For instance, a
 524higher amount of fine roots (Giardina et al. 2005) and decaying roots support a
 525higher saprophytic fungal biomass (Hobbie 2006) in soil associated with
 526deciduous trees than evergreen, which could explain why soils associated with
 527old lindens showed such a high F/B ratio. Another important factor potentially
 528explaining the similar F/B ratio under lawns and evergreen trees may be the
 529continuous disturbance present in urban parks; in particular, soil compaction
 530can lower the F/B ratio (Hedlund et al., 2003).

531 Our third hypothesis focused on plant type mediated changes in higher
 532trophic groups, such as nematodes and earthworm. GLMM analyses focusing on
 533microbial and nematode feeding guilds revealed clear responses to plant type,
 534although in some cases the response was seemingly mediated largely through
 535age and soil properties of the parks. Total microbial biomass, as well as

bacterial and fungal biomasses separately were highest under deciduous trees in old parks, whereas no such effect was detectable in younger parks. Bacterial feeding nematodes and fungal feeding nematodes were more abundant under evergreen and deciduous trees than under lawn. Nematode communities often vary according to vegetation, as a direct result of plant-provided resources and indirectly through changes in the quality and quantity of the microbial fauna controlled by plant litter and exudate production. The lower abundance of nematodes in lawns may be due to diminished predation/consumption by earthworms (Dash 1980), which were less abundant under trees than lawns. Moreover, Yates (1981) reported complementary dynamics between earthworms and bacterial feeding nematodes, suggesting that nematodes and earthworms can compete for resources, with both using microbial biomass as food. In line with other studies, total microbial and bacterial biomasses in this study were also positively correlated with % soil C. Soil % C was also correlated with nematode abundance (both FF and BF) and community structure. Taken together, this is a clear indication that, just as in non-urban soils, the availability of carbon is a key determinant of microbial and secondary consumer biomass in urban soils, but that the effects appear only over time.

Contrary to previous findings (Lauber et al. 2009; Hui et al. 2017a), the soil microbial (PLFA) community did not correlate with pH. In contrast, the gram-/gram+ ratio did. Even though these two groups are not strict functional groups, they responded differently to soil pH; and in agreement with previous studies, gram- bacteria were relatively more abundant at high pH (Wang et al. 2016). Further, our results also corroborate Hui et al. (2017a), who showed that microbial communities (characterized with high throughput sequencing methods) in these urban parks were distinguished by plant functional types and correlated with % soil N and pH.

563

4.2. Effects of park age on the soil biota

565

Our fourth hypothesis stated that time since the establishment of a park (park age as well as tree age) will drive changes in the soil microbial and invertebrate communities. Our previous study, conducted in the same parks, but using high

throughput sequencing, showed that microbial communities differed compositionally between young parks and old / intermediate parks (Hui et al., 2017a). Consistent with those results, our NMDS analyses of microbial communities separated old and intermediate parks from young parks. This suggests that the soil microbial community reaches a stable structure between 10 and 50 years after park construction. Soil microbial community composition in old parks overlapped with that in control forests. However, microbial biomass was substantially higher in forests than in old parks. This is likely due to frequent disturbances in the latter system (Pickett and Cadenasso 2009). Malmivaara-Lämsä and Fritze (2003) reported that human soil trampling in an urban boreal forest did not affect microbial community composition. Accordingly, we also observed that microbial community composition of old parks approximates the non-disturbed forest sites, albeit microbial abundance is diminished in urban parks compared to forest sites. However, for logistical reasons, we sampled the urban parks and control sites (forest) at different times. Microbial biomass can change across seasons, with peaks in the spring and autumn (Diaz-Ravina et al. 1995). Nevertheless, the control sites also had higher soil OM and %C (Setälä et al. 2016) suggesting that the observed microbial biomass differences are unlikely mere artifacts attributable to different sampling times. In addition to biomass, microbial community composition can be seasonally dynamic (Moore-Kucera and Dick 2008), but despite such potential seasonal dynamics, the communities in control and urban park sites overlapped.”.

Contrary to our expectations, the relative amount of fungi over bacteria did not increase as a function of park age. Similarly, NCR, describing the relative abundance of bacterial-feeding over fungal-feeding nematodes, remained mostly invariant across all park age classes. In successional systems, such as agricultural systems under restoration, the F/B ratio increases (Zeller 2001, Bailey 2002) as a result of changes in soil parameters (Zeller et al. 2001) and reduced disturbance (Hedlund et al. 2003). However, in urban parks the normal successional course is arrested and disturbance is continuous, possibly preventing increases in the relative abundance of fungi. Disturbance may also be the reason behind the invariant MI (Nematode Maturity Index) among the

602 parks of different ages. Interestingly, MI, which is regarded as a good indicator
 603 of ecosystem maturity, was higher in old parks compared with forest. In a
 604 study on the response of a riparian nematode community along a rural-urban
 605 transect, Pavao-Zuckerman et al. (2007) reported no differences in MI among
 606 sites, suggesting that the level of pollution was not high enough to influence MI.
 607 Also in our case, the levels of metal pollution were below ecologically relevant
 608 values (Setälä et al. 2017), but it is still unclear why MI values in forests are
 609 lower than in parks.

610 We were also unable to detect any shifts in the abundances of different
 611 nematode functional groups. Similarly to our speculations above, here too
 612 factors such as disturbance and litter removal may greatly contribute in
 613 altering resource availability to the soil biota, thus homogenizing these taxa
 614 across park ages. This could also explain the lack of response to plant traits and
 615 park age in plant-feeding nematodes, although plant root volumes likely
 616 increased substantially with time since park construction and establishment.

617 Nematode communities did respond to park age, but only in the
 618 rhizospheres of the two tree types; yet no changes in abundance with park age
 619 were observed. Nematodes can respond to changes in biotic and abiotic soil
 620 conditions (De Goede and Bongers 1994, Hanel 1995). Therefore, it is not
 621 surprising that we observed an age related shift in the nematode community
 622 only in association with trees, i.e. plant types which with time resulted in the
 623 clearest soil property changes in our study parks (see Setälä et al., 2016), and
 624 not in lawns.

625 The nematode community was also extremely responsive when we
 626 compared old parks and control forest, responding in parallel with the
 627 microbial communities as reported in Hui et al. (2017a). The nematode
 628 community structure was also correlated with changes in the soil
 629 characteristics, which in turn were modified by the different plant functional
 630 type and time. Community composition correlated with changes in abiotic soil
 631 parameters, such as pH, C, N and soil moisture. Again, this shows that
 632 nematodes are sensitive indicators (Neher 2001) of temporally mediated
 633 changes in soil characteristics. Contrary to previous findings, nematode
 634 densities were similar in rural forests and in urban parks. Pouyat et al. (1994)

suggested that a decline in nematode populations in urban soils might be linked to higher heavy metal concentrations in urban soils. On the other hand, Ohtonen (1992) found no link between metal pollution and nematode density. It is possible that pollution, and in particular metal loads, in urban soils in Helsinki and Lahti were not extreme enough to affect the soil biota.

Abundance of the anecic earthworm *L. terrestris* was lowest in young parks, increased with park age and reached a plateau in intermediate parks. The low abundance of earthworms in young parks can be attributed to two factors: i) lack of time for earthworms to colonize newly constructed habitats, and/or ii) the inhospitality of the newly built park soil as a habitat. As resources such as C and N increase in our park soils (see Setälä et al., 2016), so too does earthworm biomass. Similar trends in urban parks were observed by Smetak et al. (2007). This can have important implications since earthworms can ameliorate negative soil characteristics, compensating for and even reducing soil compaction with time. It is interesting to note that epigeic and endogeic earthworms did respond to pH variation, being more abundant at low pH than at high pH. Earthworms are quite sensitive to pH (Edwards and Bohlen, 1996). However, variation in pH among parks and plant types was rather narrow (Setälä et al., 2016) and among the earthworm species that we recorded, *Lumbricus rubellus* and *Aporrectodea caliginosa* were more abundant. Considering that pH is highly correlated with park age (Setälä et al., 2016), the significant relationship between earthworm abundance and pH could also be interpreted as the result of a relationship between park age and earthworm abundance.

Conclusions

Our data show that microbial biomass in natural forests is much higher than in urban parks, irrespective of their age. Thus, the urban greenspaces investigated unlikely approximate ecosystem properties or functions of the surrounding non-urban areas. Yet, our study provides strong evidence that deciduous trees support a greater microbial biomass with time since urban

green space establishment. Deciduous trees also had the highest relative amount of fungi over bacteria in urban parks, that evergreen trees would promote fungal rich soil microbial communities. Not only did microbial biomass increase with age in urban parks, but so too did the abundance of secondary consumers, such as earthworms. These data suggest that although urban parks do not approximate natural forests, likely as a consequence of the maintenance/disturbance regime they experience, the soil microbial and invertebrate communities respond to vegetation and edaphic shifts over time. Nevertheless, further studies are needed to assess the correlation and type of ecosystem services provided by these trophic groups.

678

679

680 Acknowledgments

681

We are grateful to Tuukka Ryyänen, for help with the field sampling. Thanks go to Anna-Lea Rantalainen for guidance with the GC-MS, and to Riikka-Juulia Koivula and Santeri Savolainen who helped solving all sorts of technical issues. Satu Tegel, Tuuli Ylikotila and Pekka Engblom provided data for the parks in Helsinki. Markku Saari provided information about parks in Lahti. The research was funded by an Academy of Finland grant (Grant number 268548).

688

689

690 References

691

Aerts R., Chapin III F.S., 2000. The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns-iii. Nutrient-limited plant growth: which nutrient is limiting?-b. How to detect N-or P-limited plant. *Advances in Ecological Research*, 30, 7-8.

696

Amossé, J., Dózsa-Farkas, K., Boros, G., Rochat, G., Sandoz, G., Fournier, B., Mitchell, E.A., Le Bayon, R., 2016. Patterns of earthworm, enchytraeid and nematode diversity and community structure in urban soils of different ages. *European Journal of Soil Biology*, 73, 46-58.

701

Bailey, V.L., Smith, J.L., Bolton, H., 2002. Fungal-to-bacterial ratios in soils investigated for enhanced C sequestration. *Soil Biology and Biochemistry*, 34, 997-1007.

705

- 706Balser, T.C., Firestone, M.K., 2005. Linking microbial community composition and
707soil processes in a california annual grassland and mixed-conifer forest.
708Biogeochemistry, 73, 395-415.
709
- 710Bardgett, R.D., Bowman, W.D., Kaufmann, R., Schmidt, S.K., 2005. A temporal
711approach to linking aboveground and belowground ecology. Trends in Ecology &
712Evolution, 20, 634-641.
713
- 714Bardgett, R.D., McAlister, E., 1999. The measurement of soil fungal: Bacterial
715biomass ratios as an indicator of ecosystem self-regulation in temperate meadow
716grasslands. Biology and Fertility of Soils, 29, 282-290.
717
- 718Bardgett, R.D., Walker, L.R., 2004. Impact of coloniser plant species on the
719development of decomposer microbial communities following deglaciation. Soil
720Biology and Biochemistry, 36, 555-559.
721
- 722Bates, D., Mächler, M., Bolker, B., Walker, S., 2014. Fitting linear mixed-effects
723models using lme4. arXiv preprint arXiv:1406.5823.
724
- 725Baxter, J.W., Pickett, S.T., Carreiro, M.M., Dighton, J., 1999. Ectomycorrhizal
726diversity and community structure in oak forest stands exposed to contrasting
727anthropogenic impacts. Canadian Journal of Botany, 77, 771-782.
728
- 729Berendse, F., 1990. Organic matter accumulation and nitrogen mineralization during
730secondary succession in heathland ecosystems. The Journal of Ecology, 78, 413-
731427.
732
- 733Bligh, E., Dyer, W., 1959. A rapid method of total lipid extraction and purification.
734Canadian Journal of Biochemistry and Physiology, 37, 911-917.
735
- 736Blouin, M., Hodson, M.E., Delgado, E.A., Baker, G., Brussaard, L., Butt, K.R., Dai,
737J., Dendooven, L., Pérès, G., Tondoh, J., 2013. A review of earthworm impact on
738soil function and ecosystem services. European Journal of Soil Science, 64, 161-
739182.
740
- 741Bolund, P., Hunhammar, S., 1999. Ecosystem services in urban areas. Ecological
742Economics, 29, 293-301.
743
- 744Bongers, T., 1990. The maturity index: An ecological measure of environmental
745disturbance based on nematode species composition. Oecologia, 83, 14-19.
746
- 747Bongers, T., Ferris, H., 1999. Nematode community structure as a bioindicator in
748environmental monitoring. Trends in Ecology & Evolution, 14, 224-228.
749
- 750Brown, S. P., Jumpponen, A., 2014. Contrasting primary successional trajectories of
751fungi and bacteria in retreating glacier soils. Molecular Ecology, 23, 481-497.
752
- 753Brzeski, M.W., 1995. Changes of the nematode fauna in the successive age classes
754of a scots pine forest. Fragmenta Faunistica, 15, 339-345.
755

- 756Bååth, E., Anderson, T. H., 2003. Comparison of soil fungal/bacterial ratios in a pH
757gradient using physiological and PLFA-based techniques. *Soil Biology and*
758*Biochemistry*, 35, 955-963.
- 759
- 760Coleman, D.C., Reid, C., Cole, C., 1983. Biological strategies of nutrient cycling in
761soil systems. *Advances in Ecological Research*, 13, 1-55.
- 762
- 763Costanza, R., d'Arge, R., de Groot, R., Farberk, S., Grasso, M., Hannon, B.,
764Limburg, K., Naeem, S., O'Neill, R.V., Paruelo, J., 1997. The value of the world's
765ecosystem services and natural capital. *Nature*, 387, 253-260
- 766
- 767Curry, J.P., 2004. Factors affecting the abundance of earthworms in soils. In:
768*Earthworm Ecology*, Edwards C. A. (Ed.), CRC press, pp. 91-113.
- 769
- 770Dash, M., Senapati, B., Mishra, C., 1980. Nematode feeding by tropical
771earthworms. *Oikos* 34, 322-325.
- 772
- 773De Goede, R.G.M., Bongers, T., 1994. Nematode community structure in relation to
774soil and vegetation characteristics. *Applied Soil Ecology* 1, 29-44.
- 775
- 776de Vries, F.T., Bloem, J., Quirk, H., Stevens, C.J., Bol, R., Bardgett, R.D., 2012.
777Extensive management promotes plant and microbial nitrogen retention in
778temperate grassland. *PLoS One* 7, e51201.
- 779
- 780de Vries, F.T., Hoffland, E., van Eekeren, N., Brussaard, L., Bloem, J., 2006.
781Fungal/bacterial ratios in grasslands with contrasting nitrogen management. *Soil*
782*Biology and Biochemistry*, 38, 2092-2103.
- 783
- 784de Vries, F.T., Thebault, E., Liiri, M., Birkhofer, K., Tsiafouli, M.A., Bjornlund, L.,
785Bracht Jorgensen, H., Brady, M.V., Christensen, S., de Ruiter, P.C., d'Hertefeldt, T.,
786Frouz, J., Hedlund, K., Hemerik, L., Hol, W.H., Hotes, S., Mortimer, S.R., Setälä,
787H., Sgardelis, S.P., Uteseny, K., van der Putten, W.H., Wolters, V., Bardgett, R.D.,
7882013. Soil food web properties explain ecosystem services across European land use
789systems. *Proceedings of the National Academy of Sciences of the United States of*
790*America*, 110, 14296-14301.
- 791
- 792Diaz-Ravina, M., Acea, M. J., Carballas, T., 1995. Seasonal changes in microbial
793biomass and nutrient flush in forest soils. *Biology and Fertility of Soils*, 19, 220-
794226.
- 795
- 796Doblas-Miranda, E., Wardle, D.A., Peltzer, D.A., Yeates, G.W., 2008. Changes in
797the community structure and diversity of soil invertebrates across the Franz Josef
798Glacier chronosequence. *Soil Biology and Biochemistry*, 40, 1069-1081.
- 799
- 800Edwards, C.A., Bohlen, P.J., 1996. *Biology and Ecology of Earthworms*. Springer
801Science & Business Media.
- 802
- 803Ferris, H., Matute, M., 2003. Structural and functional succession in the nematode
804fauna of a soil food web. *Applied Soil Ecology*, 23, 93-110.
- 805

- 806 Fox, J., 2011. Weisberg S. 2011. An r companion to applied regression. R package,
807 Sage.
- 808
- 809 Frostegård, Å, Bååth, E., 1996. The use of phospholipid fatty acid analysis to
810 estimate bacterial and fungal biomass in soil. *Biology and Fertility of Soils*, 22, 59-
811 65.
- 812
- 813 Frostegård, Å, Tunlid, A., Bååth, E., 1991. Microbial biomass measured as total
814 lipid phosphate in soils of different organic content. *Journal of Microbiological*
815 *Methods*, 14, 151-163.
- 816
- 817 Giardina, C.P., Coleman, M.D., Hancock, J.E., King, J.S., Lilleskov, E.A., Loya,
818 W.M., Pregitzer, K.S., Ryan, M.G., Trettin, C.C., 2005. The Response of
819 Belowground Carbon Allocation in Forests to Global Change. In: Anonymous Tree
820 Species Effects on Soils: Implications for Global Change. Springer, pp. 119-154.
- 821
- 822 Gower, S. T., Krankina, O., Olson, R. J., Apps, M., Linder, S., Wang, C. 2001. Net
823 primary production and carbon allocation patterns of boreal forest ecosystems.
824 *Ecological Applications*, 11, 1395-1411.
- 825
- 826 Grayston, S.J., Wang, S., Campbell, C.D., Edwards, A.C., 1998. Selective influence
827 of plant species on microbial diversity in the rhizosphere. *Soil Biology and*
828 *Biochemistry*, 30, 369-378.
- 829
- 830 Gunn, A., 1992. The use of mustard to estimate earthworm populations.
831 *Pedobiologia* 36, 65-67.
- 832
- 833 Háněl, L., 1995. Secondary successional stages of soil nematodes in cambisols of
834 South Bohemia. *Nematologica*, 41, 197-218.
- 835
- 836 Háněl, L., 2010. An outline of soil nematode succession on abandoned fields in
837 south bohemia. *Applied Soil Ecology* 46, 355-371.
- 838
- 839 Haritash, A., Kaushik, C., 2009. Biodegradation aspects of polycyclic aromatic
840 hydrocarbons (PAHs): A review. *Journal of Hazardous Materials*, 169, 1-15.
- 841
- 842 Hedlund, K., Santa Regina, I., Van der Putten, W., Lepš, J., Diaz, T., Korthals, G.,
843 Lavorel, S., Brown, V., Gormsen, D., Mortimer, S., 2003. Plant species diversity,
844 plant biomass and responses of the soil community on abandoned land across
845 Europe: Idiosyncrasy or above-belowground time lags. *Oikos*, 103, 45-58.
- 846
- 847 Hobbie, E.A., 2006. Carbon allocation to ectomycorrhizal fungi correlates with
848 belowground allocation in culture studies. *Ecology*, 87, 563-569.
- 849
- 850 Hobbie, S. E., Baker, L. A., Buyarski, C., Nidzgorski, D., & Finlay, J. C. 2014.
851 Decomposition of tree leaf litter on pavement: implications for urban water
852 quality. *Urban Ecosystems*, 17, 369-385.
- 853
- 854 Hui, N., Jumpponen, A., Francini, G., Kotze, D.J., Liu, X., Romantschuk, M.,
855 Strömmer, R., Setälä, H., 2017a. Soil microbial communities are shaped by

- 856vegetation type and park age in cities under cold climate. *Environmental*
857*Microbiology*, 19, 1281-1295.
- 858
- 859Hui, N., Liu, X., Kotze., D.J., Jumpponen, A., Francini, G., Setälä., H.,
860Ectomycorrhizal fungal communities in urban parks are similar to those in natural
861forests but shaped by vegetation and park age. 2017b *Applied and Environmental*
862*Microbiology*, AEM-01797.
- 863
- 864Ilieva-Makulec, K., Olejniczak, I., Szanser, M., 2006. Response of soil micro-and
865mesofauna to diversity and quality of plant litter. *European Journal of Soil Biology*,
86642, 244-249.
- 867
- 868Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Bönisch, G., Garnier,
869E., Westoby, M., Reich, P. B., Wright, I. J., Cornelissen, J. H. C., Violle, C. And
870Harrison, S. P., Van Bodegom, P. M., Reichstein, M., Enquist, B. J.,
871Soudzilovskaia, N. A., Ackerly, D. D., Anand, M. And Atkin, O., Bahn, M., Baker,
872T. R., Baldocchi, D., Bekker, R., Blanco, C. C., Blonder, B., Bond, W. J.,
873Bradstock, R., Bunker, D. E., Casanoves, F., Cavender-Bares, J., Chambers, J. Q.,
874Chapin Iii, F. S., Chave, J., Coomes, D., Cornwell, W. K., Craine, J. M., Dobrin, B.
875H., Duarte, L., Durka, W., Elser, J., Esser, G., Estiarte, M., Fagan, W. F., Fang, J.,
876Fernández-Méndez, F., Fidelis, A., Finegan, B., Flores, O., Ford, H., Frank, D.,
877Freschet, G. T., Fyllas, N. M., Gallagher, R. V., Green, W. A., Gutierrez, A. G.,
878Hickler, T., Higgins, S. I., Hodgson, J. G., Jalili, A., Jansen, S., Joly, C. A.,
879Kerkhoff, A. J., Kirkup, D., Kitajima, K., Kleyer, M., Klotz, S., Knops, J. M. H.,
880Kramer, K., Kühn, I., Kurokawa, H., Laughlin, D., Lee, T. D., Leishman, M., Lens,
881F., Lenz, T., Lewis, S. L., Lloyd, J., Llusià, J., Louault, F., Ma, S., Mahecha, M. D.,
882Manning, P., Massad, T., Medlyn, B. E., Messier, J., Moles, A. T., Müller, S. C.,
883Nadrowski, K., Naeem, S., Niinemets, Ü., Nöllert, S., Nüske, A., Ogaya, R.,
884Oleksyn, J., Onipchenko, V. G., Onoda, Y., Ordoñez, J., Overbeck, G., Ozinga, W.
885A., Patiño, S., Paula, S., Pausas, J. G., Peñuelas, J., Phillips, O. L., Pillar, V.,
886Poorter, H., Poorter, L., Poschlod, P., Prinzing, A., Proulx, R., Rammig, A.,
887Reinsch, S., Reu, B., Sack, L., Salgado-Negret, B., Sardans, J., Shiodera, S.,
888Shipley, B., Siefert, A., Sosinski, E., Soussana, J.-F., Swaine, E., Swenson, N.,
889Thompson, K., Thornton, P., Waldram, M., Weiher, E., White, M., White, S.,
890Wright, S. J., Yguel, B., Zaehle, S., Zanne, A. E., Wirth, C., 2011. TRY-a global
891database of plant traits. *Global Change Biology*, 17, 2905-2935.
- 892
- 893Knops, J.M., Tilman, D., 2000. Dynamics of soil nitrogen and carbon accumulation
894for 61 years after agricultural abandonment. *Ecology*, 81, 88-98.
- 895
- 896Lauber, C.L., Hamady, M., Knight, R., Fierer, N., 2009. Pyrosequencing-based
897assessment of soil pH as a predictor of soil bacterial community structure at the
898continental scale. *Applied and Environmental Microbiology*, 75, 5111-5120.
- 899
- 900Lavelle, P., Decaëns, T., Aubert, M., Barot, S., Blouin, M., Bureau, F., Margerie, P.,
901Mora, P., Rossi, J., 2006. Soil invertebrates and ecosystem services. *European*
902*Journal of Soil Biology*, 42, S3-S15.
- 903
- 904Ledin, M., 2000. Accumulation of metals by microorganisms-processes and
905importance for soil systems. *Earth-Science Reviews*, 51, 1-31.

- 906
 907Macnaughton, S.J., Jenkins, T.L., Wimpee, M.H., Cormiér, M.R., White, D.C.,
 9081997. Rapid extraction of lipid biomarkers from pure culture and environmental
 909samples using pressurized accelerated hot solvent extraction. *Journal of*
 910*Microbiological Methods*, 31, 19-27.
 911
 912Maharning, A.R., Mills, A.A., Adl, S.M., 2009. Soil community changes during
 913secondary succession to naturalized grasslands. *Applied Soil Ecology*, 41, 137-147.
 914
 915Malmivaara-Lämsä, M., Hamberg, L., Haapamäki, E., Liski, J., Kotze, D.J.,
 916Lehvävirta, S., Fritze, H., 2008. Edge effects and trampling in boreal urban forest
 917fragments—impacts on the soil microbial community. *Soil Biology and*
 918*Biochemistry*, 40, 1612-1621.
 919
 920Marschner, P., Crowley, D., Yang, C.H., 2004. Development of specific rhizosphere
 921bacterial communities in relation to plant species, nutrition and soil type. *Plant and*
 922*Soil*, 261, 199-208.
 923
 924Moore, J.C., McCann, K., de Ruiter, P.C., 2005. Modeling trophic pathways,
 925nutrient cycling, and dynamic stability in soils. *Pedobiologia*, 49, 499-510.
 926
 927Moore-Kucera, J., Dick, R.P., 2008. PLFA profiling of microbial community
 928structure and seasonal shifts in soils of a Douglas-fir chronosequence. *Microbial*
 929*Ecology*, 55, 500-511.
 930
 931Neher, Deborah A., 2001. Role of nematodes in soil health and their use as
 932indicators. *Journal of Nematology*, 33, 161-168.
 933
 934Ohtonen, R., Fritze, H., Pennanen, T., Jumpponen, A., Trappe, J., 1999. Ecosystem
 935properties and microbial community changes in primary succession on a glacier
 936forefront. *Oecologia*, 119, 239-246.
 937
 938Ohtonen, R., Ohtonen, A., Luotonen, H., Markkola, A., 1992. Enchytraeid and
 939nematode numbers in urban, polluted scots pine (*Pinus sylvestris*) stands in relation
 940to other soil biological parameters. *Biology and Fertility of Soils*, 13, 50-54.
 941
 942Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M.H.H., Oksanen, M.J.,
 943Suggests, M., 2007. The vegan package. Community ecology package. Version 1.8-
 9445.
 945
 946Orwin, K. H., Buckland, S. M., Johnson, D., Turner, B. L., Smart, S., Oakley, S.,
 947Bardgett, R. D., 2010. Linkages of plant traits to soil properties and the functioning
 948of temperate grassland. *Journal of Ecology*, 98, 1074-1083.
 949
 950Paterson, E., Osler, G., Dawson, L.A., Gebbing, T., Sim, A., Ord, B., 2008. Labile
 951and recalcitrant plant fractions are utilised by distinct microbial communities in soil:
 952Independent of the presence of roots and mycorrhizal fungi. *Soil Biology and*
 953*Biochemistry*, 40, 1103-1113.
 954

- 955Pavao-Zuckerman, M.A., Coleman, D.C., 2007. Urbanization alters the functional
956composition, but not taxonomic diversity, of the soil nematode community.
957Applied Soil Ecology, 35, 329-339.
- 958
- 959Pickett, S.T., Cadenasso, M.L., 2009. Altered resources, disturbance, and
960heterogeneity: A framework for comparing urban and non-urban soils. Urban
961Ecosystems, 12, 23-44.
- 962
- 963Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., 2007. Linear and nonlinear mixed
964effects models. R package version 3, 57.
- 965
- 966Pižl, V., 1992. Succession of earthworm populations in abandoned fields. Soil
967Biology and Biochemistry, 24, 1623-1628.
- 968
- 969Pouyat, R.V., Parmelee, R., Carreiro, M., 1994. Environmental effects of forest soil-
970invertebrate and fungal densities in oak stands along an urban-rural land use
971gradient. Pedobiologica, 38, 385-99.
- 972
- 973
- 974Raciti, S.M., Groffman, P.M., Jenkins, J.C., Pouyat, R.V., Fahey, T.J., Pickett, S.T.,
975Cadenasso, M.L., 2011. Accumulation of carbon and nitrogen in residential soils
976with different land-use histories. Ecosystems 14, 287-297.
- 977
- 978Ramirez, K.S., Leff, J.W., Barberan, A., Bates, S.T., Betley, J., Crowther, T.W.,
979Kelly, E.F., Oldfield, E.E., Shaw, E.A., Steenbock, C., Bradford, M.A., Wall, D.H.,
980Fierer, N., 2014. Biogeographic patterns in below-ground diversity in New York
981city's central park are similar to those observed globally. Proceedings of the royal
982society B, 281 20141988
- 983
- 984Setälä, H.M., Francini, G., Allen, J.A., Hui, N., Jumpponen, A., Kotze, D.J., 2016.
985Vegetation type and age drive changes in soil properties, nitrogen, and carbon
986sequestration in urban parks under cold climate. Frontiers in Ecology and Evolution,
9874, 93.
- 988
- 989Setälä, H., Francini, G., Allen, J.A., Jumpponen, A., Hui, N., Kotze, D.J., 2017.
990Urban parks provide ecosystem services by retaining metals and nutrients in soils.
991Environmental Pollution, 231, 451-461.
- 992
- 993Smetak, K., Johnson-Maynard, J., Lloyd, J., 2007. Earthworm population density
994and diversity in different-aged urban systems. Applied Soil Ecology, 37, 161-168.
- 995
- 996Sohlenius, B., 1979. A carbon budget for nematodes, rotifers and tardigrades in a
997Swedish coniferous forest soil. Ecography, 2, 30-40.
- 998
- 999Steinberg, D.A., Pouyat, R.V., Parmelee, R.W., Groffman, P.M., 1997. Earthworm
1000abundance and nitrogen mineralization rates along an urban-rural land use gradient.
1001Soil Biology and Biochemistry, 29, 427-430.
- 1002
- 1003Thomson, B.C., Ostle, N., McNamara, N., Bailey, M.J., Whiteley, A.S., Griffiths,
1004R.I., 2010. Vegetation affects the relative abundances of dominant soil bacterial taxa

- 1005and soil respiration rates in an upland grassland soil. *Microbial Ecology*, 59, 335-
1006343.
- 1007
- 1008Trofymow, J.A., Coleman, D.C., 1982. The role of bacterivorous and fungivorous
1009nematodes in cellulose and chitin decomposition. In: Freckman, D.W. (Ed.),
1010Nematodes in Soil Ecosystem. University of Texas Press, Austin, TX, pp. 117-138.
1011
- 1012Valtanen, M., Sillanpää, N., Setälä, H., 2015. Key factors affecting urban runoff
1013pollution under cold climatic conditions. *Journal of Hydrology* 529, 1578-1589.
1014
- 1015Wang, X., Wang, X., Zhang, W., Shao, Y., Zou, X., Liu, T., Zhou, L., Wan, S., Rao,
1016X., Li, Z., Fu, S., 2016. Invariant community structure of soil bacteria in subtropical
1017coniferous and broadleaved forests. *Scientific Reports*, 6, 19071.
1018
- 1019Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H.,
1020Wall, D.H., 2004. Ecological linkages between aboveground and belowground
1021biota. *Science*, 304, 1629-1633.
1022
- 1023Wardle, D.A., 2002. *Communities and Ecosystems: Linking the Aboveground and*
1024*Belowground Components* (MPB-34). Princeton University Press.
1025
- 1026White, D., Bobbie, R., King, J., Nickels, J., Amoe, P., 1979. Lipid analysis of
1027sediments for microbial biomass and community structure. In: Litchfield CD,
1028Seyfried PL (Eds.) *Methodology for biomass determination and microbial activities*
1029*in sediments*. ASTM STP 673, American Society for Testing and Materials,
1030Philadelphia, pp 87–103
1031
- 1032Xu, H., Li, S., Su, J., Nie, S., Gibson, V., Li, H., Zhu, Y., 2014. Does urbanization
1033shape bacterial community composition in urban park soils? A case study in 16
1034representative Chinese cities based on the pyrosequencing method. *FEMS*
1035*Microbiology Ecology*, 87, 182-192.
1036
- 1037Yeates, G.W., 2003. Nematodes as soil indicators: Functional and biodiversity
1038aspects. *Biology and Fertility of Soils*, 37, 199-210.
1039
- 1040Yeates, G., 1981. Soil nematode populations depressed in the presence of
1041earthworms. *Pedobiologia*, 22, 191-195.
1042
- 1043Yeates, G.W., Bongers, T., De Goede, R.G., Freckman, D.W., Georgieva, S.S.,
10441993. Feeding habits in soil nematode families and genera-an outline for soil
1045ecologists. *Journal of Nematology*, 25, 315-331.
1046
- 1047Zak, D.R., Grigal, D.F., Gleeson, S., Tilman, D., 1990. Carbon and nitrogen cycling
1048during old-field succession: Constraints on plant and microbial biomass.
1049*Biogeochemistry*, 11, 111-129.
1050
- 1051Zeller, V., Bardgett, R.D., Tappeiner, U., 2001. Site and management effects on soil
1052microbial properties of subalpine meadows: A study of land abandonment along a
1053north–south gradient in the European Alps. *Soil Biology and Biochemistry*, 33, 639-
1054649.

1055

Figure and Table Captions

1057

1058**Fig. 1** The effects of plant functional type and park age on a) total microbial
1059PLFAs, b) fungal PLFA (18:2 ω 6) and c) the PLFA fungal to bacterial ratio. Mean
1060values $\pm$ SE are presented.

1061

1062**Fig. 2** Relation between a) the C/N ratio and predicted values of total microbial
1063PLFAs, b) % soil C and predicted values of total microbial PLFAs, c) % soil N
1064and predicted values of fungal PLFA (18:2 ω 6) and d) pH and predicted values
1065of gram-/gram+ bacteria.

1066

1067**Fig. 3** NMDS plots for bacterial (1-6, on the left) and nematode (7-12, on the
1068right) communities. Microbial communities were grouped by plant functional
1069type (1, 3, 5) and age (2, 4, 6), and nematode communities were grouped by
1070plant functional type (7, 9, 11) and age (8, 10, 12) under lawn, deciduous and
1071evergreen trees in young, intermediate and old parks. Significant effects ($p <$
10720.05) are indicated with a check mark in the upper panels and significant
1073vectors are shown in the NMDS plots.

1074

1075**Fig. 4** The effects of plant functional type and park age on a) the nematode
1076channel ratio (NCR), and b) the nematode maturity index (MI). Mean values $\pm$
1077SE are presented.

1078

1079**Table 1.** GLMM results for PLFAs, nematodes and earthworms. For PLFAs the
1080response variables included total microbial PLFA, bacterial PLFA, fungal PLFAs
1081(18:2 ω 6, saprophytic fungi), fungal to bacterial ratio (fungi/bacteria) and gram + to
1082gram – ratio (gram+/gram-). For nematodes the response variables included
1083bacterial feeders, fungal feeders, plant feeders, predators and omnivorous, the
1084maturity index (MI) and the nematode channel ratio (NCR). For earthworms the
1085response variables were: *Lumbricus terrestris* and other earthworms. For each
1086variable we reported the coefficient, standard error and p-value. Significant effects
1087($p < 0.05$) are highlighted in bold and significant interactions are indicated with an
1088asterisk. Young evergreen trees are in the intercept.

1089

1090**Supplementary material, Table S2.** Means ($\pm$ 1 SD) of the following PLFAs:1091total microbial, bacterial, fungal PLFAs (18:2 ω 6, saprophytic fungi),

1092fungal/bacterial ratio, gram/gram+ ratio, nematodes: bacterial feeders, fungal

1093feeders, plant feeders, predators and omnivores, maturity index (MI),

1094nematode channel ratio (NCR), earthworms: *Lumbricus terrestris* and other

1095earthworms, for the three different park ages (young, intermediate and old)

1096and plant functional type (evergreen, deciduous, lawn). Values for Lahti old

1097parks and control forest (evergreen and deciduous trees) are also presented.

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

Table 1. GLMM results for PLFAs, nematodes and earthworms. For PLFAs the response variables included total microbial PLFA, bacterial PLFA, fungal PLFAs (8:2 ω 6, saprophytic fungi), fungal to bacterial ratio (fungi/bacteria) and gram + to gram – ratio (gram+/gram-). For nematodes the response variables included bacterial feeders, fungal feeders, plant feeders, predators and omnivorous, the maturity index (MI) and the nematode channel ratio (NCR). For earthworms the response variables were: *Lumbricus terrestris* and other earthworms. For each variable we reported the coefficient, standard error and p-value. Significant effects ($p < 0.05$) are highlighted in bold and significant interactions are indicated with an asterisk. Young evergreen trees are in the intercept.

	variable	intercept	lawn	deciduous	intermediate	old	age x functional group	pH	% soil C
PLFAs	Total microbial PLFAs	6.821	0.041	0.040	0.081	0.068			0.064
		0.220	0.092	0.092	0.103	0.100	*		0.023
		<0.001	0.447	0.437	0.791	0.675			<0.001
	Bacterial PLFAs	5.951	0.091	0.007	0.060	0.046			0.070
		0.228	0.094	0.094	0.105	0.102	*		0.023
		<0.001	0.336	0.942	0.565	0.655			<0.001
	Fungal PLFAs	3.414	0.129	-0.156	0.248	-0.320			-0.162
		0.290	0.266	0.261	0.272	0.274	*		0.085
		<0.001	0.627	0.549	0.362	0.243			0.056
	Fungi/bacteria	17.296	-4.994	8.044	-2.107	1.353			
		3.141	3.444	3.417	3.417	3.417			
		<0.001	0.147	0.019	0.537	0.692			
	Gram-/gram+	0.327	0.007	0.037	-0.064	-0.133		0.105	
		0.320	0.050	0.052	0.056	0.053		0.051	
		0.308	0.897	0.474	0.260	0.018		0.040	
Nematodes	Bacterial feeding	8.676	-2.038	-1.782	-2.121	-1.668			0.475
		2.279	0.847	0.832	0.860	0.852			0.218
		<0.001	0.016	0.321	0.013	0.050			0.029
	Fungal feeding	5.58	-1.322	-0.245	-0.35	-0.761			
		0.963	0.492	0.485	0.55	0.548			
		<0.001	0.01	0.612	0.524	0.165			
	Plant feeders	4.303	-1.030	-1.386	0.176	0.895			
		1.335	0.918	0.886	0.929	0.924			
		<0.001	0.262	0.118	0.849	0.333			
	Predators and omnivorous	-0.082	0.161	0.770	0.770	-0.109			
		1.049	0.438	0.438	0.462	0.463			
		0.937	0.712	0.104	0.095	0.813			
	MI	0.550	0.209	0.090	0.033	-0.074			
		0.119	0.059	0.060	0.064	0.064	*		
		<0.001	<0.001	0.134	0.600	0.246			

Earthworms	NCR	0.418	0.008	-0.09	-0.084	-0.012	
		0.094	0.055	0.055	0.066	0.065	
		<0.001	0.121	0.084	0.212	0.848	
	<i>Lumbricus terrestris</i>	1.920	-0.163	-0.242	1.452	1.097	-0.265
		0.725	0.320	0.319	0.406	0.404	0.123
		0.008	0.611	0.448	<0.001	0.007	0.032
	Other earthworms	3.657	0.794	0.433	0.160	0.368	-0.486
		1.355	0.190	0.196	0.226	0.217	0.200
		0.007	< 0.001	0.027	0.480	0.090	0.015

1132

Table 2 Means ($\pm$ 1 SD) of the following PLFAs: total microbial, bacterial, fungal PLFAs (18:2 ω 6, saprophytic fungi), fungal/bacterial ratio, gram/gram+ ratio, nematodes: bacterial feeders, fungal feeders, plant feeders, predators and omnivores, maturity index (MI), nematode channel ratio (NCR), earthworms: *Lumbricus terrestris* and other earthworms, for the three different park ages (young, intermediate and old) and plant functional type (evergreen, deciduous, lawn). Values for Lahti old parks and control forest (evergreen and deciduous trees) are also presented.

Variable		Young	SD	Intermediate	SD	Old	SD	Old Lahti	SD	Control	SD	
PLFA $\mu\text{g g}^{-1}$ soil dw	Total microbial PLFAs	Evergreen	380.09	132.83	526.96	208.32	417.16	93.44	318.82	64.09	842.19	319.05
		Deciduous	271.04	93.78	530.15	146.41	616.97	347.73	669.85	439.87	957.27	281.94
		Lawn	460.25	165.97	447.87	128.15	473.87	185.37				
	Bacterial PLFAs	Evergreen	172.85	66.11	223.54	88.39	181.19	50.83	160.42	46.90	799.71	306.51
		Deciduous	116.54	43.01	224.04	59.13	241.89	112.35	270.50	105.90	982.63	252.72
		Lawn	211.31	90.41	203.88	60.37	218.14	82.75				
	Fungal PLFAs	Evergreen	30.98	13.60	45.55	25.43	26.59	15.87	21.74	14.16	127.29	99.25
		Deciduous	24.65	9.00	48.12	28.67	111.03	125.14	141.06	156.35	127.97	100.76
		Lawn	34.26	12.77	23.88	11.22	26.16	16.65				
	Fungi/bacteria	Evergreen	20.33	12.39	21.17	11.73	15.96	11.25	14.05	8.43	18.39	15.22
		Deciduous	22.65	9.65	21.89	10.83	37.53	26.93	43.87	32.64	12.91	8.83
		Lawn	18.76	8.98	11.88	5.29	11.83	4.55				
	Gram+/gram-	Evergreen	2.98	0.45	2.77	0.31	2.49	1.02	3.07	1.02	3.14	0.83
		Deciduous	3.35	0.79	2.70	0.44	2.68	0.53	2.78	0.57	3.53	0.62
		Lawn	2.79	0.40	2.66	0.26	2.73	0.35				
Nematodes 10 g ⁻¹ soil	Bacterial feeders	Evergreen	139	113	100	85	93	47	64	33	74	31
		Deciduous	106	84	51	60	85	73	34	30	38	6
		Lawn	79	83	78	63	49	51				
	Fungal feeders	Evergreen	25	23	18	14	12	12	11	10	40	53
		Deciduous	24	25	20	30	13	9	12	7	13	8
		Lawn	11	11	10	14	7	6				
	Plant feeders	Evergreen	63	63	54	55	86	67	68	36	45	27
		Deciduous	37	32	43	53	48	33	47	36	16	24
		Lawn	129	283	69	55	136	285				
	Predators and omnivorous	Evergreen	9	8	12	9	7	11	4	4	19	15
		Deciduous	15	15	14	9	10	9	12	12	18	18
		Lawn	12	14	8	5	9	11				
	MI	Evergreen	2.17	0.11	2.30	0	2.35	0.23	2.42	0.03	2.15	0.20
		Deciduous	2.24	0.43	2.84	0.38	2.23	0.44	2.56	0.18	2.34	0.27
		Lawn	2.37	0.44	2.43	0.64	2.54	0.46				
	NCR	Evergreen	0.82	0.13	0.71	0.25	0.79	0.17	0.86	0.08	0.71	0.30
		Deciduous	0.77	0.19	0.85	0.12	0.88	0.09	0.70	0.17	0.76	0.12
		Lawn	0.88	0.12	0.88	0.10	0.87	0.10				
Earthworms g m ⁻²	<i>Lumbricus terrestris</i>	Evergreen	34	43	88	60	62	65	99	65	115	133
		Deciduous	15	32	63	49	66	67	102	77	97	100
		Lawn	27	30	54	67	43	37				
	Other earthworms	Evergreen	8	10	11	12	9	12	12	15	14	11
		Deciduous	9	13	19	32	19	13	25	13	50	48
		Lawn	16	16	36	44	35	40				

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151 Fig 1.

1152

1153

1154Fig 2.

1155

1156

1157Fig. 3

1158

1159

1160
1161 Fig. 4
1162
1163