<https://helda.helsinki.fi>

Influence of postoperative pain and use of NSAID on heart rate variability of dairy cows

Frondelius, Lilli

2018-02

Frondelius , L , Hietaoja , J K , Pastell , M , Hänninen , L T , Anttila , P & Mononen , J 2018 ,
' Influence of postoperative pain and use of NSAID on heart rate variability of dairy cows ' ,
pö Journal of Dairy Research , vol. 85 , no. 1 , pp. 27 29 . <https://doi.org>

<http://hdl.handle.net/10138/312764>

<https://doi.org/10.1017/S0022029917000760>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Influence of postoperative pain and use of NSAID on heart rate variability of dairy cows

Lilli Frondelius¹, Juha Hietaoja², Matti Pastell¹, Laura Hänninen³, Paula Anttila⁴, Jaakko Mononen¹

¹ Natural Resources Institute Finland (Luke), Green technology, Halolantie 31 A, 71750 Maaninka, Finland

² University of Helsinki, Faculty of Agriculture and Forestry, Department of Agricultural Sciences, P.O Box 28, 00014 University of Helsinki, Finland

³ University of Helsinki, Faculty of Veterinary Medicine, Department of Production Animal Medicine, P.O. Box 57, 00014 University of Helsinki, Finland

⁴ Maalaislääkäri Anttila, Halolantie 27, 71750 Maaninka, Finland

correspondent author: Lilli Frondelius, lilli.frondelius@luke.fi

Summary

This Research communication describes the effect of post-operative pain and NSAID treatment on heart rate variability (HRV) of dairy cows. Postoperative pain in farm animals is often left untreated. HRV could be a promising tool for pain assessment in animals. The aim of this study was to assess if postoperative state after subcutaneous surgery and NSAID treatment affect HRV in dairy cows. Nine cows were inserted with an implantable electrocardiograph logger. Cows were divided into the NSAID treatment group and the control group. The cows in the NSAID group had higher HRV than the control group, indicating a higher sympathetic activity in control animals, most likely due to untreated post-operative pain. Besides the ethical need for treating pain in production animals, ongoing pain has an adverse effect on animal productivity. Thus post-operative pain alleviation is recommended.

Key words: Dairy cattle, Heart rate variability, Pain, Nonsteroidal anti-inflammatory drug

24

25 In most cases involving farm animals, intraoperative pain during veterinary procedures is treated
26 properly, but treatment for postoperative pain is seldom given (Walker *et al.* 2011). Behaviours,
27 such as restlessness, foot stamping and changes in lying behaviour, are used for assessing farm
28 animal pain, but also physiological parameters, such as stress hormones, are assessed (Viñuela-
29 Fernández *et al.* 2007). However, both behavioural and traditional physiological approaches have
30 their limitations in sensitivity and reliability and new methods for effective pain assessment are
31 needed. Heart rate variability (HRV) can be used as an indicator of the sympathovagal balance of
32 the autonomous nervous system (Task Force 1996) and HRV is a promising tool for pain
33 assessment in farm animals (calves: Stewart *et al.* 2008 and 2010; ewes: Stubbsjøen *et al.* 2009).
34 However, results of HRV measurements are inconsistent: pain has been shown either to increase
35 sympathetic (e.g. Stewart *et al.* 2008, Stubbsjøen *et al.* 2009) or vagal (e.g. Stewart *et al.* 2010)
36 activity. Thus there is a need for greater understanding of the underlying mechanism of HRV
37 response, especially for different types of pain (e.g. somatic and visceral).

38

39 In cattle, studies on pain and HRV are mainly conducted with disbudded or castrated calves
40 (Stewart *et al.* 2008, 2010) rather than with adult animals. The aim of the study was to assess if
41 postoperative somatic pain after small-scale surgery and postoperative nonsteroidal anti-
42 inflammatory drug (NSAID) treatment affect HRV in dairy cows. This experiment was a part of a
43 larger study focusing on the effect of stress on HRV in dairy cows measured with implantable
44 technology, and approved by the National Animal Experiment Board (Permit Number: ESHL-2008-
45 08892/Ym-23).

46

47 Materials and methods

48 The study was conducted at the Natural Resources Institute Finland research barn in Maaninka,
49 Finland. The experimental animals were four pregnant heifers and seven dry cows (six Holstein-
50 Friesian and five Ayrshire). The animals were tethered in peat-bedded rubber-matted tie stalls
51 (width 120 cm, length 180 cm). They were fed daily with 0.3 kg of barley concentrate (energy
52 content 13.2 MJ ME/kgDM), 0.2 kg commercial mineral for dry cows (Tunnu-Namino, Hankkija Oy,
53 Finland), and 32 kg (multiparous cows) or 36 kg (heifers) grass silage (10.5 MJ ME/kgDM).
54
55 Implantable loggers (Bjarnason *et al.* 2013) were used to measure the electrocardiograph (ECG) of
56 the experimental animals (Fig. 1) with a sample rate of 341.3 Hz. The system contained a 50 g 70
57 mm x 45 mm x 17 mm) implant placed on the left side of the animal into a pocket made between
58 the skin and subcutaneous tissue posterior to the scapula, and a receiver box attached to the side
59 of the cow with an elastic girth.

60

61

62 Figure 1. a) Implant for ECG measurements and b) implant and receiver box after operating on the
63 cow.

64

65 The animals were operated about one month ($\pm$ 3 days) prior to the expected parturition date
66 under sedation (0.018 mg/kg xylazine hydrochloride 20 mg/ml intravenously) and local anesthesia
67 (20 – 35 ml lidocaine hydrochloride 20 mg/ml subcutaneously). The animals received three-day
68 antibiotic treatment (20 000 IU/kg penicillin benzathine and penicillin procaine 300 000 IU/ml
69 intramuscularly) post-operatively.

70

71 Heifers and cows were randomly divided into the treatment (1 heifer + 5 cows) and the control (3
72 heifers + 2 cows) groups. On the operation day the treatment group received intravenous NSAID
73 treatment (3 mg/kg ketoprofen 100 mg/ml). An oral NSAID treatment (4 mg/kg ketoprofen 160
74 mg/g) mixed in a small amount of concentrates continued for the four consecutive days (days 1 –
75 4) after the operation day. The control group did not receive any NSAID treatment, but a small
76 amount of concentrates as a control treatment on the days 1 – 4. An oral ketoprofen has been
77 effectively used previously e.g. with acute mastitis in dairy cows (Banting *et al.* 2008).

78

79 ECG of the experimental animals was measured and behaviour video recorded during days 1 – 4
80 using cameras (Axis Q1755-E and Axis 215) positioned to face downwards to the tie-stalls. The
81 body and head postures and rumination of the animals were recorded continuously twice daily,
82 05.00 – 07.00 and 22.00 – 24.00. Because the posture of the animal affects the HRV (Frondeus *et*
83 *al.* 2015) only the ECG data from continuous lying periods (sternal recumbency while head up)
84 from animals not ruminating were used for the HRV analysis. ECG data from these periods were
85 visually inspected and good quality five-minute segments were selected for further analysis. Data
86 from two control group animals were discarded because of unreadable ECG data due to
87 malfunctioning of the implants.

88

89 The ECG data were interpolated to 500Hz using a geometric interpolation method (Tiusanen *et al.*
90 2015) and the location of the S wave (Tiusanen & Pastell 2016) was used to analyze the normal-to-
91 normal (NN) intervals. We calculated heart rate (HR), the standard deviation of NN intervals
92 (SDNN), root mean square of successive NN interval differences (RMSSD) and pNN50 (Task Force
93 1996).

94
95 HRV data from the nine animals (treatment n=6, control n=3) were used for the analysis including
96 mean daily values of HRV parameters from 94 five-minute segments; 58 segments from the
97 treatment group and 36 segments from the control group, 10.4±5.87 (mean±SD) segments per
98 cow. Linear mixed models were fitted to determine the effect of independent variables –
99 treatment (control vs NSAID), time from the procedure (days 1 – 4) and the interaction between
100 these two – on HRV parameters (dependent variables) using compound symmetry as a covariance
101 structure and cow as a random effect. The statistical analyses were made with SAS for Windows
102 version 9.2 through SAS Enterprise Guide version 4.3 (SAS Institute Inc., Cary, NC, USA).

103

104 Results and discussion

105 The cows in the treatment group had higher SDNN and tended to have higher RMSSD values than
106 the animals in the placebo group (Table 1). There were no differences in HR and pNN50 between
107 two groups. The day did not have any effect on HRV values and there was no interaction between
108 the treatment and the day.

109

110 The results indicate that NSAID affects the postoperative HRV of adult dairy cows that had
111 undergone subcutaneous implantation. The control group had a lower vagal tone and a stronger
112 sympathetically controlled response (von Borell *et al.* 2007) to the postoperative state, most likely

Commented [FL1]: Table 1 near here

113 due to untreated pain. Ketoprofen might have also mitigated the inflammatory reaction towards
114 the implant due to NSAIDs anti-inflammatory and anti-pyretic effects (Donalisio *et al.* 2012).

115

116 We did not observe any effect of NSAID treatment on long-term HR. This result is similar to
117 Raekallio *et al.* (1997), who studied horses for 72 hours after arthroscopic surgery. Generally higher
118 sympathetic activity is connected to increase in HR (Von Borell *et al.* 2007) and lack of pain
119 medication increases HR (Stewart *et al.* 2008, Stubbsjøen *et al.* 2009). In these studies, however, HR
120 was measured only during the painful procedure or few hours after it. This difference in methods
121 may partly explain lack of difference in HR in our study, since we started measurements of cardiac
122 responses one day after the surgery and measured it throughout the whole four-day post-
123 operative treatment. However, it is also possible that ketoprofen was not a potent enough NSAID
124 to cause differences in HR, as in the study by Newby *et al.* (2014).

125

126 With this small number of the animals, our experiment should be regarded as a case study.
127 However, linked with other results, we suggest that HRV may be a sensitive measure for assessing
128 pain in production animals. Stubbsjøen *et al.* (2009), who also used a limited number of animals in
129 their study, found that moderate pain in sheep had an effect on HRV.

130

131 This study shows that dairy cows are subjected to postoperative pain even after small-scale
132 subcutaneous surgery. In production animals postoperative pain is often neglected (Walker *et al.*
133 2011). Besides the ethical need for treating pain, ongoing pain may also have an adverse effect on
134 animal productivity, affecting their growth and immune functions (Anil *et al.* 2005). Thus pain
135 alleviation during and after painful procedures is beneficial for the animal, the producer and the
136 consumer.

137

138 Acknowledgments

139 The authors wish to thank the staff of Luke Maaninka research barn for technical assistance and
140 Vetcare Oy for providing pharmaceuticals used in the study. This study was part of a project
141 funded by the Maaseudun kehittämisrahasto under the Finnish Ministry of Agriculture and
142 Fisheries (grant number 2872/502/2008).

143

144 References

145 **Anil L, Anil SS & Deen J** 2005 Pain detection and amelioration in animals on the farm: Issues and
146 options. *Journal of Applied Animal Welfare Science* **8** 261 – 278

147

148 **Banting A, Banting S, Heinonen K & Mustonen K** 2008 Efficacy of oral and parenteral ketoprofen
149 in lactating cows with endotoxin-induced acute mastitis. *Veterinary Record* **163** 506 – 509

150

151 **Bjarnason Á, Vuorela T, Verho J, Riistama J, Vanhala J, Lekkala J & Hyttinen J** 2013 Implantable
152 measurement system for dairy-cattle monitoring with long recording time. *Advances in Science*
153 *and Technology* **85** 33 – 38

154

155 **Donalisio C, Barbero R, Cuniberti B, Vercelli C, Casalone M, Re G** 2012 Effects of flunixin
156 meglumine and ketoprofen on mediator production in ex vivo and in vitro models of inflammation
157 in healthy dairy cows. *Journal of Veterinary Pharmacology and Therapeutics* **36** 130 – 139

158

159 **Frondeius L, Järvenranta K, Koponen T & Mononen J** 2015 The effects of body posture and
160 temperament on heart rate variability in dairy cows. *Physiology & Behavior* **139** 437 – 441

161

162 **Newby NC, Tucker CB, Pearl DL, LeBlanc SJ, Leslie KE, von Keyserlingk MAG, Duffield TF** 2014 An
 163 investigation of the effects of ketoprofen following rumen fistulation surgery in lactating dairy
 164 cows. *The Canadian Veterinary Journal* **55** 442 – 448

165

166 **Raekallio M, Taylor PM & Bennett RC** 1997 Preliminary investigations of pain and analgesia
 167 assessment in horses administered phenylbutazone or placebo after arthroscopic surgery.
 168 *Veterinary Surgery* **26** 150 – 155

169

170 **Stewart M, Stafford KJ, Dowling SK, Scafer AL & Webster JR** 2008 Eye temperature and heart
 171 rate variability of calves disbudded with or without local anaesthetic. *Physiology & Behaviour* **93**
 172 789 – 797

173

174 **Stewart M, Verkert GA, Stafford KJ, Schaefer AL & Webster JR** 2010 Noninvasive assessment of
 175 autonomic activity for evaluation of pain in calves, using surgical castration as a mode. *Journal of*
 176 *Dairy Science* **96** 3602 – 3609

177

178 **Stubsjøen SM, Flø AS, Moe RO, Janczak AM, Skjerve E, Valle PS & Zanella AJ** 2009 Exploring non-
 179 invasive methods to assess pain in sheep. *Physiology & Behaviour* **98** 640 – 648

180

181 **Task Force of The European Society of Cardiology, North American Society of Pacing and**
 182 **Electrophysiology** 1996 Heart rate variability – Standards of measurement, physiological
 183 interpretation, and clinical use. *Circulation* **93** 1043 – 1065

184

185 **Tiusanen J, Hautala MI, Ternman EM & Pastell ME** 2015 Geometrical method for interpolating S-
186 peaks from cow ECG using a microcontroller. *Biosystems Engineering* **129** 324 – 328
187

188 **Tiusanen J & Pastell M** 2016 Simple online algorithm for detecting cow's ECG beat-to-beat interval
189 using a microcontroller. *Agricultural Engineering International: CIGR Journal* **18** 411 – 418
190

191 **Viñuela-Fernández I, Jones E, Welsh EM & Fleetwood-Walker SM** 2007 Pain mechanism and their
192 implication for the management of pain in farm and companion animals. *The Veterinary Journal*
193 **174** 227 – 239
194

195 **Von Borell E, Langbein J, Deprés G, Hansen S, Leterrier C, Marchant-Forde J, Marchand-Forde R,**
196 **Minero M, Mohr E, Prunier A, Valance D & Veissier I** 2007 Heart rate variability as a measure of
197 autonomic regulation of cardiac activity for assessing stress and welfare in farm animals – A
198 review. *Physiology & Behavior* **92** 293 – 316
199

200 **Walker KA, Duffield TF & Weary DM** 2011 Identifying and preventing pain during and after
201 surgery in farm animals. *Applied Animal Behaviour Science* **135** 259 – 265
202

203 Table 1. Comparison of the linear mixed model estimates of heart rate variability (HRV) variables
204 (mean±SEM) between NSAID (cows n=6; analyzed 5-minute HRV segments n=58) and placebo
205 groups (cows n=3; analyzed 5-minute HRV segments n=36).

	NSAID	Placebo	P
HR	61.0±3.84	61.8±5.55	ns

SDNN	56.1±2.16	45.6±3.39	<0.05
RMSSD	60.8±1.96	54.3±3.09	<0.1
pNN50	6.21±1.58	4.96±2.40	ns