<https://helda.helsinki.fi>

Predictive mapping of mosquito distribution based on environmental and anthropogenic factors in Taita Hills, Kenya

Uusitalo, Ruut Jaael

2019-04

Uusitalo , R J , Siljander , M , Culverwell , C L , Mutai , N , Forbes , K M , Vapalahti , O & Pellikka , P K E 2019 , ' Predictive mapping of mosquito distribution based on environmental and anthropogenic factors in Taita Hills, Kenya ' , International Journal of Applied Earth Observation and Geoinformation , vol. 76 , pp. 84-92 . <https://doi.org/10.1016/j.jag.2018.11.004>

<http://hdl.handle.net/10138/321847>

<https://doi.org/10.1016/j.jag.2018.11.004>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Predictive mapping of mosquito distribution based on environmental and anthropogenic factors in
Taita Hills, Kenya

Ruut Uusitalo^{1,2,3}, Mika Siljander¹, C. Lorna Culverwell^{2,3,4}, Noah C. Mutai⁵, Kristian M. Forbes²,
Olli Vapalahti^{2,3,6} Petri K.E. Pellikka^{1,7,8}

¹Department of Geosciences and Geography, P.O. Box 64, FI-00014 University of Helsinki,
Finland; mika.siljander@helsinki.fi; petri.pellikka@helsinki.fi

²Department of Virology, Haartmaninkatu 3, P.O. Box 21, FI-00014 University of Helsinki,
Finland; lorna.culverwell@helsinki.fi; kristian.forbes@helsinki.fi; olli.vapalahti@helsinki.fi

³Department of Veterinary Biosciences, Agnes Sjöberginkatu 2, P.O. Box 66, FI-00014 University
of Helsinki, Finland

⁴Department of Life Sciences, Natural History Museum, Cromwell Road, London, SW5 7BD

⁵Department of Mathematics and Informatics, Taita Taveta University, P. O. Box 635-80300, Voi,
Kenya

⁶Virology and Immunology, HUSLAB, Helsinki University Hospital

⁷Helsinki Institute of Sustainability Science, University of Helsinki, Finland

⁸Institute for Atmospheric and Earth System Research, University of Helsinki, Finland

Corresponding author:

Ruut Uusitalo

University of Helsinki, Department of Geosciences and Geography, P.O. Box 64, FI-00014
ruut.uusitalo@helsinki.fi; Tel.: +358 (0)50 5911280

Abstract

Mosquitoes are vectors for numerous pathogens, which are collectively responsible for millions of
human deaths each year. As such, it is vital to be able to accurately predict their distributions,
particularly in areas where species composition is unknown. Species distribution modeling was used
to determine the relationship between environmental, anthropogenic and distance factors on the
occurrence of two mosquito genera, *Culex* Linnaeus and *Stegomyia* Theobald (syn. *Aedes*), in the
Taita Hills, southeastern Kenya. This study aims to test whether any of the statistical prediction
models produced by the Biomod2 package in R can reliably estimate the distributions of mosquitoes

in these genera in the Taita Hills; and to examine which factors best explain their presence. Mosquito collections were collected from 122 locations between January–March 2016 along transects throughout the Taita Hills. Environmental-, anthropogenic- and distance-based geospatial data were acquired from the Taita Hills geo-database, satellite- and aerial imagery and processed in GIS software. The Biomod2 package in R, intended for ensemble forecasting of species distributions, was used to generate predictive models. Slope, human population density, normalized difference vegetation index, distance to roads and elevation best estimated *Culex* distributions by a generalized additive model with an area under the curve (AUC) value of 0.791. Mean radiation, human population density, normalized difference vegetation index, distance to roads and mean temperature resulted in the highest AUC (0.708) value in a random forest model for *Stegomyia* distributions. We conclude that in the process towards more detailed species-level maps, with our study results, general assumptions can be made about the distribution areas of *Culex* and *Stegomyia* mosquitoes in the Taita Hills and the factors which influence their distribution.

Keywords: species distribution modeling; vector-borne disease; GIS; predictive mapping; mosquito, biomod2

1. Introduction

Environmental and anthropogenic disturbances such as climate change, urbanization and deforestation are crucial factors in the distribution of pathogen vectors and the emergence of diseases that they transmit (Crowl *et al.*, 2008). This is already evident as the global rise in human population densities, and correlated mean temperatures leading to land-use changes, has created new suitable breeding sites for mosquitoes (Roiz *et al.*, 2011; Campbell *et al.*, 2015). Geographic information system (GIS) and species distribution modeling (SDM) approaches have been used to understand these connections and their implications for the spread of invasive species, particularly with regard to northern regions (Neteler *et al.*, 2011). Other important areas, however, remain unexplored, including biodiversity rich but increasingly fragmented parts of Africa, where infectious diseases can be highly prevalent but access to health care is limited (Bhutta *et al.*, 2014). As a result, tools such as GIS and SDM that can estimate and predict infectious disease risks have the potential to optimize the use of limited resources and improve public health outcomes.

Due to their ability to act as vectors for a suite of pathogens, mosquitoes (Diptera, Culicidae) are among the most economically and socially important taxa on the planet (WHO, 2017). Two notable

69 genera are *Culex* Linnaeus and *Stegomyia* Theobald (following the classification of Reinert *et al.*,
70 (2009)). *Culex* is a large genus of mosquitoes, with 769 species in 26 subgenera (MTI, 2017),
71 accounting for 21.6% of all mosquito species worldwide. Species of *Culex* have an almost worldwide
72 distribution from the tropics to cool temperate regions, but do not extend into extreme northern
73 latitudes (MTI, 2017). While species from other subgenera are known to vector pathogens, subgenus
74 *Culex* (*Culex*) contains many of the significant human vector species, including those demonstrated
75 to transmit West Nile fever virus, Rift Valley fever virus and Japanese Encephalitis virus (MTI, 2017;
76 WHO, 2017). *Stegomyia* is a moderately sized mosquito genus comprising 128 species, and
77 distributed in the Afrotropical, Oriental and Australasian regions (MTI, 2017). Following human
78 dispersal, at least two species, *St. aegypti* and *St. albopicta*, are also present in Neotropical, Nearctic
79 and Palearctic regions (MTI, 2017; Paupy, *et al.*, 2009). Member species are competent vectors of
80 yellow fever virus (Huang, 1986), dengue virus serotypes 1-4, Zika virus, and Chikungunya virus,
81 among others (Huang, 1990; MTI, 2017).

82
83 SDM helps to understand the relationship between wildlife and their environments, to estimate their
84 distributions where empirical data is missing, and to forecast range expansions in the face of
85 environmental change. SDM is used in decision-making for a range of global challenges, such as
86 informing protected areas for wildlife conservation (Guisan *et al.*, 2013), the distribution potential for
87 invasive species (Václavík & Meentemeyer, 2009), and investigating the effects of climate change
88 on species that carry pathogens of public health concern such as mosquitoes (Dukes *et al.*, 2009).
89 Species distribution modeling, therefore, provides a means to understand and predict vector responses
90 to changing climate patterns in dynamic and fragmented environments.

91
92 The purpose of this study is to test whether any of the statistical prediction models produced by the
93 Biomod2 package in R, can reliably estimate the distributions of mosquitoes in genera *Culex* and
94 *Stegomyia* in the Taita Hills; to examine which factors best explain their presence; and to create maps
95 of their predicted occurrence. The target area, Taita Hills in rural south-eastern Kenya, is an
96 ecologically diverse area with little previous mosquito research, strong variability in rainfall and a
97 rapidly growing human population. While machine-learning techniques such as maximum entropy
98 modeling is often used to understand mosquito distributions (Mughini-Gras *et al.*, 2014; Sallam *et al.*,
99 *et al.*, 2016), the employed R package, the Biomod2 (Thuiller *et al.*, 2016), has important benefits for
100 estimating mosquito distributions. This is due to its ability to provide improved simulations across
101 initial conditions, test multiple model classes and parameterizations, and include unlimited number
102 of boundary conditions (Thuiller *et al.*, 2009).

relationships, and it may result in more accurate predictions than other model types (Yee *et al.*, 1991). For *Stegomyia* estimations, we focused on the RF model because of its advantages over other models, including high classification accuracy and the ability to model interactions among predictors (Cutler *et al.*, 2007). However, some uncertainties exist concerning *Stegomyia* estimations due to a higher number of absences (94 collections) than presences (28 collections) in the sampling data. Environmental, anthropogenic and distance variables were spatially autocorrelated to some extent, which may, among other things, affect the precision of coefficients (Diniz- Filho *et al.*, 2003).

In our study, the spatial resolution of environmental data was 20 meters. This resolution may have produced some bias to the study results because mosquitoes are also breeding e.g. in tree holes and water tanks in different microclimates. The lack of very high-resolution environmental data (such as the resolutions of 2 meters and 5 meters) is a general problem and not limited just to the Taita Hills. In addition, increasing the resolution of environmental data by deriving them from a high-resolution environmental data does not always improve the species distribution models (Pradervand *et al.*, 2014). Instead, very high-resolution environmental predictors should be produced by taking into account more local field measurements such as fine environmental mapping or in-situ measurements (Pradervand *et al.*, 2014; Lembrechts *et al.*, 2018). The potential solution for the lack of microclimate data could be the use of unmanned aerial vehicles (UAVs) in order to map the sampling areas in real time, and to produce higher resolution environmental data of even 3 centimeters (Kimberly *et al.*, 2014; Anderson *et al.*, 2013). Furthermore, there certainly exist other explanatory factors affecting the mosquito distributions which were not included in the modelling process including landscape fragmentation indicators. Fragmentation variables, such as distance from forest patch, patch size, distance from patch edge and the landscape metric of PPU (patches per unit) have resulted significant statistical relations with mosquito distributions (Richman *et al.*, 2018; Reiter & LaPointe 2007). For example, the use of FRAGSTATS- a spatial pattern analysis program may add value for our future mosquito habitat studies (McGarical *et al.*, 2002).

Consistent with previous research from other global locations (Sallam *et al.*, 2017; Ding *et al.*, 2017), environmental and anthropogenic variables were important determinants of *Culex* and *Stegomyia* mosquito distribution in the Taita Hills. For example the response curve for slope (Fig 3.) shows high probability for *Culex* presence for slopes less than 35°. In the Taita Hills, dwelling units are mainly absent on slopes greater than 35° (Siljander *et al.*, 2011). High probability for *Culex* presence were found with high NDVI values caused by strongly reflecting orchard trees and croplands adjacent to dwelling units in the Taita Hills. Specifically, we found that locations with high population densities,

short distances to roads, and gentle slope angles positively influenced *Culex* presence. Residential and urban areas are often recognized as important for *Culex* distributions (Reiter & LaPointe, 2007; Conley *et al.*, 2014). Rich and poor vegetation and all elevations were suitable for *Culex* mosquito presence, strengthening previous findings of its widespread distribution (MTI, 2017). However, we cannot rule out some effects of potential sample biases (e.g. time of the day and traps involved) which may have affected the mosquito spectrum captured. Sampling bias may also have been the reason for an unexpected response curve for population density (*popden*) at values from ca. 4000 to 5000 (Fig. 3).

Our study suggests, that *Stegomyia* mosquitoes prefer locations with lower human population densities, higher temperatures and solar radiation, and poor or moderate vegetation. This finding somewhat contradicts the notion that *St. aegypti* distribution is linked to growing human population (Fatima *et al.*, 2016), but is consistent with historic records of the variety of *Stegomyia* species suggesting that most members of the *Stegomyia* genus are forest dwelling, with only some adapted to breeding close to human habitation (Powell & Tabachnick 2013). Since our collections included several different *Stegomyia* species - with *St. aegypti* being the only species collected from around human habitation, and other species being restricted to the forest - the model findings are consistent with these historical reports (Powell & Tabachnick 2013). *Stegomyia* favored locations with intermediate and high temperatures, confirming the argument that *Stegomyia* mosquitoes have temperature-based limits to survival (Brady *et al.*, 2013). Overall, anthropogenic and distance factors appeared to be more important than environmental drivers for the distribution of both *Culex* and *Stegomyia* mosquitoes. We note that other variables, not considered in this study, may also affect the distribution of *Culex* and *Stegomyia* mosquitoes and that the genus-level pooling of the species will mask species distribution determinants, which will be targeted in future work with larger and more detailed datasets.

5. Conclusions

Our results affirm the utility and reliability of the Biomod2 package in R as a valid modeling method for species distributions, resulting in insights into vector-ecological interactions. Such work will be further refined with species-level identification and screening for arboviruses. With our study results, general assumptions can be made about the distribution of mosquitoes belonging to *Culex* and *Stegomyia* genera in the Taita Hills and the factors that influence their distribution. High population densities, short distance to roads and gentle slope angles were associated with the occurrence of *Culex* mosquitoes. For *Stegomyia* mosquitoes, on the other hand, low human population densities, high

temperatures and solar radiations as well as poor or moderate vegetation were suitable factors. Together, these results have implications for applying the approach for identifying risk areas and optimizing the use of limited resources for mitigation strategies.

351

Acknowledgements

We are thankful to Essi Korhonen, Joni Uusitalo, Masika Moses and Peter Mwazi for assistance with the fieldwork, and Miska Luoto, Juha Aalto, Sakari Äärilä and Ninna Malinen for thoughtful discussion on the topic. We also acknowledge the logistical support provided by the Taita Research Station of the University of Helsinki. Funding for this work was provided by the Jane and Aatos Erkko Foundation, the Ministry for Foreign Affairs of Finland TAITAGIS project (http://www.cimo.fi/programmes/hei_ici_index/programmes/hei_ici/projects/taitagis), and the Academy of Finland. The work was carried out under a research permit for Taita Research Station from the National Council for Science and Technology, Kenya (permit NCST/RCD/17/012/33).

361

References

- Adhikari, H., Heiskanen, J., Siljander, M., Maeda, E., Heikinheimo V., & Pellikka, P.K.E. (2017). Determinants of Aboveground Biomass across an Afromontane Landscape Mosaic in Kenya. *Remote Sensing*, 9(8), 827.
- Anderson, K., & Gaston, K. (2013). Lightweight unmanned aerial vehicles will revolutionize spatial ecology. *Frontiers in Ecology and the Environment*, 11(3), 138-146.
- Bhutta, Z.A., Sommerfeld, J., Lassi, Z.S., Salam, R.A., & Das, J.K. (2014). Global burden, distribution, and interventions for infectious diseases of poverty. *Infectious Diseases of Poverty*, 3, 21.
- Broberg, A., Salminen, H., Tolvanen, R., & Ylhäisi, J. (2004). Werugha - village in the heart of the Taita Hills. In Pellikka, P., Ylhäisi, J., & B. Clark (Eds.), *Taita Hills and Kenya, 2004–seminar, reports and journal of a field excursion to Kenya* (pp.108-113). Expedition reports of the Department of Geography. University of Helsinki.
- Campbell, L. P., Luther, C., Moo-Llanes, D., Ramsey, J. M., Danis-Lozano, R., & Peterson, A. T. (2015). Climate change influences on global distributions of dengue and chikungunya virus vectors. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370(1665).
- Clark, B., & Pellikka, P. (2005, June). The development of a land use change detection methodology for mapping the Taita Hills, south-east Kenya. CD Rom publication at the meeting of the Proceedings of the 31st International Symposium of Remote Sensing of the Environment, St. Petersburg, Russia.

Climate-Data.org (2018). Monthly average rainfall in Maktau. <https://en.climate-data.org/location/103844/> Accessed 30 April 2018.

Conley, A., Fuller, D., Haddad, N., Hassan, A., Gad, A., & Beier, J. (2014). Modeling the distribution of the West Nile and Rift Valley fever vector *Culex pipiens* in arid and semi-arid regions of the Middle East and North Africa. *Parasites & Vectors*, 7, (289).

Crowl, T., Crist, T., Parmenter, R., Belovsky, G., & Lugo, A. (2008). The spread of invasive species and infectious disease as drivers of ecosystem change. *Frontiers in Ecology and the Environment*, 6, 238–246.

Cutler, D.R., Edwards, T.C.Jr., Beard, K.H., Cutler, A., Hess, K.T., Gibson, J., & Lawler, J.J. (2007). Random forests for classification in ecology. *Ecology*, 88, 2783–2792.

Ding, F., Fu, J., Jiang, D., Hao, M., & Lin, G. (2017). Mapping the spatial distribution of *Aedes aegypti* and *Aedes albopictus*. *Acta Tropica*, 178, 155–162.

Diniz- Filho, J., Bini, L., & Hawkins, B. (2003). Spatial autocorrelation and red herrings in geographical ecology. *Global Ecology & Biogeography*, 12, 53–64.

Drew, C., Wiersma, Y., & Huettmann, F. (2011). *Predictive species and habitat modeling in landscape ecology*. 319p. The United States.

Dukes, J., Pontius, J., Orwig, D., Garnas, J., Rodgers, V., Brazee, N., Cooke, B., Theoharides, K., Stange, E., Harrington, R., Ehrenfeld, J., Gurevitch, J., Lerda, M., Stinson, K., Wick, R. & M. Ayres (2009). Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: What can we predict? *Canadian Journal of Forest Research*, 39, 231–248.

Environment Systems Research Institute (ESRI). (1991). ARC/INFO User's guide. Cell-based modelling with GRID. Analysis, display and management (Redlands, CA: ESRI). European Spatial Agency (ESA). (2015). Sentinel-2 user handbook. *ESA Standard Document*, 64.

Erdogan, E.H., Pellikka, P., & Clark, B. (2011). Modelling the impact of land-cover change on potential soil loss in the Taita Hills, Kenya, between 1987 and 2003 using remote-sensing and geospatial data. *International Journal of Remote Sensing*, 32(21), 5919–5945.

Fatima, S., Atif, S., Rasheed, S., Zaidi, F., & Hussain, E. (2016). Species distribution modeling of *Stegomyia aegypti* in two dengue-endemic regions of Pakistan. *Tropical Medicine and International Health*, 21, 427–436.

Fielding, A., & Bell, J. (1997). A review of methods for the assessment of prediction errors in conservation presence/ absence models. *Environmental Conservation*, 24, 38–49.

414 Fornace, K., Drakeley, C., William, T., Espino, F. & Cox, J. (2014). Mapping infectious disease
 415 landscapes: unmanned aerial vehicles and epidemiology. *Trends in Parasitology*, 30(11), 514-
 416 519.

417 Franklin, J., & Miller, J. (2010). Statistical methods – modern regression. In: Franklin, J.(ed.)
 418 *Mapping species distribution: spatial inference and prediction*. Cambridge: Cambridge
 419 University Press, (pp. 340).

420 Guisan, A., & Zimmermann, N. (2000). Predictive habitat distribution models in ecology.
 421 *Ecological Modeling*, 135(2–3), 147–186.

422 Guisan, A., Tingley, R., Baumgartner, J., Naujokaitis-Lewis, I., Sutcliffe, P., Tulloch, A., Regan, T.,
 423 Brotons, L., McDonald-Madden, E., Mantyka-Pringle, C., Martin, T., Rhodes, J., Maggini, R.,
 424 Setterfield, S., Elith, J., Schwartz, M., Wintle, B., Broennimann, O., Austin, M., Ferrier, S.,
 425 Kearney, M., Possingham, H., & Buckley, Y. (2013). Predicting species distributions for
 426 conservation decisions. *Ecology Letters*, 16(12), 1424–1435.

427 Hastie, T., Tibshirani, R., & Buja, A. (1994). Flexible discriminant analysis by optimal scoring.
 428 *Journal of the American Statistical Association*, 89(428), 1255–1270.

429 Huang, Y-M. (1986). *Aedes (Stegomyia) bromeliae* the yellow fever virus vector in East Africa.
 430 *Journal of Medical Entomology*, 23(2), 196–200.

431 Huang, Y-M. (1990). The subgenus *Stegomyia* of *Aedes* in the Afrotropical Region I. The Africanus
 432 group of species (Diptera: Culicidae). *Contributions of the American Entomological Institute*,
 433 26(1), 1–90.

434 Huang, Y-M. (2001). A pictorial key for the identification of the subfamilies of Culicidae, genera of
 435 Culicinae, and subgenera of *Aedes* of the Afrotropical Region (Diptera: Culicidae). *Proceedings*
 436 *of the Entomological Society of Washington*, 103, 1–53.

437 Huang, Y-M. (2004). The subgenus *Stegomyia* of *Aedes* in the Afrotropical Region with keys to the
 438 species (Diptera: Culicidae). *Zootaxa*, 700(1), 1–120.

439 Hutchinson, M.F. (1991). The application of thin plate splines to continent-wide data assimilation. In
 440 Data Assimilation Systems. J.D. Jasper (Ed.). Melbourne: Bureau of Meteorology. *BMRC*
 441 *Research Report*, 27, 104–113.

442 Kaplan, I., Dobert, M.K., Marvin, B.J., McLaughlin, J.L., & Whitaker, D.P. (1976). *Area Handbook*
 443 *for Kenya*. Foreign Area Studies (FAS). The American University, Washington D.C., U.S.A, (pp.
 444 472).

445 Lembrechts, J., Nijs, I., & Lenoir, J. (2018). Incorporating microclimate into species distribution
 446 models. *Ecography*.

447 Lounibos, L.P. (2002). Invasions by insect vectors of human disease. *Annual Review of Entomology*,
448 47, 233–266.

449 McGarigal, K., Cushman, S., Neel, M., & Ene, E. (2002). FRAGSTATS: Spatial pattern analysis
450 program for categorical maps. Computer software program produced at the University of
451 Massachusetts, Amherst. <https://www.umass.edu/landeco/research/fragstats/fragstats.html/>
452 Accessed 7 November 2018.

453 Moran, P (1950). Notes on continuous stochastic phenomena. *Biometrika*, 37, 17–23.

454 Mosquito taxonomic inventory (MTI). (2017). *Anopheles; Culex; Stegomyia*. [http://mosquito-](http://mosquito-taxonomic-inventory.info/simpletaxonomy/term/8577/)
455 [taxonomic-inventory.info/simpletaxonomy/term/8577/](http://mosquito-taxonomic-inventory.info/simpletaxonomy/term/8577/) Accessed 5 April 2017.

456 Mughini-Gras, L., Mulatti, P., Severini, F., Boccolini, D., Romi, R., Gioia, B., Khoury, C., Bianchi,
457 R., Montarsi, F., Patregnani, T., Bonfanti, L., Rezza, G., Capelli, G., & Busani, L. (2014).
458 Ecological niche modelling of potential West Nile virus vector mosquito species and their
459 geographical association with equine epizootics in Italy. *EcoHealth*, 11, 120.

460 Neteler, M., Roiz, D., Rocchini, D., Castellani, C., & Rizzoli, A. (2011). Terra and Aqua satellites
461 track tiger mosquito invasion: modeling the potential distribution of *Aedes albopictus* in north-
462 eastern Italy. *International Journal of Health Geographics*, 10 (49).

463 Paupy, C., Delatte, H., Bagny, L., Corbel, V. & Fontenille, D. (2009). *Aedes albopictus*, an arbovirus
464 vector: from the darkness to the light. *Microbes and Infection*, 11, 1177–1185.

465 Pellikka, P. K. E., Heikinheimo, V., Hietanen, J., Schäfer, E., Siljander, M. & Heiskanen, J. (2018).
466 Impact of land cover change on aboveground carbon stocks in Afri-montane landscape in Kenya.
467 *Applied Geography*, 94, 178–189.

468 Powell, J. R., & Tabachnick, W. J. (2013). History of domestication and spread of *Aedes aegypti* - A
469 Review. *Memórias Do Instituto Oswaldo Cruz*, 108(1), 11–17.

470 Pradervand, J., Dubuis, A., Pellissier, L., Guisan, A., & Randin, C. (2014). Very high resolution
471 environmental predictors in species distribution models: Moving beyond topography? *Progress*
472 *in Physical Geography*, 38, 79-96.

473 R Core Team. (2013). R: A language and environment for statistical computing. *R Foundation for*
474 *Statistical Computing*. Vienna, Austria. URL <http://www.R-project.org/>.

475 Reinert, J. F., Harbach, R. E., & Kitching, I. J. (2009). Phylogeny and classification of tribe Aedini
476 (Diptera: Culicidae). *Zoological Journal of the Linnean Society*, 157, 700–794.

477 Reiter, M., & LaPointe, D. (2007). Landscape factors influencing the spatial distribution and
478 abundance of mosquito vector *Culex quinquefasciatus* (Diptera: Culicidae) in a mixed
479 residential-agricultural community in Hawai ‘i. *Journal of Medical Entomology*, 44(5), 861–868.

Richman, R., Diallo, D., Diallo, M., Sall, A., Faye, O., Diagne, C., Dia, I., Weaver, S., Hanley, K., & Buenemann, M. (2018). Ecological niche modeling of *Aedes* mosquito vectors of chikungunya virus in southeastern Senegal. *Parasites & vectors*, 11(1), 255.

Roiz, D., Neteler, M., Castellani, C., Arnoldi, D., & Rizzoli, A. (2011). Climatic factors driving invasion of the tiger mosquito (*Aedes albopictus*) into new areas of Trentino, Northern Italy. *PLoS ONE*, 6(4), e1480.

Sallam, M., Xue, R., Pereira, R., & Koehler, P. (2016). Ecological niche modeling of mosquito vectors of West Nile virus in St. John's County, Florida, USA. *Parasites & Vectors*, 9, 371.

Sallam, M.F., Michaels, S.R., Riegel, C., Pereira, R.M., Zipperer, W., Lockaby, B.G., & Koehler, P.G. Spatio-Temporal Distribution of Vector-Host Contact (VHC) Ratios and Ecological Niche Modeling of the West Nile Virus Mosquito Vector, *Culex quinquefasciatus*, in the City of New Orleans, LA, USA. *International Journal of Environmental Research and Public Health*, 14(8), 892.

Service, M. (1991). *Mosquito ecology: Field sampling methods*. (2nd edition). 988 p. Elsevier.

Siljander, M., Clark, B., & Pellikka, P. (2011). A predictive modelling technique for human population distribution and abundance estimation using remote sensing and geospatial data in a rural mountainous area in Kenya. *International Journal of Remote Sensing*, 32(21), 5997–6023.

Thuiller, W., Georges, D., Engler, R., & Breiner, F. (2016). Package “biomod2”. Ensemble platform for species distribution modeling. Version 3.3-7. <https://cran.r-project.org/web/packages/biomod2/biomod2.pdf>/ Accessed 5 November 2016.

Thuiller, W., Lafourcade, B., Engler, R., & Araújo, M. (2009). BIOMOD - A platform for ensemble forecasting of species distributions. *Ecography*, 32, 369–373.

Tucker, C. (1979). Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment*, 8, 127–150. University of Helsinki (2003-2009). The Taita Hills. <http://www.helsinki.fi/science/taita/taitahills.html>/ Accessed 22 April 2016.

Virtanen, E. (2015). *Fine-resolution climate grids for species studies in data-poor regions*. Master's Thesis. University of Helsinki.

The World Health Organization (WHO). (2017). Vector-borne diseases. October 2017. Fact sheets. <http://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases/> Accessed 6 June 2018.

Yee, T., & Mitchell, N. (1991). Generalized additive models in plant ecology. *Journal of Vegetation Science*, 2, 587–602.

514 Appendix 1. Multicollinearity of the variables included in the models. All statistical significance
515 values (p-values), are marked as follows: statistically highly significant *** = <0.001, statistically
516 significant ** = <0.01, statistically significant * = <0.05 and ns = not statistically significant.
517

<i>Culex</i>					
Variable	Slope	NDVI	Population density	Distance to roads	Elevation
Slope		0.115ns	-0.294**	0.172ns	0.279**
NDVI	0.115ns		-0.226*	0.391***	0.557***
Population density	-0.294**	-0.226*		-0.26**	-0.313***
Distance to roads	0.172ns	0.391***	-0.26**		0.143ns
Elevation	0.279**	0.557***	-0.313***	0.143ns	
<i>Stegomyia</i>					
Variable	Mean radiation	NDVI	Population density	Distance to roads	Mean temperature
Mean radiation		-0.04ns	0.312***	-0.12ns	0.10ns
NDVI	-0.04ns		-0.226*	0.391***	-0.543***
Population density	0.313***	-0.226*		-0.26**	0.286**
Distance to roads	-0.12ns	0.391***	-0.26**		-0.12ns
Mean temperature	0.10ns	-0.543***	0.286**	-0.12ns	

518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534

535 Appendix 2. Variable importance presented in each model for *Culex* and *Stegomyia* estimates.

536

<i>Culex</i>								
Variable	GLM	GAM	GBM	CTA	ANN	MARS	RF	Maxent
Slope	0	0.147	0.078	0	0.147	0.385	0.200	0.432
Population density	0.658	0.764	0.678	1	0.773	0.702	0.429	0.513
NDVI	0.261	0.165	0.088	0	0	0	0.114	0.382
Distance to roads	0	0.179	0.039	0	0.092	0	0.064	0.453
Elevation	0	0.028	0.006	0	0.097	0	0.030	0.436
<i>Stegomyia</i>								
Variable	GLM	GAM	GBM	CTA	ANN	MARS	RF	Maxent
Mean radiation	-	0.041	0	0	0.289	-	0.059	0.176
Population density	-	0.575	0.680	0.966	1	-	0.206	0.357
NDVI	-	0.388	0	0	0	-	0.210	0.291
Distance to roads	-	0.337	0	0	0.215	-	0.087	0.028
Mean temperature	-	0	0.358	0	0.081	-	0.089	0.026

537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561

562 Appendix 3. A. SAC of *Culex* predictors. B. SAC of *Stegomyia* predictors. Red circles indicated a
563 significant p-value ($p < 0.05$) and indicate distances where the variable was autocorrelated.

564 A.

565 B.

566
567
568
569
570
571