<https://helda.helsinki.fi>

Vaginal microbiota in pregnancy : Role in induction of labor and seeding the neonate's microbiota?

Kervinen, Kaisa

2019-10

Kervinen , K , Kalliala , I , Glazer-Livson , S , Virtanen , S , Nieminen , P & Salonen , A 2019 , ' Vaginal microbiota in pregnancy : Role in induction of labor and seeding the neonate's microbiota? ' , Journal of Biosciences , vol. 44 , no. 5 , 116 . <https://doi.org/10.1007/s12038-019-9925-z>

<http://hdl.handle.net/10138/322072>

<https://doi.org/10.1007/s12038-019-9925-z>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Review

Vaginal microbiota in pregnancy: Role in induction of labor and seeding the neonate's microbiota?

KAISA KERVINEN¹, ILKKA KALLIALA^{1,2}, SIVAN GLAZER-LIVSON¹, SEPPO VIRTANEN¹,
PEKKA NIEMINEN¹ and ANNE SALONEN^{3*}

¹Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

²Department of Surgery and Cancer, Institute of Reproductive and Developmental Biology, Imperial College London, London, UK

³Human Microbiome Research Program, Department of Bacteriology and Immunology, University of Helsinki, Helsinki, Finland

*Corresponding author (Email, anne.salonen@helsinki.fi)

Published online 20 September 2019

Compared to other human microbiota, vaginal microbiota is fairly simple with low bacterial diversity and high relative abundance of *Lactobacillus* species. *Lactobacillus* dominance is even more pronounced during pregnancy. Genetic factors, such as ethnicity, along with environmental, individual and lifestyle factors all have an impact on vaginal microbiota composition. The composition of the vaginal microbiota appears to play an important role in pregnancy as recent studies have linked it to adverse obstetric outcomes such as preterm birth, a leading cause of neonatal morbidity and mortality worldwide. However, the same vaginal microbiota does not seem to cause the same response in all women, calling for future research to fully understand the complex host–microbiota interplay in normal and complicated pregnancies.

Keywords. *Lactobacillus*; parturition; pregnancy; vaginal microbiota

Abbreviations: BV, bacterial vaginosis; CST, community state type; pPROM, preterm premature rupture of membranes; PTB, preterm birth; SES, socioeconomic status; VMB, vaginal microbiota

1. Introduction

Human vaginal microbiota (VMB) is unique in its composition compared to the other human microbiotas or other species' VMB with its *Lactobacillus* dominance and low bacterial diversity (Belizário and Napolitano 2015; Human Microbiome Project Consortium 2012). The vaginal mucosa differs from other mucosae also with respect to epithelial structure, mucus composition and innate and adaptive immune mechanisms (Kumamoto and Iwasaki 2012). IgG rather than IgA takes a dominant role in maintaining hemostasis in the vagina during health and infection (Bard *et al.* 2002) as well as in parturition by neutralizing pathogens and activating the innate immune system via complement activation (van de Bovenkamp *et al.* 2016).

VMB varies among healthy women. Based on the dominant bacterium VMB can be clustered into at least five distinct community state types (CSTs) (Ravel *et al.* 2011):

those dominated by *Lactobacillus crispatus*, *L. gasseri*, *L. iners* or *L. jensenii* (CST I–III and CST V, respectively). CST IV lacks lactobacilli and consists of anaerobic bacteria, such as *Gardnerella*, *Atopobium*, *Sneathia* and *Prevotella* that are typical in women with, e.g., bacterial vaginosis (BV). Age, ethnicity, lifestyle habits, use of contraceptives, anti- and probiotics and the prevailing hormonal milieu, all impact the composition of VMB (Brooks *et al.* 2017; Macklaim *et al.* 2015; Ravel *et al.* 2011) (figure 1). Other environmental factors, including dietary habits, are known to influence the human gut microbiome potentially more than the host genetics (Rothschild *et al.* 2018), but the impact of diet on VMB is yet unknown and thus calls for future studies. The socioeconomic status (SES), e.g., education, is associated to VMB composition, but currently which particular SES factors are involved and to what extent they affect the VMB remains elusive (Ding and Schloss 2014).

Figure 1. Factors known to affect vaginal microbiota (VMB) composition during pregnancy. VMB composition during pregnancy is more stable compared to the non-pregnant state. Environmental, lifestyle and individual factors do, however, have an impact on VMB composition also during pregnancy. VMB's and host's crosstalk have an influence on the woman's and neonate's health.

Pregnancy is characterized by high circulating estrogen levels produced by the placenta (Siiteri and MacDonald 1966). Estrogen increases glycogen deposition on the vaginal epithelium, favoring proliferation and dominance of lactobacilli that metabolize the breakdown products of glycogen to lactic acid, leading to low pH (<4.5) prevailing in the healthy vagina (MacIntyre et al. 2015; O'Hanlon et al. 2013). The VMB composition undergoes some changes as the pregnancy progresses, but overall it is less variable and diverse, and even more *Lactobacillus* dominant than in the non-pregnant state (Aagaard et al. 2012; Goltsman et al. 2018; Romero et al. 2014a; Walther-Antônio et al. 2014). The stability is hypothesized to be due to more stable hormonal levels and potential changes in sexual habits during pregnancy (Walther-Antônio et al. 2014). Also, the lack of menstrual bleeding, shown to have a significant effect on the VMB stability (Gajer et al. 2012), might enhance VMB stability in pregnancy. The diversity of VMB is highest during the first trimester of pregnancy and diminishes during the second and third trimester (Kroon et al. 2018; Walther-Antônio et al. 2014), possibly due to yet rising estrogen levels with progressing gestational age. During pregnancy VMB may typically change from one *Lactobacillus* spp. dominant CST to another but rarely to mixed community, i.e. CST IV (Romero et al. 2014b). As in non-pregnant women (Gajer et al. 2012; Petrova et al. 2017), *L. crispatus* appears as the most distinctive to a healthy state that also promotes the stability of normal VMB, whereas *L.*

iners is more prone to convert to abnormal VMB (Verstraelen et al. 2009) (figure 2). *L. iners* has been shown to be more prevalent in older pregnant women (Walther-Antônio et al. 2014) as well as among pregnant African-American women compared to other ethnic backgrounds, similar to CST IV (Hyman et al. 2014; Tabatabaei et al. 2018).

Interestingly, as the pregnancy progresses towards term, VMB might become more similar to the non-pregnant state (Aagaard et al. 2012). Especially in *L. iners*-dominated communities a clear increase in taxonomic richness can be seen towards the end of pregnancy (Goltsman et al. 2018). The potential significance of such VMB changes, for example, to the duration of pregnancy remains unclear.

2. Microbiota and preterm birth

Most studies on VMB during pregnancy have focused on its role in preterm birth (PTB) (Aagaard et al. 2012; Brown et al. 2018; Callahan et al. 2017; Digiulio et al. 2015; Freitas et al. 2018; Goldenberg et al. 2008; Goltsman et al. 2018; Hyman et al. 2014; Kindinger et al. 2016; Kroon et al. 2018; Liu et al. 2016; Walther-Antônio et al. 2014). PTB is the leading cause of neonatal mortality and morbidity worldwide, causing around million deaths each year (Liu et al. 2016). Despite intensive research, the ultimate trigger behind spontaneous PTB remains unclear. Overall, infections are thought to cause up to half of spontaneous PTB cases (Goldenberg et al. 2008), classically as ascending infections, drawing more attention to VMB during the recent years.

The role of VMB in PTB is controversial and seems to vary depending on the ethnicity of the cohort studied (Callahan et al. 2017). Most probably genetic factors affect and alter host-microbiome interactions so that the same microbial community structure causes divergent responses in women with different ethnic backgrounds. High VMB diversity and low lactobacilli count as signs of vaginal dysbiosis are linked to PTB in Caucasian (Brown et al. 2018; Digiulio et al. 2015; Freitas et al. 2018; Kindinger et al. 2016; Tabatabaei et al. 2018), but not in African-American women (Callahan et al. 2017; Romero et al. 2014a). African-American and black women are at increased risk of PTB compared to white women (Goldenberg et al. 2008) and their VMB diversity during pregnancy appears as most diverse (Hyman et al. 2014; Tabatabaei et al. 2018). Somewhat surprisingly, in a recent study on mainly African-American women the diversity and richness of VMB decreased significantly over pregnancy in women with preterm birth, especially between the first and second trimesters (Stout et al. 2017). Increased community instability was also associated with PTB (Stout et al. 2017), which has been shown also in Caucasian women (Callahan et al. 2017).

L. crispatus is seen as a protective factor against PTB in all ethnicities (Callahan et al. 2017; Kindinger et al. 2017; Tabatabaei et al. 2018), whereas *L. iners* dominance in the second trimester might increase the risk of PTB at least in Caucasian women (Kindinger et al. 2017; Petricevic et al.

Figure 2. Community state type (CST) I, III and IV. CST I and III are the most common community state types during pregnancy. CST IV, a mixed community where lactobacilli are replaced with anaerobic bacteria, has been linked to adverse pregnancy outcomes such as preterm birth (PTB) in Caucasian women.

2014). Vaginal dysbiosis might be associated with premature cervical remodeling and PTB, most likely through activation of a local tissue inflammation process (Kindinger *et al.* 2016).

PTB has multiple clinical presentations and heterogeneous underlying etiologies. Inflammatory neutrophils and macrophages in the uterus, decidua, fetal membranes and cervix are activated during labor (Girardi 2018; Gomez-Lopez *et al.* 2013; Thomson *et al.* 1999; Young *et al.* 2002). Complement activation leads to the chemotactic recruitment of immune cells, microbial phagocytosis, antigen uptake by dendritic cells and their delivery to local lymph node for the generation of a sterile pro-inflammatory state leading to labor (Nadeau-Vallée *et al.* 2016; Romero *et al.* 2018). Furthermore, spontaneous PTB can present as preterm premature rupture of membranes (pPROM) or cervical dilatation. PTB has also iatrogenic causes (e.g. due to maternal or fetal complications). Many VMB studies have included all PTB cases, even iatrogenic, causing probable misinterpretation of the results. Considering pPROM's strong connection to intra-amniotic infection (Goldenberg *et al.* 2008) it conceivably has a stronger link to VMB than other clinical presentations. Thus, further studies are needed before definitive connections of VMB and PTB can be indicated.

3. Prolonged and post-term pregnancy

Prolonged (>41 weeks' gestation) and post-term pregnancies (>42 weeks' gestation) are associated with several adverse neonatal and maternal outcomes compared to term

pregnancies (Olesen *et al.* 2003). Primiparity, previous post-term pregnancy, male fetus, genetic predisposition, obesity, maternal age and ethnicity have all been linked to susceptibility for post-term pregnancy (Caughey *et al.* 2009). Genetic factors, both fetal and maternal, attribute up to 30–50% of prolonged pregnancies (Laursen *et al.* 2004; Oberg *et al.* 2013). Despite intensive research the exact mechanisms leading to prolonged or post-term pregnancy are yet unknown. VMB has been linked to PTB which calls attention to VMB's potential significance behind prolonged pregnancy as well. Until today this connection has not yet been studied and thus opens a very interesting field of research in determining the interplay between VMB and host immune system during induction of labor.

4. Microbiota and parturition

To date, only few studies exist on VMB during labor. This is surprising given the anticipated role of VMB in the induction of labor as well as in neonatal colonization. A study comparing the VMB composition at 36 gestational weeks and at the onset of labor found increased VMB diversity at the time of labor onset (Avershina *et al.* 2017). Bacteria related to CST IV were significantly overrepresented at labor compared to samples at 36 weeks. Although there was less intra- than interindividual variation, a shift in CST was seen in 18% of the study cohort, most often to CST IV.

It is tempting to speculate that the increase in VMB diversity prior to labor onset could associate with or even cause a local immune and inflammatory response that starts or enhances the cervical remodeling process. An increased ratio of L- to D-lactic acid, seen in *L. iners* dominant VMB compared to *L. crispatus* dominant, has been associated with increased expression of vaginal extracellular matrix metalloproteinase inducer (EMMPRIN) and activation of matrix metalloproteinase-8, the latter being an enzyme involved in remodeling the cervix prior to onset of labor (Witkin *et al.* 2013). Vaginal communities dominated by *L. crispatus* are also associated with lower levels of pro-inflammatory cytokines than those colonized by *L. iners* or bacterial vaginosis associated bacteria that also induced secretion of pro-inflammatory cytokines (e.g. IL-1 α , IL-1 β , and IL-8) from human vaginal cells *in vitro* (Anahtar *et al.* 2015). The potential role of VMB in activating the onset of parturition remains yet to be elucidated.

Recent high-resolution analyses of mother-infant microbiomes indicate that vaginal bacteria are detected in the neonate only transiently and play minor or no role in stable intestinal colonization (Ferretti *et al.* 2018; Sakwinska *et al.* 2017). However, the maternal VMB may still have a major impact on the neonate's early immunity as the exposure to high quantities of vaginal bacteria (up to 10⁹ bacteria per gram vaginal secretions; re-calculated from Virtanen *et al.* 2017 data (Virtanen *et al.* 2017)) conceivably activate and stimulate toll-like and other bacterial sensing innate immune receptors, e.g., in the airways. However, currently there is no data to support the transfer of maternal vaginal content to a neonate born via elective C-section nor can it be recommended due to the risk of transferring pathogenic bacteria or viruses to the neonate (Haahr *et al.* 2018).

5. Microbiota after pregnancy

After parturition the estrogen levels decline fast (Nott *et al.* 1976) causing a dramatical change in VMB composition, characterized by a reduction in *Lactobacillus* spp. and an increase in alpha-diversity irrespective of ethnicity, mode of delivery, or community structure during pregnancy (Digiulio *et al.* 2015; Huang *et al.* 2015; MacIntyre *et al.* 2015). This is accompanied with a decrease in community stability. These changes can be seen to prevail up to a year after giving birth (Digiulio *et al.* 2015). Accordingly, the VMB composition during the first trimester of pregnancy was shown to associate with pregnancy history, *L. crispatus* being more abundant in women with their first pregnancy and *L. iners* in women with prior childbirth or spontaneous or induced abortion (Nasioudis *et al.* 2017). Since BV related bacteria and a diverse VMB have been linked to adverse pregnancy outcomes, such as PTB and spontaneous abortion (Kroon *et al.* 2018), these findings could favor a longer interval between pregnancies. However, the length of this diverse community prevalence remains to be elucidated

before any recommendations concerning the time before subsequent conception can be given.

6. Summary

Understanding of VMB's composition and its interactions with the host during pregnancy has vastly expanded during the recent years. VMB is known to associate at least indirectly with preterm birth, while its role on labor onset or in prolonged pregnancies has received much less attention and should hence be addressed in the future studies. Similarly, the effect of diet and other lifestyle factors that vary greatly between different populations should be investigated for their potential contribution to VMB variation and intentional microbiota modulation to support women's health.

References

- Aagaard K, Riehle K, Ma J, Segata N, Mistretta T-A, Coarfa C, Raza S, Rosenbaum S, Van den Veyver I, Milosavljevic A, Gevers D, Huttenhower C, Petrosino J and Versalovic J 2012 A metagenomic approach to characterization of the vaginal microbiome signature in pregnancy. *PLoS One* **7** e36466
- Anahtar MN, Byrne EH, Doherty KE, Bowman BA, Yamamoto HS, Soumillon M, Padavattan N, Ismail N, *et al.* 2015 Cervicovaginal bacteria are a major modulator of host inflammatory responses in the female genital tract. *Immunity* **42** 965–976
- Avershina E, Slangsvold S, Simpson MR, Storror O, Johnsen R, Øien T and Rudi K 2017 Diversity of vaginal microbiota increases by the time of labor onset. *Sci. Rep.* **7** 17558
- Bard E, Riethmuller D, Biichlé S, Meillet D, Prétet JL, Mougin C and Seillès E 2002 Validation of a high sensitive immunoenzymatic assay to establish the origin of immunoglobulins in female genital secretions. *J. Immunoassay Immunochem.* **23** 145–162
- Belizário JE and Napolitano M 2015 Human microbiomes and their roles in dysbiosis, common diseases, and novel therapeutic approaches. *Front. Microbiol.* **6** 1050
- Brooks JP, Edwards DJ, Blithe DL, Fettweis JM, Serrano MG, Sheth NU, Strauss JF, Buck GA and Jefferson KK 2017 Effects of combined oral contraceptives, depot medroxyprogesterone acetate and the levonorgestrel-releasing intrauterine system on the vaginal microbiome. *Contraception* **95** 405–413
- Brown RG, Marchesi JR, Lee YS, Smith A, Lehne B, Kindinger LM, Terzidou V, Holmes E, Nicholson JK, Bennett PR and MacIntyre DA 2018 Vaginal dysbiosis increases risk of preterm fetal membrane rupture, neonatal sepsis and is exacerbated by erythromycin *BMC Med.* **16** 9
- Callahan BJ, DiGiulio DB, Goltsman DSA, Sun CL, Costello EK, Jeganathan P, Biggio JR, Wong RJ, Druzin ML, Shaw GM, Stevenson DK, Holmes SP and Relman DA 2017 Replication and refinement of a vaginal microbial signature of preterm birth in two racially distinct cohorts of US women. *Proc. Natl. Acad. Sci. USA* **114** 9966–9971

- Caughey AB, Stotland NE, Washington AE and Escobar GJ 2009 Who is at risk for prolonged and postterm pregnancy? *Am. J. Obstet. Gynecol.* **200** 683.e1–683.e5
- DiGiulio DB, Callahan BJ, Mcmurdie PJ, Costello EK, Lyell DJ, Robaczewska A, Sun CL, Goltsman DSA, Wong RJ, Shaw G, Stevenson DK, Holmes SP and Relman DA 2015 Temporal and spatial variation of the human microbiota during pregnancy. *Proc. Natl. Acad. Sci. USA* **112** 11060–11065
- Ding T and Schloss PD 2014 Dynamics and associations of microbial community types across the human body. *Nature* **509** 357–360
- Ferretti P, Pasolli E, Tett A, Asnicar F, Gorfer V, Fedi S, Armanini F, Truong DT, *et al.* 2018 Mother-to-Infant Microbial Transmission from Different Body Sites Shapes the Developing Infant Gut Microbiome. *Cell Host Microbe* **24** 133–145
- Freitas AC, Bocking A, Hill JE, Money DM and VOGUE Research Group the VR 2018 Increased richness and diversity of the vaginal microbiota and spontaneous preterm birth. *Microbiome* **6** 117
- Gajer P, Brotman RM, Bai G, Sakamoto J, Schutte UME, Zhong X, Koenig SSK, Fu L, Ma Z, Zhou X, Abdo Z, Forney LJ and Ravel J 2012 Temporal Dynamics of the Human Vaginal Microbiota. *Sci. Transl. Med.* **4** 132ra52
- Girardi G 2018 Complement activation, a threat to pregnancy. *Semin. Immunopathol.* **40** 103–111
- Goldenberg RL, Culhane JF, Iams JD, Romero R 2008 Epidemiology and causes of preterm birth. *Lancet* **371** 75–84
- Goltsman DSA, Sun CL, Proctor DM, DiGiulio DB, Robaczewska A, Thomas BC, Shaw GM, Stevenson DK, Holmes SP, Banfield JF and Relman DA 2018 Metagenomic analysis with strain-level resolution reveals fine-scale variation in the human pregnancy microbiome. *Genome Res.* **28** 1467–1480
- Gomez-Lopez N, Tanaka S, Zaeem Z, Metz GA and Olson DM 2013 Maternal circulating leukocytes display early chemotactic responsiveness during late gestation. *BMC Pregnancy Child-birth* **13** S8
- Haahr T, Glavind J, Axelsson P, Bistrup Fischer M, Bjurström J, Andrésdóttir G, Teilmann-Jørgensen D, Bonde U, Olsén Sørensen N, Møller M and Fuglsang J, Ovesen P, Petersen J, Stokholm J, Clausen T 2018 Vaginal seeding or vaginal microbial transfer from the mother to the caesarean-born neonate: a commentary regarding clinical management. *BJOG An Int. J. Obstet. Gynaecol.* **125** 533–536
- Huang Y-E, Wang Y, He Y, Ji Y, Wang L-P, Sheng H-F, Zhang M, Huang Q-T, Zhang D-J, Wu J-J, Zhong M and Zhou H-W 2015 Homogeneity of the vaginal microbiome at the cervix, posterior fornix, and vaginal canal in pregnant Chinese women. *Microb. Ecol.* **69** 407–414
- Human Microbiome Project Consortium 2012 Structure, function and diversity of the healthy human microbiome. *Nature* **486** 207–214
- Hyman RW, Fukushima M, Jiang H, Fung E, Rand L, Johnson B, Vo KC, Caughey AB, Hilton JF, Davis RW, Giudice LC, Chi Vo K, Caughey AB, Hilton JF, Davis RW and Giudice LC 2014 Diversity of the vaginal microbiome correlates with preterm birth. *Reprod. Sci.* **21** 32–40
- Kindinger LM, Bennett PR, Lee YS, Marchesi JR, Smith A, Cacciatore S, Holmes E, Nicholson JK, Teoh TG and Macintyre DA 2017 The interaction between vaginal microbiota, cervical length, and vaginal progesterone treatment for preterm birth risk. *Microbiome* **5** 6
- Kindinger LM, MacIntyre DA, Lee YS, Marchesi JR, Smith A, McDonald JAK, Terzidou V, Cook JR, Lees C, Israfil-Bayli F, Faiza Y, Tooze-Hobson P, Slack M, Cacciatore S, Holmes E, Nicholson JK, Teoh TG and Bennett PR 2016 Relationship between vaginal microbial dysbiosis, inflammation, and pregnancy outcomes in cervical cerclage. *Sci. Transl. Med.* **8** 350ra102
- Kroon Samuel J, Ravel J and Huston WM 2018 Cervicovaginal microbiota, women's health, and reproductive outcomes. *Fertil. Steril.* **110** 327–336
- Kumamoto Y and Iwasaki A 2012 Unique features of antiviral immune system of the vaginal mucosa. *Curr. Opin. Immunol.* **24** 411–416
- Laursen M, Bille C, Olesen AW, Hjelmberg J, Skytthe A and Christensen K 2004 Genetic influence on prolonged gestation: a population-based Danish twin study. *Am. J. Obstet. Gynecol.* **190** 489–494
- Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J, Lawn JE, Cousens S, Mathers C and Black RE 2016 Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet* **388** 3027–3035
- MacIntyre DA, Chandiramani M, Lee YS, Kindinger L, Smith A, Angelopoulos N, Lehne B, Arulkumaran S, Brown R, Teoh TG, Holmes E, Nicholson JK, Marchesi JR and Bennett PR 2015 The vaginal microbiome during pregnancy and the postpartum period in a European population. *Sci. Rep.* **5** 8988
- Macklaim JM, Clemente JC, Knight R, Gloor GB and Reid G 2015 Changes in vaginal microbiota following antimicrobial and probiotic therapy. *Microb. Ecol. Health Dis.* **26** 27799
- Nadeau-Vallée M, Obari D, Palacios J, Brien M-È, Duval C, Chemtob S and Girard S 2016 Sterile inflammation and pregnancy complications: a review. *Reproduction* **152** R277–R292
- Nasioudis D, Forney LJ, Schneider GM, Gliniewicz K, France M, Boester A, Sawai M, Scholl J and Witkin SS 2017 Influence of Pregnancy History on the Vaginal Microbiome of Pregnant Women in their First Trimester. *Sci. Rep.* **7** 10201
- Nott PN, Franklin M, Armitage C and Gelder MG 1976 Hormonal changes and mood in the puerperium. *Br. J. Psychiatry* **128** 379–383
- O'Hanlon DE, Moench TR and Cone RA 2013 Vaginal pH and microbicidal lactic acid when lactobacilli dominate the microbiota. *PLoS One* **8** e80074
- Oberg AS, Frisell T, Svensson AC and Iliadou AN 2013 Maternal and fetal genetic contributions to postterm birth: familial clustering in a population-based sample of 475,429 Swedish births. *Am. J. Epidemiol.* **177** 531–537
- Olesen AW, Westergaard JG and Olsen J 2003 Perinatal and maternal complications related to postterm delivery: a national register-based study, 1978–1993. *Am. J. Obstet. Gynecol.* **189** 222–227
- Petricevic L, Domig KJ, Nierscher FJ, Sandhofer MJ, Fidesser M, Krondorfer I, Husslein P, Kneifel W and Kiss H 2014 Characterisation of the vaginal Lactobacillus microbiota associated with preterm delivery. *Sci. Rep.* **4** 5136
- Petrova MI, Reid G, Vaneechoutte M and Lebeer S 2017 Lactobacillus iners: Friend or Foe? *Trends Microbiol.* **25** 182–191
- Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SSK, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket CO, Brotman RM,

- Davis CC, Ault K, Peralta L and Forney LJ 2011 Vaginal microbiome of reproductive-age women. *Proc. Natl. Acad. Sci. USA* **108** 4680–4687
- Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Bieda J, Chaemsathong P, Miranda J, Chaiworapongsa T and Ravel J 2014a The vaginal microbiota of pregnant women who subsequently have spontaneous preterm labor and delivery and those with a normal delivery at term. *Microbiome* **2** 1–15
- Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Nikita L, Galuppi M, Lamont RF, Chaemsathong P, Miranda J, Chaiworapongsa T and Ravel J 2014b The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome* **2** 4
- Romero R, Xu Y, Plazyo O, Chaemsathong P, Chaiworapongsa T, Unkel R, Than NG, Chiang PJ, Dong Z, Xu Z, Tarca AL, Abrahams VM, Hassan SS, Yeo L and Gomez-Lopez N 2018 A Role for the Inflammasome in Spontaneous Labor at Term. *Am. J. Reprod. Immunol.* **79** e12440
- Rothschild D, Weissbrod O, Barkan E, Kurilshikov A, Korem T, Zeevi D, Costea PI, Godneva A, et al. 2018 Environment dominates over host genetics in shaping human gut microbiota. *Nature* **555** 210–215
- Sakwinska O, Foata F, Berger B, Brüssow H, Combremont S, Mercenier A, Dogra S, Soh S-E, Yen JCKCK, Heong GYSYS, Lee YSS, Yap F, Meaney MJJ, Chong Y-S, Godfrey KMM and Holbrook JDD 2017 Does the maternal vaginal microbiota play a role in seeding the microbiota of neonatal gut and nose? *Benef. Microbes* **8** 763–778
- Siiteri PK and MacDonald PC 1966 Placental estrogen biosynthesis during human pregnancy. *J. Clin. Endocrinol. Metab.* **26** 751–761
- Stout MJ, Zhou Y, Wylie KM, Tarr PI, Macones GA and Tuuli MG 2017 Early pregnancy vaginal microbiome trends and preterm birth. *Am. J. Obstet. Gynecol.* **217** 356.e1–356.e18
- Tabatabaei N, Eren AM, Barreiro LB, Yotova V, Dumaine A, Allard C and Fraser WD 2019 Vaginal microbiome in early pregnancy and subsequent risk of spontaneous preterm birth: a case-control study. *BJOG* **126** 349–358
- Thomson AJ, Telfer JF, Young A, Campbell S, Stewart CJR, Cameron IT, Greer IA and Norman JE 1999 Leukocytes infiltrate the myometrium during human parturition: further evidence that labour is an inflammatory process. *Hum. Reprod.* **14** 229–236
- van de Bovenkamp FS, Hafkenscheid L, Rispens T and Rombouts Y 2016 The Emerging Importance of IgG Fab Glycosylation in Immunity. *J. Immunol.* **196** 1435–1441
- Verstraelen H, Verhelst R, Claeys G, De Backer E, Temmerman M and Vaneechoutte M 2009 Longitudinal analysis of the vaginal microflora in pregnancy suggests that *L. crispatus* promotes the stability of the normal vaginal microflora and that *L. gasseri* and/or *L. iners* are more conducive to the occurrence of abnormal vaginal microflora. *BMC Microbiol.* **9** 116
- Virtanen S, Kalliala I, Nieminen P and Salonen A 2017 Comparative analysis of vaginal microbiota sampling using 16S rRNA gene analysis. *PLoS One* **12** e0181477
- Walther-Antônio MRS, Jeraldo P, Berg Miller ME, Yeoman CJ, Nelson KE, Wilson BA, White BA, Chia N and Creedon DJ 2014 Pregnancy's stronghold on the vaginal microbiome. *PLoS One* **9** e98514
- Witkin SS, Mendes-Soares H, Linhares IM, Jayaram A, Ledger WJ and Forney LJ 2013 Influence of vaginal bacteria and D- and L-lactic acid isomers on vaginal extracellular matrix metalloproteinase inducer: implications for protection against upper genital tract infections. *MBio.* <https://doi.org/10.1128/mbio.00460-13>
- Young A, Thomson AJ, Ledingham M, Jordan F, Greer IA and Norman JE 2002 Immunolocalization of proinflammatory cytokines in myometrium, cervix, and fetal membranes during human parturition at term. *Biol. Reprod.* **66** 445–449