<https://helda.helsinki.fi>

Effects of temperature on growth and development of amphibian larvae across an altitudinal gradient in the Tibetan Plateau

Peng, Li Qing

2020-06

Peng , L Q , Tang , M , Liao , J H , Liang , S Y , Gan , L T , Hua , K J , Chen , Y , Li , H ,
Chen , W & Merilä , J 2020 , ' Effects of temperature on growth and development of
amphibian larvae across an altitudinal gradient in the Tibetan Plateau ' , Animal Biology , vol.
70 , no. 3 , pp. 239-250 . <https://doi.org/10.1163/15707563-20201196>

<http://hdl.handle.net/10138/322297>

<https://doi.org/10.1163/15707563-20201196>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Running title: temperature influences high-altitude tadpoles' growth

**Effects of temperature on growth and development of amphibian larvae across
an altitudinal gradient in the Tibetan Plateau**

Li Qing Peng¹, Min Tang¹, Jia Hong Liao¹, Shi Yuan Liang², Li Tao Gan¹, Ke Jun
Hua¹, Yan Chen¹, Hong Li², Wei Chen^{1*} & Juha Merilä³

¹Ecological Security and Protection Key laboratory of Sichuan Province, Mianyang
Normal University, Mianyang, 621000, China

² Jiangsu Key Laboratory for Biodiversity and Biotechnology, College of Life
Sciences, Nanjing Normal University, Nanjing 210023, China.

³Ecological Genetics Research Unit, Organismal and Evolutionary Biology Research
Programme, Faculty Biological and Environmental Sciences, University of Helsinki,
P.O. Box 65, FI-00014, Helsinki, Finland

*Corresponding author: Wei Chen (Wchen1949@163.com)(+8613778065872)

17

18 **Abstract**

19 Organisms living in extreme environments, such as amphibians inhabiting the Tibetan
20 Plateau, are faced with a magnitude of potentially strong selection pressures. With an
21 average elevation exceeding 4,500 m, the Tibetan Plateau is mainly characterized by
22 low temperatures, but little is known about the influence of this factor on the growth,
23 development, and behavior of amphibian larvae living in this environment. Using a
24 common garden experiment, we studied the influence of temperatures on the early
25 growth and development of tadpoles of the Tibetan brown frog (*Rana kukunoris*)
26 endemic to the eastern Tibetan plateau. We discovered that temperature had a
27 significant influence on early growth and development of the tadpoles from
28 high-temperature treatment, growing and developing faster than their siblings from
29 low-temperature treatment. However, high-altitude individuals grew faster than
30 low-altitude individuals at low temperatures, while the opposite was true at high
31 temperatures. These results support the temperature adaptation hypothesis as tadpoles'
32 growth and developmental rates were maximized at the temperatures experienced in
33 their native environments. These results suggest that variation in ambient temperature,
34 combined with evolutionary adaptation to temperature of local environments, is
35 probably one of the most critical environmental factors shaping altitudinal differences
36 of the growth and development of amphibian larvae at the Tibetan plateau.

37

38 **Key words**

39 Development; growth; *Rana kukunoris*; survival rate; temperature

Introduction

Life history theories predict that environment conditions can pose strong influences on evolution of life history traits of animals, including amphibians (Stearns, 1992; Morrison & Hero, 2003a; Liao et al., 2014; Liao et al., 2015; Zhong et al., 2018; Wang et al., 2019; Yu et al., 2019), and that environmental conditions experienced during early life can have lasting effects on later life characteristics of individuals (Burton & Metcalfe, 2014; Garcia et al., 2017). Although organisms such as amphibians can alleviate the negative fitness consequences of harsh environmental conditions through behavioral and physiological plasticity (Voituron et al., 2005; Muir et al., 2014a), there is growing evidence to suggest that temperature lead to evolutionary adaptations in amphibian life history traits, including the growth and development of their larvae (e.g. Wells, 1977; Morrison & Hero, 2003b). However, relatively little is known about the importance of this stressor, and its possible effects on the evolution of amphibian life histories, especially in harsh high-altitude environments.

Previous studies show that temperature has markedly affect amphibian life history traits including embryonic growth and development rates (Berven, 1982; Gollman & Gollman, 1996; Laugen et al., 2003), and body size at metamorphosis (Morrison & Hero, 2003a). The tadpoles from cold environments, between such high-altitude and -latitude localities, generally have longer larval periods because of slow growth rates and finally result in larger body sizes at metamorphosis than those from warmer areas (e.g. Berven, 1982; Howard & Wallace, 1985; Morrison & Hero,

2003a; Liao & Lu 2012; Liu et al. 2018). This variation in growth and development rates has been attributed mainly to environmental temperature in that it will control developmental and physiological processes of organisms (Gollman & Gollman, 1996). The study results of a large volume of literature (e.g. Levinton & Monahan, 1983; Hangartner et al., 2011; Miaud & Merilä, 2001, Muir et al., 2014b) show that amphibian life history traits including growth and development rates exhibit high adaptation to local environmental temperature where they live, they are typically maximized at the temperatures most commonly experienced in the native environments (cf. temperature local adaption hypothesis, Yamahira & Conover, 2002). However, there is some evidence that many amphibian species show counter-gradient variation in growth and development rates. Namely, although developing more slowly in the wild, individuals from cold climate localities grow and develop faster when reared in similar temperature with the conspecifics from warmer localities (Licht, 1975; Berven et al., 1979; Gollman & Gollman 1996; Laugen et al., 2003; Palo et al., 2003; Morey & Reznick, 2004; Richter-Boix et al., 2006; Lind et al., 2008). These findings have been explained in terms of selection favoring faster growth and development rates of tadpoles from colder regions to compensate for negative effects of temperatures and growth season lengths to growth and development in low temperature populations (Conover & Present, 1990; Conover & Schultz, 1995; Yamahira & Conover, 2002; Laugen et al., 2003).

High-altitude environments are characterized by low temperatures, low oxygen levels, and strong UV-B radiation (Blumthaler et al., 1997; Bickler & Buck, 2007;

Scheinfeldt & Tishkoff, 2010). This is true also in the case of the Tibetan plateau, which has an average elevation exceeding 4,500 m (Han et al., 2018). However, the varying geomorphological and climatic conditions of the Tibetan plateau may impact upon amphibian life history traits by selecting for divergence in growth and development rates among localities. Yet, the influence of temperature on the life history traits of amphibians' larvae are poorly understood (Morrison & Hero, 2003b; Ma et al., 2009), especially when it comes to the Tibetan plateau.

Rana kukunoris is endemic to the eastern Tibetan plateau (Xie et al., 2000), and the species is an explosive breeder with a pure capital breeding strategy (Lu et al., 2008; Chen & Lu, 2011; Chen et al., 2011; Chen et al., 2013a; Chen et al., 2018). The species is sexually size dimorphic with females being the larger sex (Chen et al., 2011; Feng et al., 2015). Female fecundity is positively related to body size (Lu et al., 2008) and males prefer to mate with larger females (Chen & Lu, 2011). The next spring after winter the species will begin breeding, so females invest some energy into gonads before winter, and their gonad weight decreases with increasing altitude due to relatively lower breeding energy investment at high altitudes (Chen et al., 2013b). the species has a wide altitudinal range with 2200 m-4400 m (Xie et al., 2000) and temperatures experienced naturally by the species range from 4° C to 32° C (Chen's personal observation). But so far, little is known about the effect of temperature on the growth, development and activity of their tadpoles.

The aim of this study was to investigate the impacts of temperature on the growth, development and activity of the tadpoles of *Rana kukunoris* to discover the

growth and developmental strategies of high-altitude tadpoles. Specifically, we investigated how temperature influence 1) development, 2) behavior, 3) size at metamorphosis, and 4) survival rate of *R. kukunoris* tadpoles from different altitudes until metamorphosis using controlled laboratory experiments.

Material and Methods

Collection and husbandry of animals

We collected freshly egg clutches of *R. kukunoris* from small ponds near Galitai (32°58'N; 103°20'E, 3800 m) and Hongxin (34°05'N; 102°45'E, 3200 m) in the Zoige county in Sichuan province and from a pond near Gahai (34°17'N; 102°17'E, 3500 m) in Gansu province in March 2018. In addition, clutches were also collected from a pond near Gaoqiang village (32°31'N; 96°25'E, 4200 m) in Yushu in Qinghai province in May 2018. Egg clutches were transferred to laboratory of Mianyang Normal University (31°27'N; 104°35'E, 467m) and maintained in white plastic containers (35 × 25 × 20 cm) supplied with aerated dechlorinated water until they reached Gosner stage 18 (Gosner, 1960). Room temperature was set to 24°C, a common temperature the species experiences naturally during development. To investigate the influences of temperature we carried out the following experiment using the tadpoles from egg clutches from difference altitudes.

Experimental treatments

To examine the effect of temperature on growth and development of the tadpoles, we

use a factorial design with three temperature conditions (16°C, 24°C, and 32°C) × four altitude sites (3200 m, 3500 m, 3800 m and 4200 m). All experiments were repeated 10 times. Temperatures were selected on the basis of the temperatures recorded in the field conditions, and they were maintained using automatic air conditioning. Eggs from different altitudes were housed separately, ten embryos randomly selected at stage 18 (Gosner, 1960) from the same clutch were transferred to 1 L plastic containers filled with 750 mL dechlorinated water. After hatched (stage 25), tadpoles were fed with goldfish feed (Foshan T and F Pet Food Co., Ltd.) twice a week water changes. All the water was changed out each time. The photoperiod was set to 12 h: 12 h (L: D).

Trials to quantify behavior

Body weight of metamorphs (tail completely absorbed; Gosner stage 46) was also measured to the nearest 0.001 g with an electronic balance, body length and the maximum jumping performance of metamorphs were measured with a plastic ruler, the nearest 0.5 mm. Frogs were placed on the plastic plate under the room temperature of 24°C and allowed to rest for 5s before inducing them to jump by pinching the urostyle with a pair of fine forceps (Wilson et al., 2000). This elicited escape responses from the individuals causing them to jump away from the observer. Each metamorph was stimulated to jump at least five times, with the jump that produced the greatest distance for given individual used as a measure of maximum jumping performance (Wilson et al., 2000). The test frogs were maintained at 24°C for at least

1 hour prior to experimental procedures.

Statistical analyses

We used Linear Mixed Models (LMM) to investigate the effect of temperatures on the development and behaviour of tadpoles and metamorphs. Development rate (the time (day) from stage 18 to stage 46)/ body length of metamorph frogs was used as the response variable. Specifically, we considered the temperature and altitude as fixed explanatory variables and container identity as a random factor. Altitude $\times$ temperature interaction was also considered to test if the effects of temperature treatments were similar in populations from different altitudes. Embryonic mortality, treated as a binomial response variable, was analyzed with a type III generalized linear model with logit-link function and binomial error structure (Räsänen et al., 2003). Post-hoc tests (using SNK test) were performed to investigate significant factors in the linear mixed models.

Before the analyses, all the variables including body length and jumping distance and development rate were log-transformed to meet the assumptions of normality and homoscedasticity. Normality of residuals and homogeneity of variances were validated through visual inspection of graphical model evaluation plots. To account for the complex covariance structure in our mixed models, we used REML estimation for random effects, and the Satterthwaite procedure to approximate degrees of freedom. Analyses were conducted using R studio and the “lmerTest” package. Data are presented as means $\pm$ standard deviation (SD).

Results

The influence of temperature treatments

Temperature had positive influence on the early developmental rate of the tadpoles (estimate = -35.437 ± 8.128 ; $df = 116$; $t = -4.360$; $P < 0.001$; Table 1; Fig. 1), and developmental rate also differed between individuals from different altitudes (estimate = -12.588 ± 3.122 ; $df = 116$; $t = -4.032$; $P < 0.001$). However, a significant interaction effect between altitude and temperature treatment (estimate = 4.075 ± 0.991 ; $df = 116$; $t = 4.113$; $P < 0.001$) suggested that the developmental rate of tadpoles from different altitudes differed depending on the temperature.

The results of post-hoc tests (SNK tests) showed that high-altitude tadpoles developed faster in coldest temperature treatment than low-altitude tadpoles, with an increasing development rate of 32.12% -35.77% between temperatures. Generally, in the higher temperatures low-altitude tadpoles developed faster compared to high-altitude tadpoles with a decreasing development rate of 34.09%-54.34% (Table 1; Fig. 1). The development pattern of the tadpoles was complex in higher temperatures, the development rate of tadpoles from 3200-m population and 3500-m population showed no significant difference, the tadpoles from 3800-m population showed faster development than those from 4200-m population (Fig. 1). Nonetheless, the total trend was the tadpoles from low-altitude populations (3200m and 3500m) developed faster than those from high-altitude populations (3800 m and 4200m; Table 1)

Metamorphs' jumping ability differed among temperature treatments (estimate =

15.161 $\pm$ 5.117; df = 107.013; t = 2.963; P = 0.004; Fig. 2), as well as among individuals from different altitudes (estimate = 5.776 $\pm$ 1.965; df = 107.013; t = 2.939; P = 0.004). However, a significant temperature treatment $\times$ altitude interaction (estimate = -1.809 $\pm$ 0.624; df = 107.013; t = -2.901; P = 0.005) revealed that the effect of temperature treatment depended on altitude of origin. Tadpoles from low altitude population showed greater improvement of jumping ability (39.52%) with increasing temperature, high altitude population showed relatively small increasing jumping ability (9%). This was not a case after controlling for the effect of body length (estimate = 0.628 $\pm$ 0.320; df = 115; t = 1.960; P = 0.052; temperature: estimate = 12.640 $\pm$ 5.241, df = 115; t = 2.412; p = 0.017; altitude: estimate = 4.671 $\pm$ 2.031, df = 115; t = 2.299; P = 0.023; interaction: estimate = -1.483 $\pm$ 0.641, df = 115; t = -2.314; P = 0.022), suggesting body size was an important factor influencing Metamorphs' jumping ability.

There was significant effect of temperature treatment (estimate = 4.016 $\pm$ 1.450; df = 107.012; t = 2.769; P = 0.007) and altitude of origin (estimate = 1.761 $\pm$ 0.557; df = 107.012; t = 3.161; P = 0.002) on body length of metamorphs. The temperature $\times$ altitude interaction was significant (estimate = -0.519 $\pm$ 0.177, df = 107.012; t = -2.937; P = 0.004, Fig. 3), showing that effect of temperature treatment differed depending on the altitude of origin. In general, low temperatures resulted in larger size in metamorphosis (9.4%-19.72%; Fig.3), but tadpoles at metamorphose from high- and low- altitude populations showed no significant difference in body size in high temperature treatment (Table1; Fig.3).

Mortality of tadpoles over the experimental period was independent of temperature treatments (estimate = -112.81 ± 84.88 ; $z = -1.329$; $P = 0.184$), altitude (estimate = -46.29 ± 33.00 ; $z = -1.403$; $P = 0.161$) and their combination (altitude $\times$ temperature interaction: estimate = -13.76 ± 10.38 ; $z = 1.326$; $P = 0.185$; Table 1).

Discussion

In this study, we found that temperature had a critical influence on the growth and development rate of Tibetan brown frog tadpoles. Specifically, our results showed that tadpoles from different altitudes responded differently to different temperature treatments: high-altitude individuals grew faster than low-altitude individuals at low temperatures, while at high temperatures the results were less clear. This suggests that high-altitude populations have adapted to life in cold temperatures as their growth rate was maximized at the low temperatures they experience in their native environment (cf. Levinton & Monahan, 1983; Jin et al., 2016; Mai et al., 2017).

Low temperatures act to reduce growth and developmental rates, and consequently, prolonged growth and developmental periods typically lead to metamorphosis at larger body size (Berven, 1982; Howard & Wallace, 1985; Morrison & Hero, 2003a, b; Kuparinen et al., 2010; Liao & Lu, 2010). Our results conform to this pattern since we also found that the body length of metamorphs was also influenced by temperature treatments: Tibetan brown frog tadpoles from low temperature treatments showed slower development rate and larger body size at metamorphosis.

Temperature is a critical environmental factor that has shaped most life history traits of amphibians (Stearns, 1992; Morrison & Hero, 2003a; Kuparinen et al., 2010; Liao et al., 2010; Liao et al., 2016). Our results showed that tadpoles from all populations developed fastest in the high temperature treatment. This is typical for amphibians (Abrams et al., 1996; Morrison & Hero, 2003b) and can be explained by the fact that high temperature increases metabolic enzyme activity, speed up the process of metabolism, increase food conversion efficiency and eventually contribute to the fast growth and development of tadpoles (Marian & Pandian, 1985). Similar pattern applies also to plants (Way & Oren, 2010), insects (Wu et al., 2015), fishes (Jerry & Thépot, 2015), and reptiles (Ewert, 1985). In our study, we also found that in cold temperature condition, tadpoles from high-altitude populations showed faster developmental rate compared to these from low-altitude populations, supporting the temperature adaptation hypothesis (Yamahira & Conover, 2002).

To sum up, we investigated the impacts of temperature on the growth, development and behaviour of Tibetan brown frog tadpoles, and found that temperature appears to play a critical role in their growth, development, but not jumping behavior. Evidence was also found for adaptation to cold temperature in high altitude populations, which exhibited faster growth and developmental rates in cold temperatures than those from lower altitudes. Finally, it should be noted that our study design was not optimal because only small clutches from each of the study populations were used to carry out the experiments. Hence, there was no among family level replication in our study, and the inferences are based on the assumption

that the randomly selected family from each of the populations is representative of the given population.

Acknowledgments

This study was funded by the Natural Sciences Foundation for Distinguished Young Scholar of Sichuan (grant number 2016JQ0038), Key Foundation of Sichuan Provincial Department of Education (grant number 18ZA0255), the National Sciences Foundation of China (grant number 31670392 and 31872216) and the Finnish Cultural Foundation (grant to PM) and the Academy of Finland (grants 129662, 134728, and 218343 to JM). All field and laboratory work were conducted under permission from Mianyang Normal University.

References

- Abrams, P.A., Leimar, O., Nylin, S. & Wiklund, C. (1996). The effect of flexible growth rates on optimal sizes and development times in a seasonal environment. *Am. Nat.*, 147, 381–395.
- Berven, K. (1982). The genetic basis of altitudinal variation in the wood frog *Rana sylvatica*. I. An experimental analysis of life history traits. *Evolution*, 36, 962–983.
- Berven, K.A., Gill, D.E. & Smith-Gill, S.J. (1979). Countergradient selection in the green frog, *Rana clamitans*. *Evolution*, 33, 609–623.
- Bickler, P.E. & Buck, L.T. (2007). Hypoxia tolerance in reptiles, amphibians, and

282 fishes: life with variable oxygen availability. *Annu. Rev. Physiol.*, 69, 145–170.

283 Blumthaler, M., Ambach, W. & Ellinger, R. (1997). Increase in solar UV radiation
284 with altitude. *J. Photochem. Photobiol. B-Biol.*, 39, 130–134.

285 Burton, T. & Metcalfe, N.B. (2014). Can environmental conditions experience in early
286 life influence future generations? *Proc. Biol. Sci.* 281, 20140311.

287 Chen, W. & Lu, X. (2011). Sex recognition and mate choice in male *Rana kukunoris*.
288 *Herpetol. J.*, 21, 141–144.

289 Chen, W., Peng, L.Q., Jiang, L.C., Pike, D.A., Friesen, C.R. & Brown, G. (2018).
290 High altitude frogs (*Rana kukunoris*) adopt a diversified bet-hedging strategy
291 in the face of environmental unpredictability. *Asian Herpetol. Res.*, 9, 43–49.

292 Chen, W., Tang, Z.H., Fan, X.G., Wang, X.Y. & Pike, D.A. (2013b). Maternal
293 investment increases with altitude in a frog on the Tibetan Plateau. *J. Evol.*
294 *Biol.*, 26, 2710–2715.

295 Chen, W., Wang, X.Y. & Fan, X.G. (2013a). Do anurans living in higher altitudes have
296 higher prehibernation energy storage? Investigations from a high-altitude frog.
297 *Herpetol. J.*, 23, 45–49.

298 Chen, W., Yu, T.L. & Lu, X. (2011). Age and body size of *Rana kukunoris*, a
299 high-elevation frog native to the Tibetan plateau. *Herpetol. J.*, 21, 149–151.

300 Conover, D.O. & Present, T.M.C. (1990). Countergradient variation in growth rate:
301 compensation for length of the growing season among Atlantic silversides
302 from different latitudes. *Oecologia*, 83, 316–324.

303 Conover, D.O. & Schultz, E.F. (1995). Phenotypic similarity and the evolutionary

304 significance of countergradient variation. *Trends Ecol. Evol.*, 10, 248–252.

305 Ewert, M.A. (1985). Embryology of turtles. Pp. 75–267 in C. Gans, F. Billett, and
306 P.F.A. Madereson, eds. *Biology of the Reptilia*. Wiley, New York.

307 Feng, X.Y., Chen, W., Hu, J.H. & Jiang, J.P. (2015). Variation and sexual dimorphism
308 of body size in the plateau brown frog along an altitudinal gradient. *Asian*
309 *Herpetol. Res.*, 6, 291–297.

310 Garcia, T.S., Urbina, J.C., Bredeweg, E.M., Ferrari, M.C.O. (2017). Embryonic
311 learning and developmental carry-over effects in an invasive anuran.
312 *Oecologia*, 184, 623–631.

313 Gollman, B. & Gollman, G. (1996). Geographic variation of larval traits in the
314 Australian frog *Geocrinia victoriana*. *Herpetologica*, 52, 181–187.

315 Gosner, K.L. (1960). A simplified table for staging anuran embryos and larvae with
316 notes on identification. *Herpetologica*, 16, 183–190.

317 Han, X.Z., Zhang, Q., Fan, L.Q., Yang, L. & Liu, Z.S. (2018). The Influence of
318 oxygen on the development of *Nanoranaparkeri* tadpoles. *Asian Herpetol.*
319 *Res.*, 9, 50–55.

320 Hangartner, S.B., Laurila, A. & Räsänen, K. (2011). Adaptive divergence of the moor
321 frog (*Rana arvalis*) along an acidification gradient. *BMC Evol. Biol.*, 11,
322 366–378.

323 Howard, J.H. & Wallace, R.L. (1985). Life history characteristics of populations of
324 the long-toed salamander (*Ambystoma macrodactylum*) from different
325 altitudes. *Am. Midl. Nat.*, 113, 361–372.

326 Jerry, D.R. & Thépot, V. (2015). The effect of temperature on the embryonic
 327 development of barramundi, the Australian strain of *Lates calcarifer* (Bloch)
 328 using current hatchery practices. *Aquaculture Rep.*, 12, 132–138.

329 Jin, L., Yang, S.N., Liao, W.B. & Lüpold, S. (2016). Altitude underlies variation in the
 330 mating system, somatic condition and investment in reproductive traits in male
 331 Asian grass frogs (*Fejervarya limnocharis*). *Behav. Ecol. Sociobiol.*, 70:
 332 1197-1208.

333 Kuparinen, A., Laugen, A.T., Laurila, A. & Merilä, J. (2010). Developmental
 334 threshold model challenged by temperature. *Evol. Ecol. Res.*, 12, 821–829.

335 Laugen, A.T., Laurila, A., Räsänen, K. & Merilä, J. (2003). Latitudinal countergradient
 336 variation in the common frog (*Rana temporaria*) development rates – evidence
 337 for local adaptation. *J. Evol. Biol.*, 16, 996–1005

338 Levinton, J.S. & Monahan, R. K. (1983). The latitudinal compensation hypothesis:
 339 Growth data and a model of latitudinal growth differentiation based upon
 340 energy budgets. II. Intraspecific comparisons between subspecies of
 341 *Ophryotrocha puerilis* (Polychaeta: Dorvilleidae). *Biol. Bull.*, 165, 699–707.

342 Liao, W.B. & Lu, X. (2010). Age structure and body size of the Chuanxi Tree Frog
 343 *Hyla annectans chuanxiensis* from two different elevations in Sichuan (China).
 344 *Zool. Anz.*, 248, 255-263.

345 Liao, W.B. & Lu, X. (2012). Adult body size = f (initial size + growth rate×age):
 346 explaining the proximate cause of Bergman's cline in a toad along altitudinal
 347 gradients. *Evol. Ecol.*, 26, 579-590.

- 348 Liao, W.B., Liu, W.C. & Merilä, J. (2015). Andrew meets Rensch: Sexual size
349 dimorphism and the inverse of Rensch's rule in Andrew's toad (*Bufo*
350 *andrewsi*). *Oecologia*, 177, 389-399.
- 351 Liao, W.B., Lu, X. & Jehle R. (2014). Altitudinal variation in reproductive investment
352 and trade-off between egg size and clutch size in the Andrew's Toad (*Bufo*
353 *andrewsi*). *J. Zool.*, 293, 84-91.
- 354 Liao, W.B., Luo, Y., Lou, S.L., Lu, D. & Jehle, R. (2016). Geographic variation in
355 life-history traits: growth season affects age structure, egg size and clutch size
356 in Andrew's toad (*Bufoandrewsi*). *Front. Zool.*, 13, 6.
- 357 Liao, W.B., Zhou, C.Q., Yang, Z.S., Hu, J.C. & Lu, X. (2010). Age, size and growth in
358 two populations of the dark-spotted frog *Rana nigromaculata* at different
359 altitudes in southwestern China. *Herpetol. J.*, 20, 77–82.
- 360 Licht, L.E. (1975). Comparative life history features of the western spotted frog, *Rana*
361 *pretiosa*, from low- and high- elevation populations. *Can. J. Zool.*, 53,
362 1254–1258.
- 363 Lind, M.I., Persbo, F. & Johansson, F. (2008). Pool desiccation and developmental
364 thresholds in the common frog, *Rana temporaria*. *P. Roy. Soc. Lond. B. Bio.*,
365 275, 1073–1080.
- 366 Liu, Q.,Feng, H.,Jin, L.,Mi, Z.P., Zhou Z. M. & Liao W.B.(2018). Latitudinal
367 variation in body size in *Fejervarya limnocharis* supports the inverse of
368 Bergmann's rule.68, 113–128.
- 369 Lu, X., Zeng, X.H., Du, B. & Nie, C. (2008). Reproductive ecology of *Rana*

370 *kukunoris* Nikolskii, 1918, a high-elevation frog native to the Tibetan Plateau.
371 *Herpetozoa*, 21, 67–77.

372 Ma, X.Y., Lu, X. & Merilä, J. (2009). Altitudinal decline of body size in a Tibetan
373 frog. *J. Zool.*, 279, 364–371.

374 Mai, C.L., Liu, Y.H., Jin, L., Mi, Z.P. & Liao, W.B. 2017. Altitudinal variation in
375 somatic condition and investment in reproductive traits in male Yunnan pond
376 frog (*Pelophylax pleuraden*). *Zool. Anz.*, 266, 189–195.

377 Marian, M.P. & Pandian, T.J. (1985). Effect of temperature on development, growth
378 and bioenergetics of the bullfrog tadpole *Rana tigrine*. *J. Therm. Biol.*, 10,
379 157–161.

380 Miaud, C. & Merilä, J. (2001). Local adaptation or environmental induction? Causes
381 of population differentiation in alpine amphibians. *Biota*, 2, 31–50.

382 Morey, S.R. & Reznick, D.N. (2004). The relationship between habitat permanence
383 and larval development in California spadefoot toads: field and laboratory
384 comparisons of developmental plasticity. *Oikos*, 104, 172–190.

385 Morrison, C. & Hero, J.M. (2003a). Geographic variation in life-history
386 characteristics of amphibians: a review. *J. Anim. Ecol.*, 72, 270–279.

387 Morrison, C. & Hero, J.M. (2003b). Altitudinal variation in growth and development
388 rates of tadpoles of *Litoria chloris* and *Litoria pearsoniana* in Southeast
389 Queensland, Australia. *J. Herpetol.*, 37, 59–64.

390 Muir, A.P., Biek, R. & Mable, B.K. (2014a). Behavioural and physiological
391 adaptations to low-temperature environments in the common frog, *Rana*

392 *temporaria*. *BMC Evol. Biol.*, 14, 110. doi:10.1186/1471-2148-14-110

393 Muir, A.P., Biek, R., Thomas, R. & Mable, B.K. (2014b). Local adaptation with high
394 gene flow: temperature parameters drive adaptation to altitude in the common
395 frog (*Rana temporaria*). *Mol. Ecol.*, 23, 561–574.

396 Palo, J.U., O’Hara, R.B., Laugen, A.T., Laurila, A., Primmer, C.R. & Merilä, J. (2003).
397 Latitudinal divergence of common frog (*Rana temporaria*) life history traits
398 by natural selection: evidence from a comparison of molecular and
399 quantitative genetic data. *Mol. Ecol.*, 12, 1963–1978.

400 Räsänen, K., Pakkala, M., Laurila, A. & Merilä, J. (2003). Does jelly envelope protect
401 the common frog *Rana temporaria* embryos from UV-B radiation?
402 *Herpetologica*, 59, 293–300.

403 Richter-Boix, A., Llorente, G.A. & Montori, A. (2006). A comparative analysis of the
404 adaptive developmental plasticity hypothesis in six Mediterranean anuran
405 species along a pond permanency gradient. *Evol. Ecol. Res.*, 8, 1139–1154.

406 Scheinfeldt, L.B. & Tishkoff, S.A. (2010). Living the high life: high-altitude
407 adaptation. *Genome Biol.*, 11, 133.

408 Stearns, S.C. (1992). The evolution of life histories. Oxford: Oxford University Press.

409 Voituron, Y., Joly, P., Eugène, M. & Barré, H. (2005). Freezing tolerance of the
410 European water frogs: the good, the bad, and the ugly. *Am. J. Physiol. Regul.*
411 *Integr. Comp. Physiol.*, 288, 1563–1570.

412 Wang, X., Huang, Y., Zhong, M., Yang, S., Yang, X., Jiang, J. & Hu, J. (2019).
413 Environmental stress shapes life-history variation in the swelled-vented frog

414 (*Feirana quadranus*). *Evol. Ecol.*, 33, 435–448.

415 Way, D.A. & Oren, R. (2010). Differential responses to changes in growth
416 temperature between trees from different functional groups and biomes: a
417 review and synthesis of data. *Tree Physiol.*, 30, 669–688.

418 Wells, K.D. (1977). The social behaviour of anuran amphibians. *Anim. Behav.*, 25,
419 666–693.

420 Wu, T.H., Shiao, S.F. & Okuyama, T. (2015). Development of insects under
421 fluctuating temperature: a review and case study. *J. Appl. Entomol.*, 139,
422 592–599.

423 Xie, F., Fei, L. & Ye, C.Y. (2000). On taxonomic status and relationships of *Rana*
424 *japonica* group, in China (Amphibia: Anura: Radidae). *Cultum. Herpetol.*
425 *Sinica.*, 8, 74–80.

426 Yamahira, K. & Conover, D.O. (2002). Intra- vs. interspecific latitudinal variation in
427 growth: adaptation to temperature or seasonality? *Ecology*, 83, 1252–1262.

428 Yu, T.L., Wang, D.L., Busam, M. & Deng, Y.H. (2019). Altitudinal variation in body
429 size in *Bufo minshanicus* supports Bergmann's rule. *Evol. Ecol.*, 33, 449–460.

430 Zhong, M.J. & Liao, W.B. (2018). A review for life-history traits variation in frogs
431 especially for anurans in China. *Asian Herpetol. Res.*, 9, 165–174.

432

433

434 **Tables**

435 **Table 1.** Mean (Standard deviation SD) developmental time (in days) of Tibetan
436 brown frog tadpoles from four different populations in three temperature treatments.
437 The developmental time refers to days in between Gosner (1960) stages 18 and 45.
438 Means with different letters (a, b, c, d) are significantly different at $P < 0.05$ (SNK
439 tests).

440

Figure legends

Figure 1 Effects of temperature on early development of Tibetan brown frog tadpoles from four different populations. Tadpoles were exposed to different temperature treatments from Gosner stage 18 onwards, and the developmental time is expressed as the days from stage 18 to stage 45. Standard deviation is presented by the bars. The legend indicates different altitudes.

Figure 2 Effects of temperature on jumping ability (distance) of Tibetan brown frog metamorphs from four different populations. Standard deviation is presented by the bars. The legend indicates different altitudes.

Figure 3 Effects of temperature on body length of Tibetan brown frog metamorphs from four different populations. Standard deviation is presented by the bars. The legend indicates different altitudes.

Table 1

		<i>N</i>	3200 m	3500 m	3800 m	4200 m
Parameter	Treatments					
Development time	16°C	10	137.70 ± 0.67 ^a	111.00 ± 0.47 ^b	97.60 ± 0.70 ^c	91.60 ± 1.17 ^d
	24°C	10	27.60 ± 0.97 ^a	23.60 ± 0.70 ^b	46.70 ± 0.67 ^c	29.10 ± 0.74 ^d
	32°C	10	23.80 ± 1.23 ^a	21.90 ± 1.29 ^a	45.00 ± 0.47 ^b	27.90 ± 0.74 ^c
Body length	16°C	10	1.28 ± 0.07 ^a	1.38 ± 0.06 ^b	1.36 ± 0.03 ^b	1.43 ± 0.08 ^b
	24°C	10	1.20 ± 0.03	1.18 ± 0.05	1.18 ± 0.04	1.22 ± 0.12
	32°C	10	1.16 ± 0.06	1.13 ± 0.05	1.18 ± 0.06	1.15 ± 0.10
Jumping distance	16°C	10	4.58 ± 0.62 ^a	3.71 ± 0.53 ^b	5.07 ± 0.64 ^c	5.43 ± 0.42 ^c
	24°C	10	6.36 ± 1.17 ^a	4.98 ± 0.56 ^b	4.86 ± 0.92 ^b	5.77 ± 1.41 ^c
	32°C	10	6.39 ± 1.10	5.43 ± 0.97	5.68 ± 0.98	5.92 ± 1.50
Survival rate	16°C	10	95.00 ± 10.80	95.00 ± 10.80	98.00 ± 6.32	100.00 ± 0.00
	24°C	10	97.00 ± 6.75	97.00 ± 6.75	99.00 ± 3.16	99.00 ± 3.16
	32°C	10	99.00 ± 3.16	98.00 ± 6.32	100.00 ± 0.00	97.00 ± 6.75

Figure 1

Figure 3