<https://helda.helsinki.fi>

The same but different : equally megadiverse but taxonomically variant spider communities along an elevational gradient

Malumbres-Olarte, Jagoba

2018-04

Malumbres-Olarte , J , Crespo , L , Cardoso , P , Szuts , T , Fannes , W , Pape , T & Scharff , N 2018 , ' The same but different : equally megadiverse but taxonomically variant spider communities along an elevational gradient ' , Acta Oecologica , vol. 88 , pp. 19-28 . <https://doi.org/10.1016/j.actao.2018.02.012>

<http://hdl.handle.net/10138/323344>

<https://doi.org/10.1016/j.actao.2018.02.012>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 The same but different: structural changes in megadiverse spider communities along an 2 elevational gradient

3
4 Jagoba Malumbres-Olarte^{1, 2}, Luís Crespo², Pedro Cardoso³, Tamás Szűts⁴, Wouter Fannes⁵,
5 Thomas Pape⁶, Nikolaj Scharff¹.

6
7 ¹Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen,
8 Copenhagen, Denmark.

9 ²Biodiversity Research Institute and Department of Evolutionary Biology, Ecology and Environmental Sciences,
10 Universitat de Barcelona, Av. Diagonal 645, 08028 Barcelona, Spain.

11 ³Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland.

12 ⁴Department of Biology, Savaria University Center, Eötvös Lóránd University, Szombathely, Hungary.

13 ⁵Department of Biology, Royal Museum for Central Africa, Tervuren, Belgium.

14 ⁶Biosystematic Section, Natural History Museum of Denmark, University of Copenhagen, Copenhagen, Denmark.

16 Abstract

17 Spatial variation in biodiversity is one of the key pieces of information for the delimitation and
18 prioritisation of protected areas. This information is especially important when the protected area
19 includes different climatic and habitat conditions, such as those along elevational gradients that are
20 matched by differences between communities. Here we test whether the megadiverse communities
21 of spiders along an elevational gradient change according to two diversity models – a monotonic
22 decrease or a hump-shaped pattern in species richness, while measuring compositional variation
23 along and within elevations, and testing the role of the habitat preference and the functional
24 structure on the changes. We sampled multiple spider communities using standardised and
25 optimised sampling in three forest types, each at a different elevation along a climatic gradient in
26 the Udzungwa Mountains, Eastern Arc Mountains, Tanzania. The number of species was similar
27 between plots and forest types, and therefore the pattern did not match either diversity model.
28 However, species composition changed significantly with a gradual change along elevations. This
29 elevational trend was not matched by changes in the occupancy or preference of habitat by spiders,
30 or by any major variation in the guild structure. Interestingly, we found greater percentages of
31 adults at higher elevations, which may be explained by the absence of strong seasonal changes and
32 by more stable levels of humidity. The similarity in habitat conditions between elevations may
33 explain that species richness and the functional structure of communities remain constant while
34 composition varies between elevations. If we aim to understand and conserve some of the most
35 diverse communities in the world, researchers and managers may need to place more attention to
36 small scale and microhabitat characteristics upon which communities depend.

37
38 *Key words:* altitudinal/ elevational gradient, climatic stability, COBRA protocol, species richness,
39 tropical forest.

41 Introduction

42 Whether one aims at testing the effects of environmental factors, historical events or biotic
43 interactions on biodiversity, or at understanding the scale at which species co-exist or interact, the
44 first step must be to quantify and characterise communities (Basset, 1996; Schaffers et al., 2008;
45 Whittaker et al., 2001). Likewise, elucidating the trophic web dynamics that regulate nutrient and
46 energy cycles requires knowledge of which species share a given space and time (Montoya et al.,

2006)□. Besides the theoretical repercussions, characterising communities by studying their composition and structure, and understanding their spatio-temporal variation is imperative to monitor and assess the effects of climatic and habitat changes on ecosystems and biodiversity (Barnosky et al., 2012; Sala, 2000), and to develop conservation priorities and policies (Pereira et al., 2013).

Elevational gradients have increasingly been seen as a powerful model system to disentangle the relative effects of environmental factors on biodiversity because this allows for many replicates, enables experiments and facilitates data collection, and avoids covariation between drivers (Körner, 2007; Nogués-Bravo et al., 2008; Sanders and Rahbek, 2012). Gradients in temperature and humidity along elevations are usually matched by changes in species abundance and composition as well as in functional or traits community structure (Fitzpatrick et al., 2013; Graham et al., 2014; Sundqvist et al., 2013). With regard to species richness along elevational gradients, a number of diversity models have been put forward (Graham et al., 2014), those most commonly referred to being of two kinds: models that predict a monotonic decrease in the number of species with altitude, and models that expect a hump-shaped pattern with a peak in the middle of the gradient (Nogués-Bravo et al., 2008; Rahbek, 1997). Both model types can be based on the richness-productivity hypothesis (Grytnes, 2003; MacArthur .R, 1965; Rosenzweig, 1971; Wright, 1983). However, the former may also derive from hypothesised positive relationships between species richness and area or species distribution (Rapoport's rule) (Sanders, 2002; Stevens, 1992; Willig et al., 2003), whereas the latter include Mid-Domain-effect models (Colwell and Lees, 2000; Rahbek, 1997).

The Eastern Arc Mountains (EAM) provide an ideal system to study diversity patterns, with elevational gradients from 300 to 2400 m.a.s.l. The EAM are classified as one of the World's Biodiversity Hotspots (Myers et al., 2000) and their forests are considered some of the oldest and most stable of the African continent (Loader et al., 2014), forming an 'inland archipelago'. The limited research that has been conducted in the EAM, mainly in the Udzungwa Mountains, Uluguru Mountains, and the East and West Usambara Mountains, has revealed elevated levels of biodiversity (Nyundo, 2002; Scharff, 1992; Sørensen, 2004; Sørensen et al., 2002) and in particular a remarkable endemism compared to the surrounding savanna and lowland forest. However, no comprehensive studies of the arthropod communities have been conducted yet.

Spiders are one of the most diverse groups of organisms both taxonomically and ecologically. Over 46,000 species and around 4000 genera have been described worldwide (Natural History Museum of Bern, 2017) and many thousands more await discovery. The myriad of habitats and feeding adaptations of spiders, and their role as one of the dominant groups of arthropod predators in terrestrial ecosystems (Marc et al., 1999), make them indicators of changes in other arthropod communities and of habitat disturbance (Cardoso et al., 2010; Malumbres-Olarte et al., 2013; Moretti et al., 2002; Romero and Harwood, 2010; Wise, 1993).

Here we present the first characterisation of megadiverse spider communities along elevational gradients based on optimised and standardised sampling. Our aims are: 1) to test which elevational diversity model matches best the spider species richness in the Udzungwa Mountains; 2) to measure the variation in taxonomic structure among communities within and between elevations; and 3) to test the role of functional community structure on diversity patterns along elevations.

88

89 **2. Material and methods**

90 *2.1. Study area*

91 Our study area is the Udzungwa Mountains (7.82°S, 36.70°E), Tanzania, which are recognised as a
92 priority conservation area for mammals and birds (Dinesen et al., 2001) as well as for plants (Lovett
93 et al., 1988; Lovett and Thomas, 1986)□. The Udzungwa Mountains are located in the southern part
94 of the Eastern Arc Mountains and their forests are believed to have endured through millions of

years due to long-term climatic stability (Lovett, 1993)□. Our plots lie on the eastern slopes of the Udzungwa Mountains National Park (UMNP; 12 plots) and in the Uzungwa Scarp Forest Reserve to the south (three plots) (Fig. 1, Table 1).

Due to the climatic influence of the Indian Ocean, the eastern slopes of the Udzungwa Mountains experience a high average annual rainfall of 2000 mm (Mumbi et al., 2008)□, with a heavy rainy season between March–May and a lighter rainy season between November–February (Lovett, 1996)□. There is a gradient in forest type from deciduous miombo woodland in the lowlands (300 m a.s.l.) to evergreen montane rainforest just below the highest peaks (2400 m a.s.l.), which are covered by a mosaic of bamboo and *Hagenia abyssinica* (Bruce) J.F. Gmel. woodlands (Lovett et al., 2006; Rovero et al., 2017)□. The frequent mist in the highest parts of the mountains results in high levels of precipitation through condensation and more humid conditions during the dry season (June–October).

Sampling

We set up 15 plots, five plots at each of the three target elevations (700, 1000 and 1500 m a.s.l.) (Table 1), which correspond to three different forest types (lowland forest, submontane forest and montane forest, respectively) (Lovett, 1999, 1993)□. Horizontal distances between the first and the remaining plots of same elevation were 0.1, 1, 20 and 175 km. Each plot was a 50 m x 50 m square (0.25 hectare), within which we applied the COBRA-TF sampling protocol for spiders (Malumbres-Olarte et al., 2017)□ in October–November 2014. The COBRA protocols (Conservation Oriented Biodiversity Rapid Assessment) combine samples using different sampling methods to obtain the largest possible number of species for a given amount of effort (hence optimised) (Cardoso, 2009)□. More specifically, the COBRA-TF protocol (for tropical forest spiders) includes samples from different vegetation strata/ microhabitats of a tropical forest (see Malumbres-Olarte 2017 for a full description) and may provide a reliable sample of the diversity and structure of the community at a given location (Cardoso, 2009)□.

2.2. Data analyses

We assessed the thoroughness of our sampling by evaluating rarefied species accumulation curves visually and calculating the sampling completeness using the Chao 1 species estimator (Scharff et al., 2003)□. We analysed the species diversity in the spider communities by calculating observed and estimated numbers of species (Jackknife 1 and 2, Chao 1 and 2 and ACE estimators), the percentage of adults, and the number of individuals and species of each predatory guild (as defined in Cardoso et al., 2011)□ for each plot and elevation. We applied generalised linear models, and ran ANOVAs and linear mixed-effects analyses (with intercepts for the variable that represents the horizontal distance from the first plot as the random effect) to test the difference in these variables between elevations.

To examine the similarity in species composition and relative abundance between spider communities we generated a dissimilarity matrix based on the Bray-Curtis index (Legendre and Legendre, 1998)□ and created an ordination through non-metric multidimensional scaling (NMDS) (McCune and Grace, 2002)□. We tested for differences between the 15 communities using ANOSIM (Bray-Curtis index, 999 permutations) and for changes between communities (beta diversity) at the same elevation calculating the Jaccard and Bray-Curtis averaged pair-wise dissimilarity measures. Finally, we compared the species abundance distribution (SAD) curves of the communities and the alpha parameter values of the Gambin model (Matthews et al., 2014; Ugland et al., 2007)□. We based the Gambin model on species abundances rarefied (1000 permutations) by the minimum number of adult individuals per plot. Before these three analyses, we applied the Hellinger transformation to equalise species weights (Legendre and Gallagher, 2001)□.

142 We handled and analysed all the data in R3.2.3. (R Development Core Team, 2017) using various
143 packages including *vegan* (Oksanen et al., 2017) and *BAT* (Cardoso et al., 2015).

144 3. Results

145 The sampling yielded 40,613 individuals, of which 17,191 (42.3%) were adults. We observed 631
146 species or morphospecies belonging to 54 families and estimated between 784–866 species across
147 all the plots (Fig. 2a). The number of species per plot (alpha diversity) varied between 108–143, the
148 total number of species at each elevation was similar as well as the beta diversity among the
149 communities at each elevation (Jaccard and Bray-Curtis indices) (Table 2). We obtained a sampling
150 completeness of 79% for all plots and values between 59–83% for individual plots (Table 1). The
151 species accumulation curves changed with elevation, with lower plots providing steeper curves (Fig.
152 2b, c). An ANOVA on the final slope of the curves showed significant differences between
153 elevations ($F_{2,12}=8.81$, $p<0.01$).

154 Samples contained significantly higher percentages of adults at higher elevations (binomial [logit]
155 distribution model, $z_{\text{mid}}=12.65$, $z_{\text{top}}=18.96$, $p<0.001$) (Fig. 3b) but the observed (Fig. 3c) and
156 estimated (Fig. 3d) numbers of species were similar in all the three elevations. The number of
157 species represented by one or two specimens (here referred to as ‘rare species’) decreased with
158 elevation (Fig. 3e).

159 The NMDS on abundance data showed a clear separation of the communities according to their
160 elevation, with which the first axis was highly correlated (0.96). (Fig. 4). The ANOSIM confirmed
161 significant differences between plots at the three elevations (Fig. 5). The shape of the SAD curves
162 and the values of the alpha parameter provided by the Gambin model showed that the relative
163 abundances of species in the low and mid elevation communities were similar (Fig. 6a, b). Some of
164 the high elevation communities had lightly more even SAD curves and higher alpha values.

165 As for the microhabitat allocation, a greater percentage of individuals was present in the
166 understorey vegetation at high elevations (Fig. 7a). There were more individuals and species in the
167 understorey than on the ground and in the herbaceous vegetation at all elevations (Fig. 7b). The
168 percentages of species belonging to the predatory guilds (Cardoso2011) were similar (Fig. 8a) with
169 one exceptions: there were significantly more species of sheet web weavers at high elevation
170 ($t=4.67$, $p<0.01$). The percentages of individuals of different guilds varied slightly between
171 elevations (Fig. 8b). In the mid elevation plots there were proportionally more specialist spiders
172 ($p<0.01$) and fewer sheet web weavers ($p=0.02$), and there were fewer sensing web weavers in the
173 high elevation communities ($p=0.02$) (Fig. 8b).

174

175 4. Discussion

176 This study is the largest study of spider communities so far, in terms of observed species richness,
177 habitat coverage and number of communities. Our species richness values surpass those of previous
178 studies in other tropical areas (Baldissera et al., 2012; Coddington et al., 2009; Floren and
179 Deeleman-Reinhold, 2005; Pinkus-Rendon et al., 2006; Pinto-Leite and Rocha, 2012), with the only
180 exception of the survey conducted in Peru, where 1140 morphospecies were observed (Silva-Davila
181 and Coddington, 1996). There are likely to be other spider communities that are even more species-
182 rich, perhaps in the forests of the neotropics, and the sampling design and methodology used in this
183 study can certainly help discover, survey and analyse these. The relatively high sampling
184 completeness for all plots at all three elevations testify to the thoroughness of our sampling, and
185 therefore the adequate characterisation of the spider communities (Table 2).

186 4.1. Elevational diversity model

187 Contrary to our expectations of elevational changes in diversity, the observed and estimated
188 numbers of species in each plot were very similar at the three elevations (Fig. 3a, b, Table 2), and

189 therefore the diversity patterns did not match any of the elevational models. Indeed, communities
190 have been found to track gradients along elevations (Graham et al., 2009; Machac et al., 2011;
191 Peters et al., 2016), and several possible explanations have been put forward for changes in the
192 number of species, including range size (Stevens, 1992), productivity (Hutchinson, 1959),
193 environmental filtering (Graham et al., 2009; Maglianesi et al., 2015), geographic constraints
194 (Colwell and Lees, 2000) and competition (Graham et al., 2009)□. More specifically, in the
195 Udzungwa Mountains the number of species of plants, rodents and molluscs have been found to
196 increase with elevation in a manner that fits with the mid-domain effect (Lovett et al., 2006; Stanley
197 and Hutterer, 2007; Tattersfield et al., 2006). However, our data did not show such patterns and the
198 differences among communities resided in their species composition (Figs. 4 and 5) and, to a certain
199 point, relative species abundances (Fig. 6a,b).

200 The simplest explanation for the similarity in the number of species may be that the differences in
201 climatic and habitat conditions along the modest elevation range (700-1500 m) are too small to
202 result in significant differences in the number of spider species collected with the applied sampling
203 protocol. However, we found differences of up to 5° C and 10% of humidity between the low and
204 high elevation plots, and we propose another possible explanation based on habitat availability
205 (Greenstone, 1984; Malumbres-Olarte et al., 2013; Riechert and Gillespie, 1986)□ and
206 environmental filtering (Chatzaki et al., 2005; Foord and Dippenaar-Schoeman, 2016). The habitat
207 in the forests of the Udzungwa Mountains has been considered to be homogeneous between 300-
208 1850 m (Romdal and Rahbek, 2009)□, and yet, different studies have found elevational changes in
209 species composition (Poynton et al., 2007; Romdal and Rahbek, 2009). If the number and structure
210 of available habitats for spiders are also similar at different elevations, the number of species and
211 the functional types that each elevation can contain may be similar. However, due to differences in
212 climatic adaptations between species (environmental filtering), distinct spider communities may be
213 found at different elevations. Considering the importance of three-dimensional habitat structure for
214 spiders, testing our hypotheses may require studies that incorporate detailed data on space, habitat
215 structure and climatic conditions.

216 We found that the percentage of adult spiders was greater at higher elevations, which may be the
217 result of sampling at the end of the dry season, when forests at low elevations are considerably
218 drier. Adults are individuals that have reached the reproductive stage and may require more prey
219 (especially females) for themselves and their growing gonads. Therefore, the percentage of adults of
220 most species may be higher when climatic conditions are most favourable and more prey is
221 available (Cardoso et al., 2007; Gasnier et al., 2002). This ‘favourable’ time-window is often when
222 the most limiting resource is most abundant. For instance, late spring-early summer may be the time
223 with the highest percentages of adults in Mediterranean ecosystems (Cardoso et al., 2007) because
224 combination of temperature and rain/humidity conditions is optimal for producing offspring,
225 whereas it may be in late summer in subalpine areas where water is never scarce and when the
226 average daily temperature is the highest (Malumbres-Olarte, 2011)□. However, in the absence of
227 strong seasonal changes there may not be any ‘favourable time-windows’, and the percentage of
228 adults may be stable throughout the year. This may be the case for the higher elevation plots, where
229 levels of humidity are higher (pers. obs.) and may be less influenced by dry and wet seasonal
230 cycles. To test whether this hypothesis is true, future observational studies should look for
231 associations or correlations between seasonal variability in climatic conditions, breeding seasonality
232 and variability in percentage of adults. It is also possible that the greater number of adults may be
233 the result of more intense and efficient sampling at high elevations, which would also explain the
234 lower number of rare species collected in high plots (Coddington et al., 2009)□.

235 4.2. *Variation in taxonomic and functional structures*

236 Although the alpha and beta diversity, and the species richness at each elevation remain the same at
237 all three elevations, the species composition does change. The latter could be accredited to the

changes in habitat composition and structure, and the associated adaptations and functional roles of the species. However, we did not detect any obvious trends in habitat preference or guild structure (Figs. 6, 7). It is possible that the (homogeneity of) local-scale characteristics, such as habitat structure or microclimatic conditions, explain the similarities in functional structure in all the studied communities. In connection with what we pointed out above, more data on forest microhabitats and the morphological, physiological or behavioural adaptations of the species associated with them might serve to explain our findings.

4.3. *Repercussions for conservation*

Our findings support the notion that safeguarding the biotic communities of the Udzungwa Mountains, and probably of other parts of the Eastern Arc Mountains, requires protecting areas with different ecosystems and habitats. This is of utmost importance in the Udzungwa Mountains given the current pace of vegetation degradation and fragmentation (Barelli et al., 2015) in unprotected or semi-protected areas outside the National Park, such as the Kilombero Nature Reserve and Uzungwa Scarp Forest Reserve (Hegerl et al., 2017; Rovero et al., 2015). Unlike the areas within the National Park, the natural vegetation of the surrounding lands have become severely degraded in recent decades (Brink et al., 2016), so legal protection may be necessary to conserve native vegetation, and thereby native arthropod communities associated with it. The negative effects of human disturbance on mammals have been used to point out the urgent need for better protection of the EAM forests (Rovero et al., 2017). Here we make the point that the spatial distribution of biodiversity of less iconic but more diverse taxa should also be taken into consideration, so that protected areas includes a variety of climatic conditions and habitat types that allow their continued survival.

Contribution of the authors

JMO, NS, TP and PC conceived and designed the study and the sampling. JMO, LC, PC and TS collected the samples, JMO, LC and WF generated the data from the samples, and JMO analysed the data. NS and TP provided the logistical and financial support from grants (see acknowledgements). JMO, NS, PC and TP wrote the paper and all authors approved it.

Compliance with ethical standards

This study does not infringe any bio-ethical principles.

Conflict of interest

The authors had no conflict of interest.

Acknowledgements

NS and JM- O acknowledge the Danish National Research Foundation for the funding (grant no. DNRF96) provided to the Center for Macroecology, Evolution and Climate. NS and TP also acknowledge the support by the Carlsberg Foundation (project 2012_01_0504). We thank Tanzania National Parks (TANAPA), the Tanzania Commission for Science and Technology (COSTECH) and the Tanzania Wildlife Research Institute (TAWIRI) for providing permits to do research in the Udzungwa Mountains National Park. Richard M. L. Laizzer and Aloyce Mwakisoma provided invaluable assistance in the field, and the Udzungwa Ecological Monitoring Centre (UEMC) and its staff provided logistical support. We also thank Bjørn Hermansen, Natural History Museum of Denmark, for his help with GIS data.

284 **References**

- 285 Baldissera, R., Rodrigues, E.N.L., Hartz, S.M., 2012. Metacommunity composition of web-spiders
286 in a fragmented neotropical forest: relative importance of environmental and spatial effects.
287 PLOS ONE 7, e48099. doi:10.1371/journal.pone.0048099
- 288 Barelli, C., Albanese, D., Donati, C., Pindo, M., Dallago, C., Rovero, F., Cavalieri, D., Michael
289 Tuohy, K., Christine Hauffe, H., De Filippo, C., 2015. Habitat fragmentation is associated to
290 gut microbiota diversity of an endangered primate: implications for conservation. Sci. Rep. 5,
291 14862. doi:10.1038/srep14862
- 292 Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M., Getz, W.M.,
293 Harte, J., Hastings, A., Marquet, P.A., Martinez, N.D., Mooers, A., Roopnarine, P., Vermeij,
294 G., Williams, J.W., Gillespie, R., Kitzes, J., Marshall, C., Matzke, N., Mindell, D.P., Revilla,
295 E., Smith, A.B., 2012. Approaching a state shift in Earth's biosphere. Nature 486, 52–58.
296 doi:10.1038/nature11018
- 297 Basset, Y., 1996. Local communities of arboreal herbivores in Papua New Guinea: predictors of
298 insect variables. Ecology 77, 1906–1919. doi:10.2307/2265793
- 299 Brink, A., Martínez-López, J., Szantoi, Z., Moreno-Atencia, P., Lupi, A., Bastin, L., Dubois, G.,
300 2016. Indicators for assessing habitat values and pressures for protected areas — an integrated
301 habitat and land cover change approach for the Udzungwa Mountains National Park in
302 Tanzania. Remote Sens. 8, 862. doi:10.3390/rs8100862
- 303 Cardoso, P., 2009. Standardization and optimization of arthropod inventories — the case of Iberian
304 spiders. Biodivers. Conserv. 18, 3949–3962. doi:10.1007/s10531-009-9690-7
- 305 Cardoso, P., Arnedo, M.A., Triantis, K.A., Borges, P.A. V., 2010. Drivers of diversity in
306 Macaronesian spiders and the role of species extinctions. J. Biogeogr. 37, 1034–1046.
307 doi:10.1111/j.1365-2699.2009.02264.x
- 308 Cardoso, P., Pekár, S., Jocqué, R., Coddington, J.A., 2011. Global patterns of guild composition
309 and functional diversity of spiders. PLOS ONE 6, e21710. doi:10.1371/journal.pone.0021710
- 310 Cardoso, P., Rigal, F., Carvalho, J.C., 2015. BAT – Biodiversity Assessment Tools, an R package
311 for the measurement and estimation of alpha and beta taxon, phylogenetic and functional
312 diversity. Methods Ecol. Evol. 6, 232–236. doi:10.1111/2041-210X.12310
- 313 Cardoso, P., Silva, I., De Oliveira, N.G., Serrano, a. R.M., 2007. Seasonality of spiders (Araneae)
314 in Mediterranean ecosystems and its implications in the optimum sampling period. Ecol.
315 Entomol. 32, 516–526. doi:10.1111/j.1365-2311.2007.00894.x
- 316 Chatzaki, M., Lymberakis, P., Markakis, G., Mylonas, M., 2005. The distribution of ground spiders
317 (Araneae, Gnaphosidae) along the altitudinal gradient of Crete, Greece: Species richness,
318 activity and altitudinal range. J. Biogeogr. 32, 813–831. doi:10.1111/j.1365-
319 2699.2004.01189.x

320 Coddington, J.A., Agnarsson, I., Miller, J.A., Kuntner, M., Hormiga, G., 2009. Undersampling bias:
 321 the null hypothesis for singleton species in tropical arthropod surveys. *J. Anim. Ecol.* 78, 573–
 322 584. doi:10.1111/j.1365-2656.2009.01525.x

323 Colwell, R., Lees, D., 2000. The mid-domain effect: geometric constraints on the geography of
 324 species richness. *Trends Ecol. Evol.* 15, 70–76. doi:10.1016/S0169-5347(99)01767-X

325 Dinesen, L., Lehmberg, T., Rahner, M.C., Fjeldså, J., 2001. Conservation priorities for the forests
 326 of the Udzungwa Mountains, Tanzania, based on primates, duikers and birds. *Biol. Conserv.*
 327 99, 223–236. doi:10.1016/S0006-3207(00)00218-4

328 Fitzpatrick, M., Sanders, N., Normand, S., Svenning, J.-C., Ferrier, S., Gove, A., Dunn, R.R., 2013.
 329 Environmental and historical imprints on beta diversity: insights from variation in rates of
 330 species turnover along gradients. *Proc. R. Soc. B Biol. Sci.* 280. doi:10.1098/rspb.2013.1201

331 Floren, A., Deeleman-Reinhold, C., 2005. Diversity of arboreal spiders in primary and disturbed
 332 tropical forests. *J. Arachnol.* 33, 323–333. doi:10.1636/05-22.1

333 Foord, S.H., Dippenaar-Schoeman, A.S., 2016. The effect of elevation and time on mountain spider
 334 diversity: a view of two aspects in the Cederberg mountains of South Africa. *J. Biogeogr.* 43,
 335 2354–2365. doi:10.1111/jbi.12817

336 Gasnier, T.R., de Azevedo, C.S., Torres-Sanchez, M.P., Höfer, H., 2002. Adult size of eight hunting
 337 spider species in Central Amazonia: temporal variations and sexual dimorphisms. *J. Arachnol.*
 338 30, 146. doi:10.1636/0161-8202(2002)030[0146:ASOEHS]2.0.CO;2

339 Graham, C.H., Carnaval, A.C., Cadena, C.D., Zamudio, K.R., Roberts, T.E., Parra, J.L., McCain,
 340 C.M., Bowie, R.C.K., Moritz, C., Baines, S.B., Schneider, C.J., Vanderwal, J., Rahbek, C.,
 341 Kozak, K.H., Sanders, N.J., 2014. The origin and maintenance of montane diversity:
 342 integrating evolutionary and ecological processes. *Ecography (Cop.)*. 37, 711–719.
 343 doi:10.1111/ecog.00578

344 Graham, C.H., Parra, J.L., Rahbek, C., McGuire, J.A., 2009. Phylogenetic structure in tropical
 345 hummingbird communities. *Proc. Natl. Acad. Sci. U. S. A.* 106 Suppl, 19673–19678.
 346 doi:10.1073/pnas.0901649106

347 Greenstone, M.H., 1984. Determinants of web spider species-diversity – vegetation structural
 348 diversity vs prey availability. *Oecologia* 62, 299–304. doi:10.1007/BF00384260

349 Grytnes, J.A., 2003. Species-richness patterns of vascular plants along seven altitudinal transects in
 350 Norway. *Ecography (Cop.)*. 26, 291–300. doi:10.1034/j.1600-0587.2003.03358.x

351 Hegerl, C., Burgess, N.D., Nielsen, M.R., Martin, E., Ciolli, M., Rovero, F., 2017. Using camera
 352 trap data to assess the impact of bushmeat hunting on forest mammals in Tanzania. *Oryx* 51,
 353 87–97. doi:10.1017/S0030605315000836

354 Hutchinson, G.E., 1959. Homage to Santa Rosalia or why are there so many kinds of animals? *Am.*
 355 *Nat.* 93, 145–159. doi:10.1086/282070

- 356 Körner, C., 2007. The use of “altitude” in ecological research. *Trends Ecol. Evol.* 22, 569–74.
357 doi:10.1016/j.tree.2007.09.006
- 358 Legendre, P., Gallagher, E.D., 2001. Ecologically meaningful transformations for ordination of
359 species data. *Oecologia* 129, 271–280.
- 360 Legendre, P., Legendre, L., 1998. *Numerical ecology, Developments in environmental modelling.*
361 Elsevier, Amsterdam.
- 362 Loader, S.P., Ceccarelli, F.S., Menegon, M., Howell, K.M., Kassahun, R., Mengistu, A.A., Saber,
363 S.A., Gebresenbet, F., S, R. De, Gower, D.J., 2014. Persistence and stability of Eastern
364 Afromontane forests: evidence from brevipitid frogs. *J. Biogeogr.* 41, 1781–1792.
365 doi:10.1111/jbi.12331
- 366 Lovett, J.C., 1999. Tanzanian forest tree plot diversity and elevation. *J. Trop. Ecol.* 15, 689–694.
367 doi:10.1017/S0266467499001108
- 368 Lovett, J.C., 1996. Elevational and latitudinal changes in tree associations and diversity in the
369 Eastern Arc mountains of Tanzania. *J. Trop. Ecol.* 12, 629–650.
370 doi:10.1017/S0266467400009846
- 371 Lovett, J.C., 1993. Eastern Arc moist forest flora, in: Lovett, J.C., Wasser, S.K. (Eds.),
372 *Biogeography and Ecology of the Rain Forests of Eastern Africa.* pp. 33–55.
- 373 Lovett, J.C., Bridson, D.M., Thomas, D.W., 1988. A preliminary list of the moist forest angiosperm
374 flora of Mwanihana Forest Reserve, Tanzania. *Ann. Missouri Bot. Gard.* 75, 874–885.
375 doi:10.2307/2399376
- 376 Lovett, J.C., Marshall, A.R., Carr, J., 2006. Changes in tropical forest vegetation along an
377 altitudinal gradient in the Udzungwa Mountains National Park, Tanzania. *Afr. J. Ecol.* 44,
378 478–490. doi:10.1111/j.1365-2028.2006.00660.x
- 379 Lovett, J.C., Thomas, D.W., 1986. The ecology of pteridophytes in the Mwanihana Forest Reserve,
380 Tanzania. *Fern Gaz.* 13, 103–107.
- 381 MacArthur, R., 1965. Patterns of Species Diversity. *Biol. Rev. Camb. Philos. Soc.* 40, 510–533.
382 doi:10.1111/j.1469-185X.1965.tb00815.x
- 383 Machac, A., Janda, M., Dunn, R.R., Sanders, N.J., 2011. Elevational gradients in phylogenetic
384 structure of ant communities reveal the interplay of biotic and abiotic constraints on diversity.
385 *Ecography (Cop.)*. 34, 364–371. doi:10.1111/j.1600-0587.2010.06629.x
- 386 Maglianesi, M. a., Blüthgen, N., Böhning-Gaese, K., Schleuning, M., 2015. Functional structure
387 and specialization in three tropical plant–hummingbird interaction networks across an
388 elevational gradient in Costa Rica. *Ecography (Cop.)*. 38, 1119–1128. doi:10.1111/ecog.01538
- 389 Malumbres-Olarte, J., 2011. Spider diversity and ecology in native tussock grasslands of the South
390 Island, New Zealand. Lincoln University, Christchurch.

391 Malumbres-Olarte, J., Scharff, N., Pape, T., Coddington, J.A., Cardoso, P., 2017. Gauging
 392 megadiversity with optimized and standardized sampling protocols: a case for tropical forest
 393 spiders. *Ecol. Evol.* 7, 494–506. doi:10.1002/ece3.2626

394 Malumbres-Olarte, J., Vink, C.J., Ross, J.G., Cruickshank, R.H., Paterson, A.M., 2013. The role of
 395 habitat complexity on spider communities in native alpine grasslands of New Zealand. *Insect*
 396 *Conserv. Divers.* 6, 124–134. doi:10.1111/j.1752-4598.2012.00195.x

397 Marc, P., Canard, A., Ysnel, F., 1999. Spiders (Araneae) useful for pest limitation and
 398 bioindication. *Agric. Ecosyst. Environ.* 74, 229–273. doi:10.1016/S0167-8809(99)00038-9

399 Matthews, T.J., Borregaard, M.K., Ugland, K.I., Borges, P.A. V, Rigal, F., Cardoso, P., Whittaker,
 400 R.J., 2014. The gambin model provides a superior fit to species abundance distributions with a
 401 single free parameter: evidence, implementation and interpretation. *Ecography (Cop.)*. 37,
 402 1002–1011. doi:10.1111/ecog.00861

403 McCune, B., Grace, J.B., 2002. Analysis of ecological communities, Analysis of ecological
 404 communities. MjM Software Design, Gleneden Beach, Oregon.

405 Montoya, J.M., Pimm, S.L., Solé, R. V., 2006. Ecological networks and their fragility. *Nature* 442,
 406 259–264. doi:10.1038/nature04927

407 Moretti, M., Conedera, M., Duelli, P., Edwards, P.J., 2002. The effects of wildfire on ground-active
 408 spiders in deciduous forests on the Swiss southern slope of the Alps. *J. Appl. Ecol.* 39, 321–
 409 336. doi:10.1046/j.1365-2664.2002.00701.x

410 Mumbi, C.T., Marchant, R., Hooghiemstra, H., Wooller, M.J., 2008. Late Quaternary vegetation
 411 reconstruction from the Eastern Arc Mountains, Tanzania. *Quat. Res.* 69, 326–341.
 412 doi:10.1016/j.yqres.2007.10.012

413 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J., 2000. Biodiversity
 414 hotspots for conservation priorities. *Nature* 403, 853–858. doi:10.1038/35002501

415 Natural History Museum of Bern, 2017. World Spider Catalog [WWW Document]. online
 416 <http://wsc.nmbe.ch>, version 16.5. Accessed 1st June 2017. doi:10.24436/2

417 Nogués-Bravo, D., Araújo, M.B., Romdal, T., Rahbek, C., 2008. Scale effects and human impact on
 418 the elevational species richness gradients. *Nature* 453, 216–219. doi:10.1038/nature06812

419 Nyundo, B.A., 2002. The diversity of carabid beetles (Coleoptera: Carabidae) in the Udzungwa
 420 Mountains National Park, Tanzania. PhD thesis, University of Dar es Salaam.

421 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,
 422 O’Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2017.
 423 *vegan: community ecology pack*. <https://cran.r-project.org/package=vegan>.

424 Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., Bruford,
 425 M.W., Brummitt, N., Butchart, S.H.M., Cardoso, A.C., Coops, N.C., Dulloo, E., Faith, D.P.,
 426 Freyhof, J., Gregory, R.D., Heip, C., Hoft, R., Hurr, G., Jetz, W., Karp, D.S., McGeoch, M.A.,

427 Obura, D., Onoda, Y., Pettorelli, N., Reyers, B., Sayre, R., Scharlemann, J.P.W., Stuart, S.N.,
428 Turak, E., Walpole, M., Wegmann, M., 2013. Essential biodiversity variables. *Science*. 339,
429 277–278. doi:10.1126/science.1229931

430 Peters, M.K., Hemp, A., Appelhans, T., Behler, C., Classen, A., Detsch, F., Ensslin, A., Ferger,
431 S.W., Frederiksen, S.B., Gebert, F., Haas, M., Helbig-Bonitz, M., Hemp, C., Kindeketa, W.J.,
432 Mwangomo, E., Ngereza, C., Otte, I., Röder, J., Rutten, G., Schellenberger Costa, D.,
433 Tardanico, J., Zancolli, G., Deckert, J., Eardley, C.D., Peters, R.S., Rödel, M.-O., Schleuning,
434 M., Ssymank, A., Kakengi, V., Zhang, J., Böhning-Gaese, K., Brandl, R., Kalko, E.K.V.,
435 Kleyer, M., Nauss, T., Tschapka, M., Fischer, M., Steffan-Dewenter, I., Gaston, K.J., Brown,
436 J.H., Fine, P.V.A., Rhode, K., Mittelbach, G.G., Allen, A.P., Brown, J.H., Gillooly, J.F.,
437 Hawkins, B.A., Kreft, H., Jetz, W., Hutchinson, G.E., Brown, J.H., Hurlbert, A.H., Stegen,
438 J.C., Belmaker, J., Jetz, W., Romdal, T.S., Grytnes, J.-A., Colwell, R.K., Rahbek, C., Gotelli,
439 N.J., Brehm, G., Colwell, R.K., Kluge, J., Novotny, V., Basset, Y., Wiens, J.J., Crisp, M.D.,
440 Kühnel, S., Blüthgen, N., Classen, A., Chao, A., Colwell, R.K., Lin, C.-W., Gotelli, N.J., Chao,
441 A., Chao, A., Burnham, K.P., Anderson, D.R., Burnham, K.P., Anderson, D.R., Huyvaert,
442 K.P., Grace, J.B., Colwell, R.K., Brehm, G., Cardelus, C.L., Gilman, A.C., Longino, J.T.,
443 Duan, M., Kessler, M., Rahbek, C., Rahbek, C., Wiens, J.J., Graham, C.H., Jablonski, D.,
444 Merckx, V.S.F.T., Wu, Y., Kozak, K.H., Wiens, J.J., Tittensor, D.P., Baur, B., Meier, T., Baur,
445 A., Schmera, D., McCain, C.M., Colwell, R.K., Forister, M.L., Zhou, J., Allen, A.P., Gillooly,
446 J.F., Savage, V.M., Brown, J.H., Rolland, J., Condamine, F.L., Jiguet, F., Morlon, H., Bagchi,
447 R., Johnson, D.J., Beaulieu, W.T., Bever, J.D., Clay, K., Hemp, A., Hemp, A., Trautwein,
448 M.D., Wiegmann, B.M., Beutel, R., Kjer, K.M., Yeates, D.K., Misof, B., Grafen, A., Zancolli,
449 G., Steffan-Dewenter, I., Rödel, M.-O., Hemp, C., Hemp, C., Wang, D., Atzberger, C., Eilers,
450 P.H.C., Colwell, R.K., Hurltt, G.C., Johnson, J.B., Omland, K.S., Hsieh, T.C., Ma, K.H., Chao,
451 A., Rosseel, Y., 2016. Predictors of elevational biodiversity gradients change from single taxa
452 to the multi-taxa community level. *Nat. Commun.* 7, 13736. doi:10.1038/ncomms13736

453 Pinkus-Rendon, M.A., Leon-Cortes, J.L., Ibarra-Nunez, G., 2006. Spider diversity in a tropical
454 habitat gradient in Chiapas, Mexico. *Divers. Distrib.* 12, 61–69. doi:10.1111/j.1366-
455 9516.2006.00217.x

456 Pinto-Leite, C.M., Rocha, P.L.B., 2012. Visual search for tropical web spiders: the influence of plot
457 length, sampling effort, and phase of the day on species richness. *Environ. Entomol.* 41, 1534–
458 43. doi:10.1603/EN11257

459 Poynton, J.C., Loader, S.P., Sherratt, E., Clarke, B.T., 2007. Amphibian diversity in East African
460 biodiversity hotspots: altitudinal and latitudinal patterns. *Biodivers. Conserv.* 16, 1103–1118.
461 doi:10.1007/s10531-006-9074-1

462 R Development Core Team, 2017. R: a language and environment for statistical computing. Version
463 3.4.1.

464 Rahbek, C., 1997. The relationship among area, elevation, and regional species richness in
465 Neotropical birds. *Source Am. Nat.* 149, 875–902. doi:10.1086/286028

- 466 Riechert, S.E., Gillespie, R.G., 1986. Habitat choice and utilization in web-building spiders, in:
467 Shear, W.A. (Ed.), *Spiders: Webs, Behavior and Evolution*. Stanford University Press, pp. 23–
468 48.
- 469 Romdal, T.S., Rahbek, C., 2009. Elevational zonation of afrotropical forest bird communities along
470 a homogeneous forest gradient. *J. Biogeogr.* 36, 327–336. doi:10.1111/j.1365-
471 2699.2008.01996.x
- 472 Romero, S.A., Harwood, J.D., 2010. Diel and seasonal patterns of prey available to epigeal
473 predators: evidence for food limitation in a linyphiid spider community. *Biol. Control* 52, 84–
474 90. doi:10.1016/j.biocontrol.2009.09.013
- 475 Rosenzweig, M.L., 1971. Paradox of enrichment: destabilization of exploitation ecosystems in
476 ecological time. *Science*. 171, 385–387. doi:10.1126/science.171.3969.385
- 477 Rovero, F., Mtui, A., Kitegile, A., Jacob, P., Araldi, A., Tenan, S., 2015. Primates decline rapidly in
478 unprotected forests: evidence from a monitoring program with data constraints. *PLOS ONE*
479 10, 1–13. doi:10.1371/journal.pone.0118330
- 480 Rovero, F., Owen, N., Jones, T., Canteri, E., Iemma, A., Tattoni, C., 2017. Camera trapping surveys
481 of forest mammal communities in the Eastern Arc Mountains reveal generalized habitat and
482 human disturbance responses. *Biodivers. Conserv.* 26, 1103–1119. doi:10.1007/s10531-016-
483 1288-2
- 484 Sala, O.E., 2000. Global Biodiversity Scenarios for the Year 2100. *Science*. 287, 1770–1774.
485 doi:10.1126/science.287.5459.1770
- 486 Sanders, N., 2002. Elevational gradients in ant species richness: area, geometry, and Rapoport's
487 rule. *Ecography (Cop.)*. 1, 25–32. doi:10.1034/j.1600-0587.2002.250104.x
- 488 Sanders, N.J., Rahbek, C., 2012. The patterns and causes of elevational diversity gradients.
489 *Ecography (Cop.)*. 35, 1–3. doi:10.1111/j.1600-0587.2011.07338.x
- 490 Schaffers, A.P., Raemakers, I.P., Sýkora, K. V., Ter Braak, C.J.F., 2008. Arthropod assemblages
491 are best predicted by plant species composition. *Ecology* 89, 782–794. doi:10.1890/07-0361.1
- 492 Scharff, N., 1992. The linyphiid fauna of eastern Africa (Araneae: Linyphiidae): distribution
493 patterns, diversity and endemism. *Biol. J. Linn. Soc.* 45, 117–154. doi:10.1111/j.1095-
494 8312.1992.tb00635.x
- 495 Scharff, N., Coddington, J.A., Griswold, C.E., Hormiga, G., Bjørn, P. de P., 2003. When to quit?
496 estimating spider species richness in a northern European deciduous forest. *J. Arachnol.*
497 doi:10.1636/0161-8202(2003)031[0246:WTQESS]2.0.CO;2
- 498 Silva-Davila, D., Coddington, J. a., 1996. Spiders of Pakitza (Madre de Dios, Peru): species
499 richness and notes on community structure, in: Wilson, D.E., Sandoval, A. (Eds.), *The*
500 *Biodiversity of Southeastern Peru*. Smithsonian Institution, Washington, pp. 253–311.

501 Sørensen, L., 2004. Composition and diversity of the spider fauna in the canopy of a montane forest
502 in Tanzania. *Biodivers. Conserv.* 13, 437–452. doi:10.1023/B:BIOC.00000006510.49496.1e

503 Sørensen, L.L., Coddington, J.A., Scharff, N., 2002. Inventorying and estimating subcanopy spider
504 diversity using semiquantitative sampling methods in an Afromontane forest. *Environ.*
505 *Entomol.* 31, 319–330. doi:10.1603/0046-225X-31.2.319

506 Stanley, W., Hutterer, R., 2007. Differences in abundance and species richness between shrews and
507 rodents along an elevational gradient in the Udzungwa Mountains, Tanzania. *Acta Theriol.*
508 (Warsz). 52, 261–275. doi:10.1007/BF03194222

509 Stevens, G.C., 1992. The elevational gradient in altitudinal range: an extension of Rapoport's
510 latitudinal rule to altitude. *Am. Nat.* 140, 893–911. doi:10.1086/285447

511 Sundqvist, M.K., Sanders, N.J., Wardle, D. a., 2013. Community and ecosystem responses to
512 elevational gradients: processes, mechanisms, and insights for global change. *Annu. Rev. Ecol.*
513 *Evol. Syst.* 44, 261–280. doi:10.1146/annurev-ecolsys-110512-135750

514 Tattersfield, P., Seddon, M.B., Ngereza, C., Rowson, B., 2006. Elevational variation in diversity
515 and composition of land-snail faunas in a Tanzanian forest. *Afr. J. Ecol.* 44, 47–60.
516 doi:10.1111/j.1365-2028.2006.00612.x

517 Ugland, K.I., Lamshead, P.J.D., McGill, B., Gray, J.S., Dea, N.O., Ladle, R.J., Whittaker, R.J.,
518 2007. Modelling dimensionality in species abundance distributions: description and evaluation
519 of the Gambin model. *Evol. Ecol. Res.* 9, 313–324. doi:10.1111/ecog.00861

520 Whittaker, R.J., Willis, K.J., Field, R., 2001. Scale and species richness: towards a general,
521 hierarchical theory of species diversity. *J. Biogeogr.* 28, 453–470. doi:10.1046/j.1365-
522 2699.2001.00563.x

523 Willig, M.R., Kaufman, D.M., Stevens, R.D., 2003. Latitudinal gradients of biodiversity: pattern,
524 process, scale, and synthesis. *Annu. Rev. Ecol. Evol. Syst.* 34, 273–309.
525 doi:10.1146/annurev.ecolsys.34.012103.144032

526 Wise, D.H., 1993. *Spiders in Ecological Webs*. Cambridge University Press, Cambridge.

527 Wright, D.H., 1983. Species-energy theory: an extension of species-area theory. *Oikos* 41, 496.
528 doi:10.2307/3544109

529 **Tables**
530

Plot	Latitude	Longitude	Altitude	Elevation	Sampling completeness	Observed species richness	Estimated species richness (Chao 1)
1	-7.687091	36.94129	650.212	Low	71%	115	162.6
2	-7.687496	36.94022	649.972	Low	68%	136	201.0
3	-7.684011	36.93038	1004.936	Mid	63%	119	189.3
4	-7.684797	36.93014	993.160	Mid	68%	135	199.8
5	-7.679317	36.91859	1447.860	High	72%	134	185.8
6	-7.678377	36.91857	1481.747	High	72%	124	171.6
7	-7.689854	36.93354	707.651	Low	74%	143	193.5
8	-7.686934	36.92758	977.779	Mid	59%	126	214.5
9	-7.685076	36.91458	1526.688	High	72%	135	188.5
10	-7.841573	36.86703	674.245	Low	66%	125	188.1
11	-7.840599	36.85917	1006.378	Mid	68%	110	162.2
12	-7.827731	36.84056	1551.922	High	79%	126	159.4
13	-8.503722	35.91915	659.345	Low	75%	108	143.2
14	-8.499499	35.91654	908.324	Mid	80%	116	144.7
15	-8.488775	35.90760	1531.494	High	83%	127	152.2

531 Table 1. Location and climatic features, sampling efficiency and observed number of species of sampling plots.
532
533

Elevation	Observed species richness	Estimated species richness ($\pm$ s.e.m.)	Sampling completeness	Beta Jaccard (95% CI)	Beta Bray-Curtis (95% CI)
Low (~700 m.a.s.l.)	305	398.52 $\pm$ 11.80	77%	0.68 (0.68, 0.71)	0.67 (0.67, 0.68)
Mid (~1000 m.a.s.l.)	307	418.05 $\pm$ 10.23	73%	0.70 (0.72, 0.76)	0.70 (0.69, 0.72)
High (~1500 m.a.s.l.)	311	386.77 $\pm$ 13.92	80%	0.68 (0.69, 0.73)	0.64 (0.65, 0.67)

534 Table 2. Total observed and estimated number of species (Chao 1), sampling completeness and beta diversity for each
535 of the three elevations. Confidence intervals were calculated from rarefied communities.
536

537 **Figure legends**

538 Fig. 1. Study area and plots. Close-ups show the plots in the Udzungwa Mountains National Park
539 (A and B) and in the Uzungwa Scarp Forest Reserve (C). Plots are coloured according to whether
540 they were at low (red), mid (purple) and high (blue) elevations.
541

542 Fig. 2. Number of species in the study plots over the number of collected spider individuals. a)
543 Observed and estimated number of species in all plots. b) Rarefied accumulated number of species
544 at each elevation. c) Rarefied accumulated number of species in each plot.
545

546 Fig. 3. Diversity values for the sampling plots according to elevation. a) Number of spider
547 individuals per plot. b) Percentage of adults per plot. c) Observed number of species per plot. d)
548 Estimated number of species (Chao 1 and Abundance Coverage Estimator) per plot. e) Number of
549 rare species (species represented by one or two specimens in the entire dataset) at each elevation.
550

551 Fig. 4. Abundance-based species similarity among the studied communities. The ordination is a
552 Non-Metric Multidimensional Scaling based on the Bray-Curtis similarity index values. Colours
553 represent plots at low (red), mid (purple) and high (blue) elevations.

554
555 Fig. 5. Dissimilarity values among the spider communities obtained from an ANOSIM based on
556 Bray-Curtis dissimilarity index.

557
558 Fig. 6. Species abundance distributions using the Gambin model. Curves (a) and alpha values (b)
559 for each of the 15 spider communities at the three elevations.

560
561 Fig. 7. Diversity values in each stratum of the vegetation at the three elevations. a) Percentage of
562 spider individuals on the ground (black), in herbaceous vegetation (grey) and in the understorey or
563 lower canopy (white). b) Number of species in ground (G), herbaceous (H) and understorey (U)
564 vegetations in low (red), mid (purple) and high (blue) elevation plots.

565
566 Fig. 8. Percentages of spiders of different predatory guilds in the 15 communities. a) Percentage of
567 species per plot. b) Percentage of individuals per plot.

568
569

Legend

- Forest cover
- Udzungwa Mountains National Park
- Contour lines (100 m)
- Roads
- Plots

Fig. 2

Fig. 3

Fig. 4

Steinhaus (abundance)
Distance in taxonomic structure
between communities (NMDS)

Fig. 5

ANOSIM – Plots – Bray–Curtis

Fig. 6

Fig. 7

Fig. 8