<https://helda.helsinki.fi>

Fermentation of cereal, pseudo-cereal and legume materials with *Propionibacterium freudenreichii* and *Levilactobacillus brevis* for vitamin B12 fortification

Xie, Chong

2021-02

Xie , C , Coda , R , Chamlagain , B , Edelmann , M , Varmanen , P , Piironen , V & Katina , K
2021 , ' Fermentation of cereal, pseudo-cereal and legume materials with *Propionibacterium freudenreichii* and *Levilactobacillus brevis* for vitamin B12 fortification ' , LWT-Food Science and Technology , vol. 137 , 110431 . <https://doi.org/10.1016/j.lwt.2020.110431>

<http://hdl.handle.net/10138/323577>

<https://doi.org/10.1016/j.lwt.2020.110431>

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Journal Pre-proof

Fermentation of Cereal, Pseudo-cereal and Legume Materials with *Propionibacterium freudenreichii* and *Levilactobacillus brevis* for Vitamin B12 Fortification

Chong Xie, Rossana Coda, Bhawani Chamlagain, Minnamari Edelmann, Pekka Varmanen, Vieno Piironen, Kati Katina

PII: S0023-6438(20)31419-5

DOI: <https://doi.org/10.1016/j.lwt.2020.110431>

Reference: YFSTL 110431

To appear in: *LWT - Food Science and Technology*

Received Date: 15 April 2020

Revised Date: 21 October 2020

Accepted Date: 21 October 2020

Please cite this article as: Xie, C., Coda, R., Chamlagain, B., Edelmann, M., Varmanen, P., Piironen, V., Katina, K., Fermentation of Cereal, Pseudo-cereal and Legume Materials with *Propionibacterium freudenreichii* and *Levilactobacillus brevis* for Vitamin B12 Fortification, *LWT - Food Science and Technology*, <https://doi.org/10.1016/j.lwt.2020.110431>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

CX performed the experiments and drafted the manuscript. RD, BC, ME, PV, VP and KK participated in conceiving the experiments and reviewing the manuscript.

1 **Fermentation of Cereal, Pseudo-cereal and Legume**
2 **Materials with *Propionibacterium freudenreichii* and**
3 ***Levilactobacillus brevis* for Vitamin B12 Fortification**

4 Chong Xie ^{a*}, Rossana Coda ^{a, b}, Bhawani Chamlagain ^a, Minnamari Edelmann ^a,
5 Pekka Varmanen ^a, Vieno Piironen ^a, Kati Katina ^a

6

7 ^aDepartment of Food and Nutrition, P. O. Box 27, University of Helsinki, FI-00014,
8 Helsinki, Finland

9 ^bHelsinki Institute of Sustainability Science (HELSUS), Finland.

10 *Corresponding author. Tel.: + 358 44 912 0811

11 E-mail address: chong.xie@helsinki.fi

12

13

ABSTRACT

The present study investigated the *in situ* production of vitamin B12 in eleven cereal, pseudo-cereal and legume materials by fermentation with *Propionibacterium freudenreichii* DSM 20271 and *Levilactobacillus brevis* (formerly *Lactobacillus brevis*) ATCC 14869. *P. freudenreichii* was used as the vitamin producer and *L. brevis* was selected to improve the consistency and microbial safety of the process. The study showed that more than 300 ng/g dw of vitamin B12 (daily requirement: 2.4 µg) were produced during fermentation in most of the studied brans and legumes. The highest vitamin B12 production was observed in the fermentation of the rice bran (ca. 742 ng/g dw), followed by the fermentation of buckwheat bran (ca. 631 ng/g dw). Furthermore, partial least squares (PLS) regression analysis suggested that the production of vitamin B12 was greatly influenced by the nutrient composition of the fermented raw materials. Meanwhile, *L. brevis* was found to effectively inhibit the growth of *Enterobacteriaceae* during fermentation. These results demonstrated that fermentation of cereal, pseudo-cereal and legume materials with *P. freudenreichii* and *L. brevis* is effective in fortifying plant-based food with vitamin B12.

Keywords: *Propionibacterium freudenreichii*; *Levilactobacillus brevis*; Vitamin B12;

In situ fortification; Brans

1. Introduction

Vitamin B12 (hereafter called B12) is naturally present in foods of animal origin and can be found in plant-based foods only as a consequence of fermentation or chemical fortification (Watanabe, 2007). Humans require B12 as a cofactor in cellular metabolism, and its deficiency may cause various health disorders, such as megaloblastic anemia, ataxia, and cognitive decline (Green & Miller, 2013). Globally, clinical deficiency of B12 is relatively uncommon nowadays, but subclinical deficiency of B12 is common among people of all ages, especially among people with low intake of animal products (Green et al., 2017; Smith, Warren, & Refsum, 2018). On the other hand, the current trend of replacing animal-based products with plant-based materials could result in a considerable global decrease of dietary B12 intake (Marsh, Zeuschner, & Saunders, 2011). Therefore, developing B12-fortified plant-based food products is increasing in importance for future food trends.

In situ fortification of food materials by fermentation with selected starter cultures is an economical and efficient way to provide micronutrients to the consumer. Certain *Lactobacillus* species (e.g. *Lactobacillus reuteri*) have been reported to be able to synthesize B12-like compounds (Taranto et al., 2003; Molina, Médici, de Valdez, & Taranto 2012), which on identification revealed to be mainly pseudo-vitamin B12 (Crofts et al., 2013; Santos et al., 2007), an inactive form for humans (Watanabe, 2007). *Propionibacterium freudenreichii* is a food-grade bacterium with the ability to *de novo* synthesize active form of B12 (Martens, Barg, Warren, & Jahn, 2002). However, the ability of B12 production among different strains of *P. freudenreichii* differs considerably (Hugenschmidt, Schwenninger, Gnehm & Lacroix, 2010;

Chamlagain et al. 2016). In the previous study (Xie et al., 2018), we found that wheat flour and wheat bran were promising substrates for B12 fortification using *P. freudenreichii* DSM 20271. Moreover, *P. freudenreichii* increased B12 production in wheat bran significantly during mixed culture fermentation with *Levilactobacillus brevis* ATCC 14869 (formerly *Lactobacillus brevis*) added to control the growth of indigenous microbes during the fermentation process (Xie et al., 2019).

Although wheat is one of the most widely cultivated crops, a great proportion of people around the world consume other grains as staple foods (Curtis, 2002). Moreover, for people suffering from allergy or intolerance to wheat products, other grain materials should be considered for B12 fortification by fermentation. Sorghum, millet, pseudocereals, and legumes are popular gluten-free alternatives to wheat because they can adapt to harsh environments and contain various health-promoting ingredients, such as dietary fiber and phenolic compounds (Gobbetti, De Angelis, Di Cagno, Polo, & Rizzello, 2019; Taylor, Belton, Beta, & Duodu, 2014). Cereal brans, which are by-products of the cereal milling process, contain high levels of dietary fiber, proteins, and various beneficial compounds, such as polyphenols and phytosterols (Luithui, Nisha, & Meera, 2019). The brans are, however, underutilized for food purposes because of their detrimental effect on the technological and sensory quality of products (Prückler et al., 2014). The development of innovative technologies for upgrading these by-products into the food system is widely suggested for a more resilient food chain.

Therefore, the present study aimed to investigate the *in situ* production of B12 by mixed culture fermentation with *P. freudenreichii* and *L. brevis* in various grain

materials, including cereals (oat bran, rice bran, rye bran, millet flour, and sorghum flours), pseudocereals (buckwheat bran, amaranth flour, and quinoa flour), and legumes (soybean flour, faba bean flour, and lupine flour). Acidification properties and content of acids were also monitored to elucidate the metabolism of starters during fermentation. The content of nutrient composition and cobalt, a limiting factor for the production of B12 by *P. freudenreichii* (Hugenschmidt et al., 2010), were measured and their effects on B12 production in different materials was evaluated. The main compositional factors influencing B12 production in different raw materials were studied using partial least squares regression analysis (PLS).

2. Materials and Methods

2.1 Grain materials, microbial strains, and culture preparation

Eleven kinds of grain materials used in this study were purchased from markets. Brans and grains of sorghum and millet were milled by a rotor mill (Ultra centrifugal mill ZM 200, Retsch GmbH & Co. KG, Haan, Germany) at a speed of $10,000 \times g$. The particle size of each material is shown in Supplemental Table 1. The nutrient composition of millet flour, sorghum flour, and faba bean flour was analyzed at Eurofins Scientific (Finland). The contents of fat and protein were determined following the methods of NordVal International (NMKL) 160 (1998) and NMKL 6:2003 (2003), respectively. Dietary fiber contents were determined according to the AOAC method 985.29 (2003). The contents of available carbohydrate (monosaccharides, disaccharides, and starch) were determined by subtracting the fiber, fat, ash, and protein content from the total. The nutrient composition of other materials was obtained from their nutrition fact labels. The cobalt contents were

determined by the Finnish Environment Institute using a method based on microwave-assisted digestion and inductively coupled plasma mass spectrometry (ICP-MS) quantitation as described by Nóbrega et al. (2012).

Both *P. freudenreichii* DSM 20271 and *L. brevis* ATCC 14869 cultures were cryopreserved at -80°C in 15% glycerol (v/v). Fermentation inocula were prepared as follows: *P. freudenreichii* was propagated in the yeast extract lactate (YEL) medium (Malik, Reinbold, & Vedamuthu, 1968) at 30°C for 3 d. *L. brevis* was propagated in de Man, Rogosa, and Sharpe (MRS) medium (Lab M, Lancashire, UK) at 37°C for 1 d. The bacterial cells were harvested by centrifugation ($3,200 \times g$, 10 min) and resuspended in MilliQ water prior to inoculation.

2.2 Batter preparation and fermentation

2.2.1 Screening of grain materials for *in situ* production of B12

Batters were prepared by mixing MilliQ-water and the flours. The ratio of water to flour in each batter (Table 1) was decided according to preliminary experiments to achieve a semi-liquid batter. After mixing, 30 g of batters were transferred into Falcon tubes (50 mL) and inoculated with *P. freudenreichii* (ca. $9.0 \log \text{CFU/g}$) and *L. brevis* (ca. $6.0 \log \text{CFU/g}$). The inoculation levels of *P. freudenreichii* and *L. brevis* were established in the previous studies to provide a meaningful level of B12 production without compromising the microbial safety in non-sterilized wheat bran (Xie et al., 2018; Xie et al., 2019). Triplicate cultures were prepared for each time point (0 d, 1 d and 3 d). Fermentation was carried out in a shaking (200 rpm) incubator at 25°C for 3 d. At each time point (0 d, 1 d and 3 d), three replicate samples of each batter type

were analyzed for pH value, total titratable acids (TTA), as well as, contents of lactic acid, acetic acid, propionic acid, and B12.

2.2.2 Fermentation with different levels of starter cultures

Four materials (buckwheat bran, oat bran, rice bran, and sorghum flour) were chosen for further experiments. Batters were prepared and fermented in the same way as in the screening experiments. In each material, batters were prepared in four ways as follows: spontaneously fermented (no inoculum, Control); inoculated with 9.0 log CFU/g *P. freudenreichii* and 6.0 log CFU/g *L. brevis* (P9/L6); inoculated with 9.0 log CFU/g *P. freudenreichii* and 7.0 log CFU/g *L. brevis* (P9/L7); and inoculated with 9.0 log CFU/g *P. freudenreichii* and 8.0 log CFU/g *L. brevis* (P9/L8). Different levels of *L. brevis* were tested to determine the optimal inoculation level for a meaningful B12 production together with inhibition of *Enterobacteriaceae* growth. Triplicate tubes were prepared for each time point, and at d 0, 1, and 3, samples were taken out for measurement of pH, TTA, viable cell counts, and content of B12.

2.3 Microbial cell counts

Common plate count techniques were used to determine viable cell numbers. Batters (10 g) were serially diluted in sterile sodium chloride solution (8.5 g/l) and plated on respective agar plates for the monitoring of different microbial groups. YEL plates for the determination of *P. freudenreichii* were incubated anaerobically (Anaerogen, Oxoid, Basingstoke, UK) for 4 d and aerobically for 1 d at 30 °C to allow the colonies of *P. freudenreichii* to turn brownish and distinguishable from other bacteria. Lactic acid bacteria (LAB) were counted on MRS agar (Lab M) plates

supplemented with 0.1 mg/g cycloheximide (Sigma Chemical Co., USA) and incubated at 30 °C for 48 h. The plate count agar (PCA) plates (Lab M) and violet red bile glucose agar (VRBGA) plates (Lab M) were used for the cell counts of total aerobic bacteria (TAB) and total *Enterobacteriaceae*, respectively. The PCA plates were incubated at 30 °C for 48 h, and the VRBGA plates were incubated at 37 °C for 24 h.

2.4 Determination of pH, TTA, and content of acids

A pH meter (Portamess 752 Calimatic, Knick, Berlin, Germany) was used for the measurement of pH value. TTA was determined by a titrator (EasyPlus, Mettler Toledo, Schott, Germany), and the value is expressed as the volume of 0.1 mol/L NaOH solution (ml) required to adjust the pH of 10-g samples in 90 ml Milli-Q water to 8.5.

Batter samples (1 g) were diluted (1:1~1:10, w/v) in water and centrifuged (3,200 × g, 10 min). The supernatants were filtered (0.45 µm, Pall, USA) into vials. The contents of lactic acid (LA), acetic acid (AA), and propionic acid (PA) were determined using a high-performance liquid chromatography (HPLC) method, as reported earlier (Xie et al., 2018).

2.5 Determination of B12 content in batters

B12 in batters was determined by an ultra-high performance liquid chromatography (UHPLC) method, as described by Chamlagain, Edelmann, Kariluoto, Ollilainen, and Piironen (2015), with minor modifications. Briefly, 3 g of batter samples were mixed with 15 mL of extraction buffer (8.3 mmol/L sodium

hydroxide and 20.7 mmol/L acetic acid, pH 4.5) and 100 μ L of sodium cyanide (1g/100g in water). After extraction in boiling water (30 min), cooled mixtures were incubated in a water bath (30 min, 37°C) with the addition of 300 μ L α -amylase (50 mg/ml) to allow the breakdown of starch before centrifugation ($6,900 \times g$, 10 min). Residues after centrifugation were suspended in 5 mL of extraction buffer and centrifuged again. Both supernatants were combined and adjusted to 25 mL with the extraction buffer. Finally, 10 mL of the extracts were purified using an immunoaffinity column (Easi-Extract; Glasgow, Scotland) and analyzed with a UPLC system (Waters, Milford, MA, USA) equipped with a photodiode array detector (at 361 nm) and an Acquity HSS T3 C18 column (2.1×100 mm, 1.8 μ m). The mobile phase was a gradient flow of MilliQ water and acetonitrile, both with 0.25 mg/g trifluoroacetic acid. The presence of any other corrinoids in the samples, including pseudo-vitamin B12, was followed in the chromatograms based on knowledge of their UV-absorption spectra and retention times from previous studies (Chamlagain et al., 2015; Chamlagain et al., 2018).

2.6 Statistical analysis

Statistical analysis was performed using SPSS 24.0 for Windows (IBM Corporation, NY, USA). One-way analysis of variance (ANOVA) and Tukey's *post hoc* test were used to determine significant differences in all parameters during fermentation at a *p*-value < 0.05 among the samples. Multivariate data analysis was performed by PLS, using Simca 15.0 (Umetrics AB, Malmö, Sweden).

3. Results

3.1 Nutrient and cobalt content of grain materials

The nutrient composition of the grain materials is shown in Table 1. In general, cereal and pseudocereal flours had the highest content of available carbohydrates (57–68 g/100 g), a moderate protein content (9–14 g/100 g), and low levels of lipids (5–6 g/100 g) and dietary fiber (6–10 g/100 g). Brans differed greatly in their nutrient content (available carbohydrates 26–49 g/100 g, protein 13–40 g/100 g, dietary fiber 9–39 g/100 g, and lipids 4–20 g/100 g). Legume flours had a high protein content (31–43 g/100 g). Faba bean was rich in available carbohydrates (42 g/100 g), whereas lupine and soya bean had high fat content (12–20 g/100 g). The dietary fiber content in legume flours varied from 9 g/100 g to 28 g/100 g.

The cobalt content among the grain materials varied, and the highest was found in the faba bean flour (ca. 696 ng/g dw). In the rye bran, oat bran, and lupine flour, a low level of cobalt was observed (17–29 ng/g dw). Among other materials, the cobalt contents ranged from 68 ng/g dw to 183 ng/g dw.

3.2 Acidification of batters in the screening of materials for B12 production

3.2.1 pH and TTA

As shown in Table 2, pH in the batters ranged from 5.7 (lupine flour) to 6.8 (rice bran), and TTA varied from 0.5 mL (oat bran) to 4.8 mL (lupine flour) on d 0. In the batters of millet flour and two pseudocereal flours, pH decreased rapidly to ca. 4.0 already on d 1 and a high TTA (up to 18 mL) was reached after fermentation. A similar rapid pH drop and a high TTA level were detected also in the batters of rye bran and faba bean flour during fermentation. In the buckwheat bran batter, high TTA

(ca. 15 mL on d 1 and ca. 18 mL on d 3) and a pH remaining at 4.9 during fermentation were detected. In the batters of rice bran, oat bran, sorghum flour, soybean flour, and lupine flour, pH levels were higher than 5.5 and TTA increased by only about 1 mL during the first day. On d 3, pH in batters of rice bran (5.0), soybean flour (4.7), and lupine flour (4.9) were significantly ($p < 0.05$) lower than in the batters of oat bran (5.2) and sorghum flour (5.3). Accordingly, TTA in the rice bran, soybean flour, and lupine flour (11-13 mL) was significantly ($p < 0.05$) higher than in the batters of oat bran (6.1 mL) and sorghum flour (4.4 mL).

3.2.2 Production of organic acids

During the fermentation of oat bran, rice bran, sorghum flour, and lupine flour, LA was not detected at the measured time points, whereas a drastic increase of AA and PA contents were detected from d 1 to d 3 (Fig. 1). On d 1, the buckwheat bran batter had the highest level of both AA (ca. 9 mg/g dw) and PA (ca. 13 mg/g dw) and contained ca. 28 mg/g dw of LA. On d 3, the buckwheat bran batter contained the highest amount of PA (ca. 33 mg/g dw) and ca. 19 mg/g dw of AA but no LA was detected. During the fermentation of rye bran and flours of amaranth, quinoa, millet, and faba bean, the contents of LA (40.0–98.1 mg/g dw) were much higher than the contents of AA (1.4–4.8 mg/g dw) and PA (1.6–10.4 mg/g dw).

3.3 B12 synthesis in the screening of materials for B12 production

B12 levels of 19 - 38 ng/g dw were detected in the batters immediately following inoculum with *P. freudenreichii* (Table 3). After one day of fermentation, a significant ($p < 0.05$) increase of B12 content was observed in all batters, and the highest content

was found in the buckwheat bran batter (ca. 335 ng/g dw). From d 1 to d 3, there was no significant ($p < 0.05$) increase of B12 content in the batters of rye bran, millet flour, quinoa flour, and amaranth flour. On d 3, the highest content of B12 was found in the batter of rice bran (ca. 742 ng/g dw), followed by the batters of buckwheat bran (ca. 631 ng/g dw) and soybean flour (ca. 407 ng/g dw). In the batters of oat bran, sorghum flour, faba bean flour, and lupine flour, the B12 content ranged from 265 to 343 ng/g dw on d 3.

3.4 Correlation of nutrient composition and acidification properties with B12 production

A PLS modeling was used to investigate how nutrient composition, cobalt content, and acidification properties (pH, TTA, and production of acids) correlated with B12 production on d 1 and d 3 (Fig. 2). The first and second principal components (PC1 and PC2) together explained 68% (d 1) and 64% (d 3) of the variation. The projections of each variable on the line drawn via B12 production and the origin of the plot describes the correlation of the variables and B12 production. Moreover, the distance of each variable's projection from the origin indicate its influence on B12 production. On both d 1 and d 3, the B12 content was positively correlated with high pH value (> 4.7), the content of protein, fat and dietary fiber, and production of propionic acid and acetic acid. On the other hand, B12 production was negatively correlated with high TTA content (> 17 mL), the content of the available carbohydrate, and lactic acid. The cobalt content only had a limited influence on B12 production, both on d 1 and d 3.

3.5 Batters fermented with different inoculum levels of starter cultures

Buckwheat bran, oat bran, rice bran, and sorghum flour were chosen for further experiments to study the effect of different *L. brevis* inoculum levels on the fermentation of each material.

3.5.1 Microbial counts

Propionic acid bacteria (PAB) were not detected throughout the fermentation in any control batters (Table 4). In the inoculated batters, the cell density of PAB ranged from ca. 8.7 log CFU/g to ca. 8.8 log CFU/g. The cell density of inoculated PAB increased by ca. 0.5 log cycle in the batters of oat bran, rice bran, and buckwheat bran on d 1 and remained stable thereafter. In the sorghum batters, the increase of PAB cell density during fermentation was in the range of 0.2–0.4 log cycle.

In the control batters, LAB were detected only in the buckwheat bran and sorghum flour at a cell density at ca. 2.8 log CFU/g and ca. 2.3 log CFU/g, respectively. During fermentation, cell densities of LAB increased in all batters, and inoculation of *L. brevis* led to a higher cell number of LAB compared to the control batters with no inoculum. On d 3, the cell density of LAB reached 9.0 log CFU/g in all batters with inoculation except in the sorghum batter with an inoculum of P9/L6 (ca. 8.1 log CFU/g). Supplemental Table 2 shows the cell densities of total aerobic bacteria (TAB) in all batters. In general, similar cell densities of LAB and TAB were found during the fermentation of the oat bran and the buckwheat bran.

Before fermentation, *Enterobacteriaceae* were detected only in the buckwheat bran batter and sorghum flour batter at a level of 4.4 log CFU/g and 5.4 log CFU/g,

respectively. On d 1, the cell density of *Enterobacteriaceae* increased in all the spontaneously fermented batters and ranged from 7.6 log CFU/g to 9.0 log CFU/g. In the batters of oat bran, rice bran, and buckwheat bran, a significantly ($p < 0.05$) lower cell density of *Enterobacteriaceae* was detected when *L. brevis* was inoculated than in the corresponding control batters. In the batter of sorghum flour, a significantly ($p < 0.05$) lower cell density of *Enterobacteriaceae* was detected only when the inoculation level of *L. brevis* was higher than 7 log CFU/g. At the end of fermentation, no *Enterobacteriaceae* were detected in batters of rice bran, buckwheat bran, and sorghum flour when the inoculum size of *L. brevis* was greater than 7 log CFU/g.

3.5.2 Acidification of batters

During fermentation, the control batters reached the highest pH values and the lowest TTA content and the batters with the highest inoculum ratio of *L. brevis* (P9/L8) reached the lowest pH levels and the highest TTA content (Table 5). In both oat bran and buckwheat bran batters, a significantly ($p < 0.05$) higher TTA value was observed in P9/L6 batters than in the control batters. In the rice bran batter, a significant ($p < 0.05$) increase in TTA content and a decrease of pH value was found only when the *L. brevis* inoculum was 8.0 log CFU/g. In the sorghum batter, a significant ($p < 0.05$) increase of TTA content (from ca. 4.3 ml to 9.2 ml on d 3) and a decrease of pH value (from ca. 5.3 to ca. 3.9 on d 3) was observed for each increased inoculum level of *L. brevis*.

3.5.3 Content of B12

During d 1, the B12 production did not differ with different inocula of *L. brevis* in the batters of oat bran and rice bran (Table 5). On the contrary, on d 3, B12 content of oat bran P9/L6 batter (ca. 342 ng/g dw) was significantly ($p < 0.05$) higher than that of the oat bran P9/L7 and P9/L8 batters (298 - 312 ng/g dw). In rice bran, B12 content of P9/L8 batter (ca. 675 ng/g dw) was significantly ($p < 0.05$) lower than in P9/L7 and P9/L6 batter (ca. 720 ng/g dw). In the sorghum batters, B12 content was significantly ($p < 0.05$) lower on both d 1 and d 3 when the inoculation level of *L. brevis* was increased. Notably, only ca. 67 ng/g dw of B12 was produced in P9/L8 batter of sorghum at the end of fermentation.

4. Discussion

Our previous study showed that a significant increase of B12 content was achieved in wheat bran by fermentation with *P. freudenreichii* DSM 20271 and *L. brevis* ATCC 14869 (Xie et al., 2019). The present study aimed to study whether the mixed culture fermentation approach is applicable for the *in situ* fortification of other grains with B12. Furthermore, the correlation of B12 production with nutrient composition, cobalt content, pH, and production of acids during fermentation was investigated by PLS modeling to reveal critical factors affecting vitamin synthesis.

The materials used in this study had varying content of nutrients, which lead to different acidification kinetics and a greatly different production of B12 by *P. freudenreichii* (60 - 740 ng/g dw) during fermentation. In general, batters of cereal and pseudocereal flours had a higher level of LA content and TTA value during fermentation, which was probably due to their higher level of available carbohydrates

compared to other materials (brans and legume flours) supporting the growth of microorganisms. In line with this, the PLS model showed a positive correlation of available carbohydrate content with LA production and TTA (Fig. 2).

Although LA produced by LAB is the preferential carbon source for PAB (Piveteau, 1999), a fast drop in environmental pH caused by accumulating LA can inhibit the growth and metabolism of PAB because they preferably grow at neutral pH and stop producing acids when the pH value drops below 4.5 (Ye, Shijo, Jin, & Shimizu, 1996). B12 is a coenzyme in the metabolic pathway for PA and AA production in *P. freudenreichii* (Thierry et al., 2011). The PLS model revealed a positive correlation of B12 production with the production of PA and AA in the present study (Fig. 2). In accordance, the fermentation conditions in batters with high LA contents and correspondingly with low pH values (<4.5), such as millet, amaranth, and quinoa flours, generally produced low PA, AA, and B12 yields.

Compared to cereal and pseudocereal flours, the content of available carbohydrate in brans and legume flours was generally lower, which was consequently accompanied with higher pH values in batters during fermentation. The high pH values in some batters likely resulted from the low production levels of acids, for example, in oat bran (TTA: 6.4 mL on d 3). Besides, the high pH value may also be an effect from the presence of high buffering capacity compounds, such as proteins (Lević, Prodanović, & Sredanović, 2005) and dietary fiber (Tadesse, 1986), in the materials. For instance, in the buckwheat bran batter, despite a high TTA content observed (ca. 15 mL on d 1), the pH value remained at 4.9 from d 1 to d 3. The PLS

model also showed that dietary fiber, protein, and lipid contents were positively correlated with the production of B12.

Cobalt, the central metal ion of B12, is reported to be a limiting factor for the production of B12 during the fermentation of *P. freudenreichii* in various substrates, such as whey medium (Hugenschmidt, Schwenninger, & Lacroix, 2011) and aqueous cereal-based matrices (Chamlagain et al., 2017). Moreover, during fermentation in whey medium, the production of B12 by *P. freudenreichii* increased significantly when cobalt was added up to 5 µg/mL (Hugenschmidt et al., 2011). Our previous study also showed that adding cobalt (600 ng/g dw) in durum wheat flour batter was able to increase the production of B12 from ca. 33 ng/g dw to ca. 200 ng/g dw during fermentation with *P. freudenreichii* (Xie et al., 2018). PLS modeling indicates that cobalt content was not a limiting factor for B12 production under the conditions used here. Notably, there was no significant ($p > 0.05$) difference in B12 content (ca. 332 ng/g dw vs ca. 298 ng/g dw) at the end of fermentation between materials with the lowest level (17 ng/g dw in oat bran) and the highest level (696 ng/g dw in faba bean flour) of cobalt. On the other hand, rice bran, buckwheat bran, millet, sorghum, and amaranth flours had a similar level of cobalt (157–183 ng/g dw) but with a considerably different level of B12 production (51–742 ng/g dw), and the batter with a higher pH tended to have higher B12 production. Thus, the influence of cobalt content on B12 production seemed to be less significant than that of pH during fermentation when above the level of 17 ng/g dw of cobalt measured in this study.

Fermented soybean products, such as tempeh and natto, have been found to

contain certain amounts of B12 (0.1–8.0 µg/100 g fresh weight; Watanabe., Yabuta, Bito, & Teng, 2014). Moreover, recent studies showed that the inoculation of *P. freudenreichii* as a starter culture could significantly increase B12 production (up to 1230 ng/g dw) in lupine tempeh (Signorini et al., 2018; Wolkers – Rooijackers, Endika, & Smid, 2018). In the present study, 300–400 ng/g dw of B12 was obtained after the fermentation of three legume materials, which can provide a new option for developing B12-fortified legume products.

To the best of our knowledge, this was the first study to ferment oat, rice, and buckwheat bran with *P. freudenreichii* for *in situ* production of B12 and our study shows that they seem to be very promising materials for this purpose. Moreover, as side streams of the milling process, utilization of these brans for developing B12-fortified food can contribute to a more resilient food chain by reducing grain waste. Furthermore, developing B12-fortified sorghum products can be important because they are widely consumed in Asia and Africa (Nout, 2009). People in these two continents have a higher prevalence of B12 deficiency than people in other parts of the world (Hunt, Harrington, & Robinson, 2014). Therefore, sorghum flour and oat, rice, and buckwheat bran were chosen for further experiments.

Raw cereal materials usually contain some undesirable microorganisms, such as potential pathogens from the *Enterobacteriaceae* family, which can cause food poisoning by infecting the human gastrointestinal tract or producing various endotoxins (Singh, Sharma, & Nara, 2015). LAB in cereal materials can show antagonistic activity to pathogens by the production of acids and antimicrobial

compounds or by competitive exclusion (De Vuyst & Neysens, 2005). However, a high cell density of *Enterobacteriaceae* was found in the batters of brans and sorghum flour during fermentation in the present study. *Enterobacteriaceae* cell density in the buckwheat bran control batter increased by 4 log units during the first day of fermentation and still had a cell density at ca. 5 log CFU/g at the end of fermentation. In the oat bran and rice bran control batters, the cell numbers of *Enterobacteriaceae* increased to more than 7.0 log CFU/g even though no *Enterobacteriaceae* were detected before the fermentation.

As expected, the inoculation of *L. brevis* decreased the cell density of *Enterobacteriaceae* by at least 1 log unit compared to control batters, and stronger inhibitions were observed when the initial level of LAB increased during the fermentation. However, an increase of *L. brevis* inoculation also led to a drastic decrease of B12 production in the sorghum flour batter due to the faster pH drop. The production of PA in sorghum flour batter also drastically decreased (5.9 vs 1.4 mg/g dw) when an initial level of *L. brevis* was increased from 6 log CFU/g to 8 log CFU/g (Supplemental Table 3). In the fermentation of rice bran and buckwheat bran, although the initial level of *L. brevis* was increased from 6 log CFU/g to 8 log CFU/g, the PA production levels were not significantly affected (Supplemental Table 3), possibly because of the good buffering capacity (pH > 4.5 on d 3 despite the high TTA) of the materials. Therefore, the B12 content after d 3 was still high in the rice bran (600 ng/g dw) and in the buckwheat bran (500 ng/g dw). Considering that *in situ* synthesized B12 is stable during food processing (Edelmann, Chamlagain, Santin,

Kariluoto, & Piironen 2016), consuming a food product containing 10 g dry matter of rice bran or buckwheat bran fermented with *P. freudenreichii* could provide the daily requirement of B12 (2.4 µg for adults; Institute of Medicine, 1998). On the other hand, by replacing 30% of wheat flour in a bread recipe with fermented sorghum (with ca. 120 ng/g dw of B12), four slices of bread (120 g) could, theoretically, provide more than 2.4 µg of B12.

Further research on the sensory properties of food containing these fermented batters as ingredients (e.g. baked goods and pasta) could be performed to optimize their sensory quality and increase their acceptance. Moreover, a more in-depth study of each raw material with different nutrient composition (e.g. protein, fat, and dietary fiber) and origin would be needed for a more comprehensive understanding of their full potential in B12 production. In addition, maintaining pH at different levels and adding different ingredients (e.g. protein or materials with high availability of cobalt) during fermentation should be applied in future studies to investigate how these factors influence B12 *in situ* production in grain materials.

5. Conclusion

This work demonstrated that a nutritionally relevant amount of B12 was produced in non-sterilized grain materials by fermentation with *P. freudenreichii*. The higher B12 content in grain batters with a higher pH during fermentation suggests that maintaining optimal pH for *P. freudenreichii* can increase the production of B12. The contents of some components, such as protein and dietary fiber, in grain materials were found to be positively correlated with B12 production in PLS analysis probably due to their pH buffering capacity. Meanwhile, the combination of *P. freudenreichii*

with *L. brevis* inoculated at a certain level can effectively inhibit the growth of *Enterobacteriaceae* to ensure the microbial safety of the fermentation. In conclusion, grain materials fermented with *P. freudenreichii* and *L. brevis* can be a promising alternative to produce B12-fortified grain ingredients for a wide variety of plant-based foods.

6. Acknowledgments

The authors also would like to thank the Academy of Finland (grant number 325784) and China Scholarship Council for their financial supports. Mr. Timo Sara-Aho is acknowledged for the analysis of cobalt content.

References

- AOAC International. (2003). Total dietary fiber in foods, enzymatic- Gravimetric method. In "Official Methods of Analysis of AOAC International". 17th ed. 985.29. William Horwitz ed. Gaithersburg, MD, USA.
- Chamlagain, B., Deptula, P., Edelmann, M., Kariluoto, S., Grattepanche, F., Lacroix, C., ... & Piironen, V. (2016). Effect of the lower ligand precursors on vitamin B12 production by food-grade Propionibacteria. *LWT-Food Science and Technology*, 72, 117-124.
- Chamlagain, B., Edelmann, M., Kariluoto, S., Ollilainen, V., & Piironen, V. (2015). Ultra-high performance liquid chromatographic and mass spectrometric analysis of active vitamin B12 in cells of Propionibacterium and fermented cereal matrices. *Food Chemistry*, 166, 630-638.
- Chamlagain, B., Sugito, T., Deptula, P., Edelmann, M., Kariluoto, S., Varmanen, P., & Piironen, V. (2018). *In situ* production of active vitamin B12 in cereal matrices using *Propionibacterium freudenreichii*. *Food Science & Nutrition*, 6(1), 67-76.
- Crofts, T. S., Seth, E. C., Hazra, A. B., & Taga, M. E. (2013). Cobamide structure depends on both lower ligand availability and CobT substrate specificity. *Chemistry & biology*, 20(10), 1265-1274.
- Curtis, B. C., Rajaram, S., Gómez, M. (2002). *Bread wheat : improvement and production*. Rome: Food and Agriculture Organization of the United Nations.
- De Vuyst, L., & Neysens, P. (2005). The sourdough microflora: biodiversity and metabolic interactions. *Trends in Food Science & Technology*, 16(1-3), 43-56.
- Edelmann, M., Chamlagain, B., Santin, M., Kariluoto, S., & Piironen, V. (2016). Stability of added and in situ-produced vitamin B12 in breadmaking. *Food Chemistry*, 204, 21-28.
- Gobbetti, M., De Angelis, M., Di Cagno, R., Polo, A., & Rizzello, C. G. (2020). The sourdough fermentation is the powerful process to exploit the potential of legumes, pseudo-cereals and milling by-products in baking industry. *Critical reviews in food science*

- 468 *and nutrition*, 60(13), 2158-2173
- 469 Green, R., Allen, L. H., Bjorke-Monsen, A. L., Brito, A., Gueant, J. L., Miller, J. W., . . .
- 470 Yajnik, C. (2017). Vitamin B12 deficiency. *Nature Reviews Disease Primers*, 3(1), 1-20.
- 471 Green, R., & Miller, J. (2013). Vitamin B12. In *Handbook of Vitamins*, Fifth Edition (pp.
- 472 447-490): CRC Press.
- 473 Hugenschmidt, S., Schwenninger, S. M., Gnehm, N., & Lacroix, C. (2010). Screening of a
- 474 natural biodiversity of lactic and propionic acid bacteria for folate and vitamin B12
- 475 production in supplemented whey permeate. *International Dairy Journal*, 20(12), 852-857.
- 476 Hugenschmidt, S., Schwenninger, S. M., & Lacroix, C. (2011). Concurrent high production of
- 477 natural folate and vitamin B12 using a co-culture process with *Lactobacillus plantarum*
- 478 SM39 and *Propionibacterium freudenreichii* DF13. *Process Biochemistry*, 46(5),
- 479 1063-1070.
- 480 Hunt, A., Harrington, D., & Robinson, S. (2014). Vitamin B12 deficiency. *BMJ*, 349.
- 481 Lević, J., Prodanović, O., & Sredanović, S. (2005). Understanding the buffering capacity in
- 482 feedstuffs. *Biotechnology in Animal Husbandry*, 21(5-6), 309-313.
- 483 Institute of Medicine (1998). Dietary Reference Intakes for Thiamin, Riboflavin, Niacin,
- 484 Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. National
- 485 Academy of Sciences. Washington (DC): National Academies Press (US).
- 486 Luithui, Y., Nisha, R. B., & Meera, M. S. (2019). Cereal by-products as an important
- 487 functional ingredient: effect of processing. *Journal of Food Science and Technology*, 56(1),
- 488 1-11.
- 489 Malik, A. C., Reinbold, G. W., & Vedamuthu, E. R. (1968). An evaluation of taxonomy of
- 490 propionibacterium. *Canadian Journal of Microbiology*, 14(11), 1185-1191.
- 491 Marsh, K., Zeuschner, C., & Saunders, A. (2011). Health Implications of a Vegetarian Diet.
- 492 *American Journal of Lifestyle Medicine*, 6(3), 250-267.
- 493 Martens, J. H., Barg, H., Warren, M. J., & Jahn, D. (2002). Microbial production of vitamin
- 494 B12. *Applied Microbiology and Biotechnology*, 58(3), 275-285.
- 495 Molina, V., Mélici, M., de Valdez, G. F., & Taranto, M. P. (2012). Soybean-based functional
- 496 food with vitamin B12-producing lactic acid bacteria. *Journal of Functional Foods*, 4(4),
- 497 831-836.
- 498 Nordic Committee on Food Analysis (1998). NMKL method no. 160, 1998: Fat determination
- 499 in foods
- 500 Nordic Committee on Food Analysis (2003). NMKL method no. 6: Nitrogen determination in
- 501 foods and feeds according to Kjeldahl.
- 502 Nóbrega, J. A., Pirola, C., Fialho, L. L., Rota, G., de Campos Jordão, C. E. K. M. A., & Pollo,
- 503 F. (2012). Microwave-assisted digestion of organic samples: How simple can it become?
- 504 *Talanta*, 98, 272-276.
- 505 Nout, M. J. (2009). Rich nutrition from the poorest - cereal fermentations in Africa and Asia.
- 506 *Food Microbiology*, 26(7), 685-692.
- 507 Piveteau, P. (1999). Metabolism of lactate and sugars by dairy propionibacteria: A review.
- 508 *Lait*, 79(1), 23-41.
- 509 Prückler, M., Siebenhandl-Ehn, S., Apprich, S., Höltinger, S., Haas, C., Schmid, E., &
- 510 Kneifel, W. (2014). Wheat bran-based biorefinery 1: Composition of wheat bran and
- 511 strategies of functionalization. *LWT - Food Science and Technology*, 56(2), 211-221.

- Signorini, C., Carpen, A., Coletto, L., Borgonovo, G., Galanti, E., Capraro, J., . . . Scarafoni, A. (2018). Enhanced vitamin B12 production in an innovative lupin tempeh is due to synergic effects of *Rhizopus* and *Propionibacterium* in cofermentation. *International Journal of Food Science and Nutrition*, 69(4), 451-457.
- Singh, J., Sharma, S., & Nara, S. (2015). Evaluation of gold nanoparticle based lateral flow assays for diagnosis of enterobacteriaceae members in food and water. *Food Chemistry*, 170, 470-483.
- Smith, A. D., Warren, M. J., & Refsum, H. (2018). Vitamin B12. In *Advances in food and nutrition research* (Vol. 83, pp. 215-279). Academic Press
- Tadesse, K. (1986). The effect of dietary fibre isolates on gastric secretion, acidity and emptying. *British Journal of Nutrition*, 55(3), 507-513.
- Taylor, J. R. N., Belton, P. S., Beta, T., & Duodu, K. G. (2014). Increasing the utilisation of sorghum, millets and pseudocereals: Developments in the science of their phenolic phytochemicals, biofortification and protein functionality. *Journal of Cereal Science*, 59(3), 257-275.
- Taranto, M. P., Vera, J. L., Hugenholtz, J., De Valdez, G. F., & Sesma, F. (2003). *Lactobacillus reuteri* CRL1098 produces cobalamin. *Journal of bacteriology*, 185(18), 5643-5647.
- Thierry, A., Deutsch, S. M., Falentin, H., Dalmaso, M., Cousin, F. J., & Jan, G. (2011). New insights into physiology and metabolism of *Propionibacterium freudenreichii*. *International Journal of Food Microbiology*, 149(1), 19-27.
- Watanabe, F. (2007). Vitamin B12 sources and bioavailability. *Experimental Biology and Medicine*, 232(10), 1266-1274.
- Watanabe, F., Yabuta, Y., Bito, T., & Teng, F. (2014). Vitamin B12-containing plant food sources for vegetarians. *Nutrients*, 6(5), 1861-1873.
- Wolkers – Rooijackers, J. C. M., Endika, M. F., & Smid, E. J. (2018). Enhancing vitamin B 12 in lupin tempeh by *in situ* fortification. *LWT - Food Science and Technology*, 96, 513-518.
- Xie, C., Coda, R., Chamlagain, B., Edelmann, M., Deptula, P., Varmanen, P., . . . Katina, K. (2018). In situ fortification of vitamin B12 in wheat flour and wheat bran by fermentation with *Propionibacterium freudenreichii*. *Journal of Cereal Science*, 81, 133-139.
- Xie, C., Coda, R., Chamlagain, B., & Varmanen, P. (2019). Co-fermentation of *Propionibacterium freudenreichii* and *Lactobacillus brevis* in wheat bran for *in situ* production of vitamin B12. *Frontiers in microbiology*, 10, 1541.
- Ye, K., Shijo, M., Jin, S., & Shimizu, K. (1996). Efficient production of vitamin B 12 from propionic acid bacteria under periodic variation of dissolved oxygen concentration. *Journal of Fermentation and Bioengineering*, 82(5), 484-491.

Table 1. Nutrient content (g/100 g) and cobalt content (ng/g) of the materials and dry matter in batters (%).

Materials and their origin	Protein	Fat	Fiber	Available Carbohydrates	Cobalt	Ratio with water
Rye bran, Finland	15	4	39	26	20	15%
Oat bran, Finland	18	8	14	49	17	15%
Rice bran, USA	13	20	20	33	166	15%
Sorghum flour, Burkina Faso	10	5	10	65	183	30%
Millet flour, Burkina Faso	9	5	7	68	157	30%
Buckwheat bran, Europe	40	9	9	38	183	20%
Quinoa flour, Peru	14	6	6	57	85	25%
Amaranth flour, Germany	14	6	9	66	182	25%
Faba bean flour, Finland	31	2	9	42	696	30%
Soy bean flour, China	38	20	13	18	68	20%
Lupine flour, France	43	12	28	10	29	20%

Table 2. pH values and total titratable acids (TTA; mL) of batters during fermentation in screening of grain materials for vitamin B12 production. The results are expressed as the mean value (n = 3).

Batters	pH			TTA		
	Day 0	Day 1	Day 3	Day 0	Day 1	Day 3
Rye bran	6.6 ^{ez}	3.9 ^{ay}	3.6 ^{ax}	2.3 ^{cx}	12.4 ^{dy}	17.4 ^{cz}
Oat bran	6.5 ^{dez}	5.9 ^{fy}	5.2 ^{ex}	0.5 ^{ax}	1.7 ^{ay}	6.1 ^{az}
Rice bran	6.8 ^{fz}	6.5 ^{gy}	5.0 ^{dx}	0.9 ^{ax}	1.3 ^{ay}	12.5 ^{by}
Sorghum flour	6.2 ^{cz}	5.6 ^{ey}	5.3 ^{ex}	1.8 ^{bx}	3.1 ^{by}	4.4 ^{az}
Millet flour	6.0 ^{by}	3.7 ^{ax}	3.6 ^{ax}	3.7 ^{dx}	11.6 ^{dy}	17.3 ^{cz}
Buckwheat bran	6.5 ^{dy}	4.9 ^{dx}	4.9 ^{dx}	2.6 ^{cx}	14.9 ^{ey}	18.5 ^{cdz}
Quinoa flour	6.0 ^{bz}	3.9 ^{ay}	3.6 ^{ax}	4.7 ^{ex}	18.4 ^{fy}	21.9 ^{ez}
Amaranth flour	6.3 ^{cdz}	4.2 ^{by}	3.7 ^{ax}	3.3 ^{dx}	15.5 ^{ey}	22.1 ^{ez}
Faba bean flour	6.4 ^{dz}	4.5 ^{cy}	4.3 ^{bx}	4.4 ^{ex}	14.8 ^{ey}	19.5 ^{dz}
Soy bean flour	6.7 ^{fz}	6.1 ^{fy}	4.7 ^{cx}	2.6 ^{cx}	3.3 ^{by}	13.5 ^{bz}
Lupine flour	5.7 ^{ay}	5.6 ^{ey}	4.9 ^{dx}	4.8 ^{ex}	5.7 ^{cy}	11.3 ^{bz}

Values in the same column (a-f) and same row (x-z) bearing different superscripts are significantly different ($p < 0.05$).

Table 3. Vitamin B12 content (ng/g, dw) of batters during fermentation in screening of grain materials for vitamin B12 production. The results are expressed as the mean $\pm$ standard deviation (n = 3).

Batters	Day 0	Day 1	Day 3
Rye bran	38 $\pm$ 5 ^{cx}	103 $\pm$ 7 ^{cy}	104 $\pm$ 29 ^{by}
Oat bran	36 $\pm$ 4 ^{cx}	128 $\pm$ 3 ^{dy}	332 $\pm$ 24 ^{dz}
Rice bran	37 $\pm$ 5 ^{cx}	133 $\pm$ 16 ^{dy}	742 $\pm$ 18 ^{gz}
Sorghum flour	20 $\pm$ 3 ^{ax}	149 $\pm$ 9 ^{dey}	265 $\pm$ 13 ^{cz}
Millet flour	19 $\pm$ 3 ^{ax}	58 $\pm$ 1 ^{ay}	51 $\pm$ 6 ^{ay}
Buckwheat bran	29 $\pm$ 4 ^{bx}	335 $\pm$ 76 ^{gy}	631 $\pm$ 61 ^{fz}
Quinoa flour	32 $\pm$ 4 ^{bx}	60 $\pm$ 1 ^{ay}	65 $\pm$ 5 ^{ay}
Amaranth flour	31 $\pm$ 4 ^{bx}	81 $\pm$ 13 ^{by}	78 $\pm$ 4 ^{ay}
Faba bean flour	20 $\pm$ 5 ^{cx}	155 $\pm$ 23 ^{ey}	298 $\pm$ 51 ^{cdz}
Soy bean flour	29 $\pm$ 6 ^{bx}	154 $\pm$ 2 ^{ey}	407 $\pm$ 5 ^{ez}
Lupine flour	28 $\pm$ 4 ^{bx}	246 $\pm$ 15 ^{fy}	343 $\pm$ 71 ^{dz}

Values in the same column (a-g) and same row (x-z) bearing different superscripts are significantly different (p < 0.05).

Table 4. Cell counts of propionic acid bacteria (PAB), lactic acid bacteria (LAB) and *Enterobacteriaceae* during fermentation with different level of starter cultures (log CFU/g). The results are expressed as the mean $\pm$ standard deviation (n = 3).

	PAB			LAB			<i>Enterobacteriaceae</i>		
	Day 0	Day 1	Day 3	Day 0	Day 1	Day 3	Day 0	Day 1	Day 3
Oat bran									
Control	-*	-	-	-*	7.1 $\pm$ 0.1 ^{ax}	9.1 $\pm$ 0.1 ^{ay}	-*	7.6 $\pm$ 0.1 ^{dy}	4.7 $\pm$ 0.2 ^{bx}
P9/L6	8.7 $\pm$ 0.1 ^{ax}	9.3 $\pm$ 0.1 ^{ay}	9.4 $\pm$ 0.3 ^{ay}	5.7 $\pm$ 0.1 ^{ax}	8.4 $\pm$ 0.1 ^{by}	9.2 $\pm$ 0.1 ^{az}	-	6.4 $\pm$ 0.2 ^{cy}	4.4 $\pm$ 0.2 ^{bx}
P9/L7	8.8 $\pm$ 0.1 ^{ax}	9.3 $\pm$ 0.1 ^{ay}	9.3 $\pm$ 0.1 ^{ay}	6.9 $\pm$ 0.1 ^{bx}	8.3 $\pm$ 0.1 ^{by}	9.3 $\pm$ 0.1 ^{az}	-	4.8 $\pm$ 0.2 ^{by}	3.9 $\pm$ 0.1 ^{ax}
P9/L8	8.7 $\pm$ 0.1 ^{ax}	9.2 $\pm$ 0.1 ^{ay}	9.3 $\pm$ 0.2 ^{ay}	8.1 $\pm$ 0.1 ^{cx}	9.4 $\pm$ 0.1 ^{cy}	9.3 $\pm$ 0.1 ^{ay}	-	4.3 $\pm$ 0.1 ^a	-
Rice bran									
Control	-	-	-	-	7.4 $\pm$ 0.1 ^{ax}	8.1 $\pm$ 0.2 ^{ay}	-	7.8 $\pm$ 0.1 ^{cy}	8.7 $\pm$ 0.2 ^{bx}
P9/L6	8.8 $\pm$ 0.0 ^{ax}	9.4 $\pm$ 0.0 ^{ay}	9.5 $\pm$ 0.1 ^{ay}	5.8 $\pm$ 0.1 ^{ax}	8.1 $\pm$ 0.1 ^{by}	8.9 $\pm$ 0.2 ^{bz}	-	5.4 $\pm$ 0.2 ^{by}	3.5 $\pm$ 0.4 ^{ax}
P9/L7	8.7 $\pm$ 0.2 ^{ax}	9.2 $\pm$ 0.2 ^{ay}	9.3 $\pm$ 0.0 ^{ay}	7.0 $\pm$ 0.0 ^{bx}	8.0 $\pm$ 0.2 ^{by}	9.3 $\pm$ 0.2 ^{cz}	-	4.2 $\pm$ 0.2 ^a	-
P9/L8	8.8 $\pm$ 0.1 ^{ax}	9.3 $\pm$ 0.1 ^{ay}	9.3 $\pm$ 0.2 ^{ay}	8.0 $\pm$ 0.1 ^{cx}	9.1 $\pm$ 0.1 ^{cy}	9.5 $\pm$ 0.2 ^{cz}	-	4.3 $\pm$ 0.1 ^a	-
Buckwheat bran									
Control	-	-	-	2.8 $\pm$ 0.2 ^{ax}	8.4 $\pm$ 0.0 ^{ay}	9.4 $\pm$ 0.1 ^{az}	4.4 $\pm$ 0.2 ^{ax}	8.6 $\pm$ 0.1 ^{dz}	5.4 $\pm$ 0.2 ^{by}
P9/L6	8.8 $\pm$ 0.1 ^{ax}	9.4 $\pm$ 0.1 ^{by}	9.4 $\pm$ 0.1 ^{by}	5.7 $\pm$ 0.2 ^{bx}	9.1 $\pm$ 0.1 ^{by}	9.3 $\pm$ 0.0 ^{az}	4.3 $\pm$ 0.2 ^{ax}	6.5 $\pm$ 0.2 ^{cz}	5.1 $\pm$ 0.0 ^{ay}
P9/L7	8.7 $\pm$ 0.2 ^{ax}	9.3 $\pm$ 0.1 ^{aby}	9.3 $\pm$ 0.1 ^{by}	7.2 $\pm$ 0.1 ^{cx}	9.2 $\pm$ 0.1 ^{by}	9.3 $\pm$ 0.1 ^{ay}	4.4 $\pm$ 0.1 ^{ax}	5.5 $\pm$ 0.1 ^{by}	-
P9/L8	8.7 $\pm$ 0.1 ^{ax}	9.1 $\pm$ 0.2 ^{ay}	9.0 $\pm$ 0.2 ^{ay}	8.1 $\pm$ 0.2 ^{dx}	9.1 $\pm$ 0.1 ^{by}	9.4 $\pm$ 0.2 ^{ay}	4.3 $\pm$ 0.1 ^{ax}	5.1 $\pm$ 0.2 ^{ay}	-
Sorghum flour									
Control	-	-	-	2.3 $\pm$ 0.3 ^{ax}	6.6 $\pm$ 0.1 ^{ay}	7.9 $\pm$ 0.1 ^{az}	5.4 $\pm$ 0.1 ^{ax}	9.0 $\pm$ 0.1 ^{cy}	8.9 $\pm$ 0.1 ^{ay}
P9/L6	8.7 $\pm$ 0.2 ^{ax}	8.8 $\pm$ 0.1 ^{ax}	9.1 $\pm$ 0.2 ^{by}	5.9 $\pm$ 0.0 ^{bx}	8.1 $\pm$ 0.1 ^{by}	8.1 $\pm$ 0.1 ^{ay}	5.5 $\pm$ 0.2 ^{ax}	8.8 $\pm$ 0.1 ^{cy}	9.2 $\pm$ 0.2 ^{ay}
P9/L7	8.8 $\pm$ 0.1 ^{ax}	9.0 $\pm$ 0.1 ^{ax}	8.9 $\pm$ 0.1 ^{ax}	6.8 $\pm$ 0.2 ^{cx}	8.9 $\pm$ 0.1 ^{by}	9.1 $\pm$ 0.1 ^{by}	5.4 $\pm$ 0.0 ^{ax}	7.9 $\pm$ 0.1 ^{by}	-
P9/L8	8.7 $\pm$ 0.1 ^{ax}	8.9 $\pm$ 0.1 ^{ax}	8.9 $\pm$ 0.2 ^{ax}	7.9 $\pm$ 0.1 ^{dx}	8.8 $\pm$ 0.1 ^{by}	9.0 $\pm$ 0.1 ^{by}	5.5 $\pm$ 0.1 ^{ax}	5.4 $\pm$ 0.1 ^{ax}	-

* Not detected

Control, spontaneously fermented batter; P9/L6, P9/L7 and P9/L8 mean the batters with 9.0 log CFU/g of *P. freudenreichii* and 6.0 log CFU/g, 7.0 log CFU/ and 8.0 log CFU/g of *L. brevis*, respectively.

Values in the same column and batter type bearing different superscripts (a-c) are significantly different (p < 0.05). Values in the same row bearing different superscripts (x-z) are significantly different (p < 0.05).

Table 5. pH, total titratable acids (TTA; mL) and vitamin B12 content (ng/g, dw) during fermentation with different level of starter cultures. The results of pH and TTA are expressed as the mean. The results of vitamin B12 content are expressed as the mean $\pm$ standard deviation (n = 3).

	pH			TTA			Vitamin B12		
	Day 0	Day 1	Day 3	Day 0	Day 1	Day 3	Day 0	Day 1	Day 3
Oat bran									
Control	6.5 ^{az}	6.2 ^{cy}	5.8 ^{cx}	0.5 ^{ax}	1.0 ^{ay}	2.1 ^{az}	-*	-	-
P9/L6	6.5 ^{az}	5.8 ^{by}	5.2 ^{bx}	0.5 ^{ax}	1.6 ^{by}	5.4 ^{bz}	38 $\pm$ 5 ^{ax}	119 $\pm$ 7 ^{ay}	342 $\pm$ 14 ^{bz}
P9/L7	6.5 ^{az}	5.4 ^{ay}	4.5 ^{ax}	0.5 ^{ax}	4.8 ^{cy}	6.1 ^{bz}	38 $\pm$ 5 ^{ax}	104 $\pm$ 18 ^{ay}	312 $\pm$ 3 ^{az}
P9/L8	6.5 ^{az}	5.4 ^{ay}	4.4 ^{ax}	0.5 ^{ax}	4.7 ^{cy}	6.3 ^{bz}	38 $\pm$ 5 ^{ax}	128 $\pm$ 12 ^{ay}	298 $\pm$ 10 ^{az}
Rice bran									
Control	6.8 ^{az}	6.4 ^{by}	5.3 ^{bx}	0.9 ^{ax}	1.3 ^{ay}	11.2 ^{az}	-	-	-
P9/L6	6.8 ^{az}	6.5 ^{by}	5.1 ^{abx}	0.9 ^{ax}	1.1 ^{ay}	11.9 ^{bz}	38 $\pm$ 5 ^{ax}	112 $\pm$ 16 ^{ay}	722 $\pm$ 21 ^{bz}
P9/L7	6.8 ^{az}	6.4 ^{by}	5.0 ^{ax}	0.9 ^{ax}	1.8 ^{ay}	13.1 ^{bz}	38 $\pm$ 5 ^{ax}	119 $\pm$ 24 ^{ay}	717 $\pm$ 10 ^{bz}
P9/L8	6.8 ^{az}	5.9 ^{ay}	4.9 ^{ax}	0.9 ^{ax}	3.9 ^{by}	16.4 ^{cz}	38 $\pm$ 5 ^{ax}	122 $\pm$ 3 ^{ay}	675 $\pm$ 30 ^{az}
Buckwheat bran									
Control	6.5 ^{az}	5.3 ^{cy}	4.9 ^{bx}	2.6 ^{ax}	12.4 ^{ay}	17.0 ^{az}	-	-	-
P9/L6	6.5 ^{az}	4.9 ^{bx}	4.9 ^{bx}	2.6 ^{ax}	14.3 ^{by}	17.7 ^{bz}	29 $\pm$ 4 ^{ax}	312 $\pm$ 76 ^{by}	604 $\pm$ 41 ^{bz}
P9/L7	6.5 ^{ay}	4.7 ^{ax}	4.6 ^{ax}	2.6 ^{ax}	19.1 ^{cy}	20.0 ^{cz}	29 $\pm$ 4 ^{ax}	237 $\pm$ 38 ^{ay}	572 $\pm$ 58 ^{abz}
P9/L8	6.5 ^{ay}	4.6 ^{ax}	4.6 ^{ax}	2.6 ^{ax}	19.8 ^{cy}	20.7 ^{cz}	29 $\pm$ 4 ^{ax}	256 $\pm$ 28 ^{ay}	508 $\pm$ 26 ^{az}
Sorghum flour									
Control	6.2 ^{az}	5.7 ^{by}	5.4 ^{cx}	1.8 ^{ax}	2.4 ^{ay}	3.7 ^{az}	-*	-	-
P9/L6	6.2 ^{az}	5.6 ^{by}	5.3 ^{cx}	1.8 ^{ax}	3.2 ^{ay}	4.3 ^{az}	19 $\pm$ 3 ^{ax}	151 $\pm$ 16 ^{cy}	249 $\pm$ 11 ^{cz}
P9/L7	6.2 ^{az}	4.7 ^{ay}	4.5 ^{bx}	1.8 ^{ax}	4.8 ^{by}	7.8 ^{bz}	19 $\pm$ 3 ^{ax}	88 $\pm$ 9 ^{by}	125 $\pm$ 27 ^{bz}
P9/L8	6.2 ^{az}	4.6 ^{ay}	3.9 ^{ax}	1.8 ^{ax}	5.5 ^{cy}	9.2 ^{cz}	19 $\pm$ 3 ^{ax}	68 $\pm$ 6 ^{ay}	67 $\pm$ 14 ^{ay}

* Not detected

Control, spontaneously fermented batter; P9/L6, P9/L7 and P9/L8 mean the batters with 9.0 log CFU/g of *P. freudenreichii* and 6.0 log CFU/g, 7.0 log CFU/ and 8.0 log CFU/g of *L. brevis*, respectively.

Values in the same column and batter type bearing different superscripts (a-c) are significantly different ($p < 0.05$). Values in the same row bearing different superscripts (x-z) are significantly different ($p < 0.05$).

Figure legends

Figure 1. Contents (mg/g dw) of lactic acid (■), acetic acid (▨) and propionic acid (■) during fermentation in screening of grain materials for vitamin B12 production. The results are expressed as the mean value of three replicates and error bars indicate standard deviation of total acid content.

Figure 2. Partial least square regression (PLS) loading plots showing the distribution of x variables and y variables along the first and second principal components (PC) on day 1 (A) and day 3 (B). In each plot, the x variables are pH, total titratable acid (TTA) and content of protein, fat, fiber, available carbohydrates, cobalt, lactic acid (LA), acetic acid (AA) and propionic acid (PA). The y variable is production of vitamin B12. The projection of each x variable on the line drawn via the y variable and the origin of the plot describes the correlation of the x variables and the y variable; distance of each x variable's projection from the origin indicate its influence on the y variable.

Figure 1.

Figure 2.

Journal Pre-proof

1. More than 300 ng/g of vitamin B12 was produced in various materials
2. The highest vitamin B12 production was found in the rice bran fermentation
3. Nutrient composition of materials significantly influenced vitamin B12 production
4. *L. brevis* can inhibit the growth of *Enterobacteriaceae* during fermentation

The authors declared that no conflict of interest exists in the submission of this manuscript for publication.

Journal Pre-proof