<https://helda.helsinki.fi>

First record of the bacterial endosymbiont Wolbachia for phytophagous hoverflies from genus Merodon (Diptera : Syrphidae)

Sasic Zoric, Ljiljana

2019-09

Sasic Zoric , L , Stahls , G & Dan , M 2019 , ' First record of the bacterial endosymbiont Wolbachia for phytophagous hoverflies from genus Merodon (Diptera : Syrphidae) ' , Entomological Science , vol. 22 , no. 3 , pp. 283-296 . <https://doi.org/10.1111/ens.12361>

<http://hdl.handle.net/10138/323604>

<https://doi.org/10.1111/ens.12361>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

This is the accepted version of the following article:

Ljiljana Šašić Zorić, Gunilla Ståhls, Mihajla Đan (2019) First record of the bacterial endosymbiont *Wolbachia* for phytophagous hoverflies from genus *Merodon* (Diptera, Syrphidae). Entomological Science, 22: 283-296.,

which has been published in final form at [<https://doi.org/10.1111/ens.12361>]. This article may be used for non-commercial purposes in accordance with the Wiley Self-Archiving Policy [<http://www.wileyauthors.com/self-archiving>].

**First record of the bacterial endosymbiont *Wolbachia* for phytophagous hoverflies from
genus *Merodon* (Diptera, Syrphidae)**

Ljiljana Šašić Zorić¹, Gunilla Ståhls² and Mihajla Đan³

¹ BioSense Institute, University of Novi Sad, Dr Zorana Đinđića 1, 21000 Novi Sad, Serbia, ²
Zoology Unit, Finnish Museum of Natural History, PO Box 17, 00014 University of
Helsinki, Finland, ³ Department of Biology and Ecology, Faculty of Sciences, University of
Novi Sad, Trg Dositeja Obradovića 2, 21000 Novi Sad, Serbia

Correspondence: Ljiljana Šašić Zorić, BioSense Institute, University of Novi Sad, Zorana
Đinđića 1, 21000 Novi Sad, Serbia
Email: ljiljana.sasic@dbe.uns.ac.rs

Abstract. *Wolbachia* is a widespread bacterial endosymbiont among arthropod species. It influences the reproduction of the host species and also mtDNA diversity. Until now there were only a few studies which detected *Wolbachia* infections in hoverflies (Diptera, Syrphidae), and this is the first broader study with the aim to examine the incidence of *Wolbachia* in the hoverfly genus *Merodon*. The obtained results indicate an infection rate of 96% and the presence of both *Wolbachia* supergroup A and B, which are characteristic for most of the infected arthropod species. Additionally, the presence of multiple *Wolbachia* strains in the *M. aureus* group species was detected and the mtDNA *COI* based relationships of the group were discussed in the light of infection. Finally, we discuss plant mediated horizontal transmission of *Wolbachia* strains among the studied hoverfly species.

Keywords: 16S rRNA gene, *Drimia maritima*, *Merodon aureus* group, *wsp*.

INTRODUCTION

The hoverflies (Diptera, Syrphidae) are a widely distributed insect family present almost everywhere in the world except on the Antarctic and remote oceanic islands (Thompson & Rotheray, 1988). The hoverfly genus *Merodon* Meigen, 1903 (subfamily Eristalinae) is distributed over Palaearctic and Palaetropical regions (Hurkmans 1993). The immature stages of *Merodon* species develop in and feed on underground storage organs of geophytes of the families Asparagaceae, Iridaceae and Amaryllidaceae (Andrić *et al.* 2014; Ricarte *et al.* 2008, 2017; Preradović *et al.* 2018). Adults morphologically mimic hymenopterans and feed on pollen and nectar (Hurkmans 1993). The genus comprises more than 160 species (Ståhls *et al.* 2009; Vujić *et al.* 2012), however, the real number of *Merodon* species is still unknown considering high level of diversity and detected presence of cryptic species (e.g. Milankov *et al.* 2008, 2009; Radenković *et al.* 2011; Vujić *et al.* 2012; Popović *et al.* 2015; Ačanski *et al.* 2016; Šašić *et al.* 2016, 2018; Veselić *et al.* 2017).

The *Merodon aureus* hoverfly species group is taxonomically especially challenging, as it comprises a high genetic diversity with minor or lacking differences in morphological traits (Šašić *et al.* 2016). The group comprises species morphologically close to *M. aureus* Fabricius, 1805. The members of the taxa are small sized (8-13 mm), with a short, rounded abdomen, a distinct spike on the metatrochanter in males and a characteristic structure of the male genitalia (Vujić *et al.* 2007; Radenković *et al.* 2011). Until recently, the group comprised altogether 18 previously-known and newly-discovered taxa from the Mediterranean region and southern European mountain regions (Marcos-García *et al.* 2007; Vujić *et al.* 2007; Milankov *et al.* 2008; Radenković *et al.* 2011; Speight, 2014), while new data indicates the presence of additional species (Šašić *et al.* 2016, 2018; Veselić *et al.* 2017; Radenković *et al.* 2018).

The first results about molecular diversity of *Merodon aureus* group were based on analyses of 3' and 5' fragments of the mtDNA *COI* gene and suggested the presence of multiple cryptic species complexes within the group. The morphological character states usually used in taxonomy of hoverflies mostly failed to discern these potential species (Šašić *et al.* 2016; Radenković *et al.* 2017). However, subtle differences in wing and surstylus shape were detected using geometric morphometry (see Ačanski *et al.* 2016; Šašić *et al.* 2016; Radenković *et al.* 2017). Maximum likelihood (ML) analysis of *COI* sequences including representatives of all potential complexes from *M. aureus* species group revealed deep divergences between morphologically close species (Fig. 1). The obtained molecular evidence showed clear conflict with the morphologically defined subgroups and/or species complexes (Vujić, personal communication; Šašić *et al.* 2016; Radenković *et al.* 2017).

Figure 1.

Over the last few decades, mtDNA has been the most popular marker for quantifying molecular diversity, as the marker contains a combination of technical benefits (ease of amplification), and supposed biological and evolutionary advantages such as clonality, near-neutrality and often clocklike nature of its substitution rate. However, mtDNA is not always clonal, not neutrally evolving and not clocklike, which brings into question its use in recovering recent species and population histories (e.g. Galtier *et al.* 2009). In addition to these limitations of use, the taxonomic utility of the maternally inherited mitochondrial genome could be compromised by the presence of symbiotic bacteria, which pass from a female to its offspring (Galtier *et al.* 2009). The most important of the so-called “reproductive parasites” is *Wolbachia pipientis* (Alphaproteobacteria: Rickettsiales: Rickettsiaceae), which is facultative endosymbiont estimated to have infected more than half of arthropod species (Weinert *et al.* 2015). Although other bacterial reproductive parasites are also known (*Cardinium*, *Arsenophonus*, *Rickettsia*, *Spiroplasma*), *Wolbachia* is the most abundant

endosymbiont among arthropod species and with broadest range of host reproductive phenotypes including induction of cytoplasmic incompatibility, feminisation of genetic males, parthenogenesis and male killing (Duron *et al.* 2008; Zug & Hammerstein, 2012).

According to molecular phylogenetic analyses *Wolbachia pipientis* has been divided into seventeen clades (A-R, except G which is recombinant of A and B supergroups), termed supergroups (Werren *et al.* 1995; Bandi *et al.* 1998; Vandekerckhove *et al.* 1999; Lo *et al.* 2002, 2007; Czarnetzki & Tebbe 2004; Baldo & Werren, 2007; Bordenstein *et al.* 2009; Haegeman *et al.* 2009; Ros *et al.* 2009; Augustinos *et al.* 2011; Bing *et al.* 2014; Glowska *et al.* 2015; Wang *et al.* 2016). The supergroups taxonomic status was discussed in Ramírez-Puebla *et al.* (2015) who suggested that *Wolbachia* supergroups represent separate evolutionary lineages and that they should be designated as species. They also indicated that some of the supergroups could contain more than one *Wolbachia* species. The proposed nomenclature is criticized by Lindsey *et al.* (2016) as inadequate and confusing.

Wolbachia is probably the most widespread endosymbiont of arthropods and nematodes (Charlat *et al.* 2003; Werren *et al.* 2008). Recent studies estimated 19% to 76% infection rates of *Wolbachia* among arthropod species (Jeyaparakash & Hoy, 2000; Werren & Windsor, 2000; Hilgenboecker *et al.* 2008; Werren *et al.* 2008; Simões *et al.* 2011; Weinert *et al.* 2015; Zug & Hammerstein, 2012). The evolutionary success is achieved through a variety of effects on host biology, ranging from manipulation of reproduction in favor of females to mutualistic interactions with host species. *Wolbachia* interact with host sex-determination systems and the cell cycle, and its effect on host populations can frame sexual behaviors and species diversity (Charlat *et al.* 2003). The infection is maternally inherited via infection of developing oocytes or it can be a consequence of horizontal transmission (Werren, 1997). It is most likely to find *Wolbachia* in ovaries, although it can also occur at high intensities in the

fat body and other tissues (e. g. Werren, 1997; Dobson *et al.* 1999; Albertson *et al.* 2009; Pietri *et al.* 2016).

The first test for the presence of *Wolbachia* in hoverfly species (Syrphidae) was the study of Werren and Windsor (2000) who found that the Nearctic species *Milesia virginiensis* tested negative for the presence of *ftsZ* bacterial cell-cycle gene of *Wolbachia*. In 2006, Sintupachee *et al.* found *Syrpitta rufifacies* negative for the *ftsZ*, but the species *Graptomyza brevirostris* (Eristalinae: Volucellini) tested positive for both *ftsZ* and a *Wolbachia* surface protein (*wsp*) genes. Evison *et al.* (2012) screened pollinator groups in the UK for different groups of parasites including *Wolbachia*, and among the tested species they included four species of hoverflies, *Rhingia campestris* (Eristalinae: Rhingiini), *Eristalis arbustorum* and *E. tenax* (Eristalinae: Eristalini) and *Episyrphus balteatus* (Syrphinae: Syrphini), which all were positive for the tested *CoxA* primers. However, the current molecular taxonomy of hoverflies and applying mitochondrial markers neglect the potential bias of *Wolbachia* on the results and do not contain any *Wolbachia* screening test.

In this study, we estimate the incidence of *Wolbachia* in the genus *Merodon* (Diptera, Syrphidae) using samples from recent field collections across South European countries, Austria, Romania, Turkey, Iran, Morocco and the South African Republic. We amplified and sequenced bacterial marker genes (16S rRNA gene and *wsp* gene) with the aim to assign *Wolbachia* supergroups present in *Merodon* hoverflies. We particularly focus on the screening of *M. aureus* group species in the light of the observed high mtDNA *COI* gene variability, which is incongruent with morphological invariability in several species complexes, and test for coevolution between *M. aureus* group and *Wolbachia* strains infecting the species of the group. Additionally, we performed screening of host plant bulb with the aim to prove the presence of *Wolbachia* in plant tissue and discuss potential horizontal transmission via bulb.

MATERIAL AND METHODS

Specimens analyzed

The hoverfly specimens were collected from 2012 to 2016 and identified to the species, or subgroup (*Merodon aureus* group specimens) level (by Dr Ante Vujić and according to Šašić *et al.* 2016; Radenković *et al.* 2017). The bulb of *Drimia maritima* (syn. *Urginea maritima*) which is a host plant of *M. luteihumerus* larvae was collected in March 2017. All the data about collected samples are provided in Table S1.

DNA extraction

DNA extractions of 2 - 3 legs and separately of abdomens of the hoverfly specimens was performed by using the SDS extraction protocol according to Chen *et al.* (2010). The gDNA extracted from legs were used for *Wolbachia* specific 16SrRNA gene amplification, while the gDNA extracted from abdomens were used for *Wolbachia wsp* gene amplification. The main reason for repeated gDNA extraction was low amplification success of *wsp* gene using gDNA extracted from legs, which is probably a consequence of lower amount of bacterial DNA in legs comparing to the abdomen (as previously mentioned, the highest concentration of *Wolbachia* is in reproductive tissue).

Testing for the presence of *Wolbachia*

16S rRNA gene amplification

Primary screening on *Wolbachia* presence was based on the amplification of *Wolbachia*'s 16S rRNA gene fragment. In total, 74 specimens belonging to different *Merodon* species

were screened for *Wolbachia* presence based on amplification and sequencing of bacterial 16S rRNA gene. 45 of these belong to *M. aureus* group, with fewer samples from the following species groups: five from *M. avidus* group, five from *M. nanus* group, three from *M. geniculatus* group, three from *M. albifrons* group, three from *M. constans* group, three from *M. natans* group, three from *M. nigritarsis*, three from *M. desuturinus* group, and one from the species *M. luteihumerus*. We tested 1-3 specimens per species (Table 1).

16S rRNA gene fragment was amplified using WspecF and WspecR primer pair (Werren & Windsor, 2000). Polymerase chain reactions (PCR) were carried out in 25 µl reaction volumes. The reaction mixture contained 1x Taq Buffer without MgCl₂ (ThermoScientific, Lithuania), 1.5 mM MgCl₂, 0.25 mM of each nucleotide, 1.25 U Taq polymerase (ThermoScientific, Lithuania), 7 pmol of each primer, and approximately 50-100 ng template DNA. The amplification of the bacterial 16S rRNA gene was carried out following the protocol described in Werren & Windsor (2000).

The PCR products were checked on 1.5% agarose gels and the PCR product from gDNA of *Drosophila melanogaster* extracted from line 5 from Bloomington stock center <http://flystocks.bio.indiana.edu/Reports/5.html> (project number: OI 173012) was used as a positive control. Additionally, we also used PCR reaction mixture without gDNA as negative control in order to eliminate potential contamination.

Wsp gene amplification

In addition to 16S rRNA gene, we tested *Merodon aureus* group specimens on bacterial *wsp* gene. For this purpose, we extracted additional genomic DNA from the abdomen of hoverflies (see above). We used Phire Animal Tissue Direct PCR Master Mix (ThermoScientific, Lithuania) to amplify *wsp* gene according to the manufacturer's

instructions. The same kit was used for direct *wsp* gene amplification from the tissue of *Drimia maritima* bulb and the larvae of *M. luteihumerus* discovered within the bulb. The primers used to amplify the *wsp* fragment are 136F, 691R, 81F, 522R (Zhou *et al.* 1998). The PCR was performed with three primer pair combinations: 136F/691R for *Wolbachia* supergroup A, 81F/ 522R for *Wolbachia* supergroup B, and 81F/691R for both supergroups (Zhou *et al.* 1998). Initially, we screened all samples with *wsp* primer combination for supergroup A. The samples without products were additionally tested with *wsp* primer combination for supergroup B or universal combination for both supergroups. Only amplification products with a single band on 1.5% agarose gels were used for sequencing.

Sequencing

The PCR products are enzymatically purified using exonuclease I and shrimp alkaline phosphatase enzymes. Sequencing was done in both directions using the BigDye Terminator v.3.1 cycle sequencing kit (Applied Biosystems, Foster City, CA, USA) on ABI3730xl DNA Analyzer (Applied Biosystems, Foster City, Ca, USA) at the Sequencing Service Laboratory of the Finnish Institute for Molecular Medicine (FIMM), Helsinki, Finland.

16S rRNA gene and wsp gene sequences analyses

The produced 16S rRNA gene and *wsp* gene sequences were blasted against the nucleotide collection database at the National Center for Biotechnology Information (NCBI; www.ncbi.nlm.nih.gov) using Megablast optimized for highly similar sequences. *Wsp* sequences were also checked against the *Wolbachia* MLST database (Baldo *et al.* 2006). Finally, the screening results were presented in a form of a table with marked specimens where *Wolbachia* was identified using 16S rRNA gene, *wsp* or both genes (Table 1). All

213 sequences have been submitted in GenBank under accession numbers MK184213 –
 214 MK184277 (16S rRNA gene) and MK192943 – MK192981 (*wsp*), while *wsp* sequences are
 215 additionally deposited in MLST database (Table 2, Table S1).

216 In order to place *Wolbachia* detected in *Merodon* specimens in a particular supergroup, we
 217 constructed ML tree based on 16S rRNA gene sequences. The sequences were manually
 218 aligned and the tree was constructed using RAxML 8.2.8 (Stamatakis, 2014) through the
 219 CIPRES Science Gateway web portal (Miller *et al.* 2010) and by applying the general time-
 220 reversible (GTR) evolutionary model with gamma distribution (Rodriguez *et al.* 1990), while
 221 the statistical support for the clades was assessed using the rapid bootstrap method with 1000
 222 replicates. The analysis also included 29 sequences belonging to *Wolbachia* supergroups (A,
 223 B, F, H, I, M, N, O) founded in insect hosts, which were downloaded from GenBank (see
 224 Table S2). As outgroups, we used two species of α -Proteobacteria: *Ehrlichia canis* and
 225 *Anaplasma marginale* (GenBank accession numbers: M73226, M60313), and the tree was
 226 rooted on *Anaplasma marginale*. In order to test cophylogeny between *M. aureus* group *COI*
 227 tree and *Wolbachia* 16S rRNA gene tree, we applied Procrustean Approach to Cophylogeny
 228 (PACo) in R environment (R Core Team, 2018) as described in Balbuena *et al.* (2013). As
 229 input data, we used unrooted ML trees. For 16S rRNA gene we firstly determined sequence
 230 types by using DnaSP 5 software (Librado & Rozas, 2009) which was used for an unrooted
 231 ML tree construction in RAxML 8.2.8 (Stamatakis, 2014).

232 For *wsp* sequences, alignment was performed using the L-INS-I strategy as implemented in
 233 MAFFT (Katoh & Standley, 2013) available on the EMBL-EBI bioinformatics framework
 234 (McWilliam *et al.* 2013). The total number of alleles was determined by using DnaSP 5
 235 software (Librado & Rozas, 2009). The assessments of pairwise differences, uncorrected p
 236 distance values between alleles, were conducted in MEGA7 (Kumar *et al.* 2016). This gene is
 237 not used for phylogeny reconstruction as the evolutionary signal is masked by its mosaic

nature, however, it can be used for strain typing based on a combination of four hypervariable regions (HVRs) (Baldo *et al.* 2005, 2006). Thus, for each *wsp* allele, we determined HVRs profile by checking against the *Wolbachia* MLST database (Baldo *et al.* 2006).

RESULTS

The primary screening on *Wolbachia* using 16S rRNA gene as a marker was performed on 52 species of the genus *Merodon* or 74 specimens from which ten specimens and five species tested negative. Within the *M. aureus* group, three specimens belonging to *M. sapphous* sp. n. 2, *M. aureus* sp. n. 2 and *M. balkanicus* tested negative, within *M. segetum* (*M. natans* group) as well as within *M. melanocerus* (*M. desuturinus* group) two specimens tested positive, while one was negative. *M. albifrons* (*M. albifrons* group) specimens were all negative, while in *M. nanus* (*M. nanus* group) one specimen was positive and one was negative for *Wolbachia* infection. The one tested *M. luteihumerus* specimen were also negative. The screening results are summarized in Table 1.

Table 1.

For ML tree construction we used 16S rRNA gene sequences. The aligned sequence set used in the analysis was 415bp long. All *Wolbachia* 16S rRNA gene sequences produced in this study are resolved as supergroup A, except the *Wolbachia* sequences from *M. neofasciatus* which are resolved with supergroup B sequences (Fig. 2).

Figure 2.

PACo analysis resulted in a residual sum of squares $m^2_{xy} = 0.355$, under the probability value $P=0.064$. Thus, the cophylogeny hypothesis between *M. aureus* group *COI* tree and *Wolbachia* 16S rRNA gene tree was rejected as statistically insignificant ($P>0.05$). The

relationships between *COI* sequences of *M. aureus* group specimens and corresponding *Wolbachia* endosymbionts (based on 16S rRNA gene sequences) are presented in Figure 3.

Figure 3.

In order to achieve better resolution in *Wolbachia* strain determination, we additionally tested *Merodon aureus* group specimens (41 species or 45 specimens) for the *wsp* gene product.

The amplification products were detected in 38 species (42 specimens). *M. nisi*, *M. unicolor* and *M. balkanicus* tested negative. Multiple products of *wsp* amplification were detected in *M. naxius*, *M. andriotes* and *M. puniceus*, and these amplification products were not further processed. In total 39 sequences which correspond to *M. aureus* group specimens were produced, however, four of them were discarded because of poor quality. The *wsp* gene was also amplified and sequenced for *Drimia maritima* bulb, and *M. luteihumerus* larvae (3 specimens) from the host plant bulb (Table 1).

The final *wsp* sequence matrix contained 39 sequences. The aligned sequences were 560bp long, and with gap regions (see Fig. S1). We discovered 7 different *wsp* alleles, from which the A1 was most common among *Merodon aureus* group specimens, but present also in the bulb and *M. luteihumerus* larvae (see Fig. 2). Based on Blast search results the sequences were 99% to 100% identical to previously discovered *Wolbachia* strains *wsp* sequences from different insect hosts, except A6 which is 97% identical to *wsp* sequence from *Ceutorhynchus obstrictus* (cabbage seed pod weevil) (see Table S3). A1 is identical to *Wolbachia wsp* sequences from *Formica sanguinea*, *Formica exsecta* (both ants), *Protocalliphora sialia* (birdnest blowfly), *Conotrachelus nenuphar* (plum curculio), and *Ceutorhynchus obstrictus* (cabbage seed pod weevil). A4 is identical to *Ectemnius continuus* (a wasp species) *wsp* sequence (although the query cover is 96%).

By checking the *wsp* sequences against *Wolbachia* MLST database, we found that A1 sequence is identical to *wsp* allele 311, while rest of the alleles are detected for the first time in this study and they are submitted in the database as new alleles. Additionally, the DNA sequences of all alleles were translated and HVR peptides are determined. The new HVR peptides are submitted in the aforementioned database. The WSP profiles for each of alleles are presented in Table 2.

Table 2.

The number of base differences per site between alleles (uncorrected p distances) is shown in Table 3. The analysis involved 7 nucleotide sequences of *wsp* alleles. All ambiguous positions were removed for each sequence pair. The smallest p distance has been detected between A1 and A2, while the most divergent are A3 and A7.

Table 3.

The *Wolbachia* detection success was similar when comparing PCR amplification between the two applied molecular markers (16S rRNA gene and *wsp* gene) on *Merodon aureus* group specimens for which both markers were used. In both cases 42 out of 45 analyzed specimens had amplification product, in one there was no product (*M. balkanicus* specimen), while in four we got amplification product for only one of the markers. When comparing sequence quality, 10% of *wsp* sequences had low quality and could not be used for further analysis, while all of the 16S rRNA gene sequences were good quality sequences. Low sequence quality could be due to multiple infections by different bacterial strains, but also could be caused by contamination.

DISCUSSION

The presented results indicated a markedly high incidence of *Wolbachia* infection in *Merodon* hoverflies confirmed either by one or both amplified and sequenced *Wolbachia* genes, 16S rRNA or *wsp*. In total 50 out of 52 analyzed *Merodon* species were positive for *Wolbachia* giving an infection rate of 96%. According to the estimation of Jeyaprasath and Hoy (2000), the infection rate in arthropods reaches up to 76% (48 arthropod species out of 63 tested positive) indicating a wide distribution of *Wolbachia* infection. However, it is important to point out that in this research the estimation of infection rate is based on less than third known *Merodon* species and probably deviated from the real infection rate.

Despite wide *Wolbachia* distribution among arthropod species, the study of Bailly-Bechet *et al.* (2017) conducted on 1100 species showed that most of the species acquired *Wolbachia* only recently and the most acquisition/loss events of *Wolbachia* occurred within the last million years. These events are most likely due to imperfect maternal transmission, although in some extant because of *Wolbachia* extinction from the population. However, there are some cases which indicate longterm *Wolbachia* infection. Taking into account population level events, Bailly-Bechet *et al.* (2017) estimated that mitochondria typically accumulate 4.7% substitutions per site during an infected episode, and 7.1% substitutions per site during the uninfected phase, which means that uninfected lineages acquire *Wolbachia* every 9.3 million years, while infected lineages lose their infection every 7 million years. Assuming this scenario, it is possible that *Wolbachia* acquisition/loss dynamic shapes mtDNA genealogy of the species.

In the case of *Merodon aureus* group a potential explanation for morphologically close species splitting into two main clades on *COI* tree could be a consequence of *Wolbachia*

influence on early evolution of different mtDNA lineages within the group. This means that ancient *Wolbachia* infection shaped *COI* based phylogeny of the group. However, apparently, there is no obvious pattern of coevolution of *Wolbachia* and *M. aureus* group species when comparing *Wolbachia* 16S rRNA gene tree and *COI* gene tree of hosts.

The species within complexes of *Merodon aureus* group often shared *wsp* alleles which indicates infection by the same strain. In these cases, *Wolbachia* could influence speciation if the same strain invaded different populations independently and by coupling and spreading different mtDNA haplotypes in populations. There is evidence that *wsp* detected *Wolbachia* strains could have different variants as consequence of deeper molecular variability associated with transposable elements, as found in detailed studies of *Wolbachia* variation in *Drosophila*, *Culex*, and *Hypolimnas bolina* (Duron *et al.* 2005; 2006; Riegler *et al.* 2005; Charlat *et al.* 2009). These small differences can affect the choice of mtDNA haplotypes which will be spread together with particular *Wolbachia* strain (Charlat *et al.* 2009). The theoretical modelling (Telschow *et al.* 2007) and experimental studies on many organisms (Bordenstein *et al.* 2001; Jaenike *et al.* 2006; Koukou *et al.* 2006; Miller *et al.* 2010), including both interspecific and intersemispecific analyses, show that *Wolbachia* can promote speciation in their hosts by inducing reproductive isolation, through development of either post- and/or premating mechanisms. However, more specimens per species should be tested for *Wolbachia* before any final conclusion about the influence of infection on speciation in complexes of *M. aureus* group. Additionally, cases of potential multiple infections by different *Wolbachia* strains (indicated in Table 1 as cases where sequence quality was low * or multiple bands on electrophoresis gels were detected **) deserve an in-depth study beyond the present study.

357 The sequencing of only one or two *Wolbachia* genes is unlikely to reveal much about
358 *Wolbachia* transmission between host species (Stahlhut *et al.* 2012). The characteristic
359 transmission dynamics and cases of multiple infections of the same host with different
360 *Wolbachia* strains have resulted in a freely recombining intracellular bacterial community
361 and mosaic bacterial genome structure (Klasson *et al.* 2009). The comparisons of shared
362 polymorphisms between *Wolbachia* strains confirm a mosaic structure of the *wsp* gene,
363 which is particularly prone to recombination and is under directional selection (Schulenburg
364 *et al.* 2000; Jiggins *et al.* 2001; Baldo *et al.* 2005). The frequent recombination events
365 produce a high level of sequence variability which makes *wsp* an excellent single marker for
366 distinguishing among different *Wolbachia* strains (Stahlhut *et al.* 2010). On the other hand,
367 frequent recombination disables tracing strain genealogy and makes *wsp* a bad choice for
368 studying *Wolbachia* horizontal transmission (Stahlhut *et al.* 2012). However, we found the
369 same *wsp* alleles present in both the host plant *Drimys maritima* bulb tissue and in the larval
370 specimens of *Merodon luteihumerus* acquired from the plant bulb. Although the presence of
371 *Wolbachia* DNA is not confirmation of the presence of living bacteria (see also Kolasa *et al.*
372 2017), these findings reveal a potential way of horizontal transmission mediated by plants.
373 All known early stages of *Merodon* species are found in underground storage organs of
374 geophytes (Ricarte *et al.* 2017; Preradović *et al.* 2018).

375 Plant mediated horizontal transmission of *Wolbachia* has already been hypothesized by
376 Sintupachee *et al.* (2006). They showed that four taxonomically diverse insects feeding on
377 the same host plant contained very closely related *Wolbachia* strains, suggesting the potential
378 role of host plants in *Wolbachia* horizontal transmission. Yang *et al.* (2013) also showed that
379 identical strains of *Wolbachia* are shared by two species, the gall wasp *Andricus mukaigawae*
380 and its inquiline wasp *Synergus japonicas*, which larvae feed on modified plant tissue of the
381 gall. Ahmed *et al.* (2016) found evidence for several new instances of *Wolbachia* horizontal

transmissions in Lepidoptera, and their findings suggested that specific shared food sources and shared natural enemies were possible routes of horizontal transmission. The DNA of two different *Wolbachia* strains including the one present in asparagus beetles, *Crioceris quinquepunctata* and *C. quatuordecimpunctata*, are also detected in host plant (*Asparagus*) tissues (Kolasa *et al.* 2017) once again indicating the possible route of horizontal transmission mediated by plants. This is strongly confirmed in the study by Li *et al.* (2017) where *Wolbachia* was visualized in plant tissue, both in the phloem vessels and in some spherules along the phloem. At present, however, neither the mechanisms nor processes of *Wolbachia* horizontal transmission are completely understood. For providing more solid support for the role of the host plants in *Wolbachia* transmissions in *Merodon* hoverflies, systematic screening for *Wolbachia* should be undertaken and include both the adult flies and their developmental stages, as well as the host plants.

ACKNOWLEDGEMENTS

We kindly thank Dr Ante Vujić (Novi Sad, Serbia) for specimens collecting and help in species determination. We thank Dr Mihailo Jelić for providing genomic DNA of *Drosophila melanogaster* to use as a positive control. This research was funded by the grants of Entomological Society of Helsinki and Societas pro Fauna et Flora Fennica, both Helsinki, Finland, and it was additionally supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia under Research Project No: OI173002; and the H2020 project ANTARES (No. 664387).

REFERENCES

- Ačanski J, Vujić A, Đan M, Obreht Vidaković D, Ståhls G, Radenković S (2016) Defining species boundaries in the *Merodon avidus* complex (Diptera, Syrphidae) using integrative taxonomy, with the description of a new species. *European Journal of Taxonomy* **237**, 1–25.
- Ahmed MZ, Breinholt JW, Kawahara AY (2016) Evidence for common horizontal transmission of *Wolbachia* among butterflies and moths. *BMC Evolutionary Biology* **16**, 118.
- Albertson R, Casper-Lindley C, Cao J, Tram U, Sullivan W (2009) Symmetric and asymmetric mitotic segregation patterns influence *Wolbachia* distribution in host somatic tissue. *Journal of Cell Science* **122**, 4570–4583.
- Andrić A, Šikoparija B, Obreht D *et al.* (2014) DNA barcoding applied: identifying the larva of *Merodon avidus* (Diptera: Syrphidae). *Acta Entomologica Musei Nationalis Pragae* **54**, 741–757.
- Augustinos AA, Santos-Garcia D, Dionyssopoulou E *et al.* (2011) Detection and characterization of *Wolbachia* infections in natural populations of aphids: is the hidden diversity fully unraveled?. *PloS One* **6**, e28695.
<https://doi.org/10.1371/journal.pone.0028695>
- Bailly-Bechet M, Martins-Simões P, Szöllösi GJ, Mialdea G, Sagot MF, Charlat S (2017) How long does *Wolbachia* remain on board?. *Molecular Biology and Evolution* **34**, 1183–1193.
- Balbuena JA, Míguez-Lozano R, Blasco-Costa I (2013) PACo: a novel procrustes application to cophylogenetic analysis. *PloS One* **8**, e61048.
<https://doi.org/10.1371/journal.pone.0061048>

431 Baldo L, Lo N, Werren JH (2005) Mosaic nature of the *Wolbachia* surface protein. *Journal of*
432 *Bacteriology* **187**, 5406–5418.

433 Baldo L, Hotopp JCD, Jolley KA *et al.* (2006) Multilocus sequence typing system for the
434 endosymbiont *Wolbachia pipientis*. *Applied and Environmental Microbiology* **72**,
435 7098–7110. Available from URL: <https://pubmlst.org/wolbachia/>

436 Baldo L, Werren JH (2007) Revisiting *Wolbachia* supergroup typing based on WSP: spurious
437 lineages and discordance with MLST. *Current microbiology* **55**, 81–87.

438 Bandi C, Anderson TJ, Genchi C, Blaxter ML (1998) Phylogeny of *Wolbachia* in filarial
439 nematodes. *Proceedings of the Royal Society of London. Series B, Biological Sciences*
440 **265**, 2407–2413.

441 Bing XL, Xia WQ, Gui JD, Yan GH, Wang XW, Liu SS (2014) Diversity and evolution of
442 the *Wolbachia* endosymbionts of *Bemisia* (Hemiptera: Aleyrodidae) whiteflies.
443 *Ecology and Evolution* **4**, 2714–2737.

444 Bordenstein SR, O'hara FP, Werren JH (2001) *Wolbachia*-induced incompatibility precedes
445 other hybrid incompatibilities in *Nasonia*. *Nature* **409**, 707–710.

446 Bordenstein SR, Paraskevopoulos C, Dunning Hotopp JC *et al.* (2009) Parasitism and
447 mutualism in *Wolbachia*: what the phylogenomic trees can and cannot say. *Molecular*
448 *Biology and Evolution* **26**, 231–241.

449 Charlat S, Hurst GD, Merçot H (2003) Evolutionary consequences of *Wolbachia* infections.
450 *Trends in Genetics* **19**, 217–223.

451 Charlat S, Duplouy A, Hornett EA *et al.* (2009) The joint evolutionary histories of *Wolbachia*
452 and mitochondria in *Hypolimnas bolina*. *BMC Evolutionary Biology* **9**, 64.
453 <https://doi.org/10.1186/1471-2148-9-64>

454 Chen H, Rangasamy M, Tan SY, Wang H, Siegfried BD (2010) Evaluation of five methods
 455 for total DNA extraction from western corn rootworm beetles. *PLoS One* **5**, e11963.
 456 <https://doi.org/10.1371/journal.pone.0011963>

457 Czarnetzki AB, Tebbe CC (2004) Detection and phylogenetic analysis of *Wolbachia* in
 458 Collembola. *Environmental Microbiology* **6**, 35–44.

459 Dobson SL, Bourtzis K, Braig HR *et al.* (1999) *Wolbachia* infections are distributed
 460 throughout insect somatic and germ line tissues. *Insect Biochemistry and Molecular*
 461 *Biology* **29**, 153–160.

462 Duron O, Lagnel J, Raymond M, Bourtzis K, Fort P, Weill M (2005) Transposable element
 463 polymorphism of *Wolbachia* in the mosquito *Culex pipiens*: evidence of genetic
 464 diversity, superinfection and recombination. *Molecular Ecology* **14**, 1561–1573.

465 Duron O, Fort P, Weill M (2006) Hypervariable prophage WO sequences describe an
 466 unexpected high number of *Wolbachia* variants in the mosquito *Culex pipiens*.
 467 *Proceedings of the Royal Society B: Biological Sciences* **273**, 495–502.

468 Duron O, Bouchon D, Boutin S *et al.* (2008) The diversity of reproductive parasites among
 469 arthropods: *Wolbachia* do not walk alone. *BMC Biology* **6**, 27.

470 Evison SEF, Roberts KE, Laurenson L *et al.* (2012) Pervasiveness of Parasites in Pollinators.
 471 *PLoS One* **7**, e30641. <https://doi.org/10.1371/journal.pone.0030641>

472 Galtier N, Nabholz B, Glémin S, Hurst GDD (2009) Mitochondrial DNA as a marker of
 473 molecular diversity: a reappraisal. *Molecular Ecology* **18**, 4541–4550.

474 Glowska E, Dragun-Damian A, Dabert M, Gerth M (2015) New *Wolbachia* supergroups
 475 detected in quill mites (Acari: Syringophilidae). *Infection, Genetics and Evolution* **30**,
 476 140–146.

477 Haegeman A, Vanholme B, Jacob J *et al.* (2009) An endosymbiotic bacterium in a plant-
 478 parasitic nematode: member of a new *Wolbachia* supergroup. *International Journal*
 479 *for Parasitology* **39**, 1045–1054.

480 Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren, JH (2008) How
 481 many species are infected with *Wolbachia*?-a statistical analysis of current data.
 482 *FEMS Microbiology Letters* **281**, 215–220.

483 Hurkmans W (1993) A monograph of *Merodon* (Diptera: Syrphidae). Part 1. *Tijdschrift voor*
 484 *Entomologie* **136**, 147–234.

485 Jaenike J, Dyer KA, Cornish C, Minhas MS (2006) Asymmetrical reinforcement and
 486 *Wolbachia* infection in *Drosophila*. *PLoS Biology* **4**, e325.
 487 <https://doi.org/10.1371/journal.pbio.0040325>

488 Jeyapragash A, Hoy MA (2000) Long PCR improves *Wolbachia* DNA amplification: *wsp*
 489 sequences found in 76% of sixty-three arthropod species. *Insect Molecular Biology* **9**,
 490 393–405.

491 Jiggins FM, von der Schulenburg JHG, Hurst GD, Majerus ME (2001) Recombination
 492 confounds interpretations of *Wolbachia* evolution. *Proceedings of the Royal Society*
 493 *of London. Series B, Biological Sciences* **268**, 1423–1427.

494 Katoh K, Standley DM (2013) MAFFT Multiple 723 Sequence Alignment Software Version
 495 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**,
 496 772–780.

497 Klasson L, Westberg J, Sapountzis P *et al.* (2009) The mosaic genome structure of the
 498 *Wolbachia* wRi strain infecting *Drosophila simulans*. *Proceedings of the National*
 499 *Academy of Sciences of the United States of America* **106**, 5725–5730.

500 Kolasa M, Montagna M, Mereghetti V, Kubisz D, Mazur MA, Kajtoch Ł (2017) Preliminary
 501 evidence of the horizontal transmission of *Wolbachia* between *Crioceris* leaf beetles

502 (Coleoptera: Chrysomelidae) and their *Asparagus* host plants. *European Journal of*
503 *Entomology* **114**, 446–454.

504 Koukou K, Pavlikaki H, Kiliass G, Werren JH, Bourtzis K, Alahiotis SN (2006) Influence of
505 antibiotic treatment and *Wolbachia* curing on sexual isolation among *Drosophila*
506 *melanogaster* cage populations. *Evolution* **60**, 87–96.

507 Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis
508 version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**, 1870–1874.
509 Available from URL: https://www.megasoftware.net/download_form

510 Li SJ, Ahmed MZ, Lv N *et al.* (2017) Plant-mediated horizontal transmission of *Wolbachia*
511 between whiteflies. *The ISME Journal* **11**, 1019–1028.

512 Librado P, Rozas J (2009) DnaSP v5: A software for comprehensive analysis of DNA
513 polymorphism data. *Bioinformatics* **25**, 1451–1452. Available from URL:
514 http://www.ub.edu/dnasp/index_v5.html

515 Lindsey AR, Bordenstein SR, Newton IL, Rasgon JL (2016) *Wolbachia pipientis* should not
516 be split into multiple species: A response to Ramírez-Puebla *et al.* “Species in
517 *Wolbachia*? Proposal for the designation of ‘*Candidatus Wolbachia bourtzisii*’,
518 ‘*Candidatus Wolbachia onchocercicola*’, ‘*Candidatus Wolbachia blaxteri*’,
519 ‘*Candidatus Wolbachia brugii*’, ‘*Candidatus Wolbachia taylori*’, ‘*Candidatus*
520 *Wolbachia collemboicola*’ and ‘*Candidatus Wolbachia multihospitum*’ for the
521 different species within *Wolbachia* supergroups”. *Systematic and Applied*
522 *Microbiology* **39**, 220–222.

523 Lo, N., Casiraghi M, Salati E, Bazzocchi C, Bandi C (2002) How many *Wolbachia*
524 supergroups exist? *Molecular Biology and Evolution* **19**, 341–346.

525 Lo N, Paraskevopoulos C, Bourtzis K *et al.* (2007) Taxonomic status of the intracellular
 526 bacterium *Wolbachia pipientis*. *International Journal of Systematic and Evolutionary*
 527 *Microbiology* **57**, 654–657.

528 Marcos-García MA, Vujić A, Mengual X (2007) Revision of Iberian species of the genus
 529 *Merodon* (Diptera: Syrphidae). *European journal of entomology* **104**, 531–572.

530 McWilliam H, Li W, Uludag M *et al.* (2013) Analysis Tool Web Services from the EMBL-
 531 EBI. *Nucleic Acids Research* **41**, W597–W600. Available from URL:
 532 <https://www.ebi.ac.uk/>

533 Milankov V, Ståhls G, Stamenković J, Vujić A (2008) Genetic diversity of populations of
 534 *Merodon aureus* and *M. cinereus* species complexes (Diptera, Syrphidae): integrative
 535 taxonomy and implications for conservation priorities on the Balkan Peninsula.
 536 *Conservation Genetics* **9**, 1125–1137.

537 Milankov V, Ludoški J, Ståhls G, Stamenković J, Vujić A (2009) High molecular and
 538 phenotypic diversity in the *Merodon avidus* complex (Diptera, Syrphidae): cryptic
 539 speciation in a diverse insect taxon. *Zoological Journal of the Linnean Society* **155**,
 540 819–833.

541 Miller WJ, Ehrman L, Schneider D (2010) Infectious speciation revisited: impact of
 542 symbiont-depletion on female fitness and mating behavior of *Drosophila paulistorum*.
 543 *PLoS Pathogens* **6**, e1001214. <https://doi.org/10.1371/journal.ppat.1001214>

544 Pietri JE, DeBruhl H, Sullivan W (2016) The rich somatic life of *Wolbachia*.
 545 *MicrobiologyOpen* **5**, 923–936.

546 Popović D, Ačanski J, Djan M, Obreht D, Vujić A, Radenković S (2015) Sibling species
 547 delimitation and nomenclature of the *Merodon avidus* complex (Diptera: Syrphidae).
 548 *European Journal of Entomology* **112**, 790–809.

549 Preradović J, Andrić A, Radenković S *et al.* (2018) Pupal stages of three species of the
550 phytophagous genus *Merodon* Meigen (Diptera: Syrphidae). *Zootaxa* **4420**, 229–242.

551 Radenković S, Vujić A, Ståhls G, Pérez-Bañón C, Petanidou T, Šimić S (2011) Three new
552 cryptic species of the genus *Merodon* Meigen (Diptera: Syrphidae) from the island of
553 Lesvos (Greece). *Zootaxa* **2735**, 35–56.

554 R Core Team (2018) R: A language and environment for statistical computing. R Foundation
555 for Statistical Computing, Vienna, Austria. Available from URL: [https://www.R-](https://www.R-project.org/)
556 [project.org/](https://www.R-project.org/).

557 Radenković S, Šašić Zorić L, Djan M *et al.* (2018) Cryptic speciation in the *Merodon*
558 *luteomaculatus* complex (Diptera: Syrphidae) from the eastern Mediterranean.
559 *Journal of Zoological Systematics and Evolutionary Research* **56**, 170–191.

560 Ramirez-Puebla ST, Servín-Garcidueñas LE, Ormeno-Orrillo E *et al.* (2015) Species in
561 *Wolbachia*? Proposal for the designation of ‘*Candidatus Wolbachia bourtzisii*’,
562 ‘*Candidatus Wolbachia onchocercicola*’, ‘*Candidatus Wolbachia blaxteri*’,
563 ‘*Candidatus Wolbachia brugii*’, ‘*Candidatus Wolbachia taylori*’, ‘*Candidatus*
564 *Wolbachia collemboicola*’ and ‘*Candidatus Wolbachia multihospitum*’ for the
565 different species within *Wolbachia* supergroups. *Systematic and Applied*
566 *Microbiology* **38**, 390–399.

567 Ricarte A, Marcos-García MA, Rotheray GE (2008) The early stages and life histories of
568 three *Eumerus* and two *Merodon* species (Diptera: Syrphidae) from the Mediterranean
569 region. *Entomologica Fennica* **19**, 129–141.

570 Ricarte A, Souba-Dols GJ, Hauser M, Marcos-García MÁ (2017) A review of the early
571 stages and host plants of the genera *Eumerus* and *Merodon* (Diptera: Syrphidae), with
572 new data on four species. *PloS One* **12**, e0189852.
573 <https://doi.org/10.1371/journal.pone.0189852>

574 Riegler M, Sidhu M, Miller WJ, O'Neill SL (2005) Evidence for a global *Wolbachia*
575 replacement in *Drosophila melanogaster*. *Current Biology* **15**, 1428–1433.

576 Rodriguez FJ, Oliver JL, Marin A, Medina JR (1990) The general stochastic model of
577 nucleotide substitution. *Journal of Theoretical Biology* **142**, 485–501.

578 Ros VI, Fleming VM, Feil, EJ, Breeuwer JA (2009) How diverse is the genus *Wolbachia*?
579 Multiple-gene sequencing reveals a putatively new *Wolbachia* supergroup recovered
580 from spider mites (Acari: Tetranychidae). *Applied and Environmental Microbiology*
581 **75**, 1036–1043.

582 Rowley SM, Raven RJ, McGraw EA (2004) *Wolbachia pipientis* in Australian spiders.
583 *Current Microbiology* **49**, 208–214.

584 Šašić L, Ačanski J, Vujić A *et al.* (2016) Molecular and morphological inference of three
585 cryptic species within the *Merodon aureus* species group (Diptera: Syrphidae). *PLoS*
586 *One* **11**, e0160001. <https://doi.org/10.1371/journal.pone.0160001>

587 Šašić Zorić L, Ačanski J, Đan M *et al.* (2018) Integrative taxonomy of *Merodon caeruleus*
588 complex (Diptera: Syrphidae) – evidence of cryptic speciation. *Matica Srpska Journal*
589 *for Natural Sciences*, 103-118.

590 Schulenburg JHGV, Hurst GD, Huigens TM, van Meer MM, Jiggins FM, Majerus ME
591 (2000) Molecular evolution and phylogenetic utility of *Wolbachia ftsZ* and *wsp* gene
592 sequences with special reference to the origin of male-killing. *Molecular Biology and*
593 *Evolution* **17**, 584–600.

594 Simões PM, Mialdea G, Reiss D, Sagot M-F, Charlat S (2011) *Wolbachia* detection: an
595 assessment of standard PCR protocols. *Molecular Ecology Resources* **11**, 567–572.

596 Sintupachee S, Milne JR, Poonchaisri S, Baimai V, Kittayapong P (2006) Closely related
597 *Wolbachia* strains within the pumpkin arthropod community and the potential for
598 horizontal transmission via the plant. *Microbial Ecology* **51**, 294–301.

599 Speight MCD (2014) Species accounts of European Syrphidae (Diptera), 2014. In: Speight
 600 MCD, Castella E, Sarthou JP, Vanappelghem C (eds) *Syrph the Net, the database of*
 601 *European Syrphidae*. Vol 78, Syrph the Net publications, Dublin.

602 Stahlhut JK, Desjardins CA, Clark ME *et al.* (2010) The mushroom habitat as an ecological
 603 arena for global exchange of *Wolbachia*. *Molecular Ecology* **19**, 1940–1952.

604 Stahlhut JK, Gibbs J, Sheffield CS, Smith MA, Packer L (2012) *Wolbachia* (Rickettsiales)
 605 infections and bee (Apoidea) barcoding: a response to Gerth *et al.* *Systematics and*
 606 *Biodiversity* **10**, 395–401.

607 Ståhls G, Vujic A, Pérez-Bañón C, Radenkovic S, Rojo S, Petanidou T (2009) *COI* barcodes
 608 for identification of *Merodon* hoverflies (Diptera, Syrphidae) of Lesbos Island,
 609 Greece. *Molecular Ecology Resources* **9**, 1431–1438.

610 Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
 611 large phylogenies. *Bioinformatics* **30**, 1312–1313. Available from URL:
 612 [http://bioinformatics.oxfordjournals.org/content/early/2014/01/21/bioinformatics.btu0](http://bioinformatics.oxfordjournals.org/content/early/2014/01/21/bioinformatics.btu033.abstract)
 613 [33.abstract](http://bioinformatics.oxfordjournals.org/content/early/2014/01/21/bioinformatics.btu033.abstract).

614 Telschow A, Flor M, Kobayashi Y, Hammerstein P, Werren JH (2007) *Wolbachia*-induced
 615 unidirectional cytoplasmic incompatibility and speciation: mainland-island model.
 616 *PLoS One* **2**, e701. <https://doi.org/10.1371/journal.pone.0000701>

617 Thompson FC, Rotheray GE (1998) Family Syrphidae. In: Papp L, Darvas B (eds)
 618 *Contributions to a manual of Palaearctic Diptera (with special reference to flies of*
 619 *economic importance)*, pp 81–139. Vol. 3, Science Herald, Budapest.

620 Vandekerckhove, TT, Watteyne S, Willems A, Swings JG, Mertens J, Gillis M (1999)
 621 Phylogenetic analysis of the 16S rDNA of the cytoplasmic bacterium *Wolbachia* from
 622 the novel host *Folsomia candida* (Hexapoda, Collembola) and its implications for
 623 wolbachial taxonomy. *FEMS Microbiology Letters* **180**, 279–286.

624 Veselić S, Vujić A, Radenković S (2017) Three new Eastern-Mediterranean endemic species
625 of the *Merodon aureus* group (Diptera: Syrphidae). *Zootaxa* **4254**, 401–434.

626 Vujić A, Pérez-Bañón C, Radenković S *et al.* (2007) Two new species of genus *Merodon*
627 Meigen, 1803 (Syrphidae, Diptera) from the island of Lesvos (Greece), in the eastern
628 Mediterranean. *Annales de la Société Entomologique de France* **43**, 319–326.

629 Vujić A, Radenković S, Ståhls G *et al.* (2012) Systematics and taxonomy of the *ruficornis*
630 group of genus *Merodon* Meigen (Diptera: Syrphidae). *Systematic Entomology* **37**,
631 578–602.

632 Wang GH, Jia LY, Xiao JH, Huang DW (2016) Discovery of a new *Wolbachia* supergroup in
633 cave spider species and the lateral transfer of phage WO among distant hosts.
634 *Infection, Genetics and Evolution* **41**, 1–7.

635 Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ (2015) The incidence of bacterial
636 endosymbionts in terrestrial arthropods. *Proceedings of the Royal Society of London.*
637 *Series B, Biological Sciences* **282**, 20150249

638 Werren JH (1997) Biology of *Wolbachia*. *Annual review of entomology* **42**, 587–609.

639 Werren JH, Windsor DM (2000) *Wolbachia* infection frequencies in insects: evidence of a
640 global equilibrium?. *Proceedings of the Royal Society of London. Series B, Biological*
641 *Sciences* **267**, 1277–1285.

642 Werren JH, Zhang W, Guo LR (1995) Evolution and phylogeny of *Wolbachia*: reproductive
643 parasites of arthropods. *Proceedings of the Royal Society of London. Series B,*
644 *Biological Sciences* **261**, 55–63.

645 Werren JH, Baldo L, Clark, ME (2008) *Wolbachia*: master manipulators of invertebrate
646 biology. *Nature Reviews Microbiology* **6**: 741–751.

Zhou W, Rousset F, O'Neill S. (1998) Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. *Proceedings of the Royal Society of London. Series B, Biological Sciences* **265**, 509–515.

Zug R, Hammerstein P (2012) Still a host of hosts for *Wolbachia*: Analysis of recent data suggests that 40% of terrestrial Arthropod species are infected. *PLoS One* **7**, e38544. <https://doi.org/10.1371/journal.pone.0038544>

Yang XH, Zhu DH, Liu Z, Zhao L, Su CY (2013) High levels of multiple infections, recombination and horizontal transmission of *Wolbachia* in the *Andricus mukaigawae* (Hymenoptera; Cynipidae) communities. *PloS One* **8**, e78970. <https://doi.org/10.1371/journal.pone.0078970>

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Figure S1. The sequence alignment of *wsp* alleles.

Table S1. Data on specimens screened for *Wolbachia* infection.

Table S2. The list of *Wolbachia* 16S rRNA gene sequences representing bacterial supergroups present in insect hosts.

Table S3. Blast search results for *Wolbachia* endosymbionts of *Merodon* species and *Drimia maritima* bulb *wsp* gene sequences.

670 Figure legends and table captions

671 Figure 1. Maximum likelihood tree of *Merodon aureus* group based on combined 3' and 5'

672 *COI* sequences (Šašić *et al.* 2016; Radenković *et al.* 2017; Šašić *et al.* unpublished data).

673 Bootstrap values ≥ 50 are presented near nodes.

674 Figure 2. Maximum likelihood tree based on 16S rRNA gene sequences of *Wolbachia* strains

675 present in genus *Merodon*. Bootstrap values ≥ 50 are presented near nodes. The specimens are

676 marked with DNA IDs and the host species names or GenBank accession numbers.

677 Figure 3. Phylogeny comparison between *COI* Maximum likelihood tree of *Merodon aureus*

678 group (left) and 16S rRNA gene Maximum likelihood tree of corresponding *Wolbachia*

679 endosymbionts (right). Bootstrap values ≥ 50 are presented near nodes.

680 Table 1. List of specimens tested on *Wolbachia* infection.

681 Table 2. The list of *wsp* alleles from *Merodon aureus* group host species including *Drimia*

682 *maritima* bulb and *Merodon luteihumerus*.

683 Table 3. Uncorrected p distance values (%) between *wsp* alleles.

684

Table 1. List of specimens tested on *Wolbachia* infection.

DNA ID	Species	Group		16S	wsp
		Complex	Subgroup		
AU402	<i>M. naxius</i> Vujić & Šašić, 2018	<i>luteomaculatus</i>	<i>bessarabicus</i>	+	*
AU497	<i>M. erymanthius</i> Vujić, Ačanski & Šašić, 2018			+	A
AU812	<i>M. luteomaculatus</i> Vujić, Ačanski & Šašić, 2018			+	A
AU27	<i>M. euri</i> Vujić & Radenković, 2018			+	A
AU396	<i>M. peloponnesius</i> Vujić, Radenković, Ačanski & Šašić, 2018			+	A
AU504	<i>M. andriotes</i> Vujić, Radenković & Šašić, 2018			+	*
AU443	<i>M. sapphous</i> Vujić, Perez-Banon & Radenković, 2007	<i>sapphous</i>		+	AB
AU99	<i>M. sapphous</i> sp. n. 1			+	A
AU141	<i>M. sapphous</i> sp. n. 2			-	A
AU454	<i>M. bessarabicus</i> Paramonov, 1924	<i>bessarabicus</i>		+	AB
AU82	<i>M. bessarabicus</i> sp. n. 1			+	A
AU53	<i>M. ambiguus</i> Bradescu, 1986	<i>ambiguus</i>		+	A
AU474	<i>M. ambiguus</i> sp. n. 1			+	A
AU1435	<i>M. quercetorum</i> Marcos-García, Vujić & Mengual, 2007			+	A
AU1442	<i>M. legionensis</i> Marcos-García, Vujić & Mengual, 2007			+	A
AU1432	<i>M. nisi</i> Veselić, Vujić & Radenković 2017			+	-
AU321	<i>M. unicolor</i> Strobl, 1909	<i>unicolor</i>	+	AB	
AU796	<i>M. unicolor</i> sp. n. 1		+	-	
AU710	<i>M. aureus</i> Fabricius, 1805	<i>aureus</i>	+	AB	
AU723	<i>M. aureus</i> sp. n. 1		+	A	
AU701	<i>M. aureus</i> sp. n. 1		+	A	
AU485	<i>M. aureus</i> sp. n. 2		-	A	
AU360	<i>M. cinereus</i> (Fabricius, 1794)		+	**	
AU1371	<i>M. aff. cinereus</i>	<i>cinereus</i>	+	AB	
AU530	<i>M. cinereus</i> sp. n. 1		+	A	
AU1362	<i>M. cinereus</i> sp. n. 2		+	A	
AU236	<i>M. cinereus</i> sp. n. 3		+	A	
AU1443	<i>M. cinereus</i> sp. n. 4		+	A	
AU517	<i>M. atratus</i> (Oldenberg, 1919)	<i>cinereus</i>	+	A	
AU151	<i>M. balkanicus</i> Šašić, Ačanski & Vujić, 2016		-	-	
AU144	<i>M. virgatus</i> Vujić & Radenković, 2016		+	A	
AU550	<i>M. virgatus</i>		+	A	
AU874	<i>M. aerarius</i> Rondani, 1857		+	**	
AU311	<i>M. minutus</i> Strobl, 1893	<i>chalybeus</i>	+	A	
AU752	<i>M. chalybeus</i> Wiedemann, 1822		+	A	
AU36	<i>M. dobrogensis</i> Bradescu, 1982	<i>dobrogensis</i>	+	A	
AU632	<i>M. dobrogensis</i>		+	A	
AU413	<i>M. puniceus</i> Vujić, Radenković & Péres-Bañón, 2011		+	*	
AU47	<i>M. dobrogensis</i> sp. n. 1		+	A	

AU107	<i>M. caerulescens</i> Loew, 1869	<i>caerulescens</i>			+	A
AU176	<i>M. atricapillatus</i> Šašić, Ačanski & Vujić, 2018				+	A
AU742	<i>M. pumilus</i> Macquart, 1849				+	**
AU253	<i>M. pumilus</i>				+	**
AU115	<i>M. robustus</i> Veselić, Vujić & Radenković 2017				+	A
AU326	<i>M. unguicornis</i> Strobl, 1909				+	AB
AU272	<i>M. neofasciatus</i> Ståhls & Vujić, 2018	<i>geniculatus</i> group			+	nt
AU273	<i>M. neofasciatus</i>				+	nt
AU288	<i>M. neofasciatus</i>				+	nt
AU606	<i>M. albifrons</i> Meigen, 1822	<i>albifrons</i> group			-	nt
AU611	<i>M. albifrons</i>				-	nt
AU617	<i>M. albifrons</i>				-	nt
AU620	<i>M. constans</i> (Rossi, 1794)	<i>constans</i> group			+	nt
AU621	<i>M. constans</i>				+	nt
AU622	<i>M. constans</i>				+	nt
AU772	<i>M. segetum</i> (Fabricius, 1794)	<i>natans</i> group			+	nt
AU773	<i>M. segetum</i>				-	nt
AU775	<i>M. segetum</i>				+	nt
AU1146	<i>M. avidus</i> Rossi, 1790	<i>avidus</i> group			+	nt
AU1164	<i>M. avidus</i>				+	nt
KR1	<i>M. moenium</i> (Wiedemann in Meigen, 1822)				+	nt
KR2	<i>M. moenium</i>				+	nt
KR3	<i>M. moenium</i>	<i>nanus</i> group			+	nt
N19	<i>M. nanus</i> Sack 1931				+	nt
TS213	<i>M. nanus</i>				-	nt
TS219	<i>M. telmateia</i> Hurkmans, 1987				+	nt
TS221	<i>M. telmateia</i>				+	nt
TS222	<i>M. telmateia</i>				+	nt
NG15	<i>M. nigratarsis</i> Rondani, 1845	<i>nigratarsis</i> group			+	nt
NG16	<i>M. nigratarsis</i>				+	nt
NG17	<i>M. nigratarsis</i>				+	nt
AF55	<i>M. melanocerus</i> Bezzi, 1915	<i>desuturinus</i> group			+	nt
AF57	<i>M. melanocerus</i>				+	nt
AF58	<i>M. melanocerus</i>				-	nt
Y2367	<i>M. luteihumerus</i> Marcos-García, Vujić & Mengual, 2007				-	A
Y2368	<i>M. luteihumerus</i>				nt	A
Y2369	<i>M. luteihumerus</i>				nt	A
BULB	<i>Drimia maritima</i> (L.) Stearn; bulb	host plant of <i>M. luteihumerus</i>			-	A

A - *wsp* amplified using 136F/691R primer pair specific for supergroup A; B - *wsp* amplified using 81F/522R primer pair specific for supergroup B; AB - *wsp* amplified using 81F/522R universal primer pair; * - multiple products detected using electrophoresis; ** - poor sequence quality; nt - not tested.

685

686

687

Table 2. The list of *wsp* alleles from *Merodon aureus* group host species including *Drimia maritima* bulb and *Merodon luteihumerus*.

Alleles	WSP profile*	Sequence ID (host species)
A1	311, 53, 145, 39, 18	AU497(<i>M. erymanthus</i>), AU812(<i>M. luteomaculatus</i>), AU27(<i>M. euri</i>), AU396(<i>M. peloponnesius</i>), AU82(<i>M. bessarabicus</i> sp. n. 1), AU99(<i>M. sapphous</i> sp. n. 1), AU141(<i>M. sapphous</i> sp. n. 2), AU53(<i>M. ambiguus</i>), AU723(<i>M. aureus</i> sp. n. 1), AU701(<i>M. aureus</i> sp. n. 1), AU236(<i>M. cinereus</i> sp. n. 3), AU144(<i>M. virgatus</i>), AU550(<i>M. virgatus</i>), AU752(<i>M. chalybeus</i>); AU36(<i>M. dobrogensis</i>), AU632(<i>M. dobrogensis</i>), AU47(<i>M. dobrogensis</i> sp. n. 1), AU107(<i>M. caerulescens</i>), AU176(<i>M. atricapillatus</i>); AU115(<i>M. robustus</i>), 16060(<i>Drimia maritima</i> plant bulb), Y2367(<i>M. luteihumerus</i>), Y2369(<i>M. luteihumerus</i>), Y2368(<i>M. luteihumerus</i>)
A2	731, 53, 145, 39, 18	AU311(<i>M. minutus</i>), AU1442(<i>M. legionensis</i>), AU1435(<i>M. quercetorum</i>)
A3	735, 28, 294, 39, 18	AU474(<i>M. ambiguus</i> sp. n. 1), AU443(<i>M. sapphous</i>)
A4	734, 261, 9, 271, 18	AU321(<i>M. unicolor</i>), AU710(<i>M. aureus</i>)
A5	(incomplete sequence)	AU454 (<i>M. bessarabicus</i>)
A6	733, 262, 115, 292, 62	AU326(<i>M. unguicornis</i>), AU485(<i>M. aureus</i> sp. n. 2)
A7	732, 263, 28, 31, 30	AU530(<i>M. cinereus</i> sp. n. 1), AU1362(<i>M. cinereus</i> sp. n. 2), AU517(<i>M. atratus</i>), AU1443(<i>M. cinereus</i> sp. n. 4), AU1371(<i>M. aff. cinereus</i>)

*WSP profile: *wsp*, HVR1, HVR2, HVR3, HVR4 IDs in *Wolbachia* MLST database (Baldo et al. 2006).

688

689

690

691

692

693

694

695

696

697

Table 3. Uncorrected p distance values (%) between *wsp* alleles.

	A1	A2	A3	A4	A5	A6
A1						
A2	0.198					
A3	10.474	10.672				
A4	9.486	9.684	12.548			
A5	14.500	14.250	15.777	8.252		
A6	11.858	12.055	17.984	16.206	12.069	
A7	12.253	12.055	18.379	17.391	14.778	13.477

698