<https://helda.helsinki.fi>

Synchronized proxy-based temperature reconstructions reveal mid- to late Holocene climate oscillations in High Arctic Svalbard

Luoto, Tomi P.

2018-01

Luoto , T P , Ojala , A E K , Arppe , L , Brooks , S J , Kurki , E , Oksman , M , Wooller , M J & Zajaczkowski , M 2018 , ' Synchronized proxy-based temperature reconstructions reveal mid- to late Holocene climate oscillations in High Arctic Svalbard ' , Journal of Quaternary Science , vol. 33 , no. 1 , pp. 93-99 . <https://doi.org/10.1002/jqs.3001>

<http://hdl.handle.net/10138/323609>

<https://doi.org/10.1002/jqs.3001>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Synchronized proxy-based temperature reconstructions reveal mid- to late Holocene climate oscillations in High Arctic Svalbard

Tomi P. Luoto^{1,*}, Antti E.K. Ojala², Laura Arppe³, Stephen J. Brooks⁴, Eija Kurki⁵, Mimmi Oksman⁵, Matthew J. Wooller⁶, and Marek Zajączkowski⁷

¹ Department of Biological and Environmental Science, Surfontie 9 C, 40500 University of Jyväskylä, Finland

² Geological Survey of Finland, Betonimiehenkuja 4, 02150 Espoo, Finland

³ Laboratory of Chronology, Finnish Museum of Natural History – LUOMUS, Gustaf Hällströmin katu 2a, 00014 University of Helsinki, Finland

⁴ Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

⁵ Department of Geosciences and Geography, Gustaf Hällströmin katu 2a, 00014 University of Helsinki, Finland

⁶ College of Fisheries & Ocean Science and Water & Environmental Research Center, 461 Duckering, University of Alaska Fairbanks, USA

⁷ Institute of Oceanology Polish Academy of Sciences, Powstańców Warszawy 55, 81–712 Sopot, Poland

*Corresponding author (tomi.luoto@helsinki.fi)

Running head: Mid- to late Holocene climate oscillations in High Arctic Svalbard

26 **Abstract**

27

28 Existing paleoclimate data are exceedingly sparse from southern Spitsbergen, a High Arctic region
29 predicted to experience significant environmental changes as a result of amplified warming. We
30 analyzed biotic and isotopic paleolimnological proxies to reconstruct past climate from a lacustrine
31 sediment core, with a basal age of ~5500 years BP, in southern Spitsbergen (77°N). We used fossil
32 Chironomidae assemblages to quantitatively reconstruct past mean July air temperatures and stable
33 oxygen isotope values ($\delta^{18}\text{O}$) of these fossils to estimate changes in mean annual air temperature
34 (MAT). These proxy records are strikingly similar and show that the coldest anomaly since the mid-
35 Holocene occurred between 350-50 cal a BP, during the “Little Ice Age”, whereas the warmest
36 period in the summer temperature record occurred between 5500-5000 and ~2000 cal a BP. Our
37 findings indicate that the natural long-term air temperature dynamics in our study area are most
38 likely connected to solar minima and positive feedback mechanisms from sea-surface temperature
39 maxima. The results also highlight that the recent temperature increase is unprecedented in its rate
40 with a ~2 °C increase in the summer temperatures during the past ~50 years.

41

42

43

44

45

46

47

48

49

50 *Keywords:* Arctic; Chironomidae; Paleoclimate; Spitsbergen; Stable isotopes

51 **Introduction**

52

53 Arctic regions, such as Svalbard, are highly impacted by present day climate change due to multiple
54 feedback mechanisms related to oceanic, cryospheric and atmospheric processes (Pithan and
55 Mauritsen, 2014). Arctic sea ice extent and volume have been declining (Stroeve *et al.*, 2008;
56 Polyak *et al.*, 2010), and the Arctic Ocean may become seasonally ice-free by the mid-21st century
57 (Wang and Overland, 2009). It was recently demonstrated that the Atlantic Meridional Overturning
58 Circulation (AMOC), which plays a key role in the poleward transport of heat and has a profound
59 impact on climate in northwestern Europe and the High Arctic, may have slowed due to global
60 warming (Landerer *et al.*, 2015). To improve our understanding of the Arctic climate change
61 processes, a long-term perspective of the magnitude and rate of climate fluctuations in the past is
62 necessary.

63 Arctic lakes and their sedimentary archives provide an opportunity to examine past
64 environmental conditions. Paleolimnological methods, using biotic proxies such as chironomids
65 (Insecta: Diptera), which are sensitive to temperature changes, and stable oxygen isotope signatures
66 ($\delta^{18}\text{O}$ values) reflecting hydroclimatic conditions, have been found to be very useful in tracking past
67 climate changes (Brooks, 2006; Wooller *et al.*, 2004). Fossil chironomid assemblages have been
68 used to reconstruct temperature during summer, which is their primary season for growth and
69 development (Eggermont and Heiri, 2012), whereas the isotope values of their head capsules can
70 capture the annual $\delta^{18}\text{O}$ signal of lake water derived from precipitation (Verbruggen *et al.*, 2011).
71 The $\delta^{18}\text{O}$ of head capsules of chironomid larvae are equilibrated with the $\delta^{18}\text{O}$ of lakewaters in
72 which they live. In lakes that are not substantially influenced by evaporation or glacial meltwater
73 inputs, the $\delta^{18}\text{O}$ values of lakewater are controlled by the $\delta^{18}\text{O}$ value of local precipitation and
74 temperature. In these lakes chironomid $\delta^{18}\text{O}$ values can subsequently be used to infer mean annual
75 air temperature (MAT) (Wooller *et al.*, 2004).

76 In Svalbard, quantitative paleoclimatic records are thus far sparse, especially for the
77 southern part of Svalbard, and cover a time span of ≤ 2000 years (Brooks and Birks, 2004; Isaksson
78 *et al.*, 2005; D'Andrea *et al.*, 2012). There is a pressing need for long-term paleoclimate records
79 from the area for linking climate and AMOC dynamics. In our study area, the North Atlantic
80 Current carries warm surface waters from the Atlantic to high latitudes along the western side of
81 Svalbard. Cold waters from the Arctic Ocean move southwards on the eastern side of Svalbard
82 crossing the warmer waters at the southern tip of Svalbard, where the Polar Front develops
83 (Majewski *et al.*, 2009). Our study area has a significant marine influence from the warm West
84 Spitsbergen Current and the cold Sorokap Current, and is therefore a climatically and
85 oceanographically sensitive region. For this reason, we examined a lake sediment profile from
86 southern Spitsbergen (Fig. 1) using analyses of chironomid community assemblages to reconstruct
87 mean July air temperature combined with $\delta^{18}\text{O}$ values from analyses of chironomid head capsules to
88 infer past MAT. The results of our study are important for understanding the long-term climate
89 processes and land-ocean-atmosphere interactions in this High Arctic climate change hotspot area.

91 **Methods**

92 *Study site and sampling*

93
94 Lake Svartvatnet (76°90'N, 15°66'E, 63 m a.s.l.) is located in a periglacial landscape (Arctic
95 tundra) in southern Spitsbergen, across the Hornsund fjord from the Polish Polar Station (Fig. 1).
96 This thermally stratified lake has a maximum depth of 26.5 m, surface area of ~80 ha and one outlet
97 on its eastern side. Svartvatnet is an oligotrophic lake with a surface water pH of 7.2 (measured
98 June 2013). The lake is annually ice-covered for 9-10 months with an open-season from mid-July to
99 September. At present, the mean July temperature in the area is 4.5 °C, while the MAT is -2.1 °C
100 with annual precipitation remaining below 400 mm (Marsz and Styszyńska, 2013). The lake

101 receives most of its water during the spring/summer as meltwater via a set a shallow creeks,
102 whereas the periglacial soil appears not to produce significant groundwater storage at the study site
103 (Ojala *et al.*, 2016).

104 A 167-cm sediment core (SV4C) and a parallel 149-cm core (SV4B) were taken from the
105 northern basin of Svartvatnet (Fig. 1) using a light piston corer during the summer of 2013. The
106 sediments consisted of clay gyttja with visible sand and silt/clay laminae (Ojala *et al.*, 2016). The
107 sediment sequences were measured for their physical properties, correlated and the chronology was
108 determined based on a combination of ^{137}Cs , ^{210}Pb , AMS ^{14}C (five dates) and paleomagnetic dating.
109 The dating results and the constructed age-depth model (a Bayesian P-sequence deposition model)
110 (Fig. S1) demonstrated that the SV4C sediment profile covers a 5500-year sedimentary archive with
111 an average accumulation rate of 0.3 mm a^{-1} (Ojala *et al.*, 2016). The accumulation remained
112 otherwise stable and continuous throughout the core but according to the ^{210}Pb dating, the
113 sedimentation rate increased during the 20th century. The well-established sediment chronology,
114 with all the dating results showing good agreement, provides a solid basis for paleoclimatic
115 reconstructions from the Svartvatnet sediment sequence. Detailed descriptions of the lake basin and
116 the sediment, and the associated age-depth model for the Svartvatnet succession was previously
117 published (Ojala *et al.*, 2016).

118

119 *Analyses*

120

121 Sediment subsamples (SV4C) for fossil chironomid analysis were prepared using standard methods
122 (Brooks *et al.*, 2007). The wet sediment was sieved through a 100- μm mesh and the residue was
123 examined under a stereomicroscope. Larval head capsules were extracted with fine forceps and
124 mounted permanently on microscope slides. Identification was performed under a light microscope
125 at 400x magnification. The minimum chironomid head capsule number per sample was set to 50

126 (Heiri and Lotter, 2001). Principal component analysis (PCA) was used to detect primary direction
127 of the chironomid community variance (Abdi and Williams, 2010). The PCA was run with all taxa
128 included using a square-root transformation of the species percentage data. The chironomid-based
129 mean July air temperature reconstruction (CiT) used the trimmed Norwegian weighted-averaging
130 partial least squares (WA-PLS, 3-component) calibration model (Velle *et al.*, 2011). The model has
131 a cross-validated coefficient of determination of 0.86, a root mean squared error of prediction of
132 0.70 °C and a maximum bias of 0.66 °C. To test whether the reconstruction corresponded to the
133 primary chironomid community variability, the temperatures were compared against the PCA axis 1
134 scores using Pearson product-moment correlation coefficient and the associated level of statistical
135 significance.

136 The protocol for $\delta^{18}\text{O}$ analysis of chironomid head capsules ($\delta^{18}\text{O}_{\text{chir}}$) followed Wang *et al.*
137 (2008) without the acid treatment, and was described further by Arppe *et al.* (2017) and Kurki
138 (2016). Approximately 80 μg of cleaned chironomid head capsules were weighed in silver capsules,
139 and measurements of $\delta^{18}\text{O}$ values were performed with a Finnigan ThermoQuest TC/EA coupled to
140 a Delta^{Plus} Advantage isotope ratio mass spectrometer at the Laboratory of Chronology, Finnish
141 Museum of Natural History. The $\delta^{18}\text{O}$ values are expressed relative to the oxygen isotope ratio of
142 Vienna Standard Mean Ocean Water. The international reference materials IAEA-NO3 (25.6‰),
143 IAEA-601 (23.3‰), ANU sucrose (36.4‰), baleen whale keratin BWBII (14.0‰), and an
144 internally validated IAEA-CH3 cellulose (32.6‰) were used to normalize raw $\delta^{18}\text{O}$ values. The
145 reproducibility of similar biogenic reference materials indicates a mean analytical precision (1 σ) of
146 0.5‰, with a range from 0.4 to 2.5‰ depending on signal (sample) size, which is consistent with
147 the findings of Wang *et al.* (2008). The $\delta^{18}\text{O}_{\text{chir}}$ values were used to estimate mean annual values of
148 $\delta^{18}\text{O}_{\text{lake water}}$ and MAT following Wooller *et al.* (2004) and Verbruggen *et al.* (2011). The composite
149 errors for the reconstructions, factoring in both measurement uncertainty and calibration error, were
150 calculated according to Pryor *et al.* (2014). They range from 1.4 to 3.6 ‰ (mean 1.6‰) and 1.9 to

151 4.5 °C (mean = 2.1 °C) for the $\delta^{18}\text{O}_{\text{lake water}}$ and MAT reconstructions, respectively. A more detailed
152 description of the analytical and data processing methods was described by Arppe *et al.* (2017). It
153 should be noted that the $\delta^{18}\text{O}_{\text{chir}}$ reconstruction of mean annual temperature should be considered
154 tentative owing to reasons discussed below.

155

156 **Results**

157

158 Fossil chironomid assemblages of the initial part of the sediment profile (160-90 cm, ~5500-3300
159 cal a BP) were dominated by *Micropsectra contracta*-type (Fig. 2), which corresponds to
160 morphologically and ecologically similar taxon *M. insignilobus*-type in the chironomid-based
161 calibration set used for the temperature reconstruction (Brooks and Birks, 2000, 2001). This taxon
162 complex is typical for the oligotrophic lakes in Svalbard (Luoto *et al.* 2011; Velle *et al.* 2011). In
163 the mid-part of the profile (90-50 cm, ~3200-1800 cal a BP) the cold-indicating *Hydrobaenus*
164 *lugubris*-type began to dominate but was later (at ~1800 cal a BP) replaced by another cold
165 preferring taxon *Oliveria*. In the most recent sample representing the present, several warm-water
166 chironomids, such as *Paratanytarsus austriacus*-type and *M. junci*-type, increased that has also
167 been detected in other lakes in Svalbard (Velle *et al.* 2011; Kivilä 2014).

168 In general, the chironomid PCA axis 1 scores were negative in the lower part of the record
169 but turned into positive scores after ~3500 cal a BP. Reflecting a major shift in chironomid
170 assemblages (decreases in *M. contracta*-type and *Orthocladius trigonolabis*-type and increase in *H.*
171 *lugubris*-type), chironomid PCA axis 1 scores showed a rapid change at ~3300-3200 cal a BP (~90
172 cm) and another change of similar magnitude in the most recent samples (decreases in *Oliveridia*
173 and *H. lugubris*-type and increase in *M. junci*-type) (Fig. 2). The chironomid PCA axis 1 scores
174 correlated closely ($r=0.91$, $P<0.001$) with the CiT changes during the past 5500 years (Fig. 3).
175 Furthermore, the CiT values were very similar to the $\delta^{18}\text{O}_{\text{lake water}}$ values and the $\delta^{18}\text{O}_{\text{chir}}$

176 reconstruction of annual temperatures (Fig. 3). The CiT remained at $>5^{\circ}\text{C}$ between ~ 5500 - 3500 cal
177 a BP with a progressive decline, however, with a short warm period at ~ 1500 cal a BP and very
178 warm temperatures at the surface sample. The reconstruction of $\delta^{18}\text{O}_{\text{lake water}}$ values distinctly
179 mimics the CiT reconstruction but the warmer temperatures between ~ 5500 - 3500 cal a BP
180 identified by the CiT are not as well reflected in the isotopic reconstruction. The minimum
181 temperatures were reconstructed for the period between ~ 300 - 200 cal a BP with both proxies. In
182 addition to this temperature minimum, simultaneous cold periods were found at ~ 3400 - 3200 and
183 1400 - 1200 cal a BP.

184 The surface sediment CiT of 6.6°C is overestimated compared to the observed long-term
185 mean July air temperature (1978-2014) at the study area of 4.5°C but close to the temperature
186 measured in June 2016 of 6.0°C . Therefore, when compared to the present instrumental
187 temperature record, the surface sediment CiT is within the model's prediction error of 0.7°C . The
188 observed long-term MAT in the area is -4.4°C being clearly lower than in the $\delta^{18}\text{O}$ -based
189 reconstruction for the surface sample. However, the reconstructed surface sample MAT of -0.6°C
190 is closer to the last five-year mean of -1.9°C , which remains within the reconstruction's sample-
191 specific error estimate of 3.0°C . When comparing the reconstructed $\delta^{18}\text{O}_{\text{lake water}}$ of -9.2‰ (sample-
192 specific error 2.4‰) with the observed lake water value of -9.6‰ , measured from water samples
193 during the summer 2013, the values were found to be very close.

194

195 **Discussion**

196

197 *Temperature reconstructions and climate variability*

198

199 According to our previous assessment based on contemporary aquatic communities (Luoto *et al.*,
200 2016), the most significant confounding factor for chironomid-based temperature reconstructions in

201 the study area (Hornsund) is the potential influence of bird-driven nutrient enrichment. This kind of
202 arctic eutrophication superimposed on the recent climate warming is a typical trend for smaller
203 ponds lying close to shoreline in Svalbard with increasing sizes of arctic-nesting birds in their
204 catchment (Luoto *et al.*, 2014, 2015). However, when comparing the Svartvatnet chironomid
205 stratigraphy (Fig. 2) with the modern dataset (Luoto *et al.*, 2016), the phenomenon of bird-induced
206 eutrophication is not reflected in the chironomid communities of Svartvatnet. Therefore, since our
207 study site is not significantly influenced by birds, it is probable that the long-term biotic record is
208 driven by the direct climate impact. In fact, the PCA results indicated that the chironomid
209 assemblages significantly responded to temperature changes during the past 5500 years, as the
210 chironomid PCA axis 1 scores correlated with the CiT. Furthermore, the $\delta^{18}\text{O}_{\text{lake water}}$ reconstructions
211 showed similar trends with the CiT (Fig. 3) providing evidence that both proxies were reflecting the
212 same environmental variable (temperature), though at different timescales (annual vs. July).

213 While we consider the $\delta^{18}\text{O}_{\text{lake water}}$ reconstruction a robust representation of past changes in
214 $\delta^{18}\text{O}$ values of lake Svartvatnet water and local precipitation, the interpretation of the isotope record
215 in terms of temperature variability is not straightforward. Due to uncertainties in both the present-
216 day and past $\delta^{18}\text{O}$ /MAT relationships, possible past changes in the locations and pathways of major
217 moisture sources, and the seasonal distribution of precipitation, the $\delta^{18}\text{O}_{\text{chir}}$ -based MAT
218 reconstruction is subject to some uncertainty. The presented MAT reconstruction (Fig. 3) holds true
219 only in a scenario where all the changes observed in $\delta^{18}\text{O}_{\text{lake water}}$ can be attributed to changes in
220 MAT at the site, without associated changes in the abovementioned factors. Indeed, the record of
221 $\delta^{18}\text{O}$ values, while carrying a strong temperature dependent signal, is very likely also influenced by
222 other environmental factors such as variations in sea ice extent around Spitsbergen, and should not
223 be considered solely to reflect MAT. A detailed discussion of the $\delta^{18}\text{O}$ record was previously
224 presented by Arppe *et al.* (2017).

225 The reconstructions indicated relatively warm climatic conditions at the beginning of the
226 record, between ~5500 and 3800 cal a BP. This warm period corresponds with the presence of
227 smaller-than-present glaciers in Svalbard (Humlum *et al.*, 2005) and Franz Josef Land (Lubinsky *et*
228 *al.*, 1999), and with the general timing of the Holocene Climate Optimum (HCO, ~8000-5000 cal a
229 BP) (Wanner *et al.*, 2008). Our results are also in accordance with recent sediment-based Holocene
230 glacier reconstruction from the northwestern coast of the Svalbard Archipelago (Røthe *et al.*, 2015),
231 which shows low inorganic detrital input into the proglacial Kløsa basin, suggesting that influx of
232 meltwater ceased following the possible melting of the Karlbreen glacier. Likewise, Svendsen and
233 Mangerud (1997) concluded that the Linnébreen glacier at the mouth of Isfjorden in Spitsbergen
234 had melted during this period. Based on plant macrofossil analysis, July temperatures may have
235 been up to 1.5 °C warmer compared to the late 20th century in the western coastline of Svalbard
236 (Birks, 1991). Our CiT shows that the summer temperatures between 5500 and 5000 cal a BP in
237 southern Spitsbergen were similar to present day, which is in agreement with most paleoclimate
238 records from Svalbard indicating generally warm mid-Holocene summer conditions (Lubinsky *et*
239 *al.*, 1999; Humlum *et al.*, 2005; Luoto *et al.*, 2011).

240 Our CiT record, reflecting summer conditions, shows a general Neoglacial cooling trend
241 from ~5000 cal a BP onwards (Fig 3). Svendsen and Mangerud (1997) suggested that following the
242 HCO, the Linnébreen glacier reoccupied the valley above Linnévatnet between 4600 cal a BP and
243 around 2800 cal a BP, probably in response to a decline in summer insolation and climate cooling.
244 The first one of these episodes is fairly coincidental with two minimum peaks in our CiT. On the
245 other hand, the $\delta^{18}\text{O}_{\text{chir}}$ record, which incorporates the climatic signal of the cool/cold season, does
246 not show a decline. Summer cooling, from a slightly later peak (4250-4050 cal a BP) than in our
247 record, was recorded by van der Bilt *et al.* (2015) in the Lake Hajeren proglacial sedimentary record
248 (Fig. 3). This may reflect regional differences caused by external forcing mechanisms, or slower
249 response times of small glaciers (van der Bilt *et al.*, 2015).

250 We document a significant cooling episode in both our proxy records at ~3300 cal a BP,
251 which has previously been detected in the equilibrium-line altitude (ELA) reconstructions from
252 Karlbreen glacier on the northwest coast of Svalbard (Røthe *et al.*, 2015). Although our evidence is
253 mostly based on single data point excursions (Fig. 3), a contemporaneous centennial-scale glacier
254 advance in the Hajaren catchment was also observed by van der Bilt *et al.* (2015), which was linked
255 to orbitally forced summer cooling and increased sea-ice export through the Fram Strait. In
256 addition, a cooling event in our record at 1300 cal a BP is concurrent with the ELA reconstruction
257 (Røthe *et al.*, 2015) and the Linnévatnet record (Svendsen and Mangerud, 1997), suggesting that the
258 glacier response is probably more related to summer air temperature than precipitation changes.
259 Peak erosion due to the presence of a glacier in the Lake Hajaren catchment occurred later, at
260 around 1100 cal a BP (van der Bilt *et al.*, 2015) suggesting that the lack of glacier growth prior to
261 1100 cal a BP was due to the climate being too dry or too warm in this low-lying catchment.

262 Typical for lacustrine sites in the Arctic (Miller *et al.*, 2010), the present results do not show
263 clear signs of the Medieval Climate Anomaly (MCA, ~850-1250 CE). The closest study of
264 millennial-scale climate fluctuations to our site was presented by Pawłowska *et al.* (2015), who
265 investigated mass balance and glacial activity in a marine environment at Hornsund fjord. These
266 investigators were not able to identify clear changes in environmental conditions during the MCA
267 as there was a stable and low rate of sediment accumulation and low ice rafted debris flux.

268 According to our proxies, the general late Holocene cooling trend culminated at ~300-200
269 cal a BP, coinciding with the end of the Little Ice Age (LIA). The previous ~2000/1000-year
270 paleoclimate reconstructions from Svalbard indicate that the LIA was a rather muted and short-lived
271 event in the area (Velle *et al.*, 2011; D'Andrea *et al.* 2012) compared to continental Northern
272 Europe, where it lasted for several centuries (Luoto and Nevalainen, 2015, 2016; Zawiska *et al.*,
273 2017). This is also well reflected in glacier ELA records, since the LIA is well represented in a
274 record from continental Norway (Bakke *et al.*, 2005), while less distinct in the Svalbard glacier

record (Røthe *et al.*, 2015), suggesting that short-lived climate events may not be evident in glacier growth. According to our results, the declining trend towards colder LIA conditions may have initiated as early as 700-800 cal a BP, followed by a major cooling shift around 400 cal a BP. This date is in good accordance with the composite of high-resolution proxy climate records from the Arctic by Kaufman *et al.* (2009). It also agrees with the inferred beginning of the LIA at around 1600 CE described by Pawłowska *et al.* (2015) based on the analysis of foraminiferal ancient DNA in the marine environment of Hornsund fjord. Timing of the onset of the LIA at around 700 cal a BP also agrees well with other records from western Svalbard (Svendsen and Mangerud, 1997; van der Bilt *et al.*, 2015) suggesting a regional climate shift.

Our CiT for the surface sample is 3.5 °C higher than the lowest LIA values, whereas the $\delta^{18}\text{O}_{\text{chir}}$ -based reconstruction, if interpreted to solely represent changes in MAT at the site, is >10 °C higher. However, the dramatic negative shift observed in the reconstruction of $\delta^{18}\text{O}$ values during the LIA is likely to also reflect the significantly expanded sea ice cover limiting the input of proximally derived moisture, and thus should not be interpreted purely on the basis of MAT. These results nonetheless emphasize the influence of winter months to the annual heat budget in our study area. The annual and summer temperature increase from the LIA to the present appears to have been unprecedentedly rapid over the last 5500 years and clearly reflects the most significant climate shift in the record. As in previous evidence from the circum-Arctic (Kaufman *et al.*, 2009; Luoto *et al.*, 2017), rapid recent warming reverses the general trend of cooling in our reconstructions (Fig. 3). Our CiT record indicates a temperature increase of ~2 °C for the recent decades that is equal to the measured temperature increase during the past 30 years at the Polish Polar Station since 1979 (Marsz and Styszyńska, 2013). The surface sediment CiT of 6.6 °C is higher than the observed long-term mean July air temperature (1978-2014) at the study area of 4.5 °C but close to the present day (2016) temperature of 6.0 °C confirming that the reconstructed values are realistic. Similarly, the $\delta^{18}\text{O}_{\text{chir}}$ record indicates a MAT value of -0.6 °C for the surface sample that is higher than the

300 35-year mean of 4.4 °C but closer to the last five-year mean of -1.9 °C. The recently (2013)
301 measured $\delta^{18}\text{O}_{\text{lake water}}$ value (-9.6 ‰) and the inferred $\delta^{18}\text{O}_{\text{lake water}}$ value (-9.2 ‰) are also very
302 close. Hence, the better comparability of the reconstructed values with the more recent observations
303 are most likely owing to the observed rapid temperature increase over the temperature monitoring
304 period (Marsz and Styszyńska, 2013).

305

306 *External forcing and feedback mechanisms*

307

308 Decreasing summer solar insolation may have caused the Neoglacial cooling trend in the Arctic
309 region (Wanner *et al.*, 2008). In the North Atlantic region, the Neoglacial cooling is reflected by
310 growing glaciers (Bakke *et al.*, 2010), decreasing sea surface temperatures (SSTs) (Sarnthein *et al.*,
311 2003; Berner *et al.*, 2010) and ice-core derived temperatures (Lecavalier *et al.*, 2013). In general,
312 North Atlantic surface winds and surface ocean hydrography have been influenced by variations in
313 solar output throughout the Holocene, hence, the surface hydrographic changes may have affected
314 production of North Atlantic Deep Water, potentially providing an additional mechanism for
315 amplifying the solar radiation fluctuations and transmitting them globally (Bond *et al.*, 2001).

316 The air temperature dynamics in our study appear to be externally forced by solar minima
317 and positive feedback mechanisms related to SST maxima. The cold episodes reconstructed from
318 the current biotic and isotopic records at 3400-3200 and 1400-1200 cal a BP are synchronized with
319 periods of low solar irradiation (Steinhilber *et al.*, 2009; Fig. 3), however, it is likely that
320 combination of several factors triggers the cold and warm episodes. In agreement with Bond *et al.*
321 (2001), our record shows that the LIA is not fully synchronized with the Maunder solar minimum.
322 Cooling during the LIA is significantly connected with atmospheric forcing since a strongly
323 negative NAO index phase prevailed (Trouet *et al.*, 2009; Dylmer *et al.*, 2013; Luoto and
324 Nevalainen, 2017). The warm periods in our summer temperature record, ~5400, ~2200 and ~1600

cal a BP, show correspondence with increases in summer SSTs reconstructed from the northern Norwegian Sea (Sarnthein *et al.*, 2003; Fig. 2). However, these warm phases are not apparent in the isotopic record. Though this might reflect the sensitivity of $\delta^{18}\text{O}$ values to precipitation (Langebroek *et al.*, 2011), it more likely mirrors its sensitivity to MAT. The lack of the warm MCA signal in our record is probably due to dominance of Arctic surface water in the western Barents Sea at this time, which contributes to cooler conditions than in continental regions of the Northern Hemisphere (Dylmer *et al.*, 2013).

The recent and most significant temperature increase in our record corresponds to the human-induced increase in greenhouse gases in the atmosphere since the pre-industrial age (Crowley, 2000). Although there is an increase in solar irradiance towards the present (Steinhilber *et al.*, 2009; Fig. 3), it appears to be too small to explain the fast and major increase in the temperatures at our study site. There also exists a possibility of a regional temperature response via ocean-atmosphere heat transport that has been suggested from marine sediments off Western Svalbard (Spielhagen *et al.*, 2011).

339

340 **Conclusions**

341

Changes in fossil chironomid assemblages and $\delta^{18}\text{O}$ values of chironomid head capsules document significant paleoclimate changes during the past ~5500 years in Svalbard. Both proxies showed similar temperature trends, indicating late Holocene cooling and the LIA. The recent temperature increase is unprecedented in its speed and magnitude since the mid Holocene. Although partly associated with single data point excursions, we document connections between periods of cold temperatures and solar radiation minima and warm periods and high SSTs, suggesting that both forcing/feedback mechanisms may play an important role behind the climate dynamics of our study region. The complexity of the regional climate patterns is reinforced by the local glacier influence.

350 The current record makes a valuable contribution to paleoclimate science, since no previous
351 quantitative estimates of Holocene temperature changes are available from southern Spitsbergen,
352 which is strategically located at the crossing of major warm and cold oceanic currents and where the
353 Polar Front develops. Therefore, knowledge of climate trends and the processes controlling them at
354 our study site provides significant understanding for circum-Arctic and global climate dynamics.

355

356 **Acknowledgements**

357

358 This work was supported by the Academy of Finland [QUAL-project (A.E.K. Ojala), grant number
359 259343] and the Emil Aaltonen Foundation (T.P. Luoto, grant numbers 160156, 170161). We are
360 grateful to Willem van der Bilt for providing data for Figure 1 and for the Polish Polar Station in
361 Hornsund for logistical support during the research. Constructive comments from two anonymous
362 journal reviewers helped to improve the manuscript.

363

364 **References**

365

366 Abdi H, Williams LJ. 2010. Principal component analysis. Wiley Interdisciplinary Reviews.
367 *Computational Statistics* **2**: 433–459.

368

369 Arppe L, Kurki E, Wooller MJ, Luoto TP, Zajączkowski M, Ojala AEK. 2017. A 5500-year oxygen
370 isotope record of high-arctic environmental change from southern Spitsbergen. *The Holocene* (in
371 press).

372

373 Bakke J, Dahl SO, Paasche O, Simonsen JR, Kvisvik B, Bakke K, Nesje A. 2010. A complete
 374 record of Holocene glacier variability at Austre Okstindbreen, northern Norway: an integrated
 375 approach. *Quaternary Science Reviews* **29**: 1246–1262.
 376

377 Bakke J, Lie O, Nesje A, Dahl SO, Paasche O. 2005. Utilizing physical sediment variability in
 378 glacier-fed lakes for continuous glacier reconstructions during the Holocene, northern Folgefonna,
 379 western Norway. *The Holocene* **15**: 161–176.
 380

381 Berner KS, Koç N, Godtlielsen F. 2010. High frequency climate variability of the Norwegian
 382 Atlantic Current during the early Holocene period and a possible connection to the Gleissberg
 383 cycle. *The Holocene* **20**: 245–255.
 384

385 Birks HH. 1991. Holocene vegetational history and climatic change in west Spitsbergeneplant
 386 macrofossils from Skardtjørna, an Arctic lake. *The Holocene* **1**: 209–218.
 387

388 Bond G, Kromer B, Beer J. et al. 2001. Persistent solar influence on North Atlantic climate during
 389 the Holocene. *Science* **294**: 2130–2136.
 390

391 Brooks SJ. 2006. Fossil midges (Diptera: Chironomidae) as palaeoclimatic indicators for the
 392 Eurasian region. *Quaternary Science Reviews* **25**: 1894–1910.
 393

394 Brooks SJ, Birks HJB. 2004. The dynamics of Chironomidae (Insecta: Diptera) assemblages in
 395 response to environmental change during the past 700 years on Svalbard. *Journal of*
 396 *Paleolimnology* **31**: 483–498.
 397

398 Brooks SJ, Birks HJB. 2001. Chironomid-inferred air temperatures from Late-glacial and Holocene
399 sites in north-west Europe: Progress and problems. *Quaternary Science Reviews* **20**: 1723–1741.
400

401 Brooks SJ, Birks HJB. 2000. Chironomid-inferred late-glacial and early-Holocene mean July air
402 temperatures for Krakenes Lake, western Norway. *Journal of Paleolimnology* **23**: 77–89.
403

404 Brooks SJ, Langdon PG, Heiri O. 2007. *The identification and use of Palaeoarctic Chironomidae*
405 *larvae in palaeoecology*. QRA Technical Guide No. 10, London, Quaternary Research Association,
406 276 p.
407

408 Crowley TJ. 2000. Causes of climate change over the past 1000 years. *Science* **289**: 270–277.
409

410 D'Andrea WJ, Vaillencourt DA, Balascio NL, Werner A, Roof SR, Retelle M, Bradley RS. 2012.
411 Mild Little Ice Age and unprecedented recent warmth in an 1800 year lake sediment record from
412 Svalbard. *Geology* **40**: 1007–1010.
413

414 Dylmer CV, Giraudeau J, Eynaud F, Husum K, De Vernal A. 2013. Northward advection of
415 Atlantic water in the eastern Nordic Seas over the last 3000 yr. *Climate of the Past* **9**: 1505–1518.
416

417 Eggermont H, Heiri O. 2012. The chironomid - temperature relationship: expression in nature and
418 palaeoenvironmental implications. *Biological Reviews* **87**: 430–456.
419

420 Heiri O, Lotter AF. 2001. Effect of low count sums on quantitative environmental reconstructions:
421 an example using subfossil chironomids. *Journal of Paleolimnology* **26**: 343–350.
422

423 Humlum O, Elberling B, Hormes A, Fjordheim K, Hansen OH, Heinemeier J. 2005. Late-Holocene
 424 glacier growth in Svalbard, documented by subglacial relict vegetation and living soil microbes.
 425 *The Holocene* **15**: 396–407.

426

427 Isaksson E, Divine D, Kohler J, Martma T, Pohjola V, Motoyama H, Watanabe O. 2005. Climate
 428 oscillations as recorded in Svalbard ice core $\delta^{18}\text{O}$ records between AD 1200 and 1997. *Geografiska*
 429 *Annaler: Series A, Physical Geography* **87**: 203–214.

430

431 Kaufman DS, Schneider DP, McKay NP. et al. 2009. Recent warming reverses long-term Arctic
 432 cooling. *Science* **325**: 1236–1239.

433

434 Kivilä H. 2014. *Late Holocene climate development of Bjørnøya, Svalbard, based on chironomid*
 435 *analysis*. Master's thesis, University of Helsinki, Department of Geosciences and Geography, 72 p.

436

437 Kurki E. 2016. *The oxygen isotope composition of chironomid chitin as a proxy for oxygen isotope*
 438 *composition of lake water and related climate parameters in southern Spitsbergen, Svalbard,*
 439 *during mid- and late Holocene*. Master's thesis, University of Helsinki, Department of Geosciences
 440 and Geography, 54 p.

441

442 Landerer FW, Wiese DN, Bentel K, Boening C, Watkins MM. 2015. North Atlantic meridional
 443 overturning circulation variations from GRACE ocean bottom pressure anomalies. *Geophysical*
 444 *Research Letters* **42**: 8114–8121.

445

446 Langebroek PM, Werner M, Lohmann G. 2011. Climate information imprinted in oxygen-isotopic
 447 composition of precipitation in Europe. *Earth and Planetary Science Letters* **311**: 144–154.

448

449 Lecavalier BS, Milne GA, Vinther BM, Fisher DA, Dyke AS, Simpson MJ. 2013. Revised
450 estimates of Greenland ice sheet thinning histories based on ice-core records. *Quaternary Science*
451 *Reviews* **63**: 73–82.

452

453 Lubinsky DJ, Forman SL, Miller GH. 1999. Holocene glacier and climate fluctuations on Franz
454 Josef Land, Arctic Russia, 80 °N. *Quaternary Science Reviews* **18**: 85–108.

455

456 Luoto TP, Nevalainen L. 2017. Temperature-precipitation relationship of the Common Era in
457 Northern Europe. *Theoretical and Applied Climatology*, DOI: 10.1007/s00704-017-2139-0.

458

459 Luoto TP, Nevalainen L. 2016. Quantifying climate changes of the Common Era for Finland.
460 *Climate Dynamics*, DOI: 10.1007/s00382-016-3468-x.

461

462 Luoto TP, Nevalainen L. 2015. Late Holocene precipitation and temperature changes in Northern
463 Europe linked with North Atlantic forcing. *Climate Research* **66**: 37–48.

464

465 Luoto TP, Kuhry P, Holzkämper S, Solovieva N, Self AE. 2017. A 2000-year record of lake
466 ontogeny and climate variability from the north-eastern European Russian Arctic. *The Holocene* **27**:
467 339–348.

468

469 Luoto TP, Oksman M, Ojala AEK. 2016. Invertebrate communities of the High Arctic ponds in
470 Hornsund. *Polish Polar Research* **37**: 105–119.

471

472 Luoto TP, Oksman M, Ojala AEK. 2015. Climate change and bird impact as drivers of High Arctic
 473 pond deterioration. *Polar Biology* **38**: 357–368.

474

475 Luoto TP, Brooks SJ, Salonen VP. 2014. Ecological responses to climate change in a bird-impacted
 476 High Arctic pond (Nordaustlandet, Svalbard). *Journal of Paleolimnology* **51**: 87–97.

477

478 Luoto TP, Nevalainen L, Kubischta F, Kultti S, Knudsen KL, Salonen VP. 2011. Late Quaternary
 479 ecological turnover in high arctic Lake Einstaken, Nordaustlandet, Svalbard (80° N). *Geografiska*
 480 *Annaler: Series A, Physical Geography* **93**: 337–354.

481

482 Majewski W, Szczuciński W, Zajączkowski M. 2009. Interactions of Arctic and Atlantic
 483 water-masses and associated environmental changes during the last millennium, Hornsund (SW
 484 Svalbard). *Boreas* **38**: 529–544.

485

486 Marsz AA, Styszyńska A. (eds) 2013. *Climate and climate change at Hornsund, Svalbard*. Gdynia,
 487 Gdynia Maritime University, 402 p.

488

489 Miller GH, Brigham-Grette J, Alley RB et al. 2010. Temperature and precipitation history of the
 490 Arctic. *Quaternary Science Reviews* **29**: 1679–1715.

491

492 Ojala AEK, Arppe L, Luoto TP, Wacker L, Kurki E, Zajączkowski M, Pawłowska J, Damrat M,
 493 Oksman M. 2016. Sedimentary environment, lithostratigraphy and dating of sediment sequences
 494 from Arctic lakes Revvatnet and Svartvatnet in Hornsund, Svalbard. *Polish Polar Research* **37**: 23–
 495 48.

496

497 Pawłowska J, Zajączkowski M, Łacka M, Lejzerowicz F, Esling P, Pawlowski J. 2015.
 498 Palaeoceanographic changes in Hornsund Fjord (Spitsbergen, Svalbard) over the last millennium:
 499 new insights from ancient DNA. *Climate of the Past Discussions* **11**: 1–34.
 500
 501 Pithan F, Mauritsen T. 2014. Arctic amplification dominated by temperature feedbacks in
 502 contemporary climate models. *Nature Geoscience* **7**: 181–184.
 503
 504 Polyak L, Alley RB, Andrews JT et al. 2010. History of sea ice in the Arctic. *Quaternary Science*
 505 *Reviews* **29**: 1757–1778.
 506
 507 Pryor AJ, Stevens RE, O'Connell TC, Lister JR. 2014. Quantification and propagation of errors
 508 when converting vertebrate biomineral oxygen isotope data to temperature for palaeoclimate
 509 reconstruction. *Palaeogeography, Palaeoclimatology, Palaeoecology* **412**: 99–107.
 510
 511 Røthe TO, Bakke J, Vasskog K, Gjerde M, D'Andrea WJ, Bradley RS. 2015. Arctic Holocene
 512 glacier fluctuations reconstructed from lake sediments at Mitrahalvøya, Spitsbergen. *Quaternary*
 513 *Science Reviews* **109**: 111–125.
 514
 515 Sarnthein M, Van Kreveld S, Erlenkeuser H, Grootes PM, Kucera M, Pflaumann U, Schulz M.
 516 2003. Centennial-to-millennial-scale periodicities of Holocene climate and sediment injections off
 517 the western Barents shelf, 75°N. *Boreas* **32**: 447–461.
 518
 519 Spielhagen RF, Werner K, Sørensen SA, Zamelczyk K, Kandiano E, Budeus G, Husum K,
 520 Marchitto TM, Hald M. 2011. Enhanced modern heat transfer to the Arctic by warm Atlantic water.
 521 *Science* **331**: 450–453.

522

523 Steinhilber F, Beer J, Fröhlich C. 2009. Total solar irradiance during the Holocene. *Geophysical*
524 *Research Letters* **36**: L19704.

525

526 Stroeve JC, Serreze M, Drobot S, Gearheard S, Holland M, Maslanik J, Meier W, Scambos T. 2008.
527 Arctic sea ice extent plummets in 2007. *Eos, Transactions of the American Geophysical Union* **89**:
528 13–14.

529

530 Svendsen JJ, Mangerud J. 1997. Holocene glacial and climatic variations on Spitsbergen, Svalbard.
531 *The Holocene* **7**: 45–57.

532

533 Trouet V, Esper J, Graham NE, Baker A, Scourse JD, Frank DC. 2009. Persistent positive North
534 Atlantic Oscillation mode dominated the Medieval Climate Anomaly. *Science* **324**: 78–80.

535

536 van der Bilt WG, Bakke J, Vasskog K, D'Andrea WJ, Bradley RS, Ólafsdóttir S. 2015.
537 Reconstruction of glacier variability from lake sediments reveals dynamic Holocene climate in
538 Svalbard. *Quaternary Science Reviews* **126**: 201–218.

539

540 Velle G, Kongshavn K, Birks HJB. 2011. Minimizing the edge-effect in environmental
541 reconstructions by trimming the calibration set: Chironomid-inferred temperatures from
542 Spitsbergen. *The Holocene* **21**: 417–430.

543

544 Verbruggen F, Heiri O, Reichert GJ, Blaga C, Lotter AF. 2011. Stable oxygen isotopes in
545 chironomid and cladoceran remains as indicators for lake-water $\delta^{18}\text{O}$. *Limnology and*
546 *Oceanography* **56**: 2071–2079.

547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571

Wang M, Overland JE. 2009. A sea ice free summer Arctic within 30 years? *Geophysical Research Letters* **36**: L07502.

Wang Y, Francis DR, O'Brien DM, Wooller MJ. 2008. A protocol for preparing subfossil chironomid head capsules (Diptera: Chironomidae) for stable isotope analysis in paleoclimate reconstruction and considerations of contamination sources. *Journal of Paleolimnology* **40**: 771–781.

Wanner H, Beer J, Buetikofer J. et al. 2008. Mid- to Late Holocene climate change: an overview. *Quaternary Science Reviews* **27**: 1791–1828.

Wooller MJ, Francis D, Fogel ML, Miller U, Walker IR, Wolfe AP. 2004. Quantitative paleotemperature estimates from $\delta^{18}\text{O}$ of chironomid head capsules preserved in arctic lake sediments. *Journal of Paleolimnology* 31: 267–274.

Zawiska I, Luoto TP, Nevalainen L, Tylmann W, Jensen TC, Obremaska M, Słowiński M, Woszczyk M, Schartau AK, Walseng B. 2017. Climate variability and lake ecosystem responses in western Scandinavia (Norway) during the last Millennium. *Palaeogeography, Palaeoclimatology, Palaeoecology* **466**: 231–239.

572 **Figure captions**

573

574 **Figure 1.** Map. Lake Svartvatnet (76.895°N, 15.676°E, 63 m a.s.l.) is located 12 km south of the
575 Polish Polar Station on Hornsund fjord, Svalbard.

576

577 **Figure 2.** Chironomid assemblages. Relative abundance of chironomid taxa and the principal
578 component analysis (PCA) axis 1 scores from the Svartvatnet sediment sequence.

579

580 **Figure 3.** Reconstructions. Principal component analysis (PCA) axis 1 scores for fossil chironomid
581 assemblages, chironomid-based mean July air temperature reconstruction (CiT), lake water $\delta^{18}\text{O}$
582 values derived from fossil chironomid head capsules and annual air temperature reconstruction
583 based on the isotope records in Svartvatnet. The $\delta^{18}\text{O}$ -based reconstruction, represented as
584 anomalies from the record mean, should not be considered directly as a MAT record (discussed by
585 Arppe *et al.* (2017) in more detail). The Svartvatnet data are compared with proxy records of
586 average total solar irradiance (TSI) relative to the solar cycle minimum of the year AD 1986
587 (Steinhilber *et al.*, 2009), summer sea-surface temperature from northern Norwegian Sea (Sarnthein
588 *et al.*, 2003) and glacier variability in Svalbard (van der Bilt *et al.*, 2015). Cold climate episodes are
589 marked in gray.

590

591 **Supporting Information Figure S1.** Age-depth model originally published in Ojala *et al.* (2016).
592 Dating of the topmost sediments is based on ^{210}Pb and ^{137}Cs (solid black line). Calibrated AMS ^{14}C
593 dates with 95.4% confidence levels (2s) are given in darker gray, black crosses represent their 2s
594 median probability and black dots represent paleosecular variation dates. The age-depth relationship
595 shaded with light gray is based on a Bayesian P-sequence deposition model.