<https://helda.helsinki.fi>

Determination of the intramammary dose of benzylpenicillin required to maintain an adequate concentration in the milk to inhibit Gram-positive bacteria in the clinically normal udder for 24 hr

Abbeloos, Elke

2018-10

Abbeloos , E , Pyörälä , S , Rajala-Schultz , P & Myllys , V 2018 , ' Determination of the intramammary dose of benzylpenicillin required to maintain an adequate concentration in the milk to inhibit Gram-positive bacteria in the clinically normal udder for 24 hr ' , Journal of Veterinary Pharmacology and Therapeutics , vol. 41 , no. 5 , pp. 691-698 . <https://doi.org/10.1111/jvp.12671>

<http://hdl.handle.net/10138/323998>

<https://doi.org/10.1111/jvp.12671>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Full title: Determination of the intramammary dose of benzylpenicillin required to maintain an adequate concentration in the milk to inhibit Gram-positive bacteria in the clinically normal udder for 24 hours.

Short title: Determination of the intramammary dose of penicillin

Elke Abbeloos¹, Satu Pyörälä², Päivi Rajala-Schultz², Vesa Myllys³

¹ Boehringer Ingelheim Vetmedica. Ingelheim, Germany

² Department of Production Animal Medicine, Faculty of Veterinary Medicine, University of Helsinki, Paroninkuja 20, FI-04920 Saarentaus, Finland.

³ Vetcare Ltd, P.O.Box 26, FI-04601 Mäntsälä, Finland

Corresponding author: elke.abbeloos@boehringer-ingelheim.com

Other authors: vesa.myllys@vetcare.fi

Abstract

The aim of this study was to determine the intramammary dose of benzylpenicillin required to maintain a concentration in the milk above the MIC for the Gram-positive bacteria that cause mastitis. The product used in this study was a commercially available procaine benzylpenicillin in an oily suspension with micronized particles. Three dose levels were used: 200 000, 300 000 and 600 000 IU. Concentrations of benzylpenicillin in cow milk and plasma were determined after a single intramammary dose was administered into one quarter of each of the 5 cows in each treatment group. Samples were analyzed using an HPLC-MS/MS method which was validated during the study. Concentrations in the milk were well above the MIC for the target pathogens for all doses tested. There was a linear dose-dependent increase in the mean AUCs of benzylpenicillin concentrations in plasma and milk. At the first milking, 12 hours after dosing, there was a significant difference between the mean milk benzylpenicillin concentrations in cows treated with a dose of 600 000 IU, and

those treated with 200 000 or 300 000 IU. Although this study shows a linear relationship between the dose of procaine benzylpenicillin administered and the concentration in the milk in the healthy udder, it would be useful to conduct studies on cows with mastitis to define the optimum dose and duration of intramammary treatment with benzylpenicillin.

Keywords: penicillin, intramammary, clinical mastitis, Gram-positive bacteria, β -lactam antibiotics

Introduction

Antimicrobial resistance is an emerging issue, so the use of these drugs in food-producing animals is being scrutinized worldwide (Laxminarayan, Duse et al., 2013). Globally, the most important indication for antibiotic use on dairy farms is mastitis (Pol & Ruegg, 2007; Stevens, Piepers et al., 2016). Benzylpenicillin is considered to be the first-line treatment for mastitis caused by Gram-positive bacteria because of its narrow spectrum, according to the European Union's responsible use guidelines (European_Commission, 2015).

Benzylpenicillin is a β -lactam antibiotic with bactericidal action against the major Gram-positive bacteria often found in mastitis, such as *Streptococcus* spp. and *Staphylococcus* spp. (Prescott, 2013). All β -lactam antibiotics have a time-dependent mode of action, meaning that treatment success mainly depends on the length of time that the drug remains above the MIC (Turnidge, 1998; Erskine, Wagner et al., 2003; Prescott, 2013)

Intramammary infusion is the most common route of administration for the antibiotics used to treat mastitis (Gruet, Maincent et al., 2001). The aim of this treatment is to target the actual infection site by administering the antibiotic into the udder, and for the concentration of the drug to reach levels above the MIC for the bacteria causing the infection. The active ingredient must not only distribute well into the milk phase inside the udder, but also be absorbed into the tissue if the infective agent is an invasive (and potentially intracellular)

49 pathogen, such as *Staphylococcus aureus* (Ziv, 1980a). The distribution of penicillin
50 administered via the intramammary route was investigated in an isolated perfused model
51 by Ehinger and Kietzmann (2000). Penicillin in oily suspension with small particles
52 (micronized form) was found to have the best glandular distribution compared to aqueous
53 solution or oily suspension with large particles. Absorption into the perfusate was also
54 greatest after treatment with the oily suspension with small particles, delivering the highest
55 areas under the absorption-time curves (Ehinger & Kietzmann, 2000).

56 As a result, the decision was taken to use benzylpenicillin in oily suspension with small
57 particle size to formulate a new intramammary penicillin product. According to the European
58 Medicines Agency (EMA) guidelines on the conduct of pharmacokinetic studies in target
59 animal species, the dose of the active substance needs to be determined using the
60 concentration of penicillin in the milk as a function of time to allow the therapeutic
61 concentration-time profile at the infection site in the udder to be estimated. The same
62 procedure should be applied to the plasma to estimate potential systemic absorption. Dose
63 determination studies usually require three different dose levels to be tested and are typically
64 performed in healthy animals (EMA, 2000).

65 The aim of this study was to determine the intramammary dose of benzylpenicillin required
66 to maintain a concentration in the milk above the MIC for 24 hours for Gram-positive
67 mastitis pathogens. Concentrations of benzylpenicillin in cow milk and plasma were
68 measured following a single intramammary administration of benzylpenicillin procaine at
69 three dose levels (200 000, 300 000 and 600 000 IU) into one udder quarter. The study
70 was conducted in 2005 and followed the principles of the EMA guidelines for the conduct
71 of pharmacokinetic studies in target animal species that were in force at the time (EMA,
72 2000) . The product was originally authorized for use in only a few countries, but in 2017

an identical copy of it was approved for use in fourteen countries of the European Union through the decentralized procedure.

Materials and Methods

Subjects and product administration

Fifteen clinically healthy, 2 to 5-year-old Holstein cows were selected for this study. The cows were between the 3rd and 6th month of lactation and had not received any antibiotic treatment in the 8 days preceding the trial. The average daily milk yield was 32 liters (range 22.9–43.5) at the beginning of the study. All animals were housed on the farm of origin, which was also the site of the study, but an acclimation period of 7 days was applied to allow pre-trial observation of the cows. Each cow was examined clinically once daily throughout the entire acclimation and study phase: udders were inspected and palpated, and milk was checked for the appearance of clots or flakes, and changes in color and consistency. Milk production was recorded at each milking. During the acclimation period, the cows were ranked by lactation stage, then randomly allocated into five lactation stage groups, each containing three animals, to ensure similar distribution of days in milk across the groups. Computerized random number generation was performed with Microsoft Excel® (Microsoft Corporation, Redmond, WA, USA) to randomly allocate one animal out of each lactation stage group to each of three treatment groups. More details on the animals involved in the study can be found in Table 1.

Include Table 1 here.

The test product was a registered intramammary suspension (Carepen®, Vetcare Oy, Mantsala, Finland, also registered as Ubropen®, Boehringer Ingelheim Animal Health, Germany) containing 600 000 IU (equivalent to 600 mg) of micronized procaine

benzylpenicillin per 10 mL in an oily suspension. The product was administered intramammarily into the left front quarter. All animals were treated once after the morning milking on Day 0. For Group 1 (200 000 IU penicillin) and Group 2 (300 000 IU penicillin), two-thirds and one half, respectively, of the contents of a Carepen® /Ubropen® tube was discarded before administration to obtain the correct dose of benzylpenicillin procaine per cow. The full contents of the Carepen®/Ubropen® tube was administered to animals in Group 3. Tubes were weighed before and after administration to determine the actual dose given. Teats were properly cleaned with individual antiseptic wipes before administration.

The study was carried out in compliance with French legislation on the protection of laboratory animals, and in accordance with a valid license for experiments on vertebrate animals issued by the French Ministry for Agriculture. The study passed the ethical review committee of Avogadro (Fontenilles, France), the contract research organization that was responsible for the trial (Reference code of the study: A051143).

Milk and blood sampling

Before dosing, mixed milk samples were collected from all cows on day 0 from the milk obtained during regular milking. On day 1, milk samples from treated quarters were collected at 12, 24, 48 and 72 hours after dosing. The foremilk from the treated quarter was discarded and the teat was thoroughly cleaned and disinfected before each sample was collected. Three samples of about 5 mL of milk were collected from each treated quarter at each sampling session. The cow's total milk yield was also recorded at each sampling session. Samples were transported to the analytical laboratory (Avogadro, Fontenilles, France) at about 4 °C and frozen (to about -20 °C) as soon as possible.

Blood samples were collected from the jugular or tail vein before treatment and at 3, 6, 12, 15, 18 and 24 hours after administration. Blood (10 mL) was collected in lithium heparin tubes which were kept at 4 °C until centrifugation. They were centrifuged at around 2500 g for 10 minutes at 4 °C and 1.5 mL of plasma from each sample was transferred to each of 3 propylene tubes.

Quantification of benzylpenicillin

Samples were processed by the laboratory within 5 days of arrival. A stability test was performed, indicating that benzylpenicillin was stable in plasma at -20 °C for at least 9 days in storage (max. 20% degradation). Samples were analyzed by HPLC-MS/MS after precipitation of the protein with acetonitrile and solid-phase extraction with an OASIS® HLB cartridge (Waters Corporation, Milford, MA, USA). The Lower Limit of Quantification (LLOQ) of the testing method in milk and plasma was 2 ng/mL. For comparison, the Maximum Residue Limit (MRL) in the European Union is set at 4 µg/kg (or 4 ng/g) for milk (EMA, 1999). Specificity against endogenous substances was evaluated by carefully assessing the ion chromatograms from plasma from control cows at the retention times of benzylpenicillin and phenoxymethylpenicillin (the latter being the internal standard). No interfering peaks were observed at these times on the ion chromatograms from six control samples. The HPLC-MS/MS method used to determine the benzylpenicillin in bovine milk and plasma was validated during the assay. Linearity was verified from 2 ng/mL to 400 ng/mL of benzylpenicillin in plasma. The extraction recovery from 6 spiked plasma samples showed that benzylpenicillin and phenoxymethylpenicillin were efficiently extracted from bovine plasma (mean recovery 69.8%, coefficient of variation (CV) 3.9%). The matrix effect after replicate analyses (n=2) on benzylpenicillin and phenoxymethylpenicillin from plasma was deemed negligible (-6.3% and +2.7% respectively). The precision and accuracy evaluated at 3 concentration levels of benzylpenicillin were within an acceptance range. The highest

acceptable CV was set at 20% for the quality control (QC) LLOQ and 15% for other concentrations. The highest acceptable percentage of error was set at +/- 20% for the LLOQ and +/- 15% for other concentrations. Milk QC samples would be rejected if the concentration deviated from the theoretical value by more than -30% to +10%, or if an unsatisfactory chromatographic peak was obtained, or in the event of a manipulation error. During each batch analysis, at least 4 of the 6 QC samples were within this range, and at least 5 calibration points were included in the final calculation of the calibration curve parameters. Analysis of milk and plasma specimens was deemed acceptable as the calibration curve and the quality controls included in the analytical batches were within their respective acceptance criteria.

Data analysis

Statgraphics Plus® (Statistical Graphics, Rockville, USA) was used for statistical analysis for calibration. The linearity of the response in plasma and milk was evaluated with 3 calibration curves. Each calibration curve included a blank matrix, a zero matrix and six levels of concentration (2, 5, 25, 75, 200 and 400 ng of benzylpenicillin per mL of plasma). The calibration curve was obtained by least-squares regression of the calculated response versus theoretical concentration using several regression models (untransformed, $1/x$, $1/x^2$). ANOVA, including the lack-of-fit test, was performed on the regression curve, and linearity was determined based on examination of the residuals; outliers were checked and discarded when the absolute value of the residual was above 2. Linear adjustment by the least-squares mean method with a weighting factor of $1/x^2$ was determined to be the most appropriate model to express the relationship between the response and the theoretical concentration in bovine plasma. One outlier was discarded.

Areas Under the Curve (AUC) were calculated for benzylpenicillin concentrations in blood and milk using Microsoft Excel®. Concentrations in blood and milk were summarized at each time point into means and geometric means; the latter to reduce the influence of possible outliers. Benzylpenicillin concentrations in milk were log-transformed for easier visual interpretation. Further statistical analysis was performed using SAS® software version 9.4 (SAS Institute Inc., Cary, NC, USA). Results were considered significant at alpha=0.05 level. A simple linear regression model was fitted with AUC as the outcome variable and “actual administered dose” as an explanatory variable. A non-parametric test (the Jonckheere-Terpstra test; (Terpstra & Magel, 2003) was also performed at the given time points to evaluate whether benzylpenicillin concentrations in plasma and milk differed between treatment groups. Measurements that were below the quantification limit were treated as zero in the analysis.

Results

No adverse reactions, clinical signs or changes in daily milk production were observed after treatment. The mean dose of penicillin actually administered to each treatment group is shown in Table 1.

Concentration in plasma

The mean benzylpenicillin concentrations determined in the plasma over 24 hours are presented in Figure 1. A linear dose-dependent increase was seen in the mean AUCs for benzylpenicillin in plasma (corrected by the actual administered dose in IU/kg bodyweight; $p < 0.0001$) (Figure 2). The Jonckheere-Terpstra test demonstrated that increasing the dose increased the benzylpenicillin in the plasma at 6, 12, 15 and 18 hours after dosing (p -values 0.0002; < 0.0001 ; 0.0001; and 0.0037 respectively). The concentration in the plasma dropped below the quantification limit at 24 hours in all dosage regimens (data not shown).

192 Include Figure 1 and Figure 2 here

193 Concentration in milk

194 The mean and geometric mean of benzylpenicillin concentrations in samples of milk from
195 treated quarters are presented in Table 2. Mean log concentrations of benzylpenicillin in milk
196 are presented in Figure 3. At the first milking, 12 hours after dosing, there was evidence that
197 increasing the benzylpenicillin dose administered to the mammary quarter increased milk
198 benzylpenicillin concentrations ($p=0.002$ in the Jonckheere-Terpstra test). The difference
199 was not significant at the second milking, 24 hours after dosing, and fourth milking, 48 hours
200 after dosing ($p=0.234$ and $p=0.500$ respectively). A linear dose-dependent increase was
201 seen in the mean AUCs of benzylpenicillin concentrations in milk (corrected by the actual
202 administered dose in IU/kg bodyweight; $p=0.046$) (Figure 4).

203
204 Include Table 2 and Figure 3 here

205 **Discussion and conclusion**

206 The aim of intramammary benzylpenicillin administration is to achieve concentrations of
207 the active ingredient above the MIC for the relevant pathogens. These concentrations
208 should be reached at the site of infection (in the udder) for long enough to eliminate the
209 infection, while minimizing the exposure of commensal bacteria (such as those found in
210 the gut) to the drug to reduce the risk of selecting resistant bacteria . We believe that this
211 is the first validated study on the pharmacokinetics of benzylpenicillin after intramammary
212 administration.

213 In Europe, the most relevant Gram-positive udder pathogens isolated in mastitis and treated with
214 antimicrobials are *Staphylococcus aureus* and *Streptococcus uberis* (Tenhagen, Hansen et

al., 2009; Verbeke, Piepers et al., 2014; Santman-Berends, Lam et al., 2015; Vakkamäki, Taponen et al., 2017). No clinical MIC breakpoints are available for penicillin in bovine mastitis. The epidemiological breakpoints for the susceptibility of staphylococci (including *S. aureus*), group A streptococci (*S. dysgalactiae*) and group B streptococci (*S. agalactiae*) to penicillin are defined by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) as 0.125 µg/mL. No such breakpoint is available for *S. uberis* (EUCAST, 2017), although 0.125 µg/mL is commonly accepted (Thomas, de Jong et al., 2015).

Most mastitis-causing streptococci have remained susceptible to benzylpenicillin (Hendriksen, Mevius et al., 2008; MARAN, 2008; SVARM, 2010; FINRES, 2015). Reduced susceptibility has been reported in a small proportion of *S. uberis* isolates (Haenni, Galofaro et al., 2010; FINRES, 2015). This finding probably has no clinical relevance, because the proportion of these isolates remains below 5% and their MIC is still much below the concentrations achieved by intramammary administration of penicillin (FINRES, 2015). *S. aureus* has developed penicillin resistance by producing β-lactamase. The proportion of β-lactamase-positive isolates collected from quarter milk samples of cows with mastitis in European countries varies from less than 10% to over 50% (Hendriksen, Mevius et al., 2008; Kalmus, Aasmäe et al., 2011; FINRES, 2015; Thomas, de Jong et al., 2015). Clinical mastitis caused by coagulase-negative staphylococci can also be treated with penicillin, and resistance among these is more common than in *S. aureus* (Taponen, Nykäsenoja et al., 2016). It is remarkable that Scandinavian countries which almost exclusively treat mastitis with benzylpenicillin have reported consistently very low or decreasing numbers of penicillin-resistant *S. aureus* isolates (Pitkälä, Haveri et al., 2004; Swedres-SVARM, 2014; FINRES, 2015; NORM-VET, 2016).

In the current study, penicillin concentration in the milk compartment exceeded the MIC for the relevant pathogens for at least 24 hours with all dosages. 24 hours is proposed to be a good administration interval for an intramammary antibiotic, since it requires only once a day manipulation of the teat and ensures compliance due to adaptation to different milking regimes (twice-a-day, three-times-a-day or voluntary milking systems). With the lower dose of 200 000 IU, the concentration at 24 hours was as much as 20 times greater than the MIC; with 300 000 IU, 22 times greater; and with 600 000 IU, 70 times greater. Variation in the 600 000 IU group was large, so the geometric mean, which was 35 times greater than the MIC, may describe the results more accurately.

For time-dependent antibiotics such as benzylpenicillin, maximum kill *in vitro* is achieved at 3 to 4 times the MIC, and the most important predictor for elimination of infection is the time above the MIC (Turnidge, 1998). In light of the paradoxically reduced bactericidal activity of high doses of penicillin (first described by Eagle and Musselman (1948), and known as the “Eagle effect”), our study indicates that all doses used could possibly be unnecessarily high. It must be noted that the paradoxically inferior effect of high doses has been observed with most β -lactam antibiotics, mainly with Gram-positive bacteria, and it is not always synonymous with the classic Eagle effect (Odenholt-Tornqvist, 1988; Hamilton-Miller & Shah, 1999). Penicillin has been used to treat mastitis for over 50 years, at lower doses than at present (Le Louedec, 1978) and early studies showed that doses as low as 100 000 IU (every milking) resulted in high cure rates (Edwards, 1962; Le Louedec, 1978). However, the main target pathogen at the time of those studies was *S. agalactiae*, which responded very well to treatment (Edwards, 1962).

There are more arguments to support the higher dosing regimen. Some caution is warranted when interpreting the results, as the only samples taken consisted of foremilk. Studies with another β -lactam antibiotic (cephapirin) have shown that foremilk contains higher

concentrations of the drug than subsequent milk, potentially overestimating the drug concentration reached in the entire udder (Stockler, Morin et al., 2009).

The results of the current study also reveal considerable variation in benzylpenicillin concentrations in milk between cows in the same treatment group. The same finding was reported by Bjorland, Waage et al. (1998) who compared two intramammary doses of benzylpenicillin in five cows. Moreover, one target pathogen, *S. aureus*, is known to penetrate into udder tissue (Almeida, Matthews et al., 1996), and penicillin is also transferred into this tissue in the usual pharmacokinetic manner (Ziv, 1980b). Penicillin concentrations in udder tissue were not measured in this study, so we can only speculate on the penetration of penicillin into deeper udder tissue and whether effective concentrations accumulate there. However, Ehringer and Kietzmann (2000) noticed that the concentration of benzylpenicillin in the tissue decreased exponentially with increasing vertical distance from the teat base, and they found that benzylpenicillin concentration in the tissues of the lower part of the udder was around 1/10th of that of the upper part of the udder. This could be the reason why a field trial which studied clinical mastitis caused by *S. aureus* demonstrated significantly better cure rates when intramammary treatment was supplemented with systemic penicillin, compared to intramammary treatment alone (Taponen, Jantunen et al., 2003).

Finally, the pharmacokinetics of benzylpenicillin in healthy animals could be significantly different from cows suffering from mastitis, as the pharmacokinetic properties of active substances are determined in healthy animals (EMA, 2000). Pathological changes in cows with clinical mastitis, including swelling and occlusion of the ducts in the mammary gland, may result in uneven distribution of the drug and the associated risk of lower local penicillin concentrations (Ullberg, Hansson et al., 1958; Ziv, 1980b). In an experimental *S. uberis* infection, inflammatory changes caused widespread occlusion of the mammary gland secretory system (Pedersen, Aalbaek et al., 2003). Inflammation in the udder changes the

composition of the milk and damages the blood-milk barrier, increasing the permeability of the blood and milk compartments and potentially accelerating elimination of the drug (Erskine, Wilson et al., 1995; Zhao & Lacasse, 2008).

In conclusion, our results showed a dose-dependent increase in the concentration of benzylpenicillin in plasma and milk when applied in an oily suspension with micronized particles. Although concentrations in the milk after administration of the lower doses were well above the MIC for the target pathogens, distribution of the active ingredient in the inflamed udder and the location of the pathogens inside the udder tissue must be taken into account. For this reason, the suggested dose for this product is the higher dose of 600 000 IU administered once a day. It would be useful to conduct field studies with naturally occurring intramammary infections to define the optimum dose and duration of intramammary treatment with benzylpenicillin.

Acknowledgments

The study was conducted by Avogadro, Fontenilles, France

References

- Almeida, R.A., Matthews, K.R., Cifrian, E., Guidry, A.J. & Oliver, S.P. (1996) Staphylococcus aureus Invasion of Bovine Mammary Epithelial Cells. *Journal of Dairy Science*, **79**(6), 1021-1026.
- Bjorland, J., Waage, S., Oedegaard, S., Yndestad, M. & Hormazabal, V. (1998) Concentrations of benzylpenicillin in milk of dairy cows after intramammary administration. *Norsk Veterinaertidsskrift (Norway)*.
- Eagle, H. & Musselman, A.D. (1948) The rate of bactericidal action of penicillin in vitro as a function of its concentration, and its paradoxically reduced activity at high concentrations against certain organisms. *The Journal of Experimental Medicine*, **88**(1), 99-131.
- Edwards, S. (1962) Antibiotics in the treatment of mastitis. *Antibiotics in Agriculture. M. Wood-bine, Ed*, 43-57.

- 313 Ehinger, A.M. & Kietzmann, M. (2000) Tissue distribution of benzylpenicillin after intramammary
314 administration in the isolated perfused bovine udder. *Journal of Veterinary Pharmacology and*
315 *Therapeutics*, **23**(5), 303-310.
- 316 EMA (1999). *Committee for Veterinary Medicinal Products Penethamate Summary report*. In Committee for
317 Veterinary Medical Products Penethamate Summary report. European Agency for the Evaluation of
318 Medicinal Products.
- 319 EMA (2000). *Guidelines for the conduct of pharmacokinetic studies in target animal species*. In Guidelines
320 for the conduct of pharmacokinetic studies in target animal species.
- 321 Erskine, R., Wilson, R., Tyler, J., McClure, K., Nelson, R. & Spears, H. (1995) Ceftiofur distribution in serum
322 and milk from clinically normal cows and cows with experimental *Escherichia coli*-induced mastitis.
323 *American journal of veterinary research*, **56**(4), 481-485.
- 324 Erskine, R.J., Wagner, S. & DeGraves, F.J. (2003) Mastitis therapy and pharmacology. *Veterinary Clinics of*
325 *North America: Food Animal Practice*, **19**(1), 109-138.
- 326 EUCAST, E.C.o.A.S.T. (2017). *Antimicrobial wild type distributions of microorganisms*. In Antimicrobial wild
327 type distributions of microorganisms.
- 328 European_Commission (2015). *Commission notice*. In Commission notice Ed Union, E. pp. 7-26, Official
329 Journal of the European Union.
- 330 FINRES (2015). *FINRES-Vet 2010–2012. Finnish Veterinary Antimicrobial Resistance Monitoring and*
331 *Consumption of Antimicrobial Agents*. . In FINRES-Vet 2010–2012. Finnish Veterinary Antimicrobial
332 Resistance Monitoring and Consumption of Antimicrobial Agents. . 2/2015 edn. Finnish Food Safety
333 Authority Evira, Helsinki, Finland.
- 334 Gruet, P., Maincent, P., Berthelot, X. & Kaltsatos, V. (2001) Bovine mastitis and intramammary drug delivery:
335 review and perspectives. *Advanced Drug Delivery Reviews*, **50**(3), 245-259.
- 336 Haenni, M., Galofaro, L., Ythier, M., Giddey, M., Majcherczyk, P., Moreillon, P. & Madec, J.-Y. (2010)
337 Penicillin-binding protein gene alterations in *Streptococcus uberis* isolates presenting decreased
338 susceptibility to penicillin. *Antimicrobial agents and chemotherapy*, **54**(3), 1140-1145.
- 339 Hamilton-Miller, J.M.T. & Shah, S. (1999) Effect of antibiotic concentration on the killing of *Staphylococcus*
340 *aureus* and *Enterococcus faecalis*: comparison of the novel penem, Men 10700, with other β -lactam
341 antibiotics. *Journal of Antimicrobial Chemotherapy*, **44**(3), 418-420.
- 342 Hendriksen, R.S., Mevius, D.J., Schroeter, A., Teale, C., Meunier, D., Butaye, P., Franco, A., Utinane, A.,
343 Amado, A., Moreno, M., Greko, C., Stärk, K., Berghold, C., Myllyniemi, A.-L., Wasyl, D., Sunde, M. &
344 Aarestrup, F.M. (2008) Prevalence of antimicrobial resistance among bacterial pathogens isolated
345 from cattle in different European countries: 2002–2004. *Acta Veterinaria Scandinavica*, **50**(1), 28.
- 346 Kalmus, P., Aasmäe, B., Kärssin, A., Orro, T. & Kask, K. (2011) Udder pathogens and their resistance to
347 antimicrobial agents in dairy cows in Estonia. *Acta Veterinaria Scandinavica*, **53**(1), 4.

- 348 Laxminarayan, R., Duse, A., Wattal, C., Zaidi, A.K.M., Wertheim, H.F.L., Sumpradit, N., Vlieghe, E., Hara,
349 G.L., Gould, I.M., Goossens, H., Greko, C., So, A.D., Bigdeli, M., Tomson, G., Woodhouse, W.,
350 Ombaka, E., Peralta, A.Q., Qamar, F.N., Mir, F., Kariuki, S., Bhutta, Z.A., Coates, A., Bergstrom, R.,
351 Wright, G.D., Brown, E.D. & Cars, O. (2013) Antibiotic resistance—the need for global solutions. *The*
352 *Lancet Infectious Diseases*, **13**(12), 1057-1098.
- 353 Le Louedec, C. (1978). *Efficacité des antibiotiques contre les mammites bovines staphylococciques et*
354 *streptococciques*. In *Efficacité des antibiotiques contre les mammites bovines staphylococciques et*
355 *streptococciques* pp. 63-88.
- 356 MARAN (2008). *MARAN, 2007–2008. Monitoring of Antimicrobial Resistance and Antimicrobial Usage in*
357 *The Netherlands in 2006–2008*. . In MARAN, 2007–2008. Monitoring of Antimicrobial Resistance
358 and Antimicrobial Usage in The Netherlands in 2006–2008. .
- 359 NORM-VET, N. (2016). *Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in*
360 *Norway*. In *Usage of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Norway*.
361 Norwegian Veterinary Institute, Tromsø/ Oslo.
- 362 Odenholt-Tornqvist, I. (1988) Pharmacodynamics of Beta-Lactam Antibiotics: Studies on the Paradoxical and
363 Postantibiotic Effects in Vitro and in an Animal Model. *Scandinavian Journal of Infectious Diseases*,
364 **20**(sup58), 1-55.
- 365 Pedersen, L., Aalbaek, B., Røntved, C.M., Ingvarsen, K.L., Sorensen, N., Heegaard, P.M. & Jensen, H.
366 (2003) Early pathogenesis and inflammatory response in experimental bovine mastitis due to
367 *Streptococcus uberis*. *Journal of comparative pathology*, **128**(2-3), 156-164.
- 368 Pitkälä, A., Haveri, M., Pyörälä, S., Myllys, V. & Honkanen-Buzalski, T. (2004) Bovine mastitis in Finland
369 2001-Prevalence, distribution of bacteria, and antimicrobial resistance. *J Dairy Sci*, **87**.
- 370 Pol, M. & Ruegg, P.L. (2007) Treatment Practices and Quantification of Antimicrobial Drug Usage in
371 Conventional and Organic Dairy Farms in Wisconsin. *Journal of Dairy Science*, **90**(1), 249-261.
- 372 Prescott, J.F. (2013). *Beta-lactam Antibiotics*. In *Beta-lactam Antibiotics* pp. 153-173. John Wiley & Sons,
373 Inc.
- 374 Santman-Berends, I.M.G.A., Lam, T.J.G.M., Keurentjes, J. & van Schaik, G. (2015) An estimation of the
375 clinical mastitis incidence per 100 cows per year based on routinely collected herd data. *Journal of*
376 *Dairy Science*, **98**(10), 6965-6977.
- 377 Stevens, M., Piepers, S., Supré, K., Dewulf, J. & De Vliegher, S. (2016) Quantification of antimicrobial
378 consumption in adult cattle on dairy herds in Flanders, Belgium, and associations with udder health,
379 milk quality, and production performance. *Journal of Dairy Science*, **99**(3), 2118-2130.
- 380 Stockler, R.M., Morin, D.E., Lantz, R.K., Hurley, W.L. & Constable, P.D. (2009) Effect of milk fraction on
381 concentrations of cephapirin and desacetylcephapirin in bovine milk after intramammary infusion of
382 cephapirin sodium. *Journal of Veterinary Pharmacology and Therapeutics*, **32**(4), 345-352.

383 SVARM (2010). *SVARM, 2002–2010. Swedish Veterinary Antimicrobial Resistance Monitoring*. . In SVARM,
384 2002–2010. Swedish Veterinary Antimicrobial Resistance Monitoring. .

385 Swedres-SVARM (2014). *Consumption of antibiotics and occurrence of antibiotic resistance in Sweden*. In
386 Consumption of antibiotics and occurrence of antibiotic resistance in Sweden. Public Health Agency
387 of Sweden

388 Taponen, S., Jantunen, A., Pyörälä, E. & Pyörälä, S. (2003) Efficacy of targeted 5-day combined parenteral
389 and intramammary treatment of clinical mastitis caused by penicillin-susceptible or penicillin-resistant
390 *Staphylococcus aureus*. *Acta Veterinaria Scandinavica*, **44**(1), 53.

391 Taponen, S., Nykäsenoja, S., Pohjanvirta, T., Pitkälä, A. & Pyörälä, S. (2016) Species distribution and in
392 vitro antimicrobial susceptibility of coagulase-negative staphylococci isolated from bovine mastitic
393 milk. *Acta Veterinaria Scandinavica*, **58**(1), 12.

394 Tenhagen, B.A., Hansen, I., Reinecke, A. & Heuwieser, W. (2009) Prevalence of pathogens in milk samples
395 of dairy cows with clinical mastitis and in heifers at first parturition. *Journal of Dairy Research*, **76**.

396 Terpstra, J. & Magel, R. (2003) A new nonparametric test for the ordered alternative problem. *Journal of*
397 *Nonparametric Statistics*, **15**(3), 289-301.

398 Thomas, V., de Jong, A., Moyaert, H., Simjee, S., El Garch, F., Morrissey, I., Marion, H. & Vallé, M. (2015)
399 Antimicrobial susceptibility monitoring of mastitis pathogens isolated from acute cases of clinical
400 mastitis in dairy cows across Europe: VetPath results. *International Journal of Antimicrobial Agents*,
401 **46**(1), 13-20.

402 Turnidge, J.D. (1998) The Pharmacodynamics of β -Lactams. *Clinical Infectious Diseases*, **27**(1), 10-22.

403 Ullberg, S., Hansson, E. & Funke, H. (1958) Distribution of penicillin in mastitic udders following
404 intramammary injection; an auto-radiographic study. *American Journal of Veterinary Research*,
405 **19**(70), 84-92.

406 Ungemach, F.R., Müller-Bahrndt, D. & Abraham, G. (2006) Guidelines for prudent use of antimicrobials and
407 their implications on antibiotic usage in veterinary medicine. *International Journal of Medical*
408 *Microbiology*, **296**, Supplement 2, 33-38.

409 Vakkamäki, J., Taponen, S., Heikkilä, A.-M. & Pyörälä, S. (2017) Bacteriological etiology and treatment of
410 mastitis in Finnish dairy herds. *Acta Veterinaria Scandinavica*, **59**(1), 33.

411 Verbeke, J., Piepers, P., Supré, K. & S., D.V. (2014) Pathogen-specific incidence rate of clinical mastitis in
412 Flemish dairy herds, severity, and association with herd hygiene. *J. Dairy Sci.*, **97**, 6926–6934.

413 Zhao, X. & Lacasse, P. (2008) Mammary tissue damage during bovine mastitis: causes and control. *Journal*
414 *of animal science*, **86**(13_suppl), 57-65.

415 Ziv, G. (1980) Drug selection and use in mastitis: systemic vs local therapy. *Journal of the American*
416 *Veterinary Medical Association*, **176**(10 Spec No), 1109-1115.

417 Ziv, G. (1980) Practical pharmacokinetic aspects of mastitis therapy. III. Intramammary treatment. *VM/SAC*
418 *[Veterinary Medicine & Small Animal Clinician]*, **75**(4), 657-663, 666-670.

419

420

Authors' Contribution Statement

EA summarized the results, analyzed the data and drafted the text of the manuscript. SP and PRJ drafted the text and provided expertise particularly in pharmacokinetical, pharmacodynamical and statistical aspects. VM designed and monitored the study and reviewed the text.

Conflict of Interest

Satu Pyörälä and Päivi Rajala-Schultz have no affiliations with or involvement in any organization or entity with any financial interest. Elke Abbeloos and Vesa Myllys work for companies that market and/or distribute the product described in this article.

Tables and figures

Table 1: Mean (and SD) body weight (kg), age (months), days in milk (days) and daily milk yield (kg) for each of the treatment groups and the theoretical dose of benzylpenicillin. The last column contains the actual dose of benzylpenicillin given after weighing of the intramammary tubes after administration

Treatment group	Theoretical dose of benzylpenicillin (IU)	Cow body weight (kg)		Age (months)		Days in Milk		Daily milk yield (kg)		Actual dose of benzylpenicillin (IU)	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	200 000	585.4	55.5	40.8	12.3	149.2	34.2	32.3	7.9	189 832.0	4613.0
2	300 000	622.8	74.2	46.4	14.0	156.2	23.3	32.4	5.1	289 994.0	8404.0
3	600 000	615.2	43.1	42.8	18.3	151.0	36.7	32.8	7.4	610 244.0	5829.0
Average		607.8	57.1	43.3	14.2	152.1	29.7	32.5	6.4		

Table 2: Mean, standard deviation and geometric mean of the concentration of benzylpenicillin (in ng/mL) found in milk after intramammary administration of approximately 200.000 IU, 300.000 IU or 600.000 IU of benzylpenicillin in the form of an oily suspension with micronised particle size.

Time (h)	Group 1 (200.000 IU)			Group 2 (300.000 IU)			Group 3 (600.000 IU)		
	Mean	SD (ng/mL)	GeoMean	Mean	SD (ng/mL)	GeoMean	Mean	SD (ng/mL)	GeoMean
0	5/5 samples below quantification limit			5/5 samples below quantification limit			5/5 samples below quantification limit		
12	35345.24	22572.37	30438.00	41351.25	16982.59	38558.00	103476.36	45736.28	96380.47
24	2519.24	626.19	2459.06	2843.88	1160.82	2648.71	8778.28	12314.25	4486.51
48	13.86	10.18	11.19	18.77	19.36	11.12	20.09	8.85	18.37
72	5/5 samples below quantification limit			3/5 samples below quantification limit			5/5 samples below quantification limit

447

448 **Figure 1:** Mean benzylpenicillin concentrations in ng/mL (with SEM) determined in plasma
 449 specimens after intramammary administration of approximately 200.000 IU, 300.000 IU or
 450 600.000 IU of procaine benzylpenicillin in the form of an oily suspension with micronised
 451 particle size.

Figure 2: Simple linear regression model of Mean Area Under Curve (AUC) of benzylpenicillin levels in plasma after intramammary administration of approximately 200.000 IU, 300.000 IU or 600.000 IU benzylpenicillin in the form of an oily suspension with micronised particle size. The AUC was corrected for the actual administered dose administered to each cow in IU/kg bodyweight.

461

462 **Figure 3:** Mean benzylpenicillin log concentrations (ng.mL⁻¹) (with SEM) found in quarter
463 milk samples after intramammary administration of one quarter with respectively 200.000
464 IU, 300.000 IU and 600.000 IU of benzylpenicillin in the form of an oily suspension with
465 micronized particle size.

Figure 4: Simple linear regression model of Mean Area Under Curve (AUC) of benzylpenicillin levels in milk after intramammary administration of approximately 200.000 IU, 300.000 IU or 600.000 IU benzylpenicillin in the form of an oily suspension with micronised particle size. The AUC was corrected for the actual administered dose administered to each cow in IU/kg bodyweight.