<https://helda.helsinki.fi>

Regulation of cell adhesion : a collaborative effort of integrins, their ligands, cytoplasmic actors and phosphorylation

Gahmberg, Carl G.

2019-11

Gahmberg , C G , Grönholm , M , Madhavan , S , Jahan , F , Mikkola , E , Viazmina , L & Koivunen , E 2019 , ' Regulation of cell adhesion : a collaborative effort of integrins, their ligands, cytoplasmic actors and phosphorylation ' , Quarterly Reviews of Biophysics , vol. 52 . <https://doi.org/10.1017/S0033583519000088>

<http://hdl.handle.net/10138/324608>

<https://doi.org/10.1017/S0033583519000088>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Regulation of cell adhesion – a collaborative effort of integrins, their ligands, cytoplasmic actors and phosphorylation

Carl G. Gahmberg, Mikaela Grönholm, Sudarrshan Madhavan, Farhana Jahan, Esa Mikkola, Larisa Viazmina and Erkki Koivunen

Molecular and Integrative Biosciences Research Program, Faculty of Biological and Environmental Sciences, University of Helsinki, Viikinkaari 9 C, 00014 Helsinki, Finland

Corresponding author: Carl G. Gahmberg, Faculty of Biological and Environmental Sciences, University of Helsinki, Viikinkaari 9 C, 00014 Helsinki, Finland.

carl.gahmberg@helsinki.fi +358 505399439

Running title: Regulation of integrin activity

Word count: 14753

Abstract

Integrins are large heterodimeric type 1 membrane proteins expressed in all nucleated mammalian cells. Eighteen α -chains and eight β -chains can combine to form 24 different integrins. They are cell adhesion proteins, which bind to a large variety of cellular and extracellular ligands. Integrins are required for cell migration, hemostasis, translocation of cells out from the blood stream and further movement into tissues, but also for the immune response and tissue morphogenesis. Importantly, integrins are not usually active as such, but need activation to become adhesive. Integrins are activated by outside-in activation through integrin ligand binding, or by inside-out activation through intracellular signaling. An important question is how integrin activity is regulated, and this topic has recently drawn much attention. Changes in integrin affinity for ligand binding is due to allosteric structural alterations, but equally important are avidity changes due to integrin clustering in the plane of the plasma membrane. Recent studies have partially solved how integrin cell surface structures change during activation. The integrin cytoplasmic domains are relatively short, but by interacting with a variety of cytoplasmic proteins in a regulated manner, the integrins acquire a number of properties important not only for cell adhesion and movement, but also for cellular signaling. Recent work has shown that specific integrin phosphorylations play pivotal roles in the regulation of integrin activity. Our purpose in this review is to integrate the present knowledge to enable an understanding of how cell adhesion is dynamically regulated.

Keywords: integrin, cell adhesion, phosphorylation, leukocyte, LFA-1

Contents:

1. Introduction
2. Discovery of integrins
3. Integrin structure
4. Integrin external ligands
5. Structural changes in integrins upon activation
6. The integrin cytoplasmic domains contain numerous binding sites for intracellular proteins
7. Integrin β -tail binding cytoplasmic proteins
 - 7.1 Talin
 - 7.2 Kindlins
 - 7.3 Filamins
 - 7.4 The 14-3-3 proteins
 - 7.5 α -Actinin
 - 7.6 Cytohesin-1
 - 7.7 Dok1
 - 7.8 ICAP1
8. Integrin α -tail binding cytoplasmic proteins
 - 8.1 CD45 tyrosine phosphatase
 - 8.2 Rap1
 - 8.3 Calreticulin
 - 8.4 Paxillin
 - 8.5 SHARPIN
9. Preferential binding of proteins to clasped integrin cytoplasmic domains
10. Integrin activity is regulated by inside-out or outside-in signaling
11. Upstream events in integrin inside-out activation
12. Upstream events in integrin outside-in activation
13. Protein interactions at the integrin cytoplasmic domains are pivotal for cell adhesion and adhesion dependent functions
14. Integrin phosphorylation regulates integrin-cytoplasmic protein interactions

15. Cytoplasmic proteins compete for binding to integrin cytoplasmic tails
16. Cell adhesion is promoted by integrin mediated mechanotransduction
17. Integrin crosstalk
18. Sleep and fever affect leukocyte integrins
19. Integrins are potential targets for therapy
20. A model of cell adhesion and deadhesion
21. References

1. Introduction

Cell adhesion and its regulation are essential for the formation of different organs and maintenance of multicellularity. It is particularly important for the developing brain with its myriad of synaptic connections between neurons, but also for the dynamic interactions of other cells with their target structures. Equally important is cellular deadhesion for example during cellular movement and for metastasis of malignant cells.

Currently, we know a number of proteins, which are of pivotal importance for cellular interactions between cells and with the extracellular matrix. These include cadherins, which are essential for interactions of epithelial cells (Takeichi, 1977; Takeichi, 1991), selectins, which bind to the carbohydrate motif sialyl LeX on glycoproteins and glycolipids on blood endothelia and hematopoietic cells (Phillips et al., 1990; Walz et al., 1990; Polley et al., 1991), and the integrins with their ligands. Integrins bind to extracellular ligands such as fibronectin, fibrinogen and collagens, but also to cellular ligands such as the intercellular adhesion molecules (ICAM) (Rothlein et al., 1986; Patarroyo et al., 1987; Gahmberg, 1997) and the vascular adhesion molecule-1 (VCAM-1) .

Most integrins normally exist in a resting state and need some activation to become adhesive. This is particularly true for blood cell integrins, which must be strictly regulated in activity. Otherwise cell aggregation and blood clotting would take place, a reduced immunological response would ensue etc. resulting in dangerous complications.

Several reviews have been published on integrins and their structures and the integrin cellular ligands (Springer, 1990; Ruoslahti, 1991; Hynes, 1992; Schwartz et al., 1995; Gahmberg et al., 1997; Gahmberg, 1997; Hynes, 2002; Arnaout et al., 2005; Luo et al., 2007; Takada et al., 2007; Gahmberg et al., 2009; Bachmann et al. 2019). In this review we focus on the essential problem how integrin mediated adhesion and signaling are regulated. Integrins can be activated by inside-out activation or by outside-in activation (Kim et al., 2003; Abram and Lowell, 2009; Hu and Luo, 2013). It is, however, well established that cytosolic proteins interacting with the integrin intracellular tails are of pivotal importance for integrin regulation. We will here describe the most important findings dealing with the integrin – cytoplasmic interactions, and try to obtain a general view of how the integrin activities may be regulated. It is apparent, however, that integrin regulation is unusually complex and still incompletely understood. Integrin/cytoplasmic protein interactions can in theory be regulated by protein expression and degradation, accessibility, structural changes either in the integrins or the interacting proteins, and by posttranslational modifications. There are few ways to alter protein-protein interactions in a fast, economical and specific way. Phosphorylation is perhaps the most important one in many cell regulatory systems, and recent work has shown that specific phosphorylations play important roles in the regulation of adhesion. Due to the labile nature of phosphorylation and the potential expression of around 500 protein kinases and about 100 phosphatases in the mammalian genome, phosphorylation studies

are challenging. In this review we focus on what is currently known about the regulation of integrin activity. We focus on integrin regulation in leukocytes and platelets, because integrin regulation is best understood in these cells.

2. Discovery of integrins

In 1973, fibronectin was discovered (Hynes, 1973; Gahmberg and Hakomori, 1973). It was the first protein found that later turned out to be an adhesion protein. With the help of newly developed monoclonal antibodies, a leukocyte protein complex was identified, that was involved in several functions. The functions were mapped to two leukocyte proteins named Leukocyte function associated antigen -1 (LFA-1) and Macrophage-1 antigen (Mac-1) (Sanchez-Madrid et al., 1983). A genetic deficiency named Leukocyte adhesion deficiency (LAD) with impaired T cell functions, inability to synthesize immunoglobulins by B lymphocytes, and an increased susceptibility to life threatening infections was later described, and the cells were shown to lack the leukocyte cell surface proteins mentioned above (Arnaout et al., 1984; Springer et al., 1984). In 1982, Patarroyo and coworkers showed that phorbol esters induced leukocyte aggregation (Patarroyo et al., 1982). After screening with more than one hundred monoclonal antibodies, reacting with leukocyte cell surfaces, the phorbol ester induced leukocyte aggregation (adhesion), could be inhibited by a single monoclonal antibody (60.3) (Patarroyo et al., 1985a; Patarroyo et al., 1985b). The adhesion receptors identified were named Leu-Cam, and they turned out to be the same protein complex described above to be absent in the genetic disease called Leukocyte adhesion deficiency- I (LAD- I). Simultaneously, Ruoslahti and coworkers prepared fibronectin fragments, which replicated the activity of fibronectin (RGD) (Pierschbacher and Ruoslahti, 1984), and used them to identify the cellular receptor (Pytela et al., 1985). Sequencing of the fibroblast and leukocyte receptors showed that they were related and the common name for them "integrin" was coined by Hynes (Hynes, 1987). This name was chosen to illustrate the fact that the receptors integrate the cell surface with the interior cytoskeleton and vice versa. Later work then showed that the integrins form a receptor superfamily, and the integrins were grouped according to their β -polypeptides (Fig. 1).

Fig.1

Lower animals have few integrins and for example in the nematode *Caenorhabditis elegans* there are only two. Most cells express several types of integrins reflecting their different binding specificities and functions. Absence of $\beta 1$ integrins is lethal, whereas absence or defective functions of the $\beta 2$ integrin chain results in LAD- I, with life-threatening infections in the childhood. Defects in the $\alpha 1 \beta 3$ integrin may result in Glanzmann's thrombasthenia. However, many integrin gene knockouts in the mouse, have only mild or no effects possibly due to compensatory effects of other members of the integrin family. Selective absence of leukocyte $\alpha 4 \beta 1$, LFA-1 and Mac-1 results in increased leukocytosis and susceptibility to microbial infections (Kanwar et al., 2001; Scott et al., 2003; Ghosh et al., 2006; Chen and Sheppard, 2007).

3. Integrin structure

The integrins are heterodimers formed by two membrane spanning polypeptides, α - and β -chains (Fig. 2). They are glycoproteins, and the structures of the $\beta 2$ -integrin N-glycosidic carbohydrate

chains have been determined (Asada et al., 1991). Interestingly, they contain the sialyl LeX structure and bind E-selectin (Kotovuori et al., 1993).

Fig. 2

No complete integrin has been crystallized, and the structure elucidated, but the structure of the external domains has been elucidated for a few integrins. The α -chains of nine integrins contain an inserted I-domain, the structure of which was first determined (Lee et al., 1995). It contains a conserved metal ion-dependent adhesion site (MIDAS), which is needed for ligand binding. When the structure of the first integrin external part was determined ($\alpha V\beta 3$), it came as a surprise that the NH₂-terminal ligand binding region was turned back towards the membrane (Xiong et al., 2001). This structure does not bind ligands, or if so very weakly. Later, more integrin surface domains and parts of the transmembrane domains have been structurally determined (Xiong et al., 2009; Xie et al., 2010; Chen et al., 2010; Dong et al., 2012; Chen et al., 2012; Sen et al., 2013; Springer and Sen, 2016; Sen et al., 2018; Moore et al., 2018). It is now well established that integrins can exist in at least three different conformations: 1) bent towards the membrane, 2) extended, but with a closed ligand binding site and 3) extended with the binding site open. An extended open integrin has more than 1000 times higher ligand affinity than the resting bent form (Li and Springer, 2017). The affinities of different integrins may, however, be different even when expressed in the same cell. Using negative staining and electron microscopy whole integrins have been visualized, and one can see the different conformational states of the integrins (Fig. 3). The binding sites of different monoclonal antibodies, used for studies on $\beta 2$ integrin function are shown in Fig. 4 . The Kim127 antibody (extended closed) (Robinson et al., 1992), Mab24 (extended open) (Dransfield et al., 1992), and the 7E4 adhesion inhibitory antibody for the leukocyte LFA-1 (Nortamo et al., 1988). The Pac1 antibody reacts with the extended open form of the platelet integrin $\alpha IIb\beta 3$ (Shattil et al., 1987).

Fig. 3

Fig. 4

The transmembrane domains form α -helices, and are evidently closely associated through GxxxG interactions in the transmembrane domains (Kim et al., 2009), similar to that found in red cell glycophorin (MacKenzie et al., 1997).

The integrins have relatively short cytoplasmic tails, 20-60 amino acids long (an exception is the integrin $\beta 4$ polypeptide), but they are nevertheless important for integrin function (Ylänné et al., 1995) (Fig. 5). A number of proteins bind to the integrin cytoplasmic domains, and this is particularly true for the integrin β -chains. Much fewer cytoplasmic proteins seem to bind to the α -chain cytoplasmic domains. On the other hand, integrin external ligands bind to the α -chain I (A)-domain or to a combinatory binding site formed by the α - and β -chains. As discussed below, both the α - and the β -chains have important regulatory roles.

Fig. 5

It is obvious that only a few cytoplasmic proteins can simultaneously bind to the intracellular portions of the integrins. This means that the interactions must be both structurally and temporally regulated. These interactions are often short lived. For example, during cellular

migration, there must be a continuous formation and release of bonds between integrins and external ligands. The same must be true for integrin cytosolic interactions. The integrin interactions with intracellular molecules must not be direct. A number of cytosolic proteins bind to integrins indirectly through the integrin-bound proteins. Such adhesion complexes may be very large as evident from proteomic studies (Zaidel-Bar et al., 2007; Byron et al., 2011; Needham et al., 2019).

The cytoplasmic domains of the integrin β -chains are homologous with a proximal HDR(R/K)E sequence, two functionally important NXX(Y/F) motifs and a TTT/TTV/STF sequence between them. For the most part the α -chain cytoplasmic sequences are diverse, with the exception of the well preserved GFFKR motif in the proximal part of the domains (Fig. 5). A salt bridge is formed between the arginine residue of the GFFKR motif in the α -chains and an aspartic acid (D731 in $\beta 2$) in the β -chains (Hughes et al., 1996; Vinogradova et al., 2002).

4. Integrin external ligands

Integrins bind to a variety of ligands, which include cellular receptors, extracellular matrix proteins, soluble proteins in various body fluids, and microbial proteins and carbohydrates. The topic has been extensively reviewed, and the reader is advised to read the following reviews (Yamada and Olden, 1978; Hynes and Yamada, 1982; Yamada, 1983; Ruoslahti, 1988; Ruoslahti, 1996; Gahmberg, 1997). Many $\beta 1$ -integrins, such as $\alpha 5\beta 1$, bind to extracellular matrix proteins like fibronectin (Ruoslahti, 1991; Hynes, 1992) and $\alpha 1\beta 1$, $\alpha 2\beta 1$, $\alpha 10\beta 1$ to different types of collagens (Knight et al., 2000; Zeltz and Gullberg, 2016). Exceptions are the $\alpha 4\beta 1$, $\alpha 9\beta 1$ and $\alpha 4\beta 7$ integrins, which also can bind to the membrane protein vascular cell adhesion molecule-1 (VCAM-1). The subset of $\beta 1$ integrins binding various laminin isoforms includes $\alpha 3\beta 1$, $\alpha 6\beta 1$ and $\alpha 7\beta 1$, but also $\alpha 6\beta 4$ (Humphries et al., 2006; Takada et al., 2007). The platelet integrin $\alpha IIb\beta 3$ interacts for example with proteins that are important in blood coagulation, such as fibrinogen. Several integrins recognize the arginine-glycine-aspartic acid (RGD) or the leucine-aspartic acid-valine (LDV) sequences in the ligands. For example $\alpha 4\beta 1$ and $\alpha 9\beta 1$ bind to fibronectin via LDV, and $\alpha 5\beta 1$ via RGD.

In contrast, the $\beta 2$ -integrins interact for the most part with cellular ligands. The intercellular adhesion molecules (ICAMs) and vascular adhesion molecule-1 (VCAM-1) are type I membrane proteins belonging to the immunoglobulin superfamily. They span the cell membrane once, and their cytoplasmic domains interact with cytoskeletal proteins. Five ICAM molecules are present in mammals with different expressions. ICAM-1 and ICAM-3 have five immunoglobulin (Ig)-domains, ICAM-2 and ICAM-4 have two, and the most complex one, ICAM-5, has nine Ig-domains (Gahmberg, 1997). In most cases, the integrin binding sites are in the NH₂-terminal Ig-domain with exception of Mac-1, the binding site of which is in domain-3 of ICAM-1 (Diamond et al., 1991). The ICAM binding activity is regulated by expression. ICAM-1 is easily up-regulated by cytokines and during inflammation, whereas the other ones are more stably expressed. ICAM-1 and ICAM-2 are present in leukocytes and endothelial cells, and ICAM-3 in leukocytes. ICAM-4 is erythroid cell specific (Bailly et al., 1995), and ICAM-5 is confined to the dendrites and the cell bodies of central neurons (Yoshihara et al., 1994; Tian et al., 1997; Gahmberg et al., 2008). LFA-1 binds to all ICAMs, but not to extracellular matrix molecules. The other $\beta 2$ -integrins, Mac-1, $\alpha X\beta 2$ and $\alpha D\beta 2$

are less restricted in binding and may bind to soluble host proteins, but also to microbial molecules.

5. Structural changes in integrins upon activation

In resting cells, the integrins have their NH₂-terminal ligand-binding domains turned towards the membrane. Upon activation, the integrins extend, and may open their ligand binding site (Nishida et al., 2006). In the I-domain integrins, the binding site is present in the I- domain. Upon activation the α 7-helix moves down. In the other integrins the binding site is formed by a combination of the β -propeller of the α -chains and the β I domain of the β -chains. Upon activation the β I α 7-helix moves towards the β I interface with the hybrid domain, resulting in an outward movement of the hybrid domain, and exposure of the binding site. The cytoplasmic domains also move apart enabling new interactions of cytoplasmic proteins with the integrin tails (Springer and Dustin, 2012). These findings show that there exists a closely regulated connection between the integrin external parts and the inner domains. How the allosteric structural alterations are generated remains incompletely understood.

The fact that cells express many different integrins shows that cells need them for several different functions (Schwartz et al., 1995; Arnaout et al., 2005; Luo et al., 2007; Hogg et al., 2011; Huttenlocher and Horwitz, 2011; Mitroulis et al., 2015; Kourtzelis et al., 2017; Fagerholm et al., 2019). Their activity must be regulated, and often so in different ways. Therefore, results obtained for one integrin may not apply to another one. The cellular environment, including molecular neighbors is certainly also important.

Experimental results indicate that the β -chains are most important for the regulation of integrin activity, but they also show that the α -chains closely cooperate with the β -chains, and reciprocally that the β -chains regulate the ligand binding activity of the α -chains.

The integrin cytoplasmic tails are relatively unstructured, and their conformations and movements are restricted by interactions with cytosolic and cytoskeletal proteins. Interactions with the inner leaflet of the lipid membrane are also important.

6. The integrin cytoplasmic domains contain numerous binding sites for intracellular proteins

The integrins are not only mere adhesion proteins, although their primary functions are based on their capacity to adhere. They are pivotal components of machineries for cellular movement, cell spreading, phagocytosis, metastasis of malignant cells, blood coagulation, brain and immune synapse formation etc. Furthermore, they act, perhaps due to their "sticky nature", as receptors for various microbes (Gbarah et al., 1991; Wickham et al., 1993; Jackson et al., 2000; Campadelli-Fiume, 2016). Their ability to signal in two directions across the plasma membrane makes them unique. The integrin β -chains are quite well conserved, which indicates that their functions are quite similar. The α -chains are much more diverse, enabling more specific functions. Rather few proteins have been found to bind to the α -chain cytoplasmic domains, and at least some of them act as negative regulators of integrin activity.

Fig. 6

Fig. 6 shows the sequence of the LFA-1 cytoplasmic tails, and the binding sites for some of the intracellular proteins that interact with α L or β 2. The α -chain cytoplasmic domain contains about 60 amino acids, and the β -chain domain about 45. This means that the β -chain cytoplasmic segment is about 180 Å long if extended. In fact, some part of it forms an α -helix, and some parts are more extended. This means that at most 5-6 average size proteins could bind to it at the same time. Therefore, the interactions of cytoplasmic proteins with the β -chain, must be strictly regulated. On the other hand, several proteins may indirectly bind to the integrins through the integrin tail binding proteins, and form adhesomes. In fact it has been estimated by proteomics that hundreds of proteins can interact with the integrins (Zaidel-Bar et al., 2007; Byron et al., 2011).

In the following part of the review, we briefly describe cytoplasmic proteins that are involved in the regulation of integrin activity. Most of these proteins have been subjects of previous reviews, and the reader is referred to them for additional information (Sampath et al., 1998; Kim et al., 2003; Abram and Lowell, 2009; Moser et al., 2009b; Anthis and Campbell, 2011; Calderwood et al., 2013; Klapholz and Brown, 2017). Thereafter we discuss how the proteins, by interacting with the integrin cytoplasmic domains, may regulate integrin binding and signaling activity.

7. Integrin β -tail binding cytoplasmic proteins

7.1 Talin

Talin-1 and -2 are large cytosolic 270 kDa proteins consisting of a head region and a long rod (BurrIDGE and Connell, 1983; Critchley, 2009; Calderwood et al., 2013; Klapholz and Brown, 2017). The two talins are functionally quite similar, but have different cellular distributions. The talin head is related to the FERM (4.1, ezrin, radixin, moesin) proteins, which are known to bind to several integral membrane proteins. The rod binds to the actin cytoskeleton. Talin can be cleaved by the Ca^{2+} -activated protease calpain, and this reaction may be important for the inside-out activation of integrins (Garragher et al., 1999; Glading et al., 2002; Franco et al., 2004; Franco and Huttenlocher, 2005). The involvement of talin in cell adhesion has mainly been studied in platelets and leukocytes (Calderwood et al., 1999; Calderwood et al., 2013; Goult and Schwartz, 2018; Sun et al., 2019). The talin head binds to the proximal NXXY/F sequence in the β -chains through its phosphotyrosine binding domain (PTB) in the F3 subdomain, but also to an α -helix in the proximal part of the cytoplasmic domain (Tadokoro et al., 2003). Talin-1 knockout mice die early during embryonic development, which shows that the protein is essential. In vitro experiments show that the purified talin head can activate integrins by forcing the cytoplasmic parts apart, and it has been proposed to be the final component in the activation cascade (Kim et al., 2012). The talin rod may bind to several F-actin molecules, further increasing the clustering of integrins, and thus up-regulating adhesion by an increase in avidity. Talin interacts with phosphatidylinositol 4,5-bisphosphate at the inner surface of the plasma membrane, and this interaction is considered to be important for adhesion (Chinthalapudi et al., 2018).

7.2 Kindlins

The kindlin family of proteins consists of three members (Ussar et al., 2006). Kindlin-1 is mainly expressed in epithelia, kindlin-2 is ubiquitously expressed, and kindlin-3 expression is restricted to hematopoietic cells. Genetic absence of kindlin-1 results in a skin disease with blisters, an absence of kindlin-2 is lethal, and a deficiency of kindlin-3 results in the LAD III syndrome (Malinin et al., 2009; Moser et al., 2009a; Svensson et al., 2009). Like the talins, the kindlins contain a FERM domain, but they lack a rod domain. Kindlin-1 may bind to the proximal NPXY sequence (Gao et al., 2019) whereas the others bind to the distal integrin NPXY/F motif and the preceding TTT/TTV/STF sequence (Morrison et al., 2013; Li et al., 2017). Whereas the talins can directly activate integrins by binding to the proximal NXXY/F sequence, the kindlins cannot do that, but they have to cooperate with talins and other proteins (Moser et al., 2009b). Integrins cannot extend when kindlin is absent. Kindlin-3 is difficult to express and purify, and therefore kindlin-2 has been used for molecular studies. Structural analysis shows that when it is bound to integrin tails, it combines two binding motifs (Kammerer et al., 2017; Li et al., 2017). Importantly, kindlin-2 forms a dimer and this is necessary for cell adhesion (Ye et al., 2013a). Kindlin-3 is phosphorylated on T482 or S484. Mutation of the phosphorylated residues to alanine decreased binding and cell spreading on fibrinogen through the α IIb β 3 integrin (Bialkowska et al., 2015). Furthermore, kindlin-2 activated the small G protein Rac1, and recruited the cytoskeletal Arp2/3 complex, enabling the formation of an actin network (Böttcher et al., 2017). Taken together, the functional basis of kindlin-mediated activation of integrins is quite different from that of talin. Talin and kindlins work together in regulating adhesion. Interestingly, the minimal amount of kindlin-3 needed for platelet and leukocyte adhesion is only 5% of the normal levels. A gradual increase in kindlin-3, however, improved the integrin avidity in leukocytes (Bennett, 2015; Klapproth et al., 2015a).

7.3 Filamins

The filamin family consists of three members (Stossel et al., 2001). Filamin A is a large homodimeric protein of about 280 kDa, linked at the COOH-terminals making a V-shaped structure. The monomers contain 24 Ig-domain like β -sheet repeats. It has been shown that domain-21 binds to integrin β 2-chains between the two NP(L/K)F motifs (Liu et al., 2015a). This means that its binding site overlaps with those of talin and 14-3-3 proteins (Kiema et al., 2006; Takala et al., 2008; Chatterjee et al., 2018b). Filamin also has a binding site in the α -chains (Fig. 6) (Liu et al., 2015b). The migfilin protein binds directly to filamin (Tu et al., 2003; Takafuta et al., 2003). Filamin is considered to be an integrin inhibitor, and migfilin displaced filamin from integrin β -chains resulting in activation of β 1, β 2 and β 3 integrins (Das et al., 2011). Interestingly, filamin phosphorylation was induced by G protein-coupled receptors by protein kinase A (Tirupula et al., 2015). The phosphorylation on S2152 affected talin and kindlin-3 binding (Ithychanda et al., 2015; Waldt et al., 2018).

7.4 The 14-3-3 proteins

The 14-3-3 proteins got their peculiar name from their presence in chromatography fractions, obtained during their isolation. They are dimeric serine/threonine-phosphate binding proteins (Muslin et al., 1996), but may also bind to nonphospho-proteins, although more weakly (Petosa et al., 1998). They bind with high affinity to the pTTT phosphorylated β 2-chain. The binding was also influenced by the sequence surrounding the phosphorylation site (Campbell, 2013). 14-3-3

proteins also interact with the β 1- and β 3-cytoplasmic sequences 776-790 in a phosphorylation dependent manner (Han et al., 2001).

7.5 α -Actinin

α -Actinin is a cytoskeletal protein formed by two identical 100 kDa polypeptides. It mediates the binding of proteins to the actin cytoskeleton, among them ICAMs and integrins. α -actinin binds to the integrin β -chain upstream of the proximal NXXY/P sequence (Fig. 6) (Pavalko and LaRoche, 1993). Using conformation specific monoclonal antibodies Stanley and coworkers showed that the extended, but closed LFA-1, which binds the monoclonal antibody KIM127 epitope, is at the leading edge of a migrating cell and bound to the cytoskeleton through α -actinin (Stanley et al., 2008).

7.6 Cytohesin-1

Cytohesin-1 is a G protein exchange factor. The binding site is at the WKA motif in the most proximal part of the β 2 cytoplasmic chain. Expression of a β 2-chain mutated at this position, reduced adhesion and cell spreading in inside-out activation (Kolanus, 2007). Outside-in signaling in Jurkat T cells, expressing this β 2 mutant, resulted in threonine phosphorylation of the protein, whereas signaling through the TCR did not (Perez et al., 2003).

7.7 Dok1

The adaptor protein Dok1 contains a pleckstrin domain and a PTB domain. It binds to the integrin β -chains through the proximal NXXY/F motif with a higher affinity than that of talin (Oxley et al., 2008). The binding increased when Y747 in β 3 was phosphorylated. Dok1 bound strongly to the phosphorylated S745 and S756 in β 2, but poorly to pT758 (Gupta et al., 2015).

7.8 ICAP-1

ICAP-1 is a β 1 integrin binding protein, which binds to the valine residue close to the proximal NPIY sequence and competes with talin and kindling-3 for binding. It regulates the formation of focal adhesions (Chang et al., 1997).

8. Integrin α -tail binding cytoplasmic proteins

Rather few cytosolic proteins have been shown to bind to integrin α -chains.

8.1 CD45 tyrosine phosphatase

The CD45 tyrosine phosphatase, first identified as “common leukocyte antigen” (Gahmberg et al., 1976; Andersson and Gahmberg, 1978; Trowbridge, 1978; Thomas, 1989), is a major cell surface glycoprotein in leukocytes, which due to alternative splicing of the cell surface part expresses several isoforms. It contains two phosphatase domains in the cytoplasm, but only the proximal one is catalytically active. CD45 has an important function in the dephosphorylation of the COOH-terminal tyrosine phosphate present in cytoplasmic Src family tyrosine kinase family members including Src, Lck and Fyn. The dephosphorylation activates the Src family kinases (Mustelin et al., 1989; Mustelin et al., 1992). The effect of CD45 on the Src family kinases is, however,

complicated. In Lck, Y394 is required for full activity, and the phosphate at this residue can also be removed by CD45 (Courtney et al., 2017; Philipsen et al., 2017). Furthermore Lck has a phosphosite at Y192, and mutation of this residue to alanine inhibited the association of CD45 with Lck. This resulted in inability to dephosphorylate the COOH-terminal phosphate in the kinases (Courtney et al., 2018). Therefore, there must exist a balance between the COOH- terminal inhibitory phosphorylation, the activity promoting one on Y394 and the phosphorylation on Y192. CD45 is also a component of the immunological synapse, where it is peripherally located (Johnson et al., 2000; Chang et al., 2016).

CD45 binds to the LFA-1 α -chain at the COOH-terminal amino acid sequence 1123-1145 (Geng et al., 2005). Importantly, it does not bind to α M or α X, the cytoplasmic domains of which are shorter, and quite different from the COOH-terminal domain of α L. Full length CD45 cytoplasmic domain bound relatively poorly to LFA-1, whereas the isolated membrane proximal part, including the first phosphatase domain, showed strong binding. The binding to α L did not affect the phosphatase activity. LFA-1 and CD45 may form a complex in the immunological synapse. Interestingly, a soluble cytoplasmic form of CD45 is found in plasma and it inhibits the activation of T cells (Puck et al., 2017). It is derived from neutrophils and monocytes by proteolysis.

8.2 Rap1

Rap1 is a small G protein important in T cell adhesion (Bivona et al., 2004). Cellular activation results in localization of the protein to the plasma membrane by attachment of a geranyl-geranyl lipid. Rap1 binds weakly to the F0 domain of talin, but the interaction is stabilized when Rap1b in platelets is membrane- anchored (Zhu et al., 2017). Rap1 has two downstream effectors. In leukocytes Rap1 binds to RIAM (Rap1-interacting adapter molecule), which in turn binds to talin and activates adhesion through β 2 integrins (Calderwood, 2015; Su et al., 2015; Lagarrigue et al., 2016). RIAM is normally auto-inhibited by its N-terminal segment. The auto-inhibition can be released by phosphorylation of Y45 by FAK (focal adhesion kinase) (Chang et al., 2019). Upon TCR activation in Jurkat cells RIAM translocated to the plasma membrane, and this could be inhibited by FAK inhibitors. The Rap1 binding protein, RAPL, is implicated in leukocyte adhesion, and binds to lysines K1097 and K1099 in the α L chain. This binding is critical for LFA-1 conformational changes, and binding to ICAM-1 (Tohyama et al., 2003). In contrast, α 4 β 1 binding to VCAM-1 was only weakly affected by RIAM deficiency (Klapproth et al., 2015b), and platelet adhesion was not affected by RIAM deficiency (Stritt et al., 2015; Bromberger et al., 2018). Recent work shows that Rap1 binds to the F1 domain in the talin head and activates the α IIb β 3 integrin. Interestingly, the F1 domain activity is due to the formation of an helix when bound to membrane lipids (Gingras et al., 2019). Rap1 binds to RAPL and affects leukocyte adhesion, whereas no platelet effects have been reported (Katagiri et al., 2003). RAPL acts as an integrin transporter whereby LFA-1 localizes to the leading edge of migrating T cells.

The results show that different integrins and cells use Rap1, RIAM, RapL and talin in different ways. Interestingly, the S/T Mst1 kinase is bound to RAPL, but its possible function is still not understood (Katagiri et al., 2006).

8.3 Calreticulin

The cytosolic protein calreticulin (Krause and Michalak, 1997) binds the KXGFFKR conserved motif found in the proximal part of integrin α -chains (Coppolino et al., 1995). Treatment of the Jurkat T cell line with phorbol esters increased the calreticulin association to the $\alpha 2\beta 1$ integrin. Integrins regulate Ca^{++} transport into cells and crosslinking of the VLA-4 integrin with monoclonal antibodies induced Ca^{++} spikes, whereas knockout of calreticulin resulted in abrogation of extracellular Ca^{++} transport into cells (Coppolino et al., 1997).

8.4 Paxillin

Paxillin binds to a sequence around E983 and Y991 in the integrin $\alpha 4$ chain. Mutational analysis of amino acid residues in this region showed that S998A mutation resulted in enhanced binding of paxillin, whereas the S988D mutation abrogated binding and cell migration, but increased cell spreading (Liu and Ginsberg, 2000). Very recent work indicates that kindlin-3 recruits leupaxin to podosomes and regulates paxillin phosphorylation and thereby podosome turnover (Klapproth et al., 2019).

8.5 SHARPIN

SHARPIN (SHANK-Associated RH Domain Interactor) negatively regulates $\beta 1$, $\beta 2$ and $\beta 3$ integrins and binds to the three amino acids N-terminal to the conserved GFFKR sequence in integrin α -chains (Rantala et al., 2011; Pouwels et al., 2013). Recent work shows that it binds even more strongly to the $\beta 1$ chain and more weakly to the $\beta 3$ chain. Importantly, the binding site involves the proximal NPIY sequence. Talin binds to the same site and binding of the talin head domain was out-competed by SHARPIN (Gao et al. 2019). Furthermore, SHARPIN binds to kindlin-1, but not to kindlin-2 or -3. Importantly, SHARPIN may in this way bring kindlin-1 to the $\beta 1$ binding site and inhibit integrin activation. SHARPIN interacted with activated Rap1 and R-Ras, and inhibited downstream signaling (Lilja et al., 2017). SHARPIN is also involved in the formation of the ubiquitin chain complex, and it binds to the HOIP E3 ubiquitin ligase (Kasirer-Friede et al., 2019).

9. Preferential binding of proteins to clasped integrin cytoplasmic domains

In contrast to most cytosolic proteins, binding preferentially to either integrin β - or α -chains, the L-plastin (LCP1) protein prefers binding to the interacting, clasped cytoplasmic integrin domains (Tseng et al., 2018). It was shown to compete with talin for binding. Filamin is also known to bind to inactive, clasped integrins (Liu et al., 2015a).

10. Integrin activity is regulated by inside-out or outside-in signaling

Most experimental models on integrin activation are based either on T lymphocytes or platelets. In contrast to most cells, these cells are quiescent without activation, and therefore it is possible to observe changes in integrin- related functions after activating them. Lymphoblasts, fibroblasts and glial cells, for example, are continuously dividing and their integrins already activated at least to some extent.

A description of integrin signaling, leading to activation of adhesion, can be done in different ways. Here we first describe how cellular signaling takes place before reaching the integrins, how the integrins are modified, and the integrin-cytoplasmic interactions and signaling may take place downstream from the activated integrins.

In inside-out signaling, integrins become activated through binding of agonists to non-integrin receptors (Giancotti and Ruoslahti, 1999; Hynes, 2002; Gahmberg et al., 2009; Hogg et al., 2011). These can be cell surface proteins, such as the T cell receptor (TCR), or chemokine receptors, but it is also possible that activation takes place by binding of intracellular proteins due to changes in the environment, for example high temperature. In outside-in signaling, ligands are bound directly to the integrins, which then signal into the cell. In integrin crosstalk, an activated or inhibited integrin communicates with other integrins in a given cell by intracellular signaling. Integrin inside-out activation, resulting in ligand binding, may subsequently be followed by outside-in signaling, due to interaction of external ligands with the integrins.

Interestingly, T cells in the blood stream can migrate either upstream or downstream depending on to which ligand they bind. When the cells bind to ICAM-1 they migrate upstream, but when bound to VCAM-1, they migrate downstream (Dominguez et al., 2015). At higher shear rates the cells migrate upstream even in the presence of small amounts of ICAM-1 even in the presence of substantial levels of VCAM-1. The results show that outside-in signaling acts in different ways depending on which ligand-integrin interactions take place.

During the past few years, it has become increasingly apparent that integrins can be regulated by mechanical forces acting on integrins. In leukocyte trafficking the integrins can form slip bonds, which may be broken by shear force, but also catch bonds, which in contrast become stronger under shear force (Sun et al., 2016). In both inside-out and outside-in signaling, conformational changes in the integrins occur, enabling them to become adhesive (Springer and Sen, 2016). The distance from the ligand binding site to the lipid membrane is in the range of 200 Å for the extended integrins, and it is intriguing to speculate how the information flows between the adhesion sites and the cytoplasmic domains. In addition to the conformational changes, resulting in increased integrin affinity for ligands, an important feature is the clustering of the integrins in the plane of the membrane. This in turn will increase integrin avidity. In this case, the role of the actin cytoskeleton is of pivotal importance (Li and Springer, 2017). Obviously, a combination of both affinity and avidity increase results in highly efficient integrin binding.

A number of studies show that the hybrid domain in the external part of the β -chains swings out upon activation, concomitantly with a down- movement of the $\alpha 7$ helix in the α -chain I domain or in the β I-like domain in the integrins lacking I domains. The external integrin structure then opens, and this is coupled to a separation of the cytoplasmic domains of the α - and β -chains (Vinogradova et al., 2002; Kim et al., 2003; Tng et al., 2004). Mn^{2+} ions can activate the binding capacity for external ligands by stabilizing the extended open conformation (Sen et al., 2018), but the ion-induced activity is not seen inside the cell as separation of the cytoplasmic tails. These facts show that integrin signaling is unusually complex. Furthermore, we should realize that integrins are different, belonging to several subclasses. Therefore, the mechanisms regulating their functions are different, in spite of many common features.

The cytoplasmic domains of the integrin β polypeptides are homologous, indicating that their functions are relatively similar. Opposite to that, the α -chains are quite different in structure. This may reflect the fact that the α -chains are primarily responsible for binding of different ligands, which needs specificity, whereas the β -chains are more involved in integrin activity regulation.

Below we try to describe what we currently know about integrin activity regulation, but we realize that we still are in the very beginning of an understanding of this fascinating, but very complex subject. The reader is encouraged to get familiar with previous original work. However, during the past few years we have seen a dramatic development in the understanding of cell adhesion and its multifaceted features.

11. Upstream events in integrin inside-out activation

T lymphocytes are excellent model cells for the study of integrin activity regulation. One reason is that we know a lot about T cells, due to their importance in immunology. Activation can take place through the TCR or chemokine receptors. Each T cell has a TCR of unique specificity due to rearrangement of the DNA during development. When T cells encounter cells presenting antigen peptides bound to class I or class II transplantation antigens, the T cells must recognize whether the complex is self or novel. Further activation then needs the assembly of a TCR signaling complex for downstream signaling (Courtney et al., 2018). Associated with the TCR are the CD3 molecules, which are pivotal for signaling. An efficient way to induce extensive signaling, and circumvent specific T cell antigen restricted signaling, is by crosslinking the CD3 proteins and the associated TCR with anti-CD3 antibodies (Fig. 7). The CD3 molecules contain in their cytoplasmic domains sequence motifs called Immune-receptor-Tyrosine-based-Activation-Motif (ITAMs), which act as substrates for the cytoplasmic tyrosine kinase Lck. The phosphorylated ITAM then binds to the Zap70 tyrosine kinase through its SH2 domains. Zap70 becomes activated and phosphorylates the linker for activation of T cells (LAT). This results in the binding of phospholipase $\text{C}\gamma 1$. Then phosphatidylinositol-3 kinase is recruited and phosphorylates phosphatidylinositol 4,5-bisphosphate (PIP₂) to yield PIP₃. This brings in turn the ITK kinase to the complex, which phosphorylates PLC $\gamma 1$. PLC $\gamma 1$ then hydrolyzes PIP₂ resulting in IP₃ and diacylglycerol (DAG). IP₃ then binds to its intracellular receptor resulting in the release of Ca^{++} . Cellular Ca^{++} can also increase upon activation through the binding of the cytoplasmic protein calreticulin to the conserved KXFFKR motif in the proximal part of integrin α -chains. The Ca^{++} increase was indistinguishable in thapsigargin treated wild type and calreticulin knockout cells (Coppolino et al., 1997). Because the drug releases Ca^{++} by inhibiting the endoplasmic reticulum Ca^{++} -ATPase, the Ca^{++} must come from extracellular sources (Coppolino et al., 1997).

Fig. 7

DAG activates protein kinase C enzymes (PKC) (Castagna et al., 1982; Nishizuka, 1984; Newton, 1995), which phosphorylate the integrin $\beta 2$ -chain (Fagerholm et al., 2002b). The above scheme is based on studies of T cell activation, which obviously results in several molecular changes in the cells, among which integrin activation is only one.

On the external face of the TCR, the immunological synapse is formed (Fig. 8) (Grakoui et al., 1999; Bromley et al., 2001). The TCR is located in the center with LFA-1 in the periphery. PKC θ localizes to the central supramolecular activation cluster, where it mediates effector T cell

activation (Kong and Altman, 2013). CD45 is more peripherally located. CD45 activates Lck and the related tyrosine kinases by releasing the COOH-terminal tyrosine phosphate. The Csk kinase in turn inactivates Lck by phosphorylating the COOH-terminal of Lck, resulting in an intramolecular loop between the SH2 domain of Lck and its COOH-terminal. On the other hand, phosphorylation of Y394 in Lck is needed for Lck activation and also this phosphate can be released by CD45. Most probably the accessibility of CD45 to Lck is important in the regulation of the synapse and the subsequent influence on integrin activity (Fagerholm et al., 2002a).

Fig. 8

Calcineurin is an important serine/threonine phosphatase known to act on T cell NFAT transcription factors. It is activated by calmodulin, which is up-regulated by cytoplasmic Ca^{++} (Ye et al., 2013b). Calmodulin activates leukocyte adhesion (Fagerholm et al., 2001). Intracellular Ca^{++} is important in the regulation of integrin activity, and interestingly the intracellular Ca^{++} was increased by leukocyte integrin activation (Altieri et al., 1992). Further studies showed that the β -chain of LFA-1 is critical for Ca^{++} regulation. The phenylalanine in the distal NPKF sequence was required for Ca^{++} signaling and subsequent integrin activation (Sirim et al., 2001). Inhibition of calcineurin by cyclosporine A or FK506 is widely used in the treatment of transplantation rejection. It was recently reported that calcineurin is in fact recruited to the TCR complex, where it released the inhibitory phosphate on S59 in Lck (Dutta et al., 2017). The treatment inhibited LFA-1, which indicates that the effects of the drug are at least in part due to inhibition of T cell adhesion.

Chemokines, leukotrienes and kallikreins are deposited on endothelial cells lining the vasculature and trigger the activation of rolling leukocytes, including T cells (Alon and Feigelson, 2012; Vestweber, 2015) (Fig. 9). The chemokine receptors are G-protein-coupled receptors and induce signaling by activating PLCs, which results in generation of inositol-1,4,5-trisphosphate and DAG (Zarbock et al., 2011; Strazza et al., 2017). The former compound releases Ca^{++} from intracellular stores and DAG activates PKCs. Chemokines are also stored within endothelial cells and are released during leukocyte migration through the vessel walls (Shulman et al., 2012).

Fig. 9

The small G protein Rap1 is activated by its exchange factors such as CALDAG-GEFI by TCR or chemokine induced signaling cascades. Rap1 is brought to the membrane by RapL and in migrating cells to the leading edge (Hogg et al., 2011). Activated Rap1 activates LFA-1, whereas its down regulation inhibits T cell adhesion. Importantly, inhibition of the interaction with RIAM, and in RIAM depleted mice, the T and B lymphocytes could not traffic to peripheral lymph nodes, due to down regulation of LFA-1 and VLA-4 activities. In contrast, macrophages showed no defect in adhesion or spreading. The results show that different blood cells differently regulate their integrins.

The Del-1 (developmental endothelial locus-1) protein is a natural inhibitor of leukocyte/ endothelial interactions. It is a soluble protein that becomes attached to blood vessel endothelial proteoglycans or the $\alpha V\beta 3$ integrin through its RGD sequence (Choi et al., 2008; Schurpf et al., 2012; Mitroulis et al., 2014). It competes with ICAMs for the binding to LFA-1 and Mac-1 and therefore inhibits leukocyte binding to the ligands. Del-1 is strongly expressed in the brain and lung, but almost absent from the liver.

12. Upstream events in integrin outside-in activation

Outside-in activation of integrins takes place through agonist binding to the external domains of integrins (Abram and Lowell, 2009). It may also be a consequence of inside-out signaling, and it is then difficult to distinguish the two types of signaling. The two types of signaling may even take place at the same time. Mn^{2+} is an efficient activator of adhesion if it binds to the MIDAS site on the external aspect of cells (Gailit and Ruoslahti, 1988), but it is probably, however, not physiologically important for integrin activity. Soluble ligands, such as recombinant ICAM-2 lacking the transmembrane and cytoplasmic domains (Kotovuori et al., 1999), integrin binding peptides (Li et al., 1993a), integrin antibodies and immobilized fibrinogen have been used to demonstrate outside-in signaling in leukocytes and platelets (Shattil et al., 1998; Tan et al., 2001). Both integrin ligand binding activating and inhibitory antibodies can induce outside-in signaling, whereas antibodies binding to integrins, but which do not affect ligand binding, do not induce signaling (Grönholm et al., 2016).

Fig. 10

Outside-in signaling is in part different from that of inside-out signaling. The most studied integrin in this respect is the platelet integrin $\alpha IIb\beta 3$ (Fig. 10). A number of ligands including fibrinogen, von Willebrand factor and fibronectin bind to the integrin through the classical arginine-glycine-aspartic acid (RGD) sequence (Durrant et al., 2017). Fibrinogen also binds through the KQAGDV sequence in the α -chain (Humphries et al. 2006). Using peptides interacting with the αIIb transmembrane domain Fong et al. (2016) induced activation by activation of Src and Syk. The integrin became clustered through the signaling. Importantly, Glanzmann's thrombasthenia is due to mutations in this integrin. An early event in $\alpha IIb\beta 3$ activation is the up-regulation of Src tyrosine kinase activity. Src is highly expressed in platelets and binds by the SH3 domain to the $\beta 3$ cytoplasmic domain COOH-terminal RGT motif (Wu et al., 2015). Src phosphorylates Y747 and Y759 in the NXXY sequences of $\beta 3$ (Jenkins et al., 1998). Incidentally, the corresponding residues in $\beta 2$ are phenylalanines, which cannot be phosphorylated. Lck may have a similar function as Src in lymphocytes, but its target sites must be different. Y735 in $\beta 2$ is a potential phosphorylation site, and a Y735A mutation resulted in decreased internalization of LFA-1 in transfected cells (Bleijs et al., 2001). Src activation results in phosphorylation of the Syk tyrosine kinase and its binding to the $\beta 3$ polypeptide (Obergfell et al., 2002). Src also activates PLC γ 2 (Wonerow et al., 2003), and its activation can also be supported by the Bruton's tyrosine kinase (Shattil and Newman, 2004). PLC γ 2 catalyzes the formation of IP $_3$ and DAG, resulting in an increase of intracellular Ca^{++} and activation of PKCs.

FAK is also activated upon $\alpha IIb\beta 3$ activation, initially by autophosphorylation of Y397, which provides a docking site for Src, which then phosphorylates FAK on multiple sites and promotes its activity. The homolog Pyk2 may have similar effects in cells lacking FAK, such as macrophages (Okigaki et al., 2003). FAK may be important in integrin outside-in activation and in colon cancer epithelial cells, FAK inhibition decreased metastasis through integrins (Downey-Biechler et al., 2019).

13. Protein interactions at the integrin cytoplasmic domains are pivotal for cell adhesion and adhesion-dependent functions

Integrins are involved in a number of different cellular functions, but they are all associated with cell adhesion. This is true for cell movement, cellular cytotoxicity, phagocytosis, antibody production, synaptic transmission and as receptors for microbes. Most if not all of these functions involve signaling across the plasma membrane. The integrin cytoplasmic domains are 20-60 amino acids long (an exception is $\beta 4$, which is longer), and they have no enzymatic activity. Therefore, to assert functions, it is necessary to assemble complexes of different signaling and interacting proteins at their cytoplasmic domains. Mammals have up to 24 integrins with different cellular distributions. A given cell may contain quite a few different integrins making it possible on one hand to express several different functions, but on the other hand, it has also become necessary to regulate the integrins in a useful way. Furthermore, the integrin activities must be coordinated temporally in order to function in a beneficial manner. One such example is cell migration. If some integrin activates movement, whereas others at the same time would act as brakes, the cell would be unable to move.

Currently, we are aware of many important interactions between integrins and cytoplasmic proteins, but we are still far from understanding how the regulatory mechanisms work. A lot of excellent science has been done on this topic, but one should emphasize that much of what currently is known, has been done with different cells, different integrins and under different conditions. Certainly, integrins are individuals and facts applying to one integrin or integrin subfamily, may not be true for others.

In the following, we focus on the role of $\beta 2$ -integrins in leukocyte adhesion, and when appropriate to some extent on other integrins. Leukocyte integrins are essential for normal life, which is evident from the leukocyte adhesion deficiency syndromes. LAD-I is due to absence or inhibitory mutations in the $\beta 2$ -polypeptide, LAD-II is due to deficiency in selectin function and in LAD-III the integrin-binding protein, kindlin-3, is malfunctioning. Proteomic studies of integrin interacting proteins have identified a large number of proteins (Zaidel-Bar et al., 2007; Byron et al., 2011; Needham et al., 2019). Several of these are members of adhesion complexes, adhesomes, and not directly interacting with integrins. In Fig. 6 the best characterized interacting proteins with LFA-1 are shown, and their binding sites indicated. A number of reviews are available on integrins and their cytoplasmic interactions, but recent progress has been substantial, and this fact merits an up-to-date review.

14. Integrin phosphorylation regulates integrin-cytoplasmic protein interactions

In early studies, Nishizuka and coworkers identified a group of enzymes named protein kinase Cs, which were found to be important for cell growth, morphology and lipid metabolism. The group also found that phorbol esters strongly activate the kinases and have a similar role as the physiological lipid DAG (Castagna et al., 1982; Nishizuka, 1984). A total of 11 PKCs were identified and they could be divided into three groups. PKCs α , βI , βII and γ are regulated by Ca^{++} .

The second group includes PKCs δ , ϵ , η , θ and μ , and lack the C2 domain, which is thought to bind Ca^{++} . The atypical PKCs are ξ and λ . The last two do not respond to phorbol esters (Newton, 1995).

PKCs were first considered to be oncoproteins, because of their activation by phorbol esters, which act as tumor promoters. Surprisingly, more recent work indicates that they are in fact tumor suppressors, because they are commonly inactivated in tumors (Newton and Brognard, 2017). We will below discuss the possibility that restoration of PKC activity may in fact be beneficial in cancer treatment due to their agonist functions on integrins.

Because phorbol esters could induce leukocyte aggregation, and PKCs were known to act as receptors for these compounds, several research groups thought that it would be important to study the possibility that the adhesion receptors are phosphorylated. Using ^{32}P -labelled leukocytes, it was shown that the α -chain of LFA-1 was phosphorylated in resting cells, whereas the β -chain was not (Chatila et al., 1989; Buyon et al., 1990, Valmu et al., 1991) (Fig. 5). After activation of ^{32}P -labelled cells with phorbol esters, the β 2-chain became phosphorylated. The major phosphorylated residue was S756. However, the mutation S756A did not affect leukocyte adhesion, whereas mutations of the threonine triplet T758-760 to alanines abrogated adhesion (Hibbs et al., 1991). However, no phosphorylation of T758-760 was observed, and the possible role of phosphorylation of LFA-1 remained enigmatic. We then found that upon activation of T cells, there is in fact strong threonine phosphorylation on the β 2-chain, which was observed after the use of the phosphatase inhibitor okadaic acid (Valmu and Gahmberg, 1995). The only threonines in the cytoplasmic domain of the β 2-chain are T758, T759 and T760, and these must therefore be the targets of phosphate labeling.

We then used the integrin β 2 cytoplasmic tail as bait, and PKC β , PKC β II and PKC δ were identified as phosphorylating enzymes (Fagerholm et al., 2002b). With purified PKCs, the most efficient enzymes phosphorylating β 2 peptides were PKC α , PKC η , PKC β I, PKC β II and PKC δ . Most of them phosphorylated both serine and threonine residues in β 2, and S745, S756, T758, T759 and T760 became phosphorylated.

Detailed studies in vivo showed that the β 2-chain is phosphorylated on S745, S756, T758 and weakly on T759. Y735 may also be phosphorylated, but direct proof for this is not available. T758 was found to be most strongly labeled (Hilden et al., 2003). The RACK1 protein acts as a receptor for PKCs (Lillenthal and Chang, 1998), and it binds to the β 2-chain proximal region (Fig. 5). This could be a mechanism for bringing PKCs close to the integrins and phosphorylate them.

The phosphorylation of T758 in T cell LFA-1, resulted in binding of the dimeric 14-3-3 protein, followed by binding of the adaptor protein Tiam1, activation of the small G-protein Rac1, activation of cell adhesion, and increased expression of the activation marker CD69 (Fagerholm et al., 2002b; Fagerholm et al., 2005; Nurmi et al., 2007; Grönholm et al., 2011). The activation induced increased binding of LFA-1 to the actin cytoskeleton (Valmu et al., 1999).

The α -chain phosphorylations were not considered to be important, because they were constitutively phosphorylated in resting cells, and the phosphorylation did not seem to change upon activation. There is, however, a dynamic phosphorylation and dephosphorylation of the α -chains, because radioactive phosphate labeling of α L was stronger in cells treated with okadaic

acid than in its absence (Valmu and Gahmberg, 1995). We determined the phosphorylation site in the α L-chain, and it turned out to be S1140. Surprisingly, mutation of this residue to alanine completely abrogated cell adhesion (Fagerholm et al., 2005). Corresponding mutations gave the same result for α M and α X (Fagerholm et al., 2006; Uotila et al., 2013). α D has not been studied in this respect. We tried to find proteins interacting with α L in a phosphorylation-dependent manner, but were not able to find any. We then made the exciting discovery that in LFA-1 and α X β 2, the T758 phosphorylation on the β 2-chain depends on α -chain phosphorylation (Jahan et al., 2018). Mutations of α L S1140A and α X S1158A inhibited the important β 2-T758 phosphorylation, which is necessary for integrin activation. The mutation S1140D had a similar effect on T758 phosphorylation as the α L wt chain, indicating that the negative charge is important.

Phosphorylation of other integrins has been less studied. T788 and T789 in the β 1 chain, which correspond to T758-T59 in β 2 are likewise necessary for ligand binding, because mutations to alanine reduced binding to fibronectin, whereas mutations to aspartic acid behaved as wt β 1. It induced, however, more focal contacts and less migration (Nilsson et al., 2006). In the α V β 5 integrin, the S759 and S762 residues were phosphorylated by the p21-activated kinase 4. Mutation of these residues abolished cell migration (Li et al., 2010).

The VLA-4 (α 4 β 1) integrin is an important leukocyte integrin, but it is also expressed in many other cells. It is phosphorylated on S988 in the α -chain and the S988A mutation inhibited cell spreading, but increased cell migration. The S988A mutation prevented the binding of paxillin, whereas the S988D mutation restored the activity to resemble that of wt integrin (Han et al., 2003). The phosphorylated α 4 in VLA-4 localized to the leading edge of migrating cells (Lim et al., 2008).

In the platelet α IIb β 3 integrin, the β 3 chain is phosphorylated on T753, which is the last threonine residue in the TST triplet between the NPXY sequences. It was phosphorylated upon treatment with the phosphatase inhibitor calyculin A, and the phosphorylation inhibited platelet adhesion upon activation by outside-in activation with fibrinogen (Lerea et al., 1999; Kirk et al., 2000).

The effect of outside-in activation on integrin phosphorylation of α IIb β 3 has shown that the tyrosines in the NPXY sequences become phosphorylated in platelets by the Src kinase (Jenkins et al., 1998; Law et al., 1999; Wu et al., 2015; Dutta et al., 2017).

In addition to integrins, a number of other cellular proteins become phosphorylated upon cellular activation including several cytoplasmic proteins, which are important for cell adhesion. These include talin (Ratnikov et al., 2005), filamin (Pons et al., 2017), Rap1 (Takahashi et al., 2013), paxillin (Klapproth et al., 2019) and kindlin-3 (Bialkowska et al., 2015). The filamin domain-21 is poorly phosphorylated on S2152 in vitro by protein kinase A, but upon ligand binding the filamin phosphorylation strongly increased (Ithychanda et al., 2015; Tirupula et al., 2015). The possible importance of these phosphorylations remains poorly understood.

15. Cytoplasmic proteins compete for binding to integrin cytoplasmic tails

Both inside-out and outside-in signaling are accompanied by changes in cell adhesion, and it is well established that the intracellular integrin domains are of pivotal importance. In order to describe how integrins are regulated we focus on LFA-1.

Fig. 6 shows the binding sites of cytoplasmic proteins known to bind to the α - and β -chains of LFA-1, and which affect integrin functions. The binding sites in the cytoplasmic domains have been determined by deleting parts of the chains or by mutational analysis. The $\beta 2$ cytoplasmic domain is about 50 amino acids long, and several of the potential binding proteins are relatively large proteins. The interacting proteins have distinct functions needed for the cells. Therefore, regulation of binding must be important. One way of regulating protein binding to the β -chain is by phosphorylation, which strongly affects the binding of cytosolic proteins (Takala et al., 2008; Gupta et al., 2015; Chatterjee et al., 2016; Chatterjee et al., 2018a). Another way is accessibility. In the resting state there is a salt bridge between D731 in the $\beta 2$ -chain and R995 in αL chain. Deletion of the distal part of α -chains, which makes the salt bridge impossible, activates integrins (Liu et al., 2015b). Phosphorylation on serine has been shown to compete with salt bridge formation in various protein-protein interactions (Skinner et al., 2017), and could be a potential mechanism to change integrin cytoplasmic interactions.

In resting cells, the cytoplasmic tails are closely associated restricting protein binding. Some proteins, however, interact well with the $\beta 2$ -chain in resting cells. Filamin has been shown to bind to L753-P764 in the $\beta 2$ -chain (Kiema et al., 2006; Takala et al., 2008; Chatterjee et al., 2018b). Interestingly, it also binds to integrin α -chains (Liu et al., 2015a). Filamin could in this way form an integrin cytoplasmic domain clasp. Recent work shows that filamin is phosphorylated by the Ndr2 protein kinase on S2152 in vivo after activation of the TCR (Waltdt et al., 2018). The activation through the TCR results in the release of filamin from LFA-1. Structural analysis showed how the integrin binding motif in filamin domain-21 interacts with the $\beta 2$ -cytoplasmic domain (Takala et al., 2008). When T758 is phosphorylated, the phosphorylated peptide cannot be accommodated within the filamin pocket and it is released. Filamin is an integrin inhibitor, because it occupies sites important for binding of activating proteins. It should not, however, be considered only as an inhibitor. Its interactions may also organize the integrin cytoplasmic tails into a structure facilitating activation when appropriate.

Dok1 is a cytosolic protein binding in the resting state to the proximal NPLY sequence in $\beta 3$ (Oxley et al., 2008). The binding region partially overlaps with that of talin. Talin bound more strongly to the unphosphorylated sequence than Dok1, but upon activation and tyrosine phosphorylation of the NPLY sequence, the binding of Dok1 strongly increased (Oxley et al., 2008). Dok1 bound weakly to $\beta 2$ integrin cytoplasmic peptides, but when the S745 and S756 residues were phosphorylated, the binding increased (Gupta et al., 2015). When cells were activated through the TCR or chemokine receptors, the high affinity of 14-3-3 proteins for the phosphorylated site outcompeted talin binding from its distal binding site, and enabled a molecular switch. In $\beta 3$ the binding from the phosphorylated tyrosine in the NPLY sequence switched to a complex of $\beta 3/14-3-3/Dok1$. In $\beta 2$, When S756 was phosphorylated in $\beta 2$, the complex $\beta 2/14-3-3/Dok1$ was formed (Chatterjee et al., 2016; Chatterjee et al., 2018a). Activation through the TCR does not result in phosphorylation of S756, but it becomes phosphorylated by phorbol ester treatment. In the $\beta 2$

integrin Mac-1, S756 phosphorylation regulates Rap1 activity, which is important for phagocytosis (Lim et al., 2011).

Kindlin-3 is confined to hematopoietic cells, and its genetic absence results in the LAD-III syndrome due to defects in adhesion (Moser et al., 2009; Svensson et al., 2009; Malinin et al., 2009). Studies on kindlin-2 shows that its expression is regulated by the Smurf1 ubiquitin kinase (Wei et al., 2017), and kindlin-3 expression may also be regulated by the same mechanism. Like talin the kindlins contain a FERM domain. Kindlins-2 and -3 function as coactivators of integrins together with talin, but how the cooperation takes place has remained incompletely understood (Calderwood et al., 2013). Kindlin-1 may be inhibitory. The recent finding that kindlins form dimers may favor integrin cluster formation, resulting in increased avidity (Kammerer et al., 2017; Li et al., 2017). Talin cannot do that, but may form dimers in an antiparallel way through its COOH-terminals. Kindlin-2 is most easily expressed and purified of the kindlins. It has therefore been used for studies on kindlin structure and interactions. It directly binds to actin, which could explain that clustering occurs (Bledzka et al., 2016). In addition kindlin-2 binds paxillin and the Arp2/3 complex (Böttcher et al., 2017). Paxillin is known to bind to integrin α -chains, and this fact indicates that kindlins could inhibit the formation of clasped integrin α - and β -chains. Through the paxillin and Arp2/3 interactions, kindlins could activate Rac1, and in this way induce membrane protrusions. This could also happen through the phosphorylation of the β 2-chain on T758, which results in high affinity 14-3-3 binding and release of filamin. This in turn is followed by binding of the adaptor protein Tiam1 and activation of Rac1 (Grönholm et al., 2011; O'Toole et al., 2011) (Fig. 11).

Fig. 11

α -Actinin binds to the β 2-chain between the two talin binding sites. It is an essential component of the actin cytoskeleton. Recent studies have shown that when LFA-1 and α X β 2 are activated, phosphorylation of the α -chains is needed to enable the phosphorylation of T758 on the β 2-chain (Jahan et al., 2018). When both integrin chains were phosphorylated, α -actinin got access to the β -chain and bound more efficiently. The mutation S1140A in LFA-1 inhibited α -actinin binding to the β -chain and cell adhesion. Interestingly, the S1140D mutation restored α -actinin binding. The results indicate that when both the α - and β -chain are phosphorylated, the cytoplasmic clasp opens, perhaps due to repulsion between the cytoplasmic tails.

The Cbl-b protein is an adaptor protein, but it is also a ubiquitin ligase, and it affects cell adhesion. Deficiency of the protein resulted in increased phosphorylation of T758 in β 2. This resulted in increased binding of 14-3-3 proteins and up-regulation of adhesion (Choi et al., 2008b). It is possible that SHARPIN and Cbl-b interact, because Cbl-b is known to bind the ubiquitin ligase HOIP (Zhou et al., 2008).

The Crk proteins, Crk and CrkL, are small adaptor proteins containing SH2 and SH3 domains and they are known to be involved in T cell trafficking (Roy et al., 2018). They are active in outside-in signaling, because T cells with knock out of Crk and CrkL adhered poorly to coated ICAM-1. Under shear flow there was no difference in cell arrest in wt cells, but knock out cells did not spread. The leading edge was mobile in wt cells, whereas cells lacking the proteins did not form a leading edge. The Crk proteins promoted integrin mediated PI3K signaling, but AKT phosphorylation was

abrogated in knockout cells. Furthermore, the phosphorylation of the interacting C-cbl and Cbl-b proteins decreased. Possibly, the Crk proteins promote phosphorylation by facilitating contacts with tyrosine kinases.

The cytohesins are GEFs for the ARF family of small G proteins (Kolanus, 2007). Cytohesin-1 is known to bind to the N-terminal cytoplasmic sequence (WKA) in the integrin $\beta 2$ -chain. This results in activation of LFA-1. Cytohesin-3 resembles cytohesin-1 in inducing cell spreading, and cytohesin-2 regulates cell movement (Davies et al., 2014). Interestingly, the binding motif for cytohesin-1 in the $\beta 2$ -chain resembles that of SHARPIN (WKV) for the α -chain.

Recently, it was found that the *Aggregatibacter actinomycetemcomitans* leukotoxin (LtxA) activates LFA-1 (Nygren et al., 2018). The toxin binds to the LFA-1 external part, but surprisingly it also binds to the LFA-1 α - and β -chain cytoplasmic domains with high affinity. The affinities for αM and αX are considerably lower. The binding site in the α -chain cytoplasmic domain is in the proximal region (KVGFFKRNLKEKMEAGRGPNGI), whereas the binding to the $\beta 2$ -chain is in the middle region, because the sequence W747NN increased the binding as compared to the $\beta 2$ NH2 terminal W723-Q746 sequence. Treatment with phorbol esters decreased FRET of LFA-1 αL -CFP and $\beta 2$ -YFP as expected, but treatment with the toxin increased FRET, which indicates that the cytoplasmic domains became more close to each other. Possibly, the toxin interfered with the formation of the salt bridge between the αL and $\beta 2$ -chains, and therefore phorbol ester treatment of toxin-exposed cells no longer resulted in integrin phosphorylation. LtxA also prevented Rap1 and the formation of the Rap1/RapL complex. The lysines K998 and K1000 in the α -chain of LFA-1 are important for the binding of RapL, and this site overlaps with that of LtxA.

L-plastin (LCP1) is an actin binding protein related to α -actinin, which binds to the clasped $\alpha M \beta 2$ and keeps the integrin inactive (Tseng et al., 2018). It bound less efficiently to isolated αM and $\beta 2$ polypeptides. Interestingly, the increased adhesion of LCP1 depleted cells was inhibited by treatment with phorbol esters. LCP1 may also affect integrin trafficking, because the amount of surface expressed $\alpha M \beta 2$ increased in LCP1 depleted cells, although the total amount of $\alpha M \beta 2$ remained unchanged.

16. Cell adhesion is promoted by integrin-mediated mechanotransduction

Cells and organs live in an environment of extracellular matrix, and this is also true for blood cells at early stages of differentiation. Integrins bind to extracellular ligands and during cellular movement in the blood stream or in extracellular tissues, there may arise mechanical tension between integrin bound external molecules and the intracellular actin cytoskeleton. This integrin mediated force induces important effects on cellular signaling and molecular changes in the interacting cells (Nordenfelt et al., 2016; Sun et al., 2016; Huse, 2017).

In migrating cells, various integrin ligands, such as the ICAMs, VCAM-1 and fibronectin form nascent adhesion sites at the leading edge through "catch bonds". Here the integrins are active and induce focal adhesion or corresponding sites, and F-actin bundles force the cell further. The mechanosensitive proteins talin and vinculin are important and "slip bonds" are generated. Whereas the talin head binds to the integrin β -chains, the actin binding sites in the talin rod are

important and by stretching the rod, vinculin binding sites become exposed. Kindlins and α -actinin are also important in mechanotransduction, but they have been less studied in this connection. Kindlin-3 may be important for the $\alpha 4 \beta 1$ mediated adhesion under shear flow (Lu et al., 2016).

In leukocytes catch bonds are formed initially between E-selectin on endothelial cells and sialyl LeX oligosaccharides on leukocyte glycoconjugates. L-selectin on leukocytes may also bind to endothelial sialyl LeX ligands (Gong et al., 2017). The rolling leukocytes then engage endothelial chemokines, which induce inside-out activation of integrins and strong adhesion due to catch bonds (Morikis et al., 2017). During the selectin/ligand induced rolling the LFA-1 molecules extend, but remain closed. Due to the selectin catch bonds and chemokine activation, inside-out activation occurs, and the integrins become fully active with an open binding site. The leukocytes then move into tissues either between endothelial cells or through them. After that leukocytes utilize local chemokines to migrate further (Shulman et al., 2012). Actually, the TCR interaction with peptide-bound transplantation antigens also involves catch bonds (Huse, 2017). Interestingly, by studying single integrin attachment to a fibronectin substrate, Strohmeyer and coworkers (Stromeyer et al., 2017) found that $\alpha 5 \beta 1$ sensed load to induce adhesion within a second, and after disrupting the bond a slower adhesion reaction followed. This was due to catch bond formation.

Recent studies have shown that leukocytes can move upstream on ICAM-1 coated surfaces in fluid flow, and downstream when coated with VCAM-1 (Buffone et al., 2018). LFA-1 is the key receptor responsible of the upstream movement. Interestingly, the VLA-4/VCAM-1 interaction did not need any stimulation, whereas the LFA-1/ICAM-1 interaction must be activated. Possibly, there is a crosstalk between the integrins with LFA-1 as the dominating molecule (see below).

17. Integrin crosstalk

Due to the fact that most cells express several integrins, it is important that they can be regulated in a useful manner for the function of the cells. Communication between integrins, is needed for example when cells migrate.

Porter and Hogg (1997) observed that activation of LFA-1 in T cells down-regulated VLA-4 ($\alpha 4 \beta 1$), and $\alpha 5 \beta 1$ to a smaller extent. Activation through the TCR or with phorbol esters resulted in decreased binding to VCAM-1 and fibronectin. Then it was shown that ligation of $\alpha V \beta 3$ inhibited $\alpha 5 \beta 1$ (Blystone et al., 1999). The treatment prevented the activation of the calcium/calmodulin-dependent protein kinase II (CamKII), and up-regulation of CamKII overcame the $\alpha V \beta 3$ mediated suppression. Later work showed that $\beta 1$ blocking antibodies inhibited $\alpha V \beta 3$ mediated binding to $\alpha 4$ laminin. The treatment increased $\beta 3$ phosphorylation on S752. PKA does not phosphorylate $\beta 3$ on this site, but may act on the phosphatase inhibitor-1, which inhibits the phosphatase PP1 (Gonzales et al., 2008).

We studied how the activation of $\beta 2$ integrins in T cells results in inactivation of VLA-4. Jurkat T cells lacking LFA-1 bound strongly to the VLA-4 ligand VCAM-1. When the cells were transfected with αL , the cells bound well, but when LFA-1 was activated using the SDF-1 chemokine, the

binding to VCAM-1 was inhibited (Uotila et al., 2014). Phosphorylation of S1140 on α L was necessary, as the mutation S1140A prevented the crosstalk. Likewise, phosphorylation of T758 was essential. It was further shown that the signaling from β 2- pT758 through the 14-3-3/Tiam1/Rac1 signaling route (Fig. 11) was necessary for the crosstalk. The signaling resulted in dephosphorylation of T788/789 on β 1.

Monoclonal antibodies are potentially useful for the treatment of T cell mediated functions. We studied whether such antibodies affect integrin crosstalk in T cells. It turned out that treatment of cells with LFA-1 activating antibodies, and some inhibitory antibodies, which induce T758 phosphorylation on β 2, resulted in dephosphorylation of T788/789 on the β 1 chain of VLA-4 (see Fig. 4). The signaling took place through the 14-3-3/Tiam1/Rac1 pathway. By these antibody treatments, it became possible: 1) to activate LFA-1 and inhibit VLA-4, 2) inhibit both LFA-1 and VLA-4, 3) inhibit LFA-1 but not VLA-4 or 4) bind to LFA-1, but without any effect on either LFA-1 or VLA-4 activity (Grönholm et al., 2016).

The results show that intracellular signaling between integrins takes place through specific integrin phosphorylations. Presently, we do not know how extensive the crosstalk is, or whether other mechanisms exist to mediate communication between integrins.

18. Sleep and fever affect leukocyte integrins

Drugs such as isoproterenol and histaminergic receptor agonists are known to affect integrins. These may act through G α s-coupled agonists like epinephrine, norepinephrine, prostaglandins E2 and D2 or adenosine to inhibit integrin activation (Chigaev et al., 2008; Chigaev et al., 2011; Chigaev et al., 2014). During sleep, the levels of G α s agonists such as serotonin and catecholamines decrease, resulting in activation of T cell adhesion (Dimitrov et al., 2019).

It is a well known fact that fever promotes survival during infection. Incubation of T cells at 40 C induced the heat shock protein 90 (Hsp90), resulting in T cell integrin activation (Lin et al., 2019). Hsp90 was shown to bind to the α 4 polypeptide through the ENRRDSWSY(991) motif (Fig. 5) and mutation to alanine of R985, W989 or Y991 abrogated Hsp90 binding. The α 4 β 7 integrin was up-regulated, resulting in increased adhesion to VCAM-1. β 2 integrins were not affected. Thermal stress increased the binding of talin and kindlin-3 to α 4 integrins. When mice were infected with *Salmonella typhi* murium the body temperature rose to 40 C, and the mice with the R985A/R985A mutation in α 4 integrins showed higher lethality than wt mice. There were much fewer T cells in the small intestines of the mutated mice, which shows that the recruitment of cells was inhibited (Lin et al., 2019).

19. Integrins are potential targets for therapy

Monoclonal antibodies to leukocyte integrins are potentially useful for treatment of chronic inflammatory diseases such as psoriasis, rheumatoid arthritis, ischemia/reperfusion injury, contact dermatitis and asthma (Mitroulis et al., 2015). A positive clinical response is often seen for example in the treatment of psoriasis (Krueger et al., 2000). A problem is that harmful side

effects may appear, due to wipe out of important leukocyte functions. The statins form an interesting group of drugs. These drugs are commonly used for the treatment of hypercholesterolemia (Corsini et al., 1995). The statins inhibit the 3-hydroxy-3-methylglutaryl coenzyme-A reductase, which is a key enzyme in the biosynthesis of cholesterol. The lovastatin and simvastatin drugs blocked the interaction of LFA-1 with ICAM-1 by binding to the I-domain (Weitz-Schmidt et al., 2001; Frenette, 2001). Because the statins are relatively safe without appreciable side effects, they are potentially useful in clinical practice where inhibition of leukocyte functions is wanted. It could also be important to develop drugs that activate leukocyte adhesion. For the treatment of malignant diseases, an up-regulation of T cell and natural killer cell functions could be important. Activation could be achieved with integrin-activating antibodies or β 2-integrin-activating peptides (Tng et al., 2004; Li et al., 1993a; Grönholm et al., 2016).

In the future it could become possible to further improve in cancer treatment by immune checkpoint therapy (Sharma and Allison, 2015). The CTLA-4 and PD-1 T cell molecules induce inhibitory pathways in T cells. Their normal functions are to keep T cells unreactive to self tissues. CTLA-4 and PD-1 inhibit T cell killing of malignant tumors due to T cell mediated suppression. Whereas PD-1 inhibits Akt phosphorylation by preventing CD28-mediated activation of PI3K (Parry et al., 2005; Buchbinder and Desai, 2016), the PP2A phosphatase is important for the CTLA-4 mediated effect. By use of antibodies to inhibit the inhibitory molecules, excellent results have been obtained in the treatment of melanoma. Promising results have also been obtained with lung cancer and kidney cancer. Interestingly, engaging CTLA-4 with antibodies up-regulated LFA-1, resulting in increased adhesion and integrin capping. The effect was even stronger using a combination of CD3 and CTLA-4 antibodies. The treatments induced signaling and activation of the small G protein Rap1 (Gatta et al., 2002; Schneider et al., 2005).

20. A model of cell adhesion and deadhesion

So far we have reviewed the topic of cell adhesion and integrins with a focus on T lymphocytes, but also to some extent discussed integrins in platelets and other cells.

It is certainly evident from the above text that we currently know quite a lot about integrin structure and allosteric changes during activation, and how dominant integrins such as LFA-1, may regulate other integrins. We can distinguish between the unique abilities of integrins to signal both inside-out and outside-in across the cell plasma membrane. We are also aware of the fact that a large number of cytoplasmic proteins interact with the intracellular domains of integrins, and the binding sites have been mapped with detailed precision. Proteomic studies show that there are still an enormous number of proteins, binding either directly or indirectly to integrin cytoplasmic domains, and certainly some of them may turn out to be essential for function. The ability to bind several different proteins to the cytoplasmic domains provides means to regulate functions in a number of ways in different cells, within subdomains in the same cell and at different times. This ability is quite different from that of most other type I membrane proteins,

which have more restricted functions, such as transport through the membrane or membrane spanning tyrosine kinases or phosphatases.

Knockout experiments and mutational analyses show that most of the cytoplasmic proteins dealt with above are important for integrin functions and, likewise, several different integrin cytoplasmic motifs are of pivotal importance.

Furthermore, some interacting proteins are essential for adhesion and allosteric signaling through the integrin external domains to activate or inhibit binding, whereas others may be important for integrin localization, cellular movement and intracellular signaling. Because the integrins have many different functions, it is important that they can switch interactions, and in this way cellular functions.

Fig. 12 α 4 β 1

After describing what currently is known, let us now look at how T cell integrins could become activated using LFA-1 as a model (Fig. 12). In inside-out activation, the LFA-1 molecule is activated either through the TCR or through chemokine receptors. After proximal signaling, talin is associated with the β 2 polypeptide, PKCs become activated and phosphorylate T758 in the β 2-chain. At this point a large proportion of S1140 in the α -chain is phosphorylated. There is a turnover of the α -chain phosphorylation at S1140, which is observed by treatment with okadaic acid (Valmu and Gahmberg, 1995). If the α -chain phosphorylation site is without phosphate, the β -chain cannot be phosphorylated, and the integrin remains inactive (Jahan et al. 2018). When both chains are phosphorylated, the cytoplasmic part of the clasped integrin opens up and filamin is released due to high affinity 14-3-3 binding. Structural work showed that the phosphorylated β -chain cannot be accommodated within the filamin pocket (Takala et al., 2008). The 14-3-3 proteins bind to the phosphorylated chain with a K_d of $\sim 5 \mu\text{M}$ (Chatterjee et al. 2016). The open cytoplasmic domain is now able to bind α -actinin (Jahan et al., 2018). Talin is then released from the NPLF site, but remains associated with the β - chain through the proximal binding site, and by interaction with the lipid membrane (Gingras et al., 2019). Dok-1 is bound to the other half of the 14-3-3 dimer, and makes a complex of pT758/14-3-3/Dok-1/kindlin-3 (Chatterjee et al., 2018a). Both 14-3-3 and kindlin-3 are dimers and the proteins bind to the actin cytoskeleton. This results in clustering of integrins leading to increased avidity. Furthermore, signaling through the pT758/14-3-3/Tiam1/Rac1(GTP) route activates the cytoskeleton allowing protrusions at the plasma membrane (Grönholm et al., 2011). Simultaneously, the bound talin directly activates the integrin by inducing an allosteric structural alteration transmitted through the membrane resulting in an increase of the integrin affinity for external ligands such as ICAM-1. The cycle of phosphorylation and dephosphorylation of S1140, combined with an activation signal through the TCR or chemokine receptors, in turn enables T758 phosphorylation and dephosphorylation of the β -chain and an integrin activation cycle.

It should be pointed out that integrins differ particularly in their cytoplasmic parts, and the same regulatory mechanisms certainly do not apply to all. The $\alpha\text{M}\beta$ 2 integrin has a relatively short α -chain cytoplasmic domain as compared to the others in the integrin subfamily (Fig. 5). This fact may be the reason that S756, which is phosphorylated in phorbol ester treated cells, is more accessible to β -chain interacting cytoplasmic proteins. Rap1 interacts with the phosphorylated

S756 in $\alpha M\beta 2$, recruits talin and positions it for subsequent functions such as phagocytosis in myeloid cells (Lim et al., 2011).

In outside-in activation, the beginning of activation is different involving early integrin phosphorylation by tyrosine kinases. In leukocytes Lck is important, whereas Src is involved in $\alpha IIb\beta 3$ activation. Further signaling then results in integrin phosphorylation as described above.

In the resting state the integrins are clasped by inhibitory cytosolic proteins such as filamin, SHARPIN and others. These are outcompeted early during activation due to lower affinity for integrin binding, ~ 0.5 mM for filamin (Chattarjee et al., 2018b), as compared to that of 14-3-3 and subsequent activators.

With the exception of PKCs involved in T758 phosphorylation of the $\beta 2$ -chain, little is currently known about which protein kinases and phosphatases are involved in integrin regulation, and this subject certainly needs much further effort.

The outlined scheme is for some part hypothetical and needs further experimental work to become more complete. However, we now have a way to proceed to enable us to get a deeper understanding of this exciting and important subject in cell biology and medicine.

Financial support

The current studies were supported by the Finnish Medical Society, the Magnus Ehrnrooth Foundation, the Liv och hälsa Foundation, the Finnish Society of Science and Letters, and the Wilhelm och Else Stockmann Foundation.

21. References

- Abram CL and Lowell CA (2009) The ins and outs of leukocyte integrin signaling. *Annual Review of Immunology* 27, 339-362.
- Alon R and Feigelson SW (2012) Chemokine-triggered leukocyte arrest: force-regulated bi-directional integrin activation in quantal adhesive contacts. *Current Opinion of Cell Biology* 24, 670-676.
- Altieri DC, Stamnes SJ and Gahmberg CG (1992) Regulated Ca^{++} signalling through leukocyte CD11b/CD18 integrin. *Biochemical Journal* 288, 465-473.
- Andersson LC and Gahmberg CG (1978) Surface glycoproteins of human white blood cells. *Blood* 52, 57-67.
- Anthis NJ and Campbell ID (2011) The tail of integrin activation. *Trends in Biochemical Sciences* 36, 191-198.
- Arnaout MA, Mahalingam B and Xiong JP (2005) Integrin structure, allostery, and bidirectional signaling. *Annual Review of Cell and Developmental Biology* 21, 381-410.
- Arnaout MA, Spits H, Terhorst C, Pitt J and Todd RF (1984) Deficiency of a leukocyte surface glycoprotein (LFA-1) in 2 patients with MO1 deficiency – effects of cell activation on Mo1 LFA-1

surface expression in normal and deficient leukocytes. *Journal of Clinical Investigation* 74, 1291-1300.

Asada M, Furukawa K, Kantor C, Gahmberg CG and Kobata A (1991) Structural study of the sugar chains of human leukocyte cell adhesion molecules CD11/CD18. *Biochemistry* 30, 1561-1571.

Bachmann M, Kukkurainen S, Hytönen VP and Wehrle-Haller B (2019) Cell adhesion by integrins. *Physiological Reviews* 99, 1655-1699.

Bailly P, Tontti E, Hermand P, Cartron JP and Gahmberg CG (1995) The red cell LW blood group protein is an intercellular adhesion molecule which binds to CD11/CD18 leukocyte integrins. *European Journal of Immunology* 25, 3316-3320.

Bennett JS (2015) A little bit of kindlin goes a long way. *Blood* 126, 2535-2536.

Bialkowska K, Byzova TV and Plow EF (2015) Site-specific phosphorylation of kindlin-3 protein regulates its capacity to control cellular responses mediated by integrin $\alpha\text{IIb}\beta 3$. *Journal of Biological Chemistry* 290, 6226-6242.

Bivona TG, Wiener HH, Ahearn IM, Silletti J, Chiu VK and Philips MR (2004) Rap1 up-regulation and activation on plasma membrane regulates T cell adhesion. *Journal of Cell Biology* 164, 461-470.

Bleijds DA, van Duijnhoven GCF, van Vliet SJ, Thijssen JPH, Figdor CG and van Kooyk Y (2001) A single amino acid in the cytoplasmic domain of the $\beta 2$ integrin lymphocyte function-associated antigen-1 regulates avidity-dependent inside-out signaling. *Journal of Biological Chemistry* 276, 10338-10346.

Bledzka K, Bialkowska K, Sossey-Alaoui K, Vaynberg J, Pluskota E, Qin J and Plow EJ (2016) Kindlin-2 directly binds actin and regulates integrin outside-in signaling. *Journal of Cell Biology* 213, 97-108.

Blystone SD, Slater SE, Williams MP, Crow MT and Brown EJ (1999) A molecular mechanism of integrin crosstalk: $\alpha\text{V}\beta 3$ suppression of calcium/calmodulin-dependent protein kinase II regulates $\alpha 5\beta 1$ function. *Journal of Cell Biology* 145, 889-897.

Böttcher RT, Veelders M, Rombaut P, Faix J, Theodosiou M, Stradal TE, Rottner K, Zent R, Herzog F and Fässler R (2017) Kindlin-2 recruits paxillin and Arp2/3 to promote membrane protrusions during initial cell spreading. *Journal of Cell Biology* doi 10.1083/jcb.201701176.

Bromberger T, Klapproth S, Rohwedder I, Zhu L, Mittmann L, Reichel CA, Sperandio M, Qin J and Moser M (2018) Direct Rap1/talin interaction regulates platelet and neutrophil integrin activity in mice. *Blood* 132, 2754-2762.

Bromley SK, Burack WR, Johnson KG, Somersalo K, Sims TN, Sumen C, Davis MM, Shaw AS, Allen PM and Dustin ML (2001) The immunological synapse. *Annual Review of Immunology* 19, 375-396.

Buchbinder EI and Desai A (2016) CTLA-4 and PD-1 pathways. Similarities, differences and implications of their inhibition. *American Journal of Clinical Oncology*. 39, 98-106.

- Buffone A, Anderson NR and Hammer DA (2018) Migration against the direction of flow is LFA-1-dependent in human hematopoietic stem and progenitor cells. *Journal of Cell Science* 131, jcs205575. doi:10.1242/jcs.205575.
- Burridge K and Connell LA (1983) New protein of adhesion plaques and ruffling membranes. *Journal of Cell Biology* 97, 359-367.
- Buyon JP, Slade SG, Reibman J, Abramson JB, Philips MR, Weissmann G and Winchester R (1990) Constitutive and induced phosphorylation of the α - and β -chains of the CD11/CD18 leukocyte integrin family. *Journal of Immunology* 144, 191-197.
- Byron A, Humphries JD, Bass MD, Knight D and Humphries MJ (2011) Proteomic analysis of integrin adhesion complexes. *Science Signaling* 4, 167 pt2.
- Calderwood DA (2015) The Rap1-RIAM pathway prefers β 2 integrins. *Blood* 126, 2658-2659.
- Calderwood DA, Campbell ID and Critchley DR (2013) Talins and kindlins: partners in integrin-mediated adhesion. *Nature Reviews Molecular Cell Biology* 14, 503-517
- Calderwood DA, Zent R, Grant R, Rees DJ, Hynes RO and Ginsberg MH (1999) The talin head domain binds to integrin- β subunit cytoplasmic tails and regulates integrin activation. *Journal of Biological Chemistry* 274, 28071-28074.
- Campadelli-Fiume G, Collins-McMillen D, Gianni T and Yurochko AD (2016) Integrins as herpesvirus receptors and mediators of the host signalosome. *Annual Review of Virology* 3, 215-236.
- Campbell ID (2013) Characterization of 14-3-3- ζ interactions with integrin tails. *Journal of Molecular Biology* 425, 3060-3072.
- Carragher NO, Levkau B, Ross R and Raines EW (1999) Degraded collagen fragments promote rapid disassembly of smooth muscle focal adhesions that correlates with cleavage of pp125(FAK), paxillin, and talin. *Journal of Cell Biology* 147, 619-629.
- Castagna M, Takai Y, Kaibuchi K, Sano K, Kikkawa U and Nishizuka Y (1982) Direct activation of calcium-activated, phospholipid-dependent protein-kinase by tumor-promoting phorbol esters. *Journal of Biological Chemistry* 257, 7847-7851.
- Chang DD, Wong C, Smith H and Liu J (1997) ICAP-1, a novel β 1 integrin cytoplasmic domain-associated protein, binds to a conserved and functionally important NPXY sequence motif of β 1 integrin. *Journal of Cell Biology* 138, 1149-1157.
- Chang VT, Fernandes RA, Ganzinger KA, Lee SF, Siebold C, McColl J, Jönsson P, Palayret M, Harlos K, Coles CH, Jones EY, Lui Y, Huang E, Gilbert RJC, Klenerman D, Aricescu AR and Davis SJ (2016) Initiation of T cell signaling by CD45 segregation at 'close contacts'. *Nature Immunology* 17, 574-582.
- Chang Y-C, Su W, Cho E-a, Zhang H, Huang Q, Philips MR and Wu J (2019) Molecular basis for autoinhibition of RIAM regulated by FAK in integrin activation. *Proceedings of the National Academy of Sciences of the United States of America* 116, 3524-3529

- Chatila TA, Geha RS and Arnaout MA (1989) Constitutive and stimulus-induced phosphorylation of CD11/CD18 leukocyte adhesion molecules. *Journal of Cell Biology* 109, 3435-3444.
- Chatterjee D, D'Souza A, Zhang Y, Bin W, Tan S-M and Bhattacharjya S (2018a) Interaction analyses of 14-3-3 ζ , Dok1, and phosphorylated integrin β cytoplasmic tails reveal a bi-molecular switch in integrin regulation. *Journal of Molecular Biology* 430, 4419-4430.
- Chatterjee D, Zhiping LL, Tan S-M and Bhattacharjya S (2016) Interaction analyses of the integrin β 2 cytoplasmic tail with the F3 FERM domain of talin and 14-3-3 ζ reveal a ternary complex with phosphorylated tail. *Journal of Molecular Biology* 428, 4129-4142.
- Chatterjee D, Zhiping LL, Tan S-M and Bhattacharjya S (2018b) NMR structure, dynamics and interactions of the integrin β 2 cytoplasmic tail with filamin domain IgFLNa21. *Scientific Reports* 8, 5490. doi:1038/s41598-018-23866-6.
- Chen C and Sheppard D (2007) Identification and molecular characterization of multiple phenotypes in integrin knockout mice. *Methods in Enzymology* 426, 291-305.
- Chen X, Xie C, Nishida N, Li Z, Walz T and Springer TA (2010) Requirement of open headpiece conformation for activation of leukocyte integrin α X β 2. *Proceedings of the National Academy of Sciences of the United States of America* 107, 14727-14732.
- Chen X, Yu YM, Mi LZ, Waltz T and Springer TA (2012) Molecular basis for complement recognition by integrin alpha(X) beta (2). *Proceedings of the National Academy of Sciences of the United States of America* 109, 4586-4591.
- Chigaev A, Smagley Y and Sklar LA (2011) Nitric oxide/cGMP pathway signaling actively down-regulates α 4 β 1-integrin affinity: an unexpected mechanism for inducing cell de-adhesion. *BMC Immunology* 12, 28.
- Chigaev A, Smagley Y and Sklar LA (2014) Carbon monoxide down-regulates α 4 β 1 integrin-specific ligand binding and cell adhesion: a possible mechanism for cell mobilization. *BMC Immunology* 15, 55.
- Chigaev A, Waller A, Amit O and Sklar LA (2008) G α phs-coupled receptor signaling actively down-regulates alpha4beta1-integrin affinity: a possible mechanism for cell-deadhesion. *BMC Immunology* 9, 26.
- Chinthalapudi K, Rangarajan ES and Izard T (2018) The interaction of talin with the cell membrane is essential for integrin activation and focal adhesion formation. *Proceedings of the National Academy of Sciences of the United States of America* 115, 10339-10344.
- Choi EY, Chavakis E, Czabanka MA, Langer HF, Fraemohs L, Economopoulou M, Kundu RK, Orlandi A, Zheng YY, Prieto DA, Ballantyne CM, Constant SL, Aird WC, Papayannopoulou T, Gahmberg CG, Udey MC, Vajkoczy O, Quartermous T, Dimmeler S, Weber C and Chavakis T (2008a) Del-1, an endogenous leukocyte-endothelial adhesion inhibitor, limits inflammatory cell recruitment. *Science* 322, 1101-1104.

Choi EY, Orlova VV, Fagerholm SC, Nurmi SM, Zhang L, Ballantyne CM, Gahmberg CG and Chavakis T (2008b) regulation of LFA-1-dependent inflammatory cell recruitment by Cbl-b and 14-3-3 proteins. *Blood* 111, 3607-3614.

Coppolino M, Leung-Hagesteijn C, Dedhar S and Wilkins J (1995) Inducible interaction of integrin $\alpha 2\beta 1$ with calreticulin. Dependence on the activation state of the integrin. *Journal of Biological Chemistry* 270, 23132-23138.

Coppolino MG, Woodside MJ, Demaurex N, Grinstein S, St-Arnaud R and Dedhar S (1997) Calreticulin is essential for integrin-mediated calcium signalling and adhesion. *Nature* 386, 843-847.

Corsini A, Maggi FM and Catapano AL (1995) Pharmacology of competitive inhibitors of HMG-CoA reductase. *Pharmacological Research* 31, 9-27.

Courtney AH, Amacher JF, Kadlecsek TA, Mollenauer MN, Au-Yeung BB, Kuriyan J and Weiss A (2017) A phosphosite within the SH2 domain of Lck regulates its activation by CD45. *Molecular Cell* 67, 498-511.

Courtney AH, Lo W-L and Weiss A (2018) TCR signaling: mechanisms of initiation and propagation. *Trends in Biochemical Sciences* 43, 108-123.

Critchley DR (2009) Biochemical and structural properties of the integrin-associated cytoskeletal protein talin. *Annual Review of Biophysics* 38, 235-254.

Das M, Ithychanda SS, Qin J and Plow EF (2011) Migfilin and filamin as regulators of integrin activation in endothelial cells and neutrophils. *PLoS One* 6, e26355

Davies JC, Tamaddon-Jahromi S, Jannoo R and Kanamalapudi V (2014) Cytohesin 2/ARF6 regulates preadipocyte migration through the activation of ERK 1/2. *Biochemical Pharmacology* 15, 651-660.

Diamond MS, Staunton DE, Marlin SD and Springer TA (1991) Binding of the integrin Mac-1 (CD11B/CD18) to the 3rd immunoglobulin-like domain of ICAM-1 (CD54) and its regulation by glycosylation. *Cell* 65, 961-971.

Dimitrov S, Lange T, Gouttefangeas C, Jensen ATR, Szczepanski M, Lehnholz J, Soekadar S, Rammensee H-G, Born J and Besedovsky L (2019) Gas-coupled receptor signaling and sleep regulate integrin activation of human antigen-specific T cells. *Journal of Experimental Medicine* 216, 517-526.

Dominguez GA, Anderson NR and Hammer DA (2015) The direction of migration of T-lymphocytes under flow depends upon which adhesion receptors are engaged. *Integrative Biology* 7, 345-355.

Dong XC, Mi LZ, Zhu JH, Wang W, Hu O, Luo BH and Springer TA (2012) Alpha(v)beta(3) integrin crystal structures and their functional implications. *Biochemistry* 51, 8814-8828.

Downey-Biechler C, Craig DH, More SK and Basson MD (2019) Inside-out signaling through FAK-integrin axis may regulate circulating cancer cell metastatic adhesion. *Proceedings of the National Academy of Sciences of the United States of America* 116, 19795-19796.

- Dransfield I, Cabanas C, Craig A and Hogg N (1992) Divalent-cation regulation of the function of the leukocyte integrin LFA-1. *Journal of Cell Biology* 116, 219-226.
- Durrant TN, van den Bosch M and Hers I (2017) Integrin $\alpha\text{IIb}\beta 3$ outside-in signaling. *Blood* 130, 1607-1619.
- Dutta D, Barr VA, Akpan I, Mittelstadt PR, Singha LI, Samelson LE and Ashwell JD (2017) Recruitment of calcineurin to the TCR positively regulates T cell function. *Nature Immunology* 18, 196-204.doi:10.1038/ni.3640.
- Fagerholm SC, Guenther C, Asens ML, Savinko T and Uotila LM (2019) Beta2-integrins and interacting proteins in leukocyte trafficking, immune suppression, and immunodeficiency disease. *Frontiers in Immunology* 10, article 254.
- Fagerholm S, Hilden TJ and Gahmberg CG (2002a) Lck tyrosine kinase is important for activation of the CD11a/CD18 integrins in human T lymphocytes. *European Journal of Immunology* 32, 1670-1678.
- Fagerholm SC, Hilden TJ, Nurmi SM and Gahmberg CG (2005) Specific integrin α and β chain phosphorylations regulate LFA-1 activation through affinity-dependent and -independent mechanisms. *Journal of Cell Biology* 171, 705-715.
- Fagerholm S, Morrice N, Gahmberg CG and Cohen P (2002b) Phosphorylation of the cytoplasmic domain of the integrin CD18 chain by protein kinase C isoforms in leukocytes. *Journal of Biological Chemistry* 277, 1728-1738.
- Fagerholm S, Prescott A, Cohen P and Gahmberg CG (2001) An essential role for calmodulin in regulating human T cell aggregation. *FEBS Letters* 491, 131-136.
- Fagerholm SC, Varis M, Stefanidakis M, Hilden TJ and Gahmberg CG (2006) α -chain phosphorylation of the human leukocyte CD11b/CD18 (Mac-1) integrin is pivotal for integrin activation to bind ICAMs and leukocyte extravasation in vivo. *Blood* 108, 3379-3386.
- Fong KP, Zhu H, Span LM, Moore DT, Yoon K, Tamura R, Yin H, DeGrado WF and Bennett JS (2016) Directly activating the integrin $\alpha\text{IIb}\beta 3$ initiates outside-in signaling by causing $\alpha\text{IIb}\beta 3$ clustering. *Journal of Biological Chemistry* 291, 11706-11716.
- Franco SJ and Huttenlocher A (2005) Regulating cell migration: Calpains make the cut. *Journal of Cell Science* 118, 829-838.
- Franco SJ, Rodgers MA, Perrin BJ, Han J, Bennin DA, Critchley DR and Huttenlocher A (2004) Calpain-mediated proteolysis of talin regulated adhesion dynamics. *Nature Cell Biology* 6, 977-983.
- Frenette PS (2001) Locking a leukocyte integrin with statins. *The New England Journal of Medicine* 345, 1419-1421.
- Gahmberg CG (1997) Leukocyte adhesion. CD11/CD18 integrins and intercellular adhesion molecules. *Current Opinion in Cell Biology* 9, 643-650.
- Gahmberg CG, Fagerholm SC, Nurmi SM, Chavakis T, Marchesan S and Grönholm M (2009) Regulation of integrin activity and signalling. *Biochimica et Biophysica Acta* 1790, 431-444.

- Gahmberg CG and Hakomori S (1973) Altered growth behavior of malignant cells associated with changes in externally labeled glycoprotein and glycolipid. *Proceedings of the National Academy of Sciences of the United States of America* 70, 3329-3333.
- Gahmberg CG, Häyry P and Andersson LC (1976) Characterization of surface glycoproteins of mouse lymphoid cells. *Journal of Cell Biology* 68, 642-653.
- Gahmberg CG, Tian L, Ning L and Nyman-Huttunen H (2008) ICAM-5 – a novel two-faceted adhesion molecule in the mammalian brain. *Immunology Letters* 117, 131-135
- Gahmberg CG, Tolvanen M and Kotovuori P (1997) Leukocyte adhesion. Structure and function of human leukocyte β 2-integrins and their cellular ligands. *European Journal of Biochemistry* 245, 215-232.
- Gallit J and Ruoslahti E (1988) Regulation of the fibronectin receptor affinity by divalent-cations. *Journal of Biological Chemistry* 263, 12927-12932
- Gao J, Bao Y, Ge S, Sun P, Sun J, Liu J, Chen F, Han L, Cao Z, Qin J, White GC, Xu Z and Ma Y-Q (2019) Sharpin suppresses β 1-integrin activation by complexing with the β 1 tail and kindlin-1. *Cell Communication and Signaling* doi.org/10.1186/s12964-019-0407-6.
- Garragher NO, Levkau B, Ross R and Raines EW (1999) Degraded collagen fragments promote rapid disassembly of smooth muscle focal adhesions that correlates with cleavage of pp125(FAK), paxillin, and talin. *Journal of Cell Biology* 147, 619-630.
- Gatta L, Calviello G, di Nicuolo F, Pace L, Ubaldi V, Doria G and Pioli C (2002) Cytotoxic T lymphocyte-associated antigen-4 inhibits integrin-mediated stimulation. *Immunology* 107, 209-2016.
- Gbarah A, Gahmberg CG, Ofek I, Jacobi U and Sharon N (1991) Identification of the leukocyte adhesion molecules CD11 and CD18 as receptors for type 1 fimbriated (Mannose specific) *Escherichia coli*. *Infection and Immunity* 59, 4524-4530.
- Geng X, Tang R-H, Law SKA and Tan S-M (2005) Integrin CD11a cytoplasmic tail interacts with the CD45 membrane-proximal protein tyrosine phosphatase domain 1. *Immunology* 115, 347-357.
- Ghosh S, Chackerian AA, Parker CM, Ballantyne CM and Behar SM (2006) The LFA-1 adhesion molecule is required for protective immunity during pulmonary *Mycobacterium tuberculosis* infection. *Journal of Immunology* 176, 4914-4922.
- Giancotti FG and Ruoslahti E (1999) Integrin signaling. *Science* 285, 1028-1032.
- Gingras AR, Lagarrique F, Cuevas MN, Valadez AJ, Zorovich M, McLaughlin W, Lopez-Ramirez MA, Seban N, Ley K, Kiosses WB and Ginsberg MH (2019) Rap1 binding and a lipid-dependent helix in talin F1 domain promote integrin activation in tandem. *Journal of Cell Biology* doi.org/10.1083/jcb.201810061.
- Glading A, Lauffenburger D and Wells A (2002) Cutting to the chase: Calpain proteases in cell motility. *Trends in Cell Biology* 12, 46-54.

Gong Y, Zhang Y, Feng S, Liu X, Lu S and Long M (2017) Dynamic distributions of P- and E-selectins to beta2-integrin-induced neutrophil transmigration. *FASEB Journal* 31, 212-223.

Gonzales AM, Claiborne J and Jones CR (2008) Integrin cross-talk in endothelial cells is regulated by protein kinase A and protein phosphatase 1. *Journal of Biological Chemistry* 283, 31849-31860.

Goult BT and Schwartz MA (2018) Talin as a mechanosensitive signaling hub. *Journal of Cell Biology* 217, 3776-3784.

Grakoui A, Bromley SK, Sumen C, Davis MM, Shaw AS, Allen PM and Dustin ML (1999) The immunological synapse: a molecular machine controlling T cell activation. *Science* 285, 221-227.

Grönholm M, Jahan F, Bryushkova EA, Madhavan S, Aglialoro F, Hinojosa LS, Uotila LM and Gahmberg CG (2016) LFA-1 integrin antibodies inhibit leukocyte $\alpha 4\beta 1$ -mediated adhesion by intracellular signaling. *Blood* 128, 1270-1281.

Grönholm M, Jahan F, Marchesan S, Karvonen U, Aatonen M, Narumanchi S and Gahmberg CG (2011) TCR-induced activation of LFA-1 involves signaling through Tiam1. *Journal of Immunology* 187, 3613-3619.

Gupta S, Chit JC-Y, Feng C, Bhunia A, Tan S-M and Bhattacharjya S (2015) An alternative phosphorylation switch in integrin $\beta 2$ (CD18) tail for Dok1 binding. *Scientific Reports* 5, 11630, doi:10.1038/srep11630.

Han DC, Rodriguez LG and Guan JL (2001) Identification of a novel interaction between integrin $\beta 1$ and 14-3-3 β . *Oncogene* 20, 346-357.

Han J, Rose DM, Woodside DG, Goldfinger LE and Ginsberg MH (2003) Integrin $\alpha 4\beta 1$ -dependent T cell migration requires both phosphorylation and dephosphorylation of the $\alpha 4$ cytoplasmic domain to regulate the reversible binding of paxillin. *Journal of Biological Chemistry* 278, 34845-34853.

Hibbs ML, Jakes S, Stacker SA, Wallace RW and Springer TA (1991) The cytoplasmic domain of the integrin lymphocyte function-associated antigen 1 β subunit: sites required for binding to intercellular adhesion molecule 1 and the phorbol ester-stimulated phosphorylation site. *Journal of Experimental Medicine* 174, 1227-1238.

Hilden TJ, Valmu L, Kärkkäinen S and Gahmberg CG (2003) Threonine phosphorylation sites in the $\beta 2$ and $\beta 7$ leukocyte integrin polypeptides. *Journal of Immunology* 170, 4170-4177.

Hogg N, Patzak I and Willenbrock F (2011) The insider's guide to leukocyte integrin signaling and function. *Nature Reviews Immunology* 11, 416-426.

Hu P and Luo B-H (2013) Integrin bi-directional signaling across the plasma membrane. *Journal of Cellular Physiology* 228, 306-312.

Hughes PE, Diaz-Gonzales F, Leong L, Wu C, McDonald JA, Shattil SJ and Ginsberg MH (1996) Breaking the integrin hinge. A defined structural constraint regulates integrin signaling. *Journal of Biological Chemistry* 271, 6571-6574.

Humphries JD, Byron A and Humphries MJ (2006) Integrin ligands. *Journal of Cell Science* 119, 3901-3903.

- Huse M (2017) Mechanical forces in the immune system. *Nature Reviews Immunology* 17, 679-690.
- Huttenlocher A and Horwitz AR (2011) Integrins in cell migration. *Cold Spring Harbor Perspectives in Biology* 2011;3;a005074.
- Hynes RO (1973) Alteration of cell-surface proteins by viral transformation and by proteolysis. *Proceedings of the National Academy of Sciences of the United States of America* 70, 3170-3174.
- Hynes RO (1987) Integrins: a family of cell surface receptors. *Cell* 48, 549-554.
- Hynes RO (1992) Integrins – versatility, modulation, and signaling in cell-adhesion. *Cell* 69, 11-25.
- Hynes RO (2002) Integrins: bidirectional, allosteric signaling machines. *Cell* 110, 673-687.
- Hynes RO and Yamada KM (1982) Fibronectins-multifunctional modular glycoproteins. *Journal of Cell Biology* 95, 369-377.
- Ithychanda SS, Fang X, Mohan ML, Zhu L, Tirupula KC, Prasad SVN, Wang Y-X, Karnik SS and Qin J (2015) A mechanism of global shape-dependent recognition and phosphorylation of filamin by protein kinase A. *Journal of Biological Chemistry* 290, 8527-8538.
- Jackson T, Sheppard D, Denyer M, Blakemore W and King AMQ (2000) The epithelial integrin $\alpha\text{v}\beta 6$ is a receptor for foot-and-mouth disease virus. *Journal of Virology* 74, 4949-4956.
- Jahan F, Madhavan S, Rolova T, Viazmina L, Grönholm M and Gahmberg CG (2018) Phosphorylation of the α -chain in the integrin LFA-1 enables $\beta 2$ -chain phosphorylation and α -actinin binding required for cell adhesion. *Journal of Biological Chemistry* 293, 12318-12330.
- Jenkins AL, Nannizzi-Alaimo L, Silver D, Sellers JR, Ginsberg MH, Law DA and Phillips DR (1998) Tyrosine phosphorylation of the $\beta 3$ cytoplasmic domain mediates integrin-cytoskeletal interactions. *Journal of Biological Chemistry* 273, 13878-13885.
- Johnson KG, Bromley SK, Dustin ML and Thomas ML (2000) A supramolecular basis for CD45 tyrosine phosphatase regulation in sustained T cell activation. *Proceedings of the National Academy of Sciences of the United States of America* 97, 10138-10143.
- Kammerer P, Aretz J and Fässler R (2017) Lucky kindlin: a cloverleaf at the integrin tail. *Proceedings of the National Academy of Sciences of the United States of America* 114, 9234-9236.
- Kanwar S, Smith CW, Shardonofsky FR and Burns AR (2001) The role of Mac-1 (CD11b/CD18) in antigen-induced airway eosinophilia in mice. *American Journal of Respiratory Cell and Molecular Biology* 25, 170-177.
- Kasirer-Friede A, Tjahjono W, Eto K and Shattil SJ (2019) SHARPIN at the nexus of integrin, immune, and inflammatory signaling in human platelets. *Proceedings of the National Academy of Sciences of the United States of America*. doi 10.1073/pnas.1819156116.
- Katagiri K, Imamura M and Kinashi T (2006) Spatiotemporal regulation of the kinase Mst1 by binding protein RAPL is critical for lymphocyte polarity and adhesion. *Nature Immunology* 7, 919-928.

- Katagiri K, Maeda A, Shimonaka M and Kinashi T (2003) RAPL, a Rap1-binding molecule that mediates Rap1-induced adhesion through spatial regulation of LFA-1. *Nature Immunology* 4, 741-748.
- Kiema T, Lad Y, Jiang P, Oxley CL, Baldassarre M, Wegener KL, Campbell ID, Ylänne J and Calderwood DA (2006) The molecular basis of filamin binding to integrins and competition with talin. *Molecular Cell* 21, 337-347.
- Kim C, Lau T-L, Ulmer TS and Ginsberg MH (2009) Interactions of platelet integrin α IIb and β 3 transmembrane domains in mammalian cell membranes and their role in integrin activation. *Blood* 113, 4747-4753.
- Kim CK, Ye F, Hu X and Ginsberg MH (2012) Talin activates integrins by altering the topology of the β transmembrane domain. *Journal of Cell Biology* 197, 605-611.
- Kim M, Carman CV and Springer TA (2003) Bidirectional transmembrane signaling by cytoplasmic domain separation in integrins. *Science* 301, 1720-1725.
- Kirk RI, Sanderson MR and Lerea KM (2000) Threonine phosphorylation of the β 3 integrin cytoplasmic tail, at a site recognized by PDK1 and Akt/PKB in vitro, regulates Shc binding. *Journal of Biological Chemistry* 275, 30901-30906.
- Klapholz B and Brown NH (2017) Talin-the master of integrin adhesions. *Journal of Cell Science* 130, 2435-2446.
- Klapproth S, Bromberger T, Turk C, Kruger M and Moser M (2019) A kindlin-3-leupaxin-paxillin signaling pathway regulates podosome stability. *Journal of Cell Biology* 218, 3436.
doi:10.1083/jcb.201903109.
- Klapproth S, Moretti FA, Zeiler M, Ruppert R, Breithaupt U, Mueller S, Haas R, Mann M, Sperandio M, Fässler R and Moser M (2015a) Minimal amounts of kindlin-3 suffice for basal platelet and leukocyte functions in mice. *Blood* 126, 2592-2600.
- Klapproth S, Sperandio M, Pinheiro EM, Prunster M, Soehnlein O, Gertler FB, Fässler R and Moser M (2015b) Loss of the Rap1 effector RIAM results in leukocyte adhesion deficiency due to impaired β 2 integrin function in mice. *Blood* 126, 2704-2712.
- Knight CG, Morton LF, Peachey AR, Tuckwell DS, Farndale RW and Barnes MJ (2000) The collagen-binding A-domains of integrins α 1 β 1 and α 2 β 1 recognize the same specific amino acid sequence, GFOGER, in native (triple-helical) collagens. *Journal of Biological Chemistry* 275, 35-40.
- Koivunen E, Ranta TM, Annala A, Taube S, Uppala A, Jokinen M, van Willigen G, Ihanus E and Gahmberg CG (2001) Inhibition of β 2 integrin-mediated leukocyte cell adhesion by leucine-leucine-glycine motif-containing peptides. *Journal of Cell Biology* 153, 905-916.
- Kolanus W (2007) Guanine nucleotide exchange factors of the cytohesin family and their roles in signal transduction. *Immunological Reviews* 218, 102-113.

Kondo N, Ueda Y, Kita T, Ozawa M, Tomiyama T, Yasuda K, Lim D-S and Kinashi T (2017) NDR1-dependent regulation of kindlin-3 controls high-affinity LFA-1 binding and immune synapse organization. *Molecular and Cellular Biology* 37, e00424-16.

Kong K-F and Altman A (2013) In and out of the bull's eye: protein kinase Cs in the immunological synapse. *Trends in Immunology* 34, 234-242.

Kotovuori A, Pessa-Morikawa T, Kotovuori P, Nortamo P and Gahmberg CG (1999) ICAM-2 and a peptide from its binding domain are efficient activators of leukocyte adhesion and integrin affinity. *Journal of Immunology* 162, 6613-6620.

Kotovuori P, Tontti E, Pigott R, Shepherd M, Kiso M, Hasegawa A, Renkonen R, Nortamo P, Altieri DC and Gahmberg CG (1993) The vascular E-selectin binds to the leukocyte integrins CD11/CD18. *Glycobiology* 3, 131-136.

Kourtzelis I, Mitroulis I, von Renesse J, Hajishengallis G and Chavakis T (2017) From leukocyte recruitment to resolution of inflammation: the cardinal role of integrins. *Journal of Leukocyte Biology* 102, 677-683.

Krause K-H and Michalak M (1997) Calreticulin. *Cell* 88, 439-443.

Savinko T, Guenther C, Uotila LM, Asens ML, Yao S, Tojkander S and Fagerholm SC (2018) Filamin A is required for optimal T cell integrin-mediated force transmission, flow adhesion, and T cell trafficking. *Journal of Immunology* 200, 3109-3116.

Krueger J, Gottlieb A, Miller B, Dedrick R, Garovoy M and Walicke P (2000) Anti-CD11a treatment for psoriasis concurrently increases circulating T-cells and decreases plaque T-cells, consistent with inhibition of cutaneous T-cell trafficking. *Journal of Investigative Dermatology* 115, 333.

Lagarrique F, Kim C and Ginsberg MH (2016) The Rap1-RIAM-talin axis of integrin activation and blood cell function. *Blood* 128, 479-487.

Law DA, DeGuzman FR, Heiser P, Ministri-Madrid K, Killeen N and Phillips DR (1999) Integrin cytoplasmic tyrosine motif is required for outside-in α IIb β 3 signalling and platelet function. *Nature* 401, 808-811.

Lee J-O, Rieu P, Arnaout MA and Liddington R (1995) Crystal structure of the A domain from the α subunit of integrin CR3 (CD11b/CD18) *Cell* 80, 631-638.

Lerea KM, Cordero KP, Sakariassen KS, Kirk RI and Fried VA (1999) Phosphorylation sites in the integrin β 3 cytoplasmic domain in intact platelets. *Journal of Biological Chemistry* 274, 1914-1919.

Li H, Deng Y, Sun K, Yang H, Liu J, Wang M, Zhang Z, Lin J, Wu C, Wei Z and Yu C (2017) Structural basis of kindlin-mediated integrin recognition and activation. *Proceedings of the National Academy of Sciences of the United States of America* 114, 9349-9354.

Li J and Springer TA (2017) Integrin extension enables ultrasensitive regulation by cytoskeletal force. *Proceedings of the National Academy of Sciences of the United States of America* 114, 4685-4690.

- Li J and Springer TA (2017) Energy landscape differences among integrins establish the framework for understanding activation. *Journal of Cell Biology* 217, 397-412.
- Li R, Nortamo P, Kantor C, Kovanen P, Timonen T and Gahmberg CG (1993a) A leukocyte integrin binding peptide from intercellular adhesion molecule-2 stimulates T cell adhesion and natural killer cell activity. *Journal of Biological Chemistry* 268, 21474-21477.
- Li Z, Zhang H, Lundin L, Thullberg M, Liu Y, Wang Y, Claesson-Welsh L and Strömblad S (2010) p21-activated kinase 4 phosphorylation of integrin $\beta 5$ Ser-759 and Ser-762 regulates cell migration. *Journal of Biological Chemistry* 285, 23699-23710.
- Lilienthal J and Chang DD (1998) Rack1, a receptor for activated protein kinase C, interacts with integrin β subunit. *Journal of Biological Chemistry* 273, 2379-2383.
- Lilja J, Zacharchenko T, Georgiadou M, Jacquemet G, De Franceschi N, Peuhu E, Hamidi H, Pouwels J, Martens V, Nia FH, Beifuss M, Boeckers T, Kreienkamp H-J, Barsukov IL and Ivaska J (2017) SHANK proteins limit integrin activation by directly interacting with Rap1 and R-Ras. *Nature Cell Biology* 19, 292-305.
- Lim CJ, Kain KH, Tkachenko E, Goldfinger LE, Gutierrez E, Allen MD, Groisman A, Zhang J and Ginsberg MH (2008) Integrin-mediated protein kinase A activation at the leading edge of migrating cells. *Molecular Biology of the Cell* 19, 4930-4941.
- Lim J, Hotchin NA and Caron E (2011) Ser756 of $\beta 2$ integrin controls Rap1 activity during inside-out activation of $\alpha M\beta 2$. *Biochemical Journal* 437, 461-467.
- Lin C, Zhang Y, Zhang K, Zheng Y, Lu L, Chang H, Yang H, Yang Y, Wan Y, Wang S, Yuan M, Yan Z, Zhang R, He Y, Ge G, Wu D and Chen J (2019) Fever promotes T lymphocyte trafficking via a thermal sensory pathway involving heat shock protein 90 and $\alpha 4$ integrins. *Immunity* 50, 137-151.
- Liu J, Das M, Yang J, Ithychanda SS, Yakubenko V, Plow EF and Qin J (2015a) Structural mechanism of integrin inactivation by filamin. *Nature Structural & Molecular Biology* 22, 383-391.
- Liu J, Wang Z, Thinn AMM, Ma Y-Q and Zhu J (2015b) The dual structural roles of the membrane distal region of the α -integrin cytoplasmic tail during integrin inside-out activation. *Journal of Cell Science* 128, 1718-1731.
- Liu S and Ginsberg MH (2000) Paxillin binding to a conserved sequence motif in the $\alpha 4$ integrin cytoplasmic domain. *Journal of Biological Chemistry* 275, 22736-22742.
- Lu L, Lin C, Yan Z, Wang S, Zhang Y, Wang S, Wang J, Liu C and Chen J (2016) Kindlin-3 is essential for the resting $\alpha 4\beta 1$ integrin-mediated firm cell adhesion under shear flow conditions. *Journal of Biological Chemistry* 291, 10363-10371.
- Luo BH, Carman CV and Springer TA (2007) Structural basis of integrin regulation and signaling. *Annual Review of Immunology* 25, 619-647.
- MacKenzie KR, Prestegard JH and Engelman DM (1997) A transmembrane helix dimer, structure and implications. *Science* 276, 131-133.

- MacPherson M, Savinko T, Lek HS, Prescott A and Fagerholm SC (2013) The β 2 integrin-kindlin-3 interaction is essential for T cell homing but dispensable for T- cell activation in vivo. *Blood* 122, 1428-1436.
- Malinin NL, Zhang L, Choi J, Ciocea A, Razorenova O, Ma Y-Q, Podrez EA, Tosi M, Lennon DP, Caplan AI, Shurin SB, Plow EF and Byzova TV (2009) A point mutation in KINDLIN3 ablates activation of three integrin subfamilies in humans. *Nature Medicine* 15, 313-318.
- Mitroulis J, Alexaki VI, Kourtzelis J, Ziogas A, Hajishengallis G and Chavakis T (2015) Leukocyte integrins: role in leukocyte recruitment and as therapeutic targets in inflammatory disease. *Pharmacology & Therapeutics* 147, 123-135.
- Moore, TI, Aaron J, Chew T-L and Springer TA (2018) Measuring integrin conformational change on the cell surface with super-resolution microscopy. *Cell Reports* 22, 1903-1912.
- Morikis VA, Chase S, Wun T, Chaikof EL, Magnani JL and Simon SI (2017) Selectin catch –bonds mechanotransduce integrin activation and neutrophil arrest on inflamed endothelium under shear flow. *Blood* 130, 2101-2110.
- Morrison VL, MacPherson M, Savinko T, Lek HS, Prescott A and Fagerholm SC (2013) The β 2 integrin-kindlin-3 interaction is essential for T cell homing but dispensable for T- cell activation in vivo. *Blood* 122, 1428-1436.
- Moser M, Bauer M, Schmid S, Ruppert R, Schmidt S, Sixt M, Wang H-V, Sperandio M and Fässler R (2009a) Kindlin-3 is required for β 2 integrin-mediated leukocyte adhesion to endothelial cells. *Nature Medicine* 15, 300-305
- Moser M, Legate KR, Zent R and Fässler R (2009b) The tail of integrins, talin and kindlins. *Science* 324, 895-899.
- Muslin AJ, Tanner JW, Allen PM and Shaw AS (1996) Interaction of 14-3-3 with signaling proteins is mediated by the recognition of phosphoserine. *Cell* 84, 889-897.
- Mustelin T, Coggeshall KM and Altman A (1989) Rapid activation of the T-cell tyrosine protein kinase pp56lck by the CD45 phosphotyrosine phosphatase. *Proceedings of the National Academy of Sciences of the United States of America* 86, 6302-6306.
- Mustelin T, Pessa-Morikawa T, Autero M, Gassman M, Gahmberg CG, Andersson LC and Burn P (1992) Regulation of the pp59fyn tyrosine kinase by the CD45 phosphotyrosine phosphatase. *European Journal of Immunology* 22, 1173-1178.
- Needham EJ, Parker BL, Burykin T, James DE and Humphrey SJ (2019) Illuminating the dark phosphoproteome. *Science Signaling* 12, eaau8645.
- Newton AC (1995) Protein kinase C: structure, function, and regulation. *Journal of Biological Chemistry* 270, 28495-2849
- Newton AC and Brognard J (2017) Reversing the paradigm: protein kinase C as a tumor suppressor. *Trends in Pharmacological Sciences* 38, 438-447.

- Nilsson S, Kaniowska D, Brakebusch C, Fässler R and Johansson S (2006) Threonine 788 in integrin subunit $\beta 1$ regulates integrin activation. *Experimental Cell Research* 312, 844-853.
- Nishida N, Xie C, Shimaoka M, Cheng Y, Walz T and Springer TA (2006) Activation of leukocyte $\beta 2$ integrins by conversion from bent to extended conformations. *Immunity* 25, 583-594.
- Nishizuka Y (1984) The role of protein kinase-C in cell surface signal transduction and tumor promotion. *Nature* 308, 693-698.
- Nordenfelt P, Elliott HL and Springer TA (2016) Coordinated integrin activation by actin-dependent force during T-cell migration. *Nature Communications* 7, 13119
doi:10.1038/ncomms13119.
- Nortamo P, Patarroyo M, Kantor C, Suopanki J and Gahmberg CG (1988) Immunological mapping of the human leukocyte adhesion glycoprotein GP90 (CD18) by monoclonal antibodies. *Scandinavian Journal of Immunology* 28, 537-546.
- Nurmi SM, Autero M, Raunio AK, Gahmberg CG and Fagerholm SC (2007) Phosphorylation of the LFA-1 integrin $\beta 2$ -chain on Thr-758 leads to adhesion, Rac-1/Cdc42 activation and stimulation of CD69 expression in human T cells. *Journal of Biological Chemistry* 282, 968-975.
- Nygren P, Balashova N, Brown AC, Kieba I, Dhingra A, Boesze-Battaglia K and Lally ET (2018) *Aggregatibacter actinomycetemcomitans* leukotoxin causes activation of lymphocyte function-associated antigen 1. *Cellular Microbiology* 21, doi.org/10.1111/cmi.12967.
- Obergfell A, Eto K, Mocsai A, Buensuceso C, Moores SL, Brugge JS, Lowell CA and Shattil SJ (2002) Coordinate interactions of Csk, Src, and Syk kinases with $\alpha \text{IIb}\beta 3$ initiate integrin signaling to the cytoskeleton. *Journal of Cell Biology* 157, 265-275.
- Okigaki M, Davis C, Falasca M, Harroch S, Felsenfeld DP, Sheetz MP and Schlesinger J (2003) Pyk2 regulates multiple signaling events crucial for macrophage morphology and migration. *Proceedings of the National Academy of Sciences of the United States of America* 100, 10740-10745.
- O'Toole TE, Bialkowska K, Li X and Fox JEB (2011) Tiam1 is recruited to $\beta 1$ -integrin complexes by 14-3-3 ζ where it mediates integrin-induced Rac1 activation and motility. *Journal of Cellular Physiology* 226, 2965-2978.
- Oxley CL, Anthis NJ, Lowe ED, Vakonakis I, Campbell ID and Wegener KL (2008) An integrin phosphorylation switch. The effect of $\beta 3$ integrin tail phosphorylation on Dok1 and talin binding. *Journal of Biological Chemistry* 283, 5420-5426.
- Parry RV, Chemnitz JM, Frauwirth KA, Lanfranco AR, Braunstein I, Kobayashi SV, Linsley PS, Thompson CB and Riley JL (2005) CTLA-4 and PD-1 receptors inhibit T-cell activation by distinct mechanisms. *Molecular and Cellular Biology* 25, 9543-9553.
- Patarroyo M, Beatty PG, Fabre JW and Gahmberg CG (1985) Identification of a cell surface protein complex mediating phorbol ester-induced adhesion (binding) among human mononuclear leukocytes. *Scandinavian Journal of Immunology* 22, 171-182.

Patarroyo M, Beatty PG, Serhan CN and Gahmberg CG (1985) Identification of a cell surface glycoprotein mediating adhesion in human granulocytes. *Scandinavian Journal of Immunology* 22, 619-631.

Patarroyo M, Clark EA, Prieto J, Kantor C and Gahmberg CG (1987) Identification of a novel adhesion molecule in human leukocytes by monoclonal antibody LB-2. *FEBS Letters* 210, 127-131.

Patarroyo M, Yogeewaran G, Biberfeld P, Klein E and Klein G (1982) Morphological-changes, cell-aggregation and Cell-membrane alterations caused by phorbol 12,13-dibutyrate in human-blood lymphocytes. *International Journal of Cancer* 30, 707-717.

Pavalko FM and LaRoche SM (1993) Activation of human neutrophils induces an interaction between the integrin β 2-subunit (CD18) and the actin binding protein α -actinin. *Journal of Immunology* 151, 3795-3807.

Perez OD, Mitchell D, Jager GC, South S, Murriel C, McBride J, Herzenberg LA, Kinoshita S and Nolan GP (2003) Leukocyte functional antigen 1 lowers T cell activation thresholds and signaling through cytohesin-1 and Jun-activating binding protein 1. *Nature Immunology* 4, 1083-1094.

Petosa C, Masters SC, Bankston LA, Pohl J, Wang B, Fu H and Liddington RC (1998) 14-3-3zeta binds a phosphorylated Raf peptide and an unphosphorylated peptide via its conserved amphipathic groove. *Journal of Biological Chemistry* 273, 16305-16310.

Philipsen L, Reddycherla AV, Hartig R, Gumz J, Kästle M, Kritikos A, Poltorak MP, Prokazov Y, Turbin E, Weber A, Zuschratter W, Schraven B, Simeoni L and Muller AJ (2017) De novo phosphorylation and conformational opening of the tyrosine kinase Lck act in concert to initiate T cell receptor signaling. *Science Signaling* 10, eaaf4736.

Phillips ML, Nudelman E, Gaeta FCA, Perez M, Singhal AK, Hakomori S and Paulson JC (1990) ELAM-1 mediates cell-adhesion by recognition of a carbohydrate ligand, sialyl –LeX. *Science* 250, 1130-1132.

Pierschbacher MD and Ruoslahti E (1984) Cell attachment activity of fibronectin can be duplicated by small synthetic fragments of the molecule. *Nature* 309, 30-33.

Polley MJ, Phillips ML, Wayner E, Nudelman E, Singhal AK, Hakomori S and Paulson JC (1991) CD62 and endothelial cell-leukocyte adhesion molecule I (ELAM-1) recognize the same carbohydrate ligand, sialyl-Lewis x. *Proceedings of the National Academy of Sciences of the United States of America* 88, 6224-6228.

Pons M, Izquierdo I, Andreu-Carbo M, Garrido G, Planaguma J, Muriel O, del Pozo MA, Geli MI and Aragay AM (2017) Phosphorylation of filamin A regulates chemokine receptor CCR2 recycling. *Journal of Cell Science* 130, 490-501.

Porter JC and Hogg N (1997) Integrin cross-talk: activation of lymphocyte function-associated antigen-1 on human T cells alters α 4 β 1- and α 5 β 1-mediated function. *Journal of Cell Biology* 138, 1437-1447.

- Pouwels J, de Franceschi N, Mattila E, Potter C, Sundberg J, Hogg N, Gahmberg CG, Salmi M and Ivaska J (2013) SHARPIN regulates uropod detachment in migrating lymphocytes. *Cell Reports* 5, 619-628.
- Puck A, Hopf S, Modak M, Majdic O, Cejka P, Blumi S, Schmetterer K, Arnold-Schrauf C, Gerwien JG, Frederiksen KS, Thell E, Leitner J, Steinberger P, Aigner R, Seyerl-Jiresch M, Zlabinger GJ and Stöckl J (2017) The soluble cytoplasmic tail of CD45 (ct-CD45) in human plasma contributes to keep T cells in a quiescent state. *European Journal of Immunology* 47, 193-205.
- Pytela R, Pierschbacher MD and Ruoslahti E (1985) Identification and isolation of a 140-kd cell-surface glycoprotein with properties expected of a fibronectin receptor. *Cell* 40, 191-198.
- Rantala JK, Pouwels J, Pellinen T, Veltel S, Laasola P, Mattila E, Potter CS, Duffy T, Sundberg JP, Kallioniemi O, Askari JA, Humphries MJ, Parsons M, Salmi M and Ivaska J (2011) SHARPIN is an endogenous inhibitor of β 1-integrin activation. *Nature Cell Biology* 13, 1315-1324.
- Ratnikov B, Ptak C, Han J, Shabanowitz J, Hunt DF and Ginsberg MH (2005) Talin phosphorylation sites mapped by mass spectrometry. *Journal of Cell Science* 118, 4921-4923.
- Robinson MK, Andrew D, Rosen H, Brown D, Orllepp S, Stephens P and Butcher EC (1992) Antibody against the Leu-cam β -chain (CD18) promotes both LFA-1- and CR3-dependent adhesion events. *Journal of Immunology* 148, 1080-1085.
- Rothlein R, Dustin ML, Marlin SD and Springer TA (1986) A human intercellular adhesion molecule (ICAM-1) distinct from LFA-1. *Journal of Immunology* 137, 1270-1274.
- Roy NH, MacKay JL, Robertson TF, Hammer DA and Burkhardt JK (2018) Crk adaptor proteins mediate actin-dependent T cell migration and mechanosensing induced by the integrin LFA-1. *Science Signaling* 11, eaat3178.
- Ruoslahti E (1988) Fibronectin and its receptors. *Annual Review of Biochemistry* 57, 375-413.
- Ruoslahti E (1991) Integrins. *Journal of Clinical Investigation* 87, 1-5.
- Ruoslahti E (1996) RGD and other recognition sequences for integrins. *Annual Review of cell and developmental biology* 12, 697-715.
- Sampath R, Gallagher PJ and Pavalko FM (1998) Cytoskeletal interactions with the leukocyte integrin β 2 cytoplasmic tail. *Journal of Biological Chemistry* 273, 33588-33594.
- Sanchez-Madrid F, Simon P, Thompson S and Springer TA (1983) Mapping of antigenic and functional epitopes on the α - and β -subunits of two related mouse glycoproteins involved in cell interactions. *Journal of Experimental Medicine* 158, 586-602.
- Schneider H, Valk E, da Rocha Dias S, Wei B and Rudd CE (2005) CTLA-4 up-regulation of lymphocyte function-associated antigen 1 adhesion and clustering as an alternative basis for coreceptor function. *Proceedings of the National Academy of Sciences of the United States of America*. 102, 12861-12866.
- Schurpf T, Chen Q, Liu JH, Wang R, Springer TA and Wang JH (2012) The RGD finger of Del-1 is a unique structural feature critical for integrin binding. *FASEB Journal* 26, 3412-3420.

Schwartz MA, Schaller MD and Ginsberg MH (1995) Integrins: emerging paradigms of signal transduction. Annual Review of Cell and Developmental Biology 11, 549-599.

Scott LM, Priestley GV and Papayannopolou T (2003) Deletion of $\alpha 4$ integrins from adult hematopoietic cells reveals roles in homeostasis, regeneration, and homing. Molecular and Cellular Biology doi: 10.1128/MCB.23.24.9349-9360.2003. Sharma P and Allison JP (2015) The future of immune checkpoint therapy. Science 348, 56-61.

Sen M, Koksai AC, Yuki K, Wang J and Springer TA (2018) Ligand- and cation-induced structural alterations of the leukocyte integrin LFA-1. Journal of Biological Chemistry 293, 6565-6577.

Sen M, Yuki K and Springer TA (2013) An internal ligand-bound, metastable state of a leukocyte integrin, $\alpha X\beta 2$. Journal of Cell Biology 203, 629-642.

Sharma P and Allison JP (2015) The future of immune checkpoint therapy. Science 348, 56-61.

Shattil SJ, Cunningham M and Hoxie JA (1987) Detection of activated platelets in whole-blood using activation-dependent monoclonal antibodies and flow-cytometry. Blood 70, 307-315.

Shattil SJ, Kashiwagi H and Pampori N (1998) Integrin signaling: the platelet paradigm. Blood 91, 2645-2657.

Shattil SJ and Newman PJ (2004) Integrins: dynamic scaffolds for adhesion and signaling in platelets. Blood 104, 1606-1615.

Shulman Z, Cohen SJ, Roediger B, Kalchenko V, Jain RSJ, Grabovsky V, Klein E, Shinder V, Stoler-Barak L, Feigelson SW, Nurmi SM, Goldstein I, Hartley O, Gahmberg CG, Etzioni A, Weninger W, Ben-Baruch A and Alon R (2012) Transendothelial migration of lymphocytes mediated by endothelial vesicle stores rather than by extracellular chemokine depots. Nature Immunology 13, 67-77.

Sirim P, Zeitlmann L, Kellersch B, Falk CS, Schendel DJ and Kolanus W (2001) Calcium signaling through the $\beta 2$ -cytoplasmic domain of LFA-1 requires intracellular elements of the T cell receptor complex. Journal of Biological Chemistry 276, 42945-42956.

Skinner JJ, Wang S, Lee J, Ong C, Sommese R, Sivaramakrishnan S, Koelmel W, Hirschbeck M, Schindelin H, Kisker C, Lorenz K, Sosnick TR and Rosner MR (2017) Conserved salt-bridge competition triggered by phosphorylation regulates the protein interactome. Proceedings of the National Academy of Sciences of the United States of America 114, 13453-13458.

Springer TA (1990) Adhesion receptors of the immune-system. Nature 346, 425-434.

Springer TA and Dustin ML (2012) Integrin inside-out signaling and the immunological synapse. Current Opinion in Cell Biology 24, 107-115.

Springer TA and Sen M (2016) Leukocyte integrin $\alpha L\beta 2$ headpiece structures: the αI domain, the pocket for the internal ligand, and concerted movements of its loops. Proceedings of the National Academy of Sciences of the United States of America 113, 2940-2945.

- Springer TA, Thompson WS, Miller LJ, Schmalstieg FC and Anderson DC (1984) Inherited deficiency of the MAC-1, LFA-1, P150,95 glycoprotein family and its molecular basis. *Journal of Experimental Medicine* 160, 1901-1918.
- Stanley P, Smith A, McDowall A, Nicol A, Zicha D and Hogg N (2008) Intermediate-affinity LFA-1 binds α -actinin-1 to control migration at the leading edge of the T cell. *EMBO Journal* 27, 62-75.
- Stossel TP, Condeelis J, Cooley L, Hartwig JH, Noegel A, Schleicher M and Shapiro SS (2001) Filamins as integrators of cell mechanics and signalling. *Nature Reviews of Molecular Cell Biology* 2, 138-145.
- Strazza M, Azoulay-Alfaguter I, Peled M, Smrcka AV, Skolnik EY, Srivastava S and Mor A (2017) PLC ϵ 1 regulates SDF-1 α -induced lymphocyte adhesion and migration to sites of inflammation. *Proceedings of the National Academy of Sciences of the United States of America* 114, 2693-2698.
- Stritt S, Wolf K, Lorenz V, Vogtle T, Gupta S, Bosl MR and Nieswandt B (2015) Rap1-GTP-interacting adaptor molecule (RIAM) is dispensable for platelet integrin activation and function in mice. *Blood* 125, 219-222.
- Strohmeyer N, Bharadwaj M, Costell M, Fässler R and Muller DJ (2017) Fibronectin-bound $\alpha 5 \beta 1$ integrins sense load and signal to reinforce adhesion in less than a second. *Nature Materials* 16, 1262-1272.
- Su W, Wynne J, Pinheiro EM, Strazza M, Mor A, Montenont E, Berger J, Paul DS, Bergmeier W, Gertler FB and Philips MR (2015) Rap1 and its effector RIAM are required for lymphocyte trafficking. *Blood* 126, 2695-2703.
- Sun Z, Costell M and Fässler R (2019) Integrin activation by talin, kindlin and mechanical forces. *Nature Cell Biology* 21, 25-31.
- Sun Z, Guo SS and Fässler R (2016) Integrin-mediated mechanotransduction. *Journal of Cell Biology* 215, 445-456.
- Svensson L, Howarth K, McDowall A, Patzak I, Evans R, Ussar S, Moser M, Metin A, Fried M, Tomlinson I and Hogg N (2009) Leukocyte adhesion deficiency-III is caused by mutations in KINDLIN3 affecting integrin activation. *Nature Medicine* 15, 306-312.
- Tadokoro S, Shattil SJ, Eto K, Tai V, Liddington RC, de Pereda JM, Ginsberg MH and Calderwood DA (2003) Talin binding to integrin β tails: a final common step in integrin activation. *Science* 302, 103-106. Takafuta T, Sacki M, Fujimoto TT, Fujimura K and Shapiro SS (2003) A new member of the LIM protein family binds to filamin B and localizes at stress fibers. *Journal of Biological Chemistry* 278, 12175-12181.
- Takada Y, Ye X and Simon S (2007) The integrins. *Genome Biology* 8, 215 doi:10.1186/gb-2007-8-5-215.
- Takafuta T, Sacki M, Fujimoto TT, Fujimura K and Shapiro SS (2003) A new member of the LIM protein family binds to filamin B and localizes at stress fibers. *Journal of Biological Chemistry* 278, 12175-12181.

- Takala H, Nurminen E, Nurmi SM, Aatonen M, Strandin T, Takatalo M, Kiema T, Gahmberg CG, Ylännä J and Fagerholm SC (2008) $\beta 2$ integrin phosphorylation on Thr758 acts as a molecular switch to regulate 14-3-3 and filamin binding. *Blood* 112, 1853-1862.
- Takafuta T, Sacki M, Fujimoto TT, Fujimura K and Shapiro SS (2003) A new member of the LIM protein family binds to filamin B and localizes at stress fibers. *Journal of Biological Chemistry* 278, 12175-12181.
- Takahashi M, Dillon TJ, Liu C, Kariya Y, Wang Z and Stork PJS (2013) PKA-dependent phosphorylation of Rap1 regulates its membrane localization and cell migration. *Journal of Biological Chemistry*
- Takeichi M (1977) Functional correlation between cell adhesion properties and some cell-surface proteins. *Journal of Cell Biology* 75, 464-474.
- Takeichi M (1991) Cadherin cell adhesion receptors as a morphogenetic regulator. *Science* 251, 1451-1455.
- Tan S-M, Robinson MK, Drbal K, van Kooyk Y, Shaw JM and Law SKA (2001) The N-terminal region and the mid-region complex of the integrin $\beta 2$ subunit. *Journal of Biological Chemistry* 276, 36370-36376.
- Tian L, Lappalainen J, Autero M, Hänninen S, Rauvala H and Gahmberg CG (2008) Shedded neuronal ICAM-5 suppresses T cell activation. *Blood* 111, 3615-3625.
- Tian L, Yoshihara Y, Mizuno T, Mori K and Gahmberg CG (1997) The neuronal glycoprotein, telencephalin, is a cellular ligand for the CD11a/CD18 leukocyte integrin. *Journal of Immunology* 158, 928-936.
- Tirupula KC, Ithychanda SS, Mohan ML, Prasad SVN, Qin J and Karnik SS (2015) G protein-coupled receptors directly bind filamin A with high affinity and promote filamin phosphorylation. *Biochemistry* 54, 6673-6683.
- Tng E, Tan S-M, Ranganathan S, Cheng M and Law SKA (2004) The integrin $\alpha L\beta 2$ hybrid domain serves as a link for the propagation of activation signal from its stalk regions to the I-like domain. *Journal of Biological Chemistry* 279, 54334-54339.
- Thomas ML (1989) The leukocyte common antigen family. *Annual Review of Immunology* 7, 339-369.
- Tohyama Y, Katagiri K, Pardi R, Lu C, Springer TA and Kinashi T (2003) The critical cytoplasmic regions of the $\alpha L/\beta 2$ integrin in Rap1-induced adhesion and migration. *Molecular Biology of the Cell* 14, 2570-2582.
- Trowbridge IS (1978) Interspecies spleen-myeloma hybrid producing monoclonal antibodies against mouse lymphocyte surface glycoprotein, T200. *Journal of Experimental Medicine* 148, 313-323.
- Tseng H-y, Samarelli AV, Kammerer P, Scholze S, Ziegler T, Immier R, Zent R, Sperandio M, Sanders CR, Fässler R and Böttcher RT (2018) LCP1 preferentially binds clasped $\alpha M\beta 2$ integrin and attenuates leukocyte adhesion under flow. *Journal of Cell Science* 131, doi:10.1242/jcs.218214.

- Tu Y, Wu S, Shi X, Chen K and Wu C (2003) Migfilin and Mig-2 link focal adhesions to filamin and the actin cytoskeleton and function in cell shape modulation. *Cell* 113, 37-47.
- Uotila LM, Aatonen M and Gahmberg CG (2013) Integrin CD11c/CD18 α -chain phosphorylation is functionally important. *Journal of Biological Chemistry* 288, 33494-33499.
- Uotila LM, Jahan F, Hinojosa LS, Melandri E, Grönholm M and Gahmberg CG (2014) Specific phosphorylations transmit signals from leukocyte β 2 to β 1 integrins and regulate adhesion. *Journal of Biological Chemistry* 289, 32230-32242.
- Ussar S, Wang HV, Linder S, Fässler R and Moser M (2006) The Kindlins: subcellular localization and expression during murine development. *Experimental Cell Research* 312, 3142-3151.
- Valmu L, Autero M, Siljander P, Patarroyo M and Gahmberg CG (1991) Phosphorylation of the β -subunit of CD11/CD18 integrins by protein kinase C correlates with leukocyte adhesion. *European Journal of Immunology* 21, 2857-2862.
- Valmu L, Fagerholm S, Suila H and Gahmberg CG (1999) The cytoskeletal association of CD11/CD18 leukocyte integrins in phorbol ester activated T cells correlates with CD18 phosphorylation. *European Journal of Immunology* 29, 2107-2118.
- Valmu L and Gahmberg CG (1995) Treatment with okadaic acid reveals strong threonine phosphorylation of CD18 after activation of CD11/CD18 leukocyte integrin with phorbol esters or CD3 antibodies. *Journal of Immunology* 155, 1175-1183.
- Vestweber D (2015) How leukocytes cross the vascular endothelium. *Nature Reviews of Immunology* 15, 692-704.
- Vinogradova O, Velyvis A, Velyviene A, Hu B, Haas TA, Plow EF and Qin J (2002) A structural mechanism of integrin α IIb β 3 "inside-out" activation as regulated by its cytoplasmic face. *Cell* 110, 587-597.
- Waldt N, Seifert A, Demiray YE, Devroe E, Turk BE, Reichardt P, Mix C, Reinhold A, Freund C, Muller AJ, Schraven B, Stork O and Kliche S (2018) Filamin phosphorylation at serine 2152 by the serine/threonine kinase Ndr2 controls TCR-induced LFA-1 activation in T cells. *Frontiers in Immunology* 9, Article 2852. doi: 10.3389/fimmu.2018.02852.
- Walz G, Aruffo A, Kolanus W, Bevilacqua M and Seed B (1990) Recognition by ELAM-1 of the sialyl-LeX determinant on myeloid and tumor cells. *Science* 250, 1132-1135.
- Wei X, Wang X, Zhan J, Chen Y, Fang W, Zhang L and Zhang H (2017) Smurf1 inhibits integrin activation by controlling kindlin-2 ubiquitination and degradation. *Journal of Cell Biology* doi.org/10.1083/jcb.201609073.
- Weitz-Schmidt G, Welzenbach K, Brinkmann V, Kamata T, Kallen J, Bruns C, Cottens S, Takada Y and Hommel U (2001) Statins selectively inhibit leukocyte function antigen-1 by binding to a
- Wickham TJ, Mathias P, Cheresch DA and Nemerow GR (1993) Integrins α v β 3 and α v β 5 promote adenovirus internalization but not virus attachment. *Cell* 73, 309-319.

- Wonerow P, Pearce AC, Vaux DJ and Watson SP (2003) A critical role for phospholipase C gamma2 in alphaIIb beta3-mediated platelet spreading. *Journal of Biological Chemistry* 278, 37520-37529.
- Wu Y, Span LM, Nygren P, Zhu H, Moore DT, Cheng H, Roder H, DeGrado WF and Bennett JS (2015) The tyrosine kinase c-Src specifically binds to the active integrin α IIb β 3 to initiate outside-in signaling in platelets. *Journal of Biological Chemistry* 290, 15825-15834.
- Xie C, Zhu J, Chen X, Mi L, Nishida N and Springer TA (2010) Structure of an integrin with an α I domain, complement receptor type 4. *EMBO Journal* 29, 666-679.
- Xiong J-P, Mahalingham B, Alonso JL, Borrelli LA, Rui X, Anand S, Hyman TR, Rysiok T, Muller-Pompalla D, Goodman SL and Arnaout MA (2009) Crystal structure of the complete integrin α V β 3 ectodomain plus an α / β transmembrane fragment. *Journal of Cell Biology* 186, 589-600.
- Xiong J-P, Stehle T, Diefenbach B, Zhang R, Dunker R, Scott DI, Joachimiak A, Goodman SL and Arnaout MA (2001) Crystal structure of the extracellular segment of integrin α V β 3. *Science* 294, 339-345.
- Yamada KM (1983) Cell surface interactions with extracellular materials. *Annual Review of Biochemistry* 52, 761-799.
- Yamada KM and Olden K (1978) Fibronectins - adhesive glycoproteins of cell – surface and blood. *Nature* 275, 179-184
- Ye F, Petrich BG, Anekal P, Lefort CT, Kasirer-Friede A, Shattil SJ, Ruppert R, Moser M, Fässler R and Ginsberg MH (2013) The mechanism of kindlin-mediated activation of integrin α IIb β 3. *Current Biology* 23, 2288-2295.
- Ye F, Petrich BG, Anekal P, Lefort CT, Kasirer-Friede A, Shattil SJ, Ruppert R, Moser M, Fässler R and Ginsberg MH (2013a) The mechanism of kindlin-mediated activation of integrin α IIb β 3. *Current Biology* 23, 2288-2295.
- Ye Q, Feng Y, Yin Y, Faucher F, Currie MA, Rahman MN, Jin J, Li S, Wei Q and Jia Z ((2013b) Structural basis of calcineurin activation by calmodulin. *Cell Signaling* 25, 2661-2667.
- Ylännä J, Huuskonen J, O'Toole TE, Ginsberg MH, Virtanen I and Gahmberg CG (1995) Mutation of the cytoplasmic domain of the integrin β 3 subunit: differential effects on cell spreading, recruitment to adhesion plaques, endocytosis and phagocytosis. *Journal of Biological Chemistry* 270, 9550-9557.
- Yoshihara Y, Oka S, Nemoto Y, Watanabe Y, Nagata S, Kagamiyama H and Mori K (1994) An ICAM-related neuron glycoprotein, telencephalin, with brain segment-specific expression. *Neuron* 12, 541-553.
- Zaidel-Bar R, Itzkovitz S, Ma'ayan A, Iyengar R and Geiger B (2007) Functional atlas of the integrin adhesome. *Nature Cell Biology* 9, 858-868.
- Zarbock A, Ley K, McEver RP and Hidalgo A (2011) Leukocyte ligands for endothelial selectins: specialized glycoconjugates that mediate rolling and signaling under flow. *Blood* 118, 6743-6751.

Zhou Z-R, Gao H-C, Zhou C-J, Chang Y-G, Hong J, Song A-X, Lin D-H and Hu H-Y (2008) Differential ubiquitin binding of the UBA domains from human c-Cbl and Cbl-b: NMR structural and biochemical insights. *Protein Science* 17, 1805-1814.

Zhu L, Yang J, Bromberger T, Holly A, Lu F, Liu H, Sun K, Klapproth S, Hirbawi J, Byzova TV, Plow EF, Moser M and Qin J (2017) Structure of Rap1b bound to talin reveals a pathway for triggering integrin activation. *Nature Communications* 8:1744 doi:10.1038/s41467-017-01822-8.

Abbreviations

LFA-1, α L β 2, CD11a/CD18 (Leukocyte function associated antigen-1); (Mac-1, α M β 2, CD11b/CD18 (Macrophage-1 antigen); α X β 2, CD11c/CD18 (Integrin α X antigen); α D β 2, CD11d/CD18 (Integrin α D); ICAM (Intercellular adhesion molecule); VCAM-1 (Vascular cell adhesion molecule-1); VLA-4, α 4 β 1 (Very late antigen-4); LAD (Leukocyte adhesion deficiency disease); MIDAS (Metal ion –dependent adhesion site)); RIAM (Rap1-GTP—interacting adaptor molecule); RAPL (Rap1 ligand), I domain (Integrin ligand binding site); TCR (T cell receptor); ITAM (Immune- receptor-tyrosine-based-activation-motif); Csk (C-Src tyrosine kinase); PLC (phospholipase); PKC (Protein kinase C); DAG (Diacyl glycerol); FAK (Focal adhesion kinase); RACK1 (Receptor of activated protein kinase C 1); Dok1 (Docking protein 1); SHARPIN (SHANK associated RH domain interactor); Del-1 (Developmental endothelial locus-1); Crk (Adaptor molecule Crk); GEF (Guanine nucleotide exchange factor).

Figure legends

Fig. 1. A schematic figure of the human integrin family. The integrins are grouped according to their β -chains. Note that some β -chains can associate with more than one α -chain.

Fig. 2. Domain structures of I-domain integrins. In (A) the domains are depicted in a linear fashion. An inactive integrin is shown in (B) with the NH₂-terminal binding site turned towards the lipid membrane. In (C) the integrin is extended with the external ligand binding site closed, and the cytoplasmic tails clasped. In (D) the integrin is fully active with the ligand binding site open and the cytoplasmic tails apart. The KIM127 epitope is available for antibody binding in (C) and (D). The mab24 binds only to the active integrin in (D).

Fig. 3. Electron micrographs of negative stained α X β 2 integrin. In A is shown the inactive integrin with the ligand binding site closed. In B1 and B2 are shown extended, but not fully active integrins. B3 shows the active, open integrin. The figure was reprinted from Chen et al. (2010) with permission from the Proceedings of the US National Academy of Sciences.

Fig. 4. A schematic figure of the LFA-1 integrin with functionally important monoclonal antibody binding sites marked. See also Fig. 2.

Fig. 5. Human integrin cytoplasmic tail sequences. Known phosphorylation sites are marked in red.

Fig. 6. Protein binding sites to the cytoplasmic tails of LFA-1. The phosphorylation sites are marked in red and the conserved α -tail GFKKR sequence in blue.

Fig. 7. LFA-1 activation through the TCR. Activation takes place by activation of the CD3 associated molecules through the Lck tyrosine kinase, followed by ZAP70 kinase activation, phosphorylation of LAT, and activation of PLC γ 1. This results in the generation of DAG and IP3. DAG activates PKC enzymes, which phosphorylate T758 in the β 2 chain. IP3 activates Ca $^{2+}$ release. Calcineurin is recruited to the TCR complex, where it may release the inhibitory phosphate on S59 in Lck, activate the kinase. In β 1 and β 3 integrins, calreticulin is known to bind to the α -chains, but this has not been shown for β 2.

Fig. 8. The immunological synapse. The synapse is formed between T cells and antigen presenting cells. The specific interaction is due to TCR and MHC-peptide recognition in the center. The synapse is strengthened by LFA-1/ICAM-1 interaction in the peripheral part. The CD45 tyrosine phosphatase must be close to Lck to be able to regulate its activity.

Fig. 9. Activation of LFA-1 through chemokines. After binding of chemokines to the receptors, PLCs become activated and generate DAG and IP3. DAG activates PKCs, which phosphorylate T758 in β 2. On the other hand, the Cal-DAG G protein exchange factor activates Rap1. Rap1-GTP binds RapL, which binds to the α L-chain. In addition RIAM binds to talin and the β 2-chain.

Fig. 10. A simplified view of α IIb β 3 and LFA-1 activation by outside-in signaling. Ligands such as fibrinogen bind to the outside and activate Src family kinases, which in turn activate focal adhesion kinase and Syk by tyrosine phosphorylation. In LFA-1, the binding of external ligands induce the phosphorylation of T758 in the β -chain followed by binding of 14-3-3 proteins, Tiam1 and activation of Rac1. It remains unclear when talin and kindlin interact with the integrin, but results support the proposal that it is important in the final activation of integrins.

Fig. 11. Integrin crosstalk. Activation through the TCR or chemokine receptors activate LFA-1. This may also happen through antibody binding to LFA-1 or by outside-in activation by ICAMs. Then signaling through the β 2 T758 route results in dephosphorylation of the β 1 chain in VLA-4 resulting in its inactivation.

Fig. 12. An integrated view of LFA-1 activation. In the resting integrin, the α -chain is phosphorylated, and filamin and Dok1 are associated with the β -chain. Upon activation, T758 in the β -chain is phosphorylated resulting in the release of filamin, whereas kindlin-3, Dok1 and 14-3-3 form a complex bound to the β -chain. The phosphorylations on both the α - and β -chains open the cytoplasmic domain, and α -actinin binds to the β -chain, and connect the integrin to the actin cytoskeleton. Talin remains associated with the β 2-chain, and may induce a transmembrane allosteric activation of the LFA-1 external ligand binding site. Tiam1 activates Rac1 to the GTP form, inducing reorganization of the actin cytoskeleton and facilitating binding to the integrin. The 14-3-3, kindlin-3 and α -actinin proteins form dimers, enabling clustering of the integrins, which results in increased avidity for the ligands such as ICAM-1.