<https://helda.helsinki.fi>

Exploring individual and population eco-evolutionary feedbacks under the coupled effects of fishing and predation

Jusufovski, Dunja

2020-11

Jusufovski , D & Kuparinen , A 2020 , ' Exploring individual and population eco-evolutionary feedbacks under the coupled effects of fishing and predation ' , Fisheries Research , vol. 231 , 105713 . <https://doi.org/10.1016/j.fishres.2020.105713>

<http://hdl.handle.net/10138/324662>

<https://doi.org/10.1016/j.fishres.2020.105713>

submittedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **Exploring individual and population eco-evolutionary feedbacks under the coupled effects of**
2 **fishing and predation**

3
4
5 Dunja Jusufovski^{a*,1} and Anna Kuparinen^b
6

7 ^aOrganismal and Evolutionary Biology Research Program, University of Helsinki, P.O. Box 65,
8 00014 Helsinki, Finland; dunja.jusufovski@helsinki.fi

9 ^bDepartment of Biological and Environmental Sciences, University of Jyväskylä, P.O. Box 35, 40014
10 Jyväskylä, Finland; anna.k.kuparinen@jyu.fi

11 * Corresponding author
12
13
14
15
16
17
18

19 Running head: Eco-evolutionary feedbacks under fishing and predation
20
21
22
23

Abbreviations:

FR_{II} – Holling's functional response Type II

FR_{III} – Holling's functional response Type III

¹ Corresponding author's present address and mobile phone number: Ulica kneza Domagoja 1, 31220 Višnjevac, Croatia; +385(0)915609853

24 **Abstract**

25 Intensive fishing that selects for large and old individuals can have pervasive effects on traits directly
26 associated with the fecundity and survival of the target species. The observed reduction in fish body
27 size can result in earlier sexual maturity at a smaller body size, leading to a lower individual
28 reproductive output and population productivity in the long term. In addition, increased predation can
29 induce similar responses in age and size at maturity due to the release of intraspecific competition
30 and the lower population density. Thus, the combined impact of fisheries and predation is more
31 difficult to predict due to their competition for fish, ultimately limiting and directing the prey size
32 selectivity of each competitor. This can lead to various responses of fish on individual and population
33 levels, depending on how the impacts of fisheries and predation affect each other. To explore this
34 topic, we used an individual-based model to investigate the responses of the European hake
35 (*Merluccius merluccius*) to the coupled pressures of the prevailing fishing regime and predation by
36 bottlenose dolphins (*Tursiops truncatus*) as a study system. By simulating the growth trajectories of
37 hake with consideration to its pronounced sexual size dimorphism (SSD), we applied Holling's Type
38 II and III functional responses (FRII and FRIII, respectively) and the Rosenzweig-MacArthur
39 predator model to describe prey–predator dynamics. Our results demonstrate how the intensity,
40 duration, and prey sizes selected by predation and fisheries determine the potential for evolutionary
41 changes in hake life-history traits. Additionally, pronounced SSD in hake as well as the predation
42 type play a pivotal role in hake resilience and recovery following a period of overexploitation.
43 Different types of predation reveal the discrepancy in the intensity of competition between predators
44 and fisheries for fish, as well as predator survival, which is relevant for sustainable fisheries
45 management and effective predator conservation strategies.

46

47 Keywords: Size-selectivity; Fisheries-induced evolution; *Merluccius*; Prey–predator dynamics;
48 Functional responses

1. Introduction

Fisheries-induced alterations in fish morphology, behavior, and development have been a focus of research and discussion in fisheries science and management for the past several decades (e.g., Hendry et al., 2011; Palkovacs et al., 2012; Heino et al., 2015). A number of researchers have emphasized the negative, long-term, and potentially irreversible effects of fisheries-induced evolution (FIE) on fish stock resilience and recovery (Swain et al., 2007; Enberg et al., 2009; Neubauer et al., 2013), as well as sustainable fisheries yields (Zhou et al., 2010; Garcia et al., 2012). In particular, the size selectivity of fisheries and removal of larger and older individuals poses a persistent threat due to its pervasive, damaging effects on individual fecundity and survival (Law, 2000; Palkovacs et al., 2012; Belgrano and Fowler, 2013). Continued high harvest rates and fishing intensity, selecting for life-history traits with specific heritability, represent the main conditions that can induce an evolutionary shift in fish species (Hutchings and Kuparinen, 2019). However, difficulties in distinguishing phenotypic plasticity from evolution (reviewed in Hendry, 2016) and discerning the direction of ecology-to-evolution or evolution-to-ecology processes are the main obstacles to proving FIE in harvested fish stocks (see Hendry et al., 2011; Heino et al., 2015).

In addition to FIE, predation and other selective pressures of environmental and ecological origin occurring in marine ecosystems can potentially induce similar responses in the direction or magnitude of targeted phenotypic traits (see Reznick et al., 1990; Philips et al., 2010; Crozier and Hutchings, 2014; Ohlberger, 2013; Waples and Audzijonyte, 2016). Predation is, among others, a size-selective source of natural mortality, removing individuals according to the prey size selectivity of the predator. Moreover, the size-dependent mortality caused by predation increases with a decrease in fish body size, especially in those predators limited by gape size (Pettorelli et al., 2011). Similarly to fisheries, predation directly affects the prey's investment to reproduction (DeLong and Luhring, 2018) and its reproductive success or resilience (Lowerre-Barbieri et al., 2017) to other disturbances. This can ultimately lead to shifts in the timing of their sexual

74 maturation and alter their reproductive strategy (e.g., behavioral patterns linked to reproduction)
75 due to the removal of sexually mature individuals. While such predation-induced changes in prey
76 have been detected as shifts in size and age at maturity observed under experimental conditions
77 (Beckerman et al., 2010), changes in predators induced by the changes in prey have also been
78 observed. For instance, Shuter et al. (2015) predicted that larger size ratios of lake trout (*Salvelinus*
79 *namaycush*, predator) to cisco (*Coregonus artedii*, prey) lead to earlier maturity at a smaller size in
80 the predator species. Thus, prey characteristics as well as their adaptations to predation affect predator
81 development and abundance (Yamamichi and Miner, 2015; Fryxell et al., 2019).

82 Considering that fishing can erode size-dependent interactions, change community compositions,
83 and weaken the structural integrity of ecosystems (Palkovacs et al., 2012; Bell et al., 2018),
84 evolutionary changes in individual fish traits can reinforce these shifts and limit or reduce
85 intraspecific genotypic diversity, which ultimately diminishes fishing yields (Law and Grey, 1989;
86 Garcia et al., 2012; Belgrano and Fowler, 2013). As a result, concern has been raised regarding the
87 management of evolving fish stocks and their implications for ecosystem functioning and
88 sustainable fisheries yields. Recent managerial advice for fish populations has included the
89 application of balanced harvesting (Law and Plank, 2018), as well as the active monitoring and
90 accountability of individual trait responses to fishing and other selective pressures (Conover and
91 Munch, 2002; Palkovacs et al., 2018). Furthermore, particular attention should be given to
92 exploitation strategies for late and early maturing species (Ernande et al., 2004), as well as to fish
93 stocks already demographically and genetically altered, which require careful implementation of
94 different fishing moratoria (de Roos et al., 2006). However, these proposed measures are
95 accompanied by various operational and logistical difficulties, and they only address fisheries-
96 induced alterations in individual and population parameters, other threats such as climate change
97 notwithstanding (Fraser, 2013; Palkovacs et al., 2018).

98 In harvested ecosystems, fisheries and natural predators often compete for the same target fish in
99 areas of temporal and spatial resource overlap. To account for this, the majority of fish stock
100 assessments include predation as part of a species-specific fixed value for natural mortality that is
101 equally applied across all age and size classes of fish (Gislason et al., 2010). However, the risk of
102 being consumed by a predator decreases as a function of increasing body size, as bigger prey fish
103 become more elusive due to their size (Abrams and Ginzburg, 2000), as well as their improved
104 swimming and maneuvering skills (Scharf et al., 2000; DeLong and Luhring, 2018). From the
105 predator's viewpoint, their prey size selectivity will change and adjust according to prey availability
106 and the predator's ontogenetic development (Griffiths, 1975). While humans are considered as
107 predators, their prey selectivity and preference are not governed by the sole need to feed for
108 survival. As Darimont et al. (2015) concluded, humans are not limited in their exploitation rates or
109 techniques as natural predators are; thus, humans are considered as "super predators". As a
110 consequence of intensive harvesting, humans can alter the predation risk through direct impacts on
111 predators as well as indirectly by affecting the physical environment (see Madin et al., 2016).
112 Further negative impacts can occur through density-mediated effects (Lankau and Strauss, 2011)
113 where the removal of large predators has direct implications for the strength of trophic cascades as
114 well as food web stability (DeLong et al., 2015).

115 The European hake (*Merluccius merluccius*, Linnaeus, 1758) is a gadoid bottom-dweller
116 inhabiting the seas along the NE and central E Atlantic, the NW African coast, and the
117 Mediterranean and Black Seas, with a decreasing presence from west to east (Fig. 1; Sion et al.,
118 2019). Adult hake have a mainly piscivorous diet and exhibit strong sexual size dimorphism (SSD),
119 with males maturing earlier and having a shorter lifespan than females (Cerviño, 2014; Murua,
120 2010). Furthermore, the female body size in hake is not just positively correlated with the number
121 of produced eggs (i.e., individual fecundity), but also with the quality of the produced eggs, which
122 ensures a higher chances of offspring survival (Mehault et al., 2010). Significant fluctuations in size

123 at maturity have been observed in hake populations of the Bay of Biscay and Galician waters
124 (Domínguez-Petit et al., 2008). Although these changes have been attributed to phenotypic
125 plasticity of various strengths (see Hidalgo et al., 2014), these populations have experienced very
126 high harvest rates since the 1980s and continue to do so (ICES, 2019). Additionally, hake are
127 preyed upon by the bottlenose dolphin (*Tursiops truncatus*, Montagu, 1821), which has been
128 confirmed by dietary studies on the Irish, Bay of Biscay, and Galician, as well as the W
129 Mediterranean bottlenose dolphin populations (Blanco et al., 2001; Fernández et al., 2011;
130 Hernandez-Milian et al., 2015; Santos et al., 2007, 2014). These opportunistic predators often
131 interact with fishing activities (i.e., operational interactions; Bearzi et al., 2008), following trawlers
132 or depredating on coastal nets (e.g., López, 2006; Bearzi et al., 2010; Goetz et al., 2015). While
133 competing with hake fisheries through direct fish uptake (i.e., ecological interactions), bottlenose
134 dolphins can notably contribute to the total mortality of hake (Santos et al., 2014). Nevertheless,
135 only a few studies have attempted to quantify the fish biomass removed by these dolphin species.
136 For instance, Bearzi et al. (2010) found the annual biomass of fish removed by fisheries to be about
137 38 times greater than the fish uptake by the bottlenose dolphin population in W Greek waters.
138 Furthermore, Santos et al. (2014) estimated the predation mortalities of four resident dolphin
139 species of the NE Atlantic area preying upon European hake, with the bottlenose dolphin population
140 having the highest uptake of hake (82% of the removed hake biomass).

141 Since little is known about how fishing and predation in combination affect the ecology and
142 evolution of individual traits and, in return, alter predator abundance and fishing yields, here we
143 investigate the responses of individual traits and population parameters of the European hake under
144 the prevailing fishing selectivity and predation by the bottlenose dolphin. Gårdmark et al. (2003)
145 theoretically explored the evolutionary responses of the age at first reproduction to density-
146 independent and density-dependent predation and harvesting with regards to the number of species
147 involved and predation type (i.e., Holling's Type I and II). Other studies investigating the effects of

148 fishing and predation have explored these from the perspectives of fisheries assessment and eco-
149 evolutionary feedbacks independently, using approaches that range from individual-based to
150 ecosystem models (e.g., Melián et al., 2011; Allhoff et al., 2015; Kindsvater and Palkovacs, 2017;
151 Wood et al., 2018; Govaert et al., 2019). In this paper, we use an individual-based model to
152 examine how the competitive interactions of natural predators and fisheries affect individual fish
153 traits as well as population resilience and recovery. Similarly to ecosystem resilience (Holling,
154 1973), we observe the population resilience as the capacity of the population dynamics and
155 structure to withstand and absorb a disturbance whilst maintaining its main ecological and structural
156 characteristics, while recovery is reflected in the speed of recovery of observed parameters after the
157 disturbance has ceased. While the methodological approach allows us to account for the ecological
158 characteristics of the prey and the co-evolved prey–predator dynamics, we firstly focus our
159 investigation on the potential for FIE under the coupled effects of the size-selective pressures. Then
160 we observe the impact of predation on hake population and fisheries. Lastly, we attempt to
161 understand the dynamics between the predator and fisheries with consideration to the eco-
162 evolutionary feedbacks and disruptive fishing regime.

163

164 **2. Material and methods**

165

166 **2.1 Individual-based model**

167 In the attempt to address the current research questions, we used an individual based model (IBM),
168 which is an eco-genetic model encompassing quantitative genetics, evolutionary processes, and the
169 ecological characteristics of the fish species. The IBM approach provides an insight into fish
170 individual variability of life history traits, whilst aiding to overcome the lack of information about
171 the exact gene responsible for shaping the response to size-selective removals in hake growth.
172 Furthermore, it also allows us to observe an evolutionary change in fish life history traits in “fast
173 forward mode”. To identify the coupled impact of fishing and predation, we expanded an IBM

174 initially developed for the Atlantic cod (*Gadus morhua*; see Kuparinen et al. 2012, 2014; Kuparinen
175 and Hutchings, 2012 for details) and later parameterized for the European hake (see Jusufovski and
176 Kuparinen, 2014). While the baseline model and parameterization for hake are described
177 extensively elsewhere (see Jusufovski and Kuparinen, 2014), in the following text we provide a
178 brief overview of the main model characteristics with a focus on the novel elements: (i) size-
179 selective mortality caused by predation and (ii) SSD in hake accounted for through sex-specific
180 differences in asymptotic lengths with an individual-based prediction of the length at which a fish
181 becomes sexually mature.

182

183 **2.2 Data collection and model parameterization**

184

185 **2.2.1 Hake growth and ecological characteristics**

186 The model simulates fish growth trajectories described by the von Bertalanffy (1938) equation,
187 which calculates an individual's length at age or time step t (L_t) as: $L_t = L_\infty - (L_\infty - L_0) e^{-Kt}$. L_∞ is the
188 asymptotic length of an individual, K is the individual intrinsic growth rate (i.e., the rate at which an
189 individual approaches its L_∞), and L_0 is the fish body length at $t = 0$. L_0 is the same value for all
190 individuals and calculated from obtained data, while the length–weight relationship applied
191 corresponds to values used for the southern hake stock (ICES, 2019). We attained the final model of
192 the growth equation through the log-transformation of K and linear regression as: $\log(K) = -0.511 -$
193 $0.0041L_\infty$. For this purpose, we used hake growth parameters through the introduction of 64
194 unsexed data points, available as published data from Mediterranean study cases (see Jusufovski
195 and Kuparinen, 2014). Information on the biological characteristics and fishing regime of European
196 hake reflects the current state of the southern hake stock of the Bay of Biscay and Iberian waters
197 ICES ecoregion (ICES, 2019). Overall, we did not focus on geographical or stock-specific

198 characteristics, but rather prioritized reliability and quality in the choice of data and information that
199 is representative for the fish species in general.

200 Assuming that quantitative traits are defined by a large number of loci with a small additive
201 effect (Roff, 2002), we describe the genotype of each individual with 20 loci, assigning a value of 1
202 or 0 to each locus. Therefore, summing the values for the loci gives the genotypic value of an
203 individual, ranging from 0 (minimum value) up to 20 (the maximum value). The genetic diversity
204 then corresponds to the range of genotypic values observed with respect to the set conditions and
205 parameters (see further explanation of the tested scenarios in the section *Simulation design*).
206 Through sexual reproduction, parental genotypic information is transferred to the offspring in
207 accordance with the basic Mendelian laws of inheritance. The phenotype is then based on the
208 individual's genotype, to which phenotypic variation (i.e., normally distributed random noise with
209 mean zero) is added to account for phenotypic plasticity, whilst remaining within the realistic
210 values of heritability (0.2–0.3 for fish species; Mousseau and Roff, 1987). After the genotype is
211 transferred to the phenotype (i.e., L_∞), K is then predicted through the strong negative correlation
212 between K and L_∞ . The maximum hake age is set according to Murua (2010), so that all individuals
213 above 12 years of age have a 100% probability of dying.

214 Each individual goes through basic ecological processes depicted in Figure 2. To account for
215 SSD among hake males and females, L_∞ is increased by 41% for female hake, a value representing
216 the discrepancy between the female and male L_∞ and calculated using the RL ratio = L_{mat} / L_∞ . As
217 male and female hake sexually mature at different lengths or L_{mat} (i.e., length at maturity), L_∞ was
218 calculated separately for each sex, applying the average value of RL (0.51) obtained from the hake
219 dataset. The male and female L_{mat} applied for the calculation of the percentage difference in L_∞ was
220 32 and 45 cm, respectively (Cerviño, 2014). Apart from the average RL value used as an initial
221 individual value in all simulations (see Table 1), with each time step and for each individual, the RL
222 is attained based on the individual's L_{mat} and L_∞ values. During each reproduction event, sexually

223 mature male and female hake are randomly selected and mated, with the sex of the offspring being
224 assigned by a Bernoulli trial ($p = 0.5$). Egg production depends directly on female weight, which is
225 derived from the length–weight relationship, and the average recruit-per-spawner value attained
226 through available information on recruitment and the spawning stock biomass (Jusufovski and
227 Kuparinen, 2014). Juvenile survival is determined by a baseline juvenile mortality rate with a
228 probability of survival set to 10%, which also takes into account the known cannibalism in hake
229 (Murua, 2010). Each juvenile is then assigned its own density-dependent growth (for further details,
230 see Jusufovski and Kuparinen, 2014).

231 Other ecological processes included in the model are density-dependent growth and three
232 sources of natural mortality (Fig. 2). The density-dependent growth of each individual diminishes
233 the growth by 50% each time the fish population exceeds 90% of its carrying capacity (see
234 Jusufovski and Kuparinen, 2014 for a detailed description and logistic equation). Instantaneous
235 natural mortality or M (for individuals older than two years) and mortality due to the cost of
236 reproduction or M_{rc} (only for sexually mature individuals) are estimated through parameterization
237 and both are applied equally to male and female hake (Table 1). The fate of each individual is then
238 decided using a Bernoulli trial at each time step.

239 Fishing is described through a logistic selectivity curve with L_{50} (the length at which 50% of the
240 population is retained) set at 26.4 cm based on the prevailing fishing regime (ICES, 2019; Table 1).
241 Only individuals older than three are fished, since at this age, hake have reached or are well over
242 their permitted minimum landing size (MLS).

243

244 **2.2.2 Introduction of predation**

245 The predator population is calculated on a population level using a modification of the
246 Rosenzweig–MacArthur prey–predator model (Rosenzweig and MacArthur, 1963). Depending on

the Holling's functional response (Holling, 1959) used in the prey equations, the predator equations yield the value of the predator density (p) at each time step t as:

$$p_t = p_i e^{e_p a_p n_t / (1 + h_p a_p n_t) - d_p} \quad (1)$$

in the case of FRII, while in FRIII, p_t depends on:

$$p_t = p_i e^{(e_p a_p n_t)^2 / (1 + (h_p a_p n_t)^2) - d_p} \quad (2)$$

where p_i is the predator's initial density set for the first time step (see Table 1), after which it becomes the value of p_{t-1} for all subsequent time steps. p_t depends on n_t or the prey population density attained as a quotient of the number of prey at time t (N_t) and the prey's carrying capacity (CC) at each time step. e_p , or the predator's assimilation efficiency, the attack rate (a_p), and handling time (h_p) are fixed values estimated through parameterization. In turn, the quantity of fish eaten or, more precisely, the probability of an individual being eaten (p_{eat}) is introduced at an individual level and given as:

$$p_{eat} = a_p p_t / (1 + h_p a_p n_t) \quad (3)$$

when FRII is applied and

$$p_{eat} = a_p p_t n_t^{(2-1)} / (1 + (h_p a_p n_t)^2) \quad (4)$$

in the case of FRIII. p_t in both equations denotes the predator population density calculated by equations (1) and (2) for FRII and FRIII, respectively. The functional responses are used here as the probability of a prey fish being eaten in order to account for any uncertainty brought about by intra- and interspecific competition of the predator.

Predation is applied across the hake size range from 21 to 51 cm, set according to the average size of hake (36 cm) and the hake size range consumed by the bottlenose dolphin populations of the Ibero-Atlantic waters (Saavedra, 2017). The predator parameters a_p , h_p , and d_p were estimated (Table 1) and compared to species-specific information on the generation time and dietary requirements from published research on wild and captive bottlenose dolphins (Kastelein et al., 2003; Lockyer, 2007; Santos et al., 2014; Saavedra, 2017; Spitz et al., 2018). Since the diet

272 composition of the bottlenose dolphin is catholic (Santos et al., 2007; 2014) and its feeding is
273 opportunistic, this is accounted for in the estimation of h_p . The applied value of the predator's
274 assimilation efficiency (e_p) is in accordance with the observed range of values for toothed whales,
275 i.e. 0.90–0.95 (Lockyer, 2007). Prey and predator population parameters (p_i , N_i and C) are
276 estimated through parameterizations (see Table 1) and set in such a manner that they allow for
277 stable prey–predator dynamics throughout the simulation.

278

279 2.2.3 Simulation design

280 In order to observe and detect the independent and coupled effects of fishing and predation, as
281 well as the eco-evolutionary feedbacks under these effects, we formed eight different scenarios
282 (Figure 3). Among the tested scenarios, those without predation and a sex-specific difference in L_∞
283 were simulated as reference scenarios. The scenarios used for the comparative approach are chosen
284 with the focus on the independent and joint impacts of predation and fisheries, rather than the effect
285 of the ecological characteristics of the prey. As the stabilization of prey–predator dynamics is time-
286 consuming (a total of 3000 time steps), to minimize the computation time, we recorded *a priori* the
287 adapted hake populations for each scenario. The adapted populations were recorded during the last
288 150 time steps of a 3000 time step simulation based on the confirmation of stabilization gained
289 through preliminary simulations (output and visual confirmation using graphs). Then, to test the
290 scenario, we used the recorded populations to run simulations for 600 time steps in total and
291 simulated 30 runs for each scenario in order to obtain the representative outcomes of the individual
292 and population parameters. During these simulations in all predation scenarios, predation was
293 applied continuously, while in all eight scenarios, fishing was introduced at the 400th time step. In
294 this manner, we could effectively distinguish the impact of predation, fishing, and evolutionary
295 processes. After the fishing period, we observed effects of the given pressures during the recovery

296 time until the end of the simulation. All simulations were performed in R Program v3.6.1 (R Core
297 Team, 2019).

298 During the simulation testing stage, the introduction of fishing with an intensity of 0.15 and the
299 duration of 100 time steps revealed that each scenario had a different threshold for the fishing
300 duration (Figs S1–S3 in the Supplementary material). In other words, prey and predator population
301 resilience as well as post-fishing recovery were affected by the total mortality applied to hake. The
302 scenarios without predation demonstrated the highest resilience, enduring 100 time steps of fishing,
303 followed by a successful recovery. The reference scenarios had a lower resilience of 75 time steps
304 for the fishing duration, while the FRII scenarios endured 35 time steps of fishing. The FRIII
305 predation scenarios were the least resilient and could only endure 30 time steps of fishing. There
306 was no difference between non-evolutionary and evolutionary scenarios in the maximum sustained
307 duration of the fishing period. For a balanced scenario design, we further applied exactly 30 time
308 steps of fishing to investigate the ecological and evolutionary feedbacks under the coupled effects
309 of fishing and predation. All outputs (incl. Supplement) represent average values of extracted
310 parameters pulled across all 30 simulations for each scenario. In the results, we observe different
311 aspects of predation, fisheries, evolution, and SSD on hake growth, maturity, genotypic diversity
312 and demographic parameters with the regards to their implication for hake recovery and resilience.
313 Firstly, we explore the coupled effects of predation and fishing with respect to FIE alone focusing
314 on the differences between the evolutionary and non-evolutionary scenarios. Secondly, we explore
315 the impact of predation on hake survival as well as how predation affects the fish size availability to
316 fisheries. Lastly, we observe the interactions between the predator and hake fisheries with respect to
317 evolutionary processes, and their implications for predator survival and fishing catches.

318

319 3. Results

320 3.1 Potential for FIE under the coupled effect of predation and fishing

321 Overall results between the evolutionary scenarios and their non-evolutionary versions (e.g., the
322 evolutionary and non-evolutionary scenarios without predation and with SSD, and so on) indicate
323 mainly minor discrepancies for the stage of initial introduction of fisheries (presented in the
324 Supplement). In the individual parameters for hake such as asymptotic length (L_{∞}) and intrinsic
325 growth rate (K), all non-evolutionary scenarios reach higher and lower scale values, respectively,
326 than the same scenarios with evolution (Figs S1a and c). Similar responses in L_{∞} and K are observed
327 in simulations where an identical duration of fishing was applied (Figs 4a and b). More precisely,
328 the impact of evolutionary processes in achieving the maximum length and its effect on growth rate
329 work in favor of a higher growth rate per time step (age). Furthermore, in simulations with different
330 and identical fishing durations, the evolutionary scenarios demonstrate continuously increasing and
331 decreasing trends in all evolutionary scenarios of L_{∞} and K , respectively (Figs S1a, c and Figs 4a,
332 b). Compared to non-evolutionary scenarios, trends observed in the evolutionary scenarios of L_{∞}
333 and K indicate that the ecological characteristics of hake (SSD, lifespan, and reproduction) in
334 combination with evolutionary processes notably delay the stabilization of these individual
335 parameters, despite the hake populations applied for each scenario being fully adapted to the
336 respective scenario settings.

337 The FRIII scenarios demonstrate differences with evolution absent and present in age at maturity
338 (Fig. 5a), affecting the speed of recovery. In the absence of evolution and with FRIII predation
339 present, hake displays a slower recovery to the respective pre-fishing level, while evolutionary
340 processes provide quicker recovery after fishing cessation. In addition, in the absence of evolution,
341 hake displays a tendency for maturity at a younger age in the FRIII scenarios, although no similar
342 effects of evolutionary processes on size at maturity (Fig. 5b) are observed under FRIII predation.
343 In contrast, with only 30 time steps for the fishing period, size at maturity drops slightly (by $\approx 1\%$)

344 in the FRII scenarios, while under FRII predation, a similar reduction in age at maturity gradually
345 levels out at its pre-fishing level by the end of the simulation time. However, longer fishing periods
346 applied during the introduction of fishing induce an increase and decrease in the age and size at
347 maturity, respectively, in the FRII scenario, whereas during the reference and no-predation
348 scenarios (Figs S1b and d), age and size at maturity do not return to their pre-fishing values within
349 the 170 time steps of the recovery period. The reference scenarios, as well as the scenarios without
350 predation, endure the longest fishing periods (75 and 100 time steps in the reference and no-
351 predation scenario, respectively); this indicates that longer fishing periods have a higher potential to
352 induce persistent phenotypic changes (e.g., a long-term decrease or increase in the age and size at
353 maturity) when hake SSD is unaccounted for.

354 The observed cumulative number of juveniles compared to the present genotypes suggests
355 different genotypic range values among all scenarios (Figs S2 and 5c). Regardless of the applied
356 fishing duration, a wider genotypic range is present in all evolutionary scenarios, as opposed to their
357 respective non-evolutionary versions, which indicates a lower genotypic range due to the absence of
358 sexual reproduction. When a longer fishing period is applied during the introduction fishing (Fig.
359 S2), the range of genotypic values widens in the no-predation scenarios without evolution, while all
360 FRII scenarios, as well as the evolutionary FRIII scenario, exhibit higher numbers of hake juveniles
361 among lower genotypic values when compared to the same scenarios with a fishing period having
362 30 time steps. While the genotypic values of the current approach translate directly to phenotype, or
363 more precisely to body size, the genotypic trends observed are not reflected in the age–size
364 distribution among evolutionary and non-evolutionary scenarios, meaning that the higher age–size
365 ranges are not present in the evolutionary scenarios, as would be expected (Fig. 6). However, one
366 exception exists where, under the conditions of absent evolutionary processes, the range of the
367 average size at age is slightly higher in the non-evolutionary FRIII scenario than its evolutionary
368 version (S7 and S8, respectively, in Fig. 6).

369 Among population parameters, recruitment, and biomass with respect to the carrying capacity
370 (BM to CC) show a discrepancy between the FRIII scenarios in which evolutionary processes are
371 present and absent (Figs 7a–b). The FRIII evolutionary scenario exhibits higher scale values for
372 these parameters than the non-evolutionary scenario, which suggests higher hake survival due to the
373 survival of favorable genotypes (i.e., evolutionary rescue). A similar discrepancy between
374 evolutionary and non-evolutionary FRIII scenarios is observed in recruitment when a longer fishing
375 period is applied (S3a), where no-predation and reference scenarios indicate delayed or impeded
376 recovery to their respective pre-fishing levels. Although not presented in the Supplement, other
377 population parameters such as prey abundance and BM to CC exhibit the same trend in reference
378 scenarios and scenarios without predation in the presence of evolutionary processes and a longer
379 fishing period. Hake population level parameters (i.e., BM to CC, recruitment) in the FRIII
380 scenarios show a reciprocal relationship between hake numbers and predator density (Figs S3d and
381 8c), where higher hake biomass do not increase FRIII predator density in the presence of evolution,
382 although there are no differences in recovery among evolutionary and non-evolutionary FRIII
383 scenarios. While evolutionary rescue supports a higher hake biomass, this occurs through the
384 survival of prey sizes that are elusive to an FRIII predator, which otherwise thrives in the absence of
385 an evolving prey. These trends in hake abundance with evolution present support higher fishing
386 catches in the evolutionary FRIII scenario. While the presence or absence of evolutionary processes
387 does not affect the prey–predator density ratio among FRII and FRII scenarios (Figs S3c and 8b),
388 the female–male ratio of hake captures some discrepancies between non-evolutionary and
389 evolutionary scenarios of no-predation and reference scenarios. FRII and FRIII scenarios show
390 similar responses to the fishing period with respect to the absence and presence of evolution,
391 although there is a slight delay in response to fishing in the case of FRIII predation.

392

393 3.2 Impact of predation mortality on hake survival and fisheries

394 The discrepancy in population resilience observed in the simulation testing stage is most evident
395 among scenarios with and without predation, where hake is more susceptible to demographic
396 collapse under predation, especially in the FRIII scenario due to the lowest resilience of only 30
397 time steps (Figs S1 and S3). Independent of fishing duration, most individual and population
398 parameters of hake demonstrate different scale values due to the trait-mediated effects of predation
399 (Figs S1 and S3, as well as Figs 4–5 and 7–8). Contrary to all scenarios without predation, L_{∞}
400 shows distinctly higher values under the effect of predation (Figs S1a and 4a), which is reflected as
401 a higher K in the absence of predation (Figs S1c and 4b). As the increasing trend in L_{∞} and the
402 decreasing trend in K are also present in all scenarios without predation, this suggests that the
403 density-dependent parameters applied in all the scenarios contribute to the slow adaptation of these
404 parameters (in addition to evolutionary processes and hake ecological characteristics). The reduced
405 somatic growth rate under predation is further reflected in the lower age at maturity due to
406 potentially higher allocation to reproduction under predation pressure, whereas reference and no-
407 predation scenarios demonstrate maturity at an older age (Figs S1b and 5a). While all predation
408 scenarios indicate lower somatic growth, size at maturity is unaffected by the presence of predation,
409 with a slightly higher size at maturity in the FRII scenarios than the FRIII scenarios during the pre-
410 fishing period (Figs S1d and 5b).

411 Both predation types exhibit strong overcompensation in age and size at maturity with fishing
412 cessation, due to the release of fishing pressure (Figs 5a and b). For both age and size at maturity,
413 the recovery is faster in all scenarios without predation but with SSD accounted for. In age at
414 maturity, recovery under the reference and FRIII scenarios occurs within a narrow time period,
415 whereas FRII scenarios prolong the recovery the most. Size at maturity more distinctly
416 demonstrates the differences in recovery among scenarios, where the no-predation scenarios are
417 followed by the FRIII, FRII, and reference scenarios. However, with a longer fishing period in the

reference and the no-predation scenarios (Figs S1b and d), these individual parameters exhibit a strong phenotypic shift in the absence of predation. Comparing the two predation types, the FRIII scenarios allow for quicker recovery, indicating a lesser negative impact than FRII predation, although this could be partly implied by the slightly longer fishing period applied in the FRII scenarios.

Predation mortality and the prey size selectivity shape the genotypic diversity (Fig. 5c) and demographic structure of the hake population (Fig. 6). Reference scenarios demonstrate a very narrow range of genotypes in comparison to the genotypic range values driven by the sex-specific difference in L_{∞} and predation size selectivity. This strengthens the importance of the size discrepancy among hake females and males, and also suggests that predation drives genotypic diversity in hake, which in return reflects the wider range of the age–size distribution (Fig. 6) in all the predation scenarios. In comparison to the reference scenarios (S1–S2 in Fig. 6), the scenarios without predation (S3–S4 in Fig. 6) show an age–size distribution skewed towards larger females. However, the average size of hake is clearly higher in the FRII and FRIII scenarios (S5–S8 in Fig. 6) than in all the scenarios without predation (S1–S4 in Fig. 5). While the size at age 0 (i.e., L_0) is the same for all the scenarios, the discrepancies are firstly observed in 1- to 2-year-old hake, which are larger in size for S4–S8 than for S2–S3.

Population parameters (BM to CC and recruitment) exhibit strong overcompensation after fishing cessation in all the predation scenarios (Figs 7a and b), where hake is clearly low in biomass and other abundance-correlated parameters in the presence of predation. The strong fish removal, however, is more pronounced under FRII than FRIII predation, which can be explained by the higher efficiency of an FRII predator in consuming prey at a lower prey density than an FRIII predator, particularly as prey density in the current IBM is solely dependent on prey abundance. While biomass levels are lower in the reference and all the predation scenarios, in the no-predation scenarios, biomass exceeds the carrying capacity during the pre-fishing and recovery periods. Such

443 levels of biomass suggest weaker density-dependent processes in the conditions of the scenarios
444 without predation, which with the absence of top-down control allow the hake population to
445 outgrow its own carrying capacity. During the pre-fishing period, FRII additionally induces greater
446 fluctuation in all population parameters, regardless of fishing duration (Figs S3a and 7a–b).
447 However, during the recovery period, all the presented population parameters demonstrate the same
448 trend of recovery among scenarios as the size at maturity, where the no-predation scenarios allow
449 for the fastest recovery, followed by the FRIII and FRII scenarios, and lastly the reference
450 scenarios. The order of scenarios showing the fastest to the slowest recovery changes as the
451 duration of the fishing period is increased, as FRIII scenarios now enable faster recovery in
452 recruitment, followed by FRII scenarios, which show a delayed reduction in numbers towards the
453 end of simulation time (Fig. S3a). Furthermore, all scenarios without predation notably slow down
454 the recovery in recruitment, while in the absence of hake, SSD and predation recovery (i.e.,
455 reference scenarios) is completely disabled.

456 The ratio of female to male hake exhibits different responses to fishing depending on the
457 presence or absence of predation, particularly when observed within the overlapping time frame of
458 fishing and predation (Fig. 7c). Reference scenarios as well as scenarios without predation exhibit a
459 similar trend of a delayed response to fishing, where the proportion of hake males removed by
460 fisheries increases by the end of the fishing period. Contrary to this, FRII and FRIII predators
461 demonstrate different sex (size) selectivity during the fishing period, where an FRII predator
462 increasingly selects hake females towards the middle of the fishing period and then shifts to male
463 hake as fishing ends. An FRIII predator consumes more males as fishing starts, but this ratio of
464 hake sexes gradually shifts towards hake females and then fluctuates around the middle towards the
465 cessation of fishing. Furthermore, the predation types, with their respective targeted prey size,
466 reveal the dynamics of predation and fisheries, suggesting a higher intensity of competition between
467 fisheries and FRII predators. Particularly in case of the FRIII evolutionary scenario, the prey size

selectivity of predation is reciprocal to the size selectivity of the fisheries. When comparing Figure 7c with the same graph for a longer fishing period (S3b), the comparison of hake sexes targeted by fisheries and predation is only possible among the scenarios with the two longest durations of the fishing period, meaning the reference scenarios and the scenarios without predation. The reference scenarios with a longer fishing period continue the trend observed in the reference scenarios under a short fishing period, where the proportion of female and male hake continues to fluctuate, with increasingly greater oscillations following each shift from one sex to the other (Fig. S3b). When comparing no-predation scenarios with short and long fishing periods, the proportions of hake males and females removed by the fisheries show a similar trend as in the reference scenarios, but with much lower oscillations.

478

479 **3.3 Predator–fisheries interactions: bottlenose dolphin survival and fishing catches**

The dynamic interactions between bottlenose dolphins and fisheries are mainly steered by the availability of hake to each competitor. While the absence of a natural competitor provides a higher gain for fisheries, the fishing catches decrease as the fishing period progresses, with catch slopes becoming steeper in all scenarios (Fig. 8a). The steepest decrease in catches is observed in the reference scenarios, which fall under the catches achieved in the conditions of the evolutionary FRIII scenario by the end of the fishing period. This demonstrates that incorporating key ecological parameters associated with the trait selected by fisheries (i.e., size), as well as predation mortality, slows down the potential depletion of fish. As observed in the hake population parameters (Figs 7a–b), as well as in the female–male ratio of hake (Fig. 7c), the most intensive fish removal when comparing the two types of predation is noted in FRII predation, which also reduces fishing catches the most. However, even though predation decreases fishing catches, the smoother catch slope observed in the evolutionary FRIII scenario (Fig. 8a) suggests a lower level of unsustainable fisheries removals in the long term.

493 Considering the competitive interaction between predators and fisheries, the prevailing fishing
494 regime exerts a negative indirect effect on predator density through prey–predator interactions (Figs
495 8b and c). More precisely, fishing reduces the predator density in both predation scenarios, which is
496 also reflected in the overcompensation of the prey–predator density ratio after the cessation of
497 fishing (Fig. 8b), particularly in the FRII scenarios with a longer fishing period (Fig. S3c). While an
498 FRII predator achieves a higher density level through the co-evolved prey–predator interaction and
499 more efficient consumption of prey, FRIII predator density benefits less from such interaction,
500 particularly if evolutionary processes are accounted for (Fig. 8c). Interestingly enough, the prey–
501 predator density ratio and particularly, the predator density of the evolutionary FRIII scenario
502 demonstrates micro-oscillations between the time steps. In comparison to predation, the removal of
503 hake by fisheries is significantly higher in intensity (observed in hake population parameters) than
504 hake removal by bottlenose dolphins, which impacts on predator survival more profoundly than
505 predators affect the fisheries gain due the absence of other target and prey species (Fig. 8c).
506 Moreover, with a slightly longer fishing period applied in the FRII scenarios (Fig. S3d), predator
507 density does not fully recover to its pre-fishing levels, with or without evolutionary processes in the
508 prey.

509

510 **4. Discussion**

511 **4.1 Specific conditions dictate the potential for FIE under coupled size-selective pressures**

512 The coupled effects of predation and fishing have diverse implications for a range of fish life-
513 history traits (e.g., K , size and age at maturity), with one trait being more susceptible to
514 evolutionary change than another under specific conditions: this dictates the resilience and stability
515 of observed hake parameters, as well as affecting the speed and pattern of recovery in the
516 parameters. These specific conditions are identified here as the presence or the absence of SSD in
517 hake, the duration and intensity of both size-selective pressures, as well as the predation type.
518 Among the individual hake parameters, L_{∞} and K show resistance towards an evolutionary shift, as

519 the ecological characteristics of hake and the density-dependent processes slow down the
520 stabilization of these parameters during the evolutionary scenarios. This can be explained by the
521 formulation of SSD incorporated directly in hake L_{∞} , which, in the presence of all other mentioned
522 conditions, prolongs the time needed for the stabilization of L_{∞} and, consequently, K . Furthermore,
523 as L_{∞} is the only evolving trait, the evolution in other life-history traits is not as affected by SSD
524 due to the phenotypic rather than genetic correlations among L_{∞} and other traits. As SSD in fish
525 species is reflected in higher growth rate in males, which reach maturity earlier than females, an
526 improved approach to the description of SSD would be to introduce additional and more reliable
527 sex-specific data on hake growth parameters.

528 The length of the fishing period, removal intensity, and predation type appear crucial in inducing
529 very persistent alterations in size and age at maturity. While age at maturity shows some resilience
530 to the phenotypic shift in the FRII predation scenarios, size at maturity slightly drops after the
531 cessation of fishing in the same scenarios. With a longer fishing period, the evolutionary change for
532 both individual parameters becomes obvious in all scenarios without predation and SSD accounted
533 for, while age at maturity under the FRII scenarios also shows a lack of recovery to its pre-fishing
534 level. A longer fishing period applied in the reference and no-predation scenarios, as well as FRII
535 scenarios, indicates that fishing duration is pivotal to inducing more pronounced shifts in these
536 individual parameters. This is corroborated by studies on exploited stocks that demonstrate a
537 delayed evolutionary response to fishing after a longer period of exploitation (Devine et al., 2012).
538 Fish species naturally prone to early maturation schedules have demonstrated a higher potential for
539 reduced size at maturity under fishing pressure (Landi et al., 2015). This can only partly explain the
540 decrease in size at maturity of hake in the FRII scenarios after fishing cessation, since the expected
541 decrease in age at maturity did not occur within the same scenario settings. Fishing duration might
542 not be enough to explain the observed alterations in size and age at maturity, as the overall intensity
543 of size-selective removal caused by predation and fishing ultimately contributes to the direction and

544 magnitude of their responses. For predation pressure, the predation type is associated with the
545 predation intensity owing to the differences in consumption patterns between the two predation
546 types. The difference in the intensity of size-selective removal between the two predation types
547 supports the higher potential for FIE in the mentioned parameters when the FRII predation type is
548 applied. However, while the currently prevailing fishing size selectivity in synergy with predation
549 removal does not impoverish the genotypic diversity of hake, as observed in other fish species
550 experiencing size-selective removals (see Marty et al., 2015), the ensured phenotypic diversity
551 (owing to the phenotypic variation added to the genotypic values) could mask the potential
552 evolutionary shifts in individual life-history traits. In this case, the potential for adaptive phenotypic
553 plasticity in hake has minimized the impact of evolutionary feedbacks, which is consistent with
554 other studies addressing the phenotypic adaptations of hake to evolutionary processes under
555 harvesting pressure (Hidalgo et al., 2014). Consequently, intraspecific phenotypic diversity further
556 facilitates population demographic stability (i.e., a wide range of age–size classes) and minimizes
557 the impact of size-selective fishing.

558

559 **4.2 The importance of accounting for size-related ecological characteristics**

560 Hake population parameters demonstrate different responses to predation and fishing depending
561 on the presence or absence of SSD and evolutionary processes, as well as the predation type and
562 fishing period. The impact of predation and fishing is most intensive in the FRII scenarios (with
563 SSD in hake accounted for), while the most negative ecological impact of fishing is noted in
564 conditions where predation and SSD are not accounted for. Evolutionary processes, however, have
565 crucial implications for hake abundance and recruitment in the case of the FRIII predation type,
566 where hake survival increases, despite fishing and predation pressures, through evolutionary rescue.
567 This demonstrates that true adaptive changes in hake life-history traits have the potential to
568 counteract the negative effects of size-selective pressures (see Hendry et al., 2011). At the

individual and population levels, a short fishing duration and accounting for SSD without predation allows hake to recover faster from fishing than under conditions with predation, which supports the cases arguing for slow or even disabled recovery in some overexploited fish stocks due to high predation mortality (Hutchings and Kupařinen, 2019). With an increase in fishing duration, the implications of accounting for SSD, the presence of predation, and the predation type become more crucial to the speed of recovery in hake population parameters. Without SSD accounted for and predation, recovery is impeded or disabled, although the FRIII predation type enables faster recovery than FRII predation. Here, SSD forms the basis for recovery, which is not surprising considering the important implications of SSD for growth parameters, size and age at maturity, size-selective mortality, and ultimately, the demographic structure of many fish species (Carroll and Lowerre-Barbieri, 2019; Cerviño, 2014; Keyl et al., 2015). Furthermore, the life-history traits of shared fish prey targeted by predators and fisheries is pivotal in determining to what degree the coexistence of predators and fisheries is possible (Huss et al., 2014).

4.3 Predator individual variability: implications for fish stock assessments

Under the condition of a two-species prey–predator interaction, it is not surprising that the predation tested here profoundly impacts on fish biomass and population resilience. The current results, however, should be considered alongside the assumptions of this approach, which does not account for the individual variability, ontogenetic development stage, or catholic diet of the bottlenose dolphin. According to Navarrete and Manzur (2008), individual predator parameters (e.g., body size, diet composition, and *per capita* prey consumption) are more responsive to changes in prey for a generalist predator. However, with regards to the wide range of prey species, bottlenose dolphins have been known to demonstrate prey species preference when offered a choice (Corkeron et al., 1990). Furthermore, under low availability of other prey species, in small food webs as well as in the case of predator specialization to feed on specific prey species (Pettorelli et

al., 2011), the factors associated with opportunistic feeding in bottlenose dolphins can be less relevant in the presence of high intra- or interspecific competition. In the absence of individual variability, predation types provide a source of ecological complexity directing the response of predator parameters to fishing. Thus, with the increased level of ecological complexity from FRII to FRIII, the competition between predators and fisheries decreases. Here the consumption pattern of predators is of high importance in fish stock assessments, not just for the quantification of predation but also to avoid negative indirect effects of fisheries on marine trophic links and the environment. Moreover, as the evolving prey can facilitate the survival of a non-evolving predator (termed as “indirect evolutionary rescue” by Yamamichi and Miner [2015]), this coincides with the predator population dynamics observed in the FRII predator, but not for the FRIII. In contrast, the reciprocal size selectivity of the prey noted between fisheries and FRIII predation is similar to the fishing strategies targeting all individuals regardless of maturation status, thus reducing the potential for FIE (Ernande et al., 2004). Studies addressing the competitive interactions between marine mammals and fisheries, as well as the impact of high fishing mortality, have revealed that overexploitation is more detrimental to the sustainability of long-term fisheries yields than marine mammals (Garcia et al., 2012; Morissette et al., 2012) or the potential FIE (Hutchings and Kuperinen, 2019). Furthermore, even without a multi-species aspect, the current results predict potential long-term unsustainable outcomes for hake abundance and, consequently, for fisheries in the absence of predation, which is consistent with ecosystem-based studies that predict a decline in fisheries yields due to the lower ecosystem productivity caused by the removal of top predators (Morissette et al., 2010; Roman et al., 2014).

615

616 **5. Conclusions**

617 In this study, we demonstrated how two sources of size-selective mortality in combination with
618 the interplay between ecological and evolutionary processes shape the responses of prey parameters

619 at individual and population levels. With respect to this, the duration, intensity, and selected fish
620 size range of both predation pressures have the most prominent implications for the heritability of
621 shifts observed in hake life-history traits, which need to be accounted for when aiming to avoid FIE.
622 As different factors can determine the duration, intensity, and size selectivity of predation and
623 fisheries, the future direction in understanding the prerequisites driving the evolution of individual
624 life-history traits is to investigate which combinations, as well as thresholds of these conditions,
625 have the potential to trigger heritable changes. Since in our study the predation type was found to
626 determine the intensity and size selectivity of predation pressure, future research should address the
627 potential for evolutionary changes with an increase in ecological complexity (i.e., predation type,
628 multi-species, or even ecosystem aspect). According to the outputs of the initial introduction of
629 fisheries, incorporating more than one type of predation in fish stock assessments can aid in setting
630 more reliable limits for harvest yields and rates, weighted against the risks associated with predation
631 intensity as well as potential heritable phenotypic changes. Ultimately, understanding of how
632 predation affects fish availability for fisheries will consequently provide insights into the conditions
633 under which predators indirectly support fisheries or render fishing yields low.

634 While conservative, the current approach applies key prey ecological characteristics and
635 evolutionary processes providing insights into pivotal, baseline mechanisms in the prey–predator
636 dynamics, as well as the ecological interactions between bottlenose dolphins and hake fisheries.
637 With consideration of the implications of SSD for hake fecundity and size-selective mortality, SSD
638 combined with the predation type provides additional insights into the species-specific resilience to
639 and recovery potential under one or multiple size-selective pressures. Such prey–predator dynamics
640 has direct implications for setting sustainable reference points and successful conservation measures
641 to ensure fish survival and the gain of future fisheries. Furthermore, while SSD and FRIII predation
642 selectivity in the presence of evolutionary processes enrich the prey’s phenotypic and consequently

643 genotypic diversity, this can ensure a higher resilience to potential shifts in size at maturity, as it
644 disperses the intensity of size-selective removals by fisheries.

645 The fishing mortality and selectivity applied here resulted in a profoundly negative ecological
646 impact on hake survival, exceeding the removals and selectivity of predation mortality. Moreover,
647 indirect fishing effects extend to predator survival, as hake consumption by the bottlenose dolphin,
648 and more precisely its prey size selectivity, will depend on the age–size classes available to the
649 predator in the presence of an intensive fishing regime. The dynamics of the fishing regime and
650 predation types of the current study demonstrate how predators adapt their prey size selectivity
651 according to the size selectivity of fishing. The question is whether these predator adjustments to
652 fishing size selectivity will be sufficient to counteract the effects of overexploitation. To answer
653 this, further research is required to address the role of predation from a wider perspective of
654 complex trophic interactions or the ecosystem level. Such aspects are comprehensively included
655 within dynamic ecosystem and individual-based models, which could also explore indirect or even
656 “food-web competition” between bottlenose dolphins and hake fisheries (see ICES, 2015; Wood et
657 al., 2018). However, incorporating individual variability in both prey and predators with the
658 integration of eco-evolutionary feedbacks in a more complex trophic structure is an ambitious step
659 forward to improving the ecosystem approach to fisheries, as well as conservation efforts for natural
660 predators. While similar approaches to the current one do exist, they have not generally accounted
661 for one of the features or processes included in our model. Thus, addressing such a multilayered
662 research question requires a basic approach, where underlying mechanisms can be tested and
663 brought to light. On the other side, holistic approaches to fisheries management demand models that
664 can encapsulate such complexity whilst maintaining the applicability of their outputs to managerial
665 decision-making processes. In line with this, the management of wild fish stocks still remains a
666 multidisciplinary agenda with a strong need for a transdisciplinary approach.

667

668 **Acknowledgements**

669 This research was financially supported by the Finnish Cultural Foundation (Grant Nr. 00190412;
670 DJ), the Academy of Finland (AK), the Natural Sciences and Engineering Research Council of
671 Canada (AK), and the European Research Council (COMPLEX-FISH 770884; AK). The authors
672 would like to thank Dr Anssi Vainikka and the two anonymous reviewers for insightful comments
673 and suggestions.

674

675

676 **References**

- 677 Abrams, P.A., Ginzburg, L.R., 2000. The nature of predation : prey dependent , ratio dependent or
678 neither ? TREE 15, 337–341.
- 679 Allhoff, K.T., Ritterskamp, D., Rall, B.C., Drossel, B., Guill, C., 2015. Evolutionary food web
680 model based on body masses gives realistic networks with permanent species turnover. Sci.
681 Rep. 5, 1–12. <https://doi.org/10.1038/srep10955>
- 682 Bearzi, G., Agazzi, S., Gonzalvo, J., Bonizzoni, S., Costa, M., Petroselli, A., 2010. Biomass
683 removal by dolphins and fisheries in a Mediterranean Sea coastal area: do dolphins have an
684 ecological impact on fisheries? Aquat. Conserv. Mar. Freshw. Ecosyst. 20, 549–559.
685 <https://doi.org/10.1002/aqc.1123>
- 686 Bearzi, G., Fortuna, C.M., Reeves, R.R., 2008. Ecology and conservation of common bottlenose
687 dolphins *Tursiops truncatus* in the Mediterranean Sea. Mamm. Rev. 39, 92–123.
688 <https://doi.org/10.1111/j.1365-2907.2008.00133.x>
- 689 Beckerman, A.P., Rodgers, G.M., Dennis, S.R., 2010. The reaction norm of size and age at maturity
690 under multiple predator risk. J. Anim. Ecol. 79, 1069–1076. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2656.2010.01703.x)
691 [2656.2010.01703.x](https://doi.org/10.1111/j.1365-2656.2010.01703.x)
- 692 Belgrano, A., Fowler, C.W., 2013. How fisheries affect evolution. Science (80-.). 342, 1176–1177.

693 <https://doi.org/10.1126/science.1245490>

694 Bell, R.J., Collie, J.S., Branch, T.A., Fogarty, M.J., Minto, C., Ricard, D., 2018. Changes in the size
695 structure of marine fish communities. *ICES J. Mar. Sci.* 75, 102–112.
696 <https://doi.org/10.1093/icesjms/fsx118>

697 Carroll, J., Lowerre-Barbieri, S.K., 2019. Interactions of dimorphic growth, reproductive behavior,
698 and a size-regulated fishery: A case study using spotted seatrout *Cynoscion nebulosus*. *Mar.*
699 *Ecol. Prog. Ser.* 608, 233–245. <https://doi.org/10.3354/meps12795>

700 Cerviño, S., 2014. Estimating growth from sex ratio-at-length data in species with sexual size
701 dimorphism. *Fish. Res.* 160, 112–119. <https://doi.org/10.1016/j.fishres.2013.11.010>

702 Conover, D.O., Munch, S.B., 2002. Sustaining Fisheries Yields Over Evolutionary Time Scales.
703 *Science* (80-.). 297, 94–96. <https://doi.org/10.1126/science.1074085>

704 Corkeron, P.J., Bryden, M.M., Hedstrom, K.E., 1990. Feeding by bottlenose dolphins in association
705 with trawling operations in Moreton Bay, Australia, in: Leatherwood, S., Reeves, R.R. (Eds.),
706 *The Bottlenose Dolphin*. Academic Press Inc., San Diego, pp. 329–336.

707 Crozier, L.G., Hutchings, J.A., 2014. Plastic and evolutionary responses to climate change in fish.
708 *Evol. Appl.* 7, 68–87. <https://doi.org/10.1111/eva.12135>

709 Darimont, C.T., Fox, C.H., Bryan, H.M., Reimchen, T.E., 2015. The unique ecology of human
710 predators. *Science* (80-.). 349, 858–860. <https://doi.org/10.1126/science.aac4249>

711 De Roos, A.M., Boukal, D.S., Persson, L., 2006. Evolutionary regime shifts in age and size at
712 maturation of exploited fish stocks. *Proc. R. Soc. B Biol. Sci.* 273, 1873–1880.
713 <https://doi.org/10.1098/rspb.2006.3518>

714 DeLong, J.P., Gilbert, B., Shurin, J.B., Savage, V.M., Barton, B.T., Clements, C.F., Dell, A.I.,
715 Greig, H.S., Harley, C.D.G., Kratina, P., McCann, K.S., Tunney, T.D., Vasseur, D.A.,
716 O'Connor, M.I., 2015. The Body Size Dependence of Trophic Cascades. *Am. Nat.* 185, 354–
717 366. <https://doi.org/10.1086/679735>

718 DeLong, J.P., Luhring, T.M., 2018. Size-dependent predation and correlated life history traits alter
 719 eco-evolutionary dynamics and selection for faster individual growth. *Popul. Ecol.* 60, 9–20.
 720 <https://doi.org/10.1007/s10144-018-0608-7>

721 Devine, J.A., Wright, P.J., Pardoe, H.E., Heino, M., 2012. Comparing rates of contemporary
 722 evolution in life-history traits for exploited fish stocks. *Can. J. Fish. Aquat. Sci.* 69, 1105–
 723 1120. <https://doi.org/10.1139/f2012-047>

724 Domínguez-Petit, R., Korta, M., Saborido-Rey, F., Murua, H., Sainza, M., Piñeiro, C., 2008.
 725 Changes in size at maturity of European hake Atlantic populations in relation with stock
 726 structure and environmental regimes. *J. Mar. Syst.* 71, 260–278.
 727 <https://doi.org/10.1016/j.jmarsys.2007.04.004>

728 Enberg, K., Jørgensen, C., Dunlop, E.S., Heino, M., Dieckmann, U., 2009. Implications of
 729 fisheries-induced evolution for stock rebuilding and recovery. *Evol. Appl.* 2, 394–414.
 730 <https://doi.org/10.1111/j.1752-4571.2009.00077.x>

731 Ernande, B., Dieckmann, U., Heino, M., 2004. Adaptive changes in harvested populations:
 732 Plasticity and evolution of age and size at maturation. *Proc. R. Soc. B Biol. Sci.* 271, 415–423.
 733 <https://doi.org/10.1098/rspb.2003.2519>

734 Fraser, D.J., 2013. The emerging synthesis of evolution with ecology in fisheries science. *Can. J.*
 735 *Fish. Aquat. Sci.* 70, 1417–1428. <https://doi.org/10.1139/cjfas-2013-0171>

736 Fryxell, D.C., Wood, Z.T., Robinson, R., Kinnison, M.T., Palkovacs, E.P., 2019. Eco-evolutionary
 737 feedbacks link prey adaptation to predator performance. *Biol. Lett.* 15, 20190626.
 738 <https://doi.org/10.1098/rsbl.2019.0626>

739 Garcia, S.M., Kolding, J., Rice, J., Rochet, M.-J., Zhou, S., Arimoto, T., Beyer, J.E., Borges, L.,
 740 Bundy, A., Dunn, D., Fulton, E.A., Hall, M., Heino, M., Law, R., Makino, M., Rijnsdorp,
 741 A.D., Simard, F., Smith, A.D.M., 2012. Reconsidering the Consequences of Selective
 742 Fisheries. *Science* (80-.). 335, 1045–1047.

743 <https://doi.org/http://dx.doi.org/10.1126/science.1214594>

744 Gårdmark, A., Dieckmann, U., Lundberg, P., 2003. Life-history evolution in harvested populations:
 745 The role of natural predation. *Evol. Ecol. Res.* 5, 239–257.

746 Gislason, H., Daan, N., Rice, J.C., Pope, J.G., 2010. Size, growth, temperature and the natural
 747 mortality of marine fish. *Fish Fish.* 11, 149–158. [https://doi.org/10.1111/j.1467-](https://doi.org/10.1111/j.1467-2979.2009.00350.x)
 748 [2979.2009.00350.x](https://doi.org/10.1111/j.1467-2979.2009.00350.x)

749 Goetz, S., Read, F.L., Ferreira, M., Portela, J.M., Santos, M.B., Vingada, J., Siebert, U., Marçalo,
 750 A., Santos, J., Araújo, H., Monteiro, S., Caldas, M., Riera, M., Pierce, G.J., 2015. Cetacean
 751 occurrence, habitat preferences and potential for cetacean-fishery interactions in Iberian
 752 Atlantic waters: Results from cooperative research involving local stakeholders. *Aquat.*
 753 *Conserv. Mar. Freshw. Ecosyst.* 25, 138–154. <https://doi.org/10.1002/aqc.2481>

754 Govaert, L., Fronhofer, E.A., Lion, S., Eizaguirre, C., Bonte, D., Egas, Ma., Hendry, A.P., De Brito
 755 Martins, A., Melián, C.J., Raeymaekers, J.A.M., Ratikainen, I.I., Saether, B.E., Schweitzer,
 756 J.A., Matthews, B., 2019. Eco-evolutionary feedbacks—Theoretical models and perspectives.
 757 *Funct. Ecol.* 33, 13–30. <https://doi.org/10.1111/1365-2435.13241>

758 Griffiths, D., 1975. Prey Availability and the Food of Predators. *Ecology* 56, 1209–1214.
 759 <https://doi.org/10.2307/1936161>

760 Heino, M., Beatriz, D., Dieckmann, U., 2015. Fisheries-Induced Evolution. *Annu. Rev. Ecol. Evol.*
 761 *Syst.* 46, 461–480. <https://doi.org/10.1146/annurev-ecolsys-112414-054339>

762 Hendry, A.P., 2016. Key questions on the role of phenotypic plasticity in eco-evolutionary
 763 dynamics. *J. Hered.* 107, 25–41. <https://doi.org/10.1093/jhered/esv060>

764 Hendry, A.P., Kinnison, M.T., Heino, M., Day, T., Smith, T.B., Fitt, G., Bergstrom, C.T.,
 765 Oakeshott, J., Jørgensen, P.S., Zalucki, M.P., Gilchrist, G., Southerton, S., Sih, A., Strauss, S.,
 766 Denison, R.F., Carroll, S.P., 2011. Evolutionary principles and their practical application.
 767 *Evol. Appl.* 4, 159–183. <https://doi.org/10.1111/j.1752-4571.2010.00165.x>

768 Hernandez-Milian, G., Berrow, S., Santos, M.B., Reid, D., Rogan, E., 2015. Insights into the
769 trophic ecology of bottlenose dolphins (*Tursiops truncatus*) in Irish waters. *Aquat. Mamm.* 41,
770 226–239. <https://doi.org/10.1578/AM.41.2.2015.226>

771 Hidalgo, M., Olsen, E., , J., Saborido- Rey, F., Murua, H., Stenseth, N.C., 2014. Contrasting
772 evolutionary demography induced by fishing : the role of adaptive phenotypic plasticity. *Ecol.*
773 *Appl.* 24, 1101–1114. <https://doi.org/https://doi.org/10.1890/12-1777.1>

774 Holling, C.S., 1973. Resilience and Stability of Ecological Systems. *Annu. Rev. Ecol. Syst.* 4, 1–23.

775 Holling, C.S., 1959. The Components of Predation as Revealed by a Study of Small Mammal
776 Predation of the European Pine Sawfly. *Can. Entomol.* 91, 293–320.

777 Huss, M., de Roos, A.M., Van Leeuwen, A., Gårdmark, A., 2014. Facilitation of fisheries by natural
778 predators depends on life history of shared prey. *Oikos* 123, 1071–1080.
779 <https://doi.org/10.1111/oik.00839>

780 Hutchings, J.A., Kuparinen, A., 2019. Implications of fisheries-induced evolution for population
781 recovery: Refocusing the science and refining its communication. *Fish Fish.* 1–12.
782 <https://doi.org/10.1111/faf.12424>

783 ICES, 2019. Report of the Working Group for the Bay of Biscay and the Iberian waters Ecoregion
784 (WGBIE). <https://doi.org/http://doi.org/10.17895/ices.pub.5299>

785 Jusufovski, D., Kuparinen, A., 2014. Contrasting evolutionary and ecological management
786 objectives in the context of sustainable harvesting. *Evol. Ecol. Res.* 16.

787 Kastelein, R.A., Staal, C., Wiepkema, P.R., 2003. Food consumption, food passage time, and body
788 measurements of captive Atlantic bottlenose dolphins (*Tursiops truncatus*). *Aquat. Mamm.* 29,
789 53–66. <https://doi.org/10.1578/016754203101024077>

790 Keyl, F., Kempf, A.J., Sell, A.F., 2015. Sexual size dimorphism in three North Sea gadoids. *J. Fish*
791 *Biol.* 86, 261–275. <https://doi.org/10.1111/jfb.12579>

792 Kindsvater, H.K., Palkovacs, E.P., 2017. Predicting Eco-evolutionary Impacts of Fishing on Body

793 Size and Trophic Role of Atlantic Cod. *Copeia* 105, 475–482. <https://doi.org/10.1643/ot-16->
794 533

795 Kuparinen, A., Hardie, D.C., Hutchings, J.A., 2012. Evolutionary and ecological feedbacks of the
796 survival cost of reproduction. *Evol. Appl.* 5, 245–255. <https://doi.org/10.1111/j.1752->
797 4571.2011.00215.x

798 Kuparinen, A., Hutchings, J.A., 2012. Consequences of fisheries-induced evolution for population
799 productivity and recovery potential. *Proc. R. Soc. B Biol. Sci.* 279, 2571–2579.
800 <https://doi.org/10.1098/rspb.2012.0120>

801 Kuparinen, A., Stenseth, N.C., Hutchings, J.A., 2014. Fundamental population-productivity
802 relationships can be modified through density-dependent feedbacks of life-history evolution.
803 *Evol. Appl.*

804 Landi, P., Hui, C., Dieckmann, U., 2015. Fisheries-induced disruptive selection. *J. Theor. Biol.* 365,
805 204–216. <https://doi.org/10.1016/j.jtbi.2014.10.017>

806 Lankau, R.A., Strauss, S.Y., 2011. Newly rare or newly common: evolutionary feedbacks through
807 changes in population density and relative species abundance, and their management
808 implications. *Evol. Appl.* 4, 338–353. <https://doi.org/10.1111/j.1752-4571.2010.00173.x>

809 Law, R., 2000. Fishing, selection, and phenotypic evolution. *ICES J. Mar. Sci.* 57, 659–668.
810 <https://doi.org/10.1006/jmsc.2000.0731>

811 Law, R., Grey, D.R., 1989. Evolution of yields from populations with age-specific cropping. *Evol.*
812 *Ecol.* 3, 343–359. <https://doi.org/10.1007/BF02285264>

813 Law, R., Plank, M.J., 2018. Balanced harvesting could reduce fisheries-induced evolution. *Fish*
814 *Fish.* 19, 1078–1091. <https://doi.org/10.1111/faf.12313>

815 Lockyer, C., 2007. All creatures great and smaller: A study in cetacean life history energetics. *J.*
816 *Mar. Biol. Assoc. United Kingdom* 87, 1035–1045.
817 <https://doi.org/10.1017/S0025315407054720>

818 López, B.D., 2006. Interactions between Mediterranean bottlenose dolphins (*Tursiops truncatus*)
 819 and gillnets off Sardinia , Italy. ICES J. Mar. Sci. 63, 946–951.
 820 <https://doi.org/10.1016/j.icesjms.2005.06.012>

821 Lowerre-Barbieri, S., DeCelles, G., Pepin, P., Catalán, I.A., Muhling, B., Erisman, B., Cadrin, S.X.,
 822 Alós, J., Ospina-Alvarez, A., Stachura, M.M., Tringali, M.D., Burnsed, S.W., Paris, C.B.,
 823 2017. Reproductive resilience: a paradigm shift in understanding spawner-recruit systems in
 824 exploited marine fish. Fish Fish. 18, 285–312. <https://doi.org/10.1111/faf.12180>

825 Madin, E.M.P., Dill, L.M., Ridlon, A.D., Heithaus, M.R., Warner, R.R., 2016. Human activities
 826 change marine ecosystems by altering predation risk. Glob. Chang. Biol. 22, 44–60.
 827 <https://doi.org/10.1111/gcb.13083>

828 Marty, L., Dieckmann, U., Ernande, B., 2015. Fisheries-induced neutral and adaptive evolution in
 829 exploited fish populations and consequences for their adaptive potential. Evol. Appl. 8, 47–63.
 830 <https://doi.org/10.1111/eva.12220>

831 Mehault, S., Domínguez-Petit, R., Cerviño, S., Saborido-Rey, F., 2010. Variability in total egg
 832 production and implications for management of the southern stock of European hake. Fish.
 833 Res. 104, 111–122. <https://doi.org/10.1016/j.fishres.2010.03.019>

834 Melián, C.J., Vilas, C., Baldó, F., González-Ortegón, E., Drake, P., Williams, R.J., 2011. Eco-
 835 evolutionary Dynamics of Individual-Based Food Webs, in: Advances in Ecological Research.
 836 Elsevier Ltd., pp. 225–268. <https://doi.org/10.1016/B978-0-12-386475-8.00006-X>

837 Morissette, L., Christensen, V., Pauly, D., 2012. Marine Mammal Impacts in Exploited Ecosystems:
 838 Would Large Scale Culling Benefit Fisheries? PLoS One 7.
 839 <https://doi.org/10.1371/journal.pone.0043966>

840 Morissette, L., Kaschner, K., Gerber, L.R., 2010. Ecosystem models clarify the trophic role of
 841 whales off Northwest Africa. Mar. Ecol. Prog. Ser. 404, 289–302.
 842 <https://doi.org/10.3354/meps08443>

843 Mousseau, T.A., Roff, D.A., 1987. Natural selection and the heritability of fitness components.
844 Heredity (Edinb). 59, 181.

845 Murua, H., 2010. The Biology and Fisheries of European Hake , *Merluccius merluccius* , in the
846 North-East Atlantic, 1st ed, Advances in Marine Biology. Elsevier Ltd.
847 <https://doi.org/10.1016/B978-0-12-381015-1.00002-2>

848 Navarrete, S.A., Manzur, T., 2008. Individual- And population-level responses of a keystone
849 predator to geographic variation in prey. Ecology 89, 2005–2018. [https://doi.org/10.1890/07-](https://doi.org/10.1890/07-1231.1)
850 1231.1

851 Neubauer, P., Jensen, O.P., Hutchings, J.A., Baum, J.K., 2013. Resilience and recovery of
852 overexploited marine populations. Science (80-.). 340, 347–349.
853 <https://doi.org/10.1126/science.1230441>

854 Ohlberger, J., Thackeray, S.J., Winfield, I.J., Maberly, S.C., Vøllestad, L.A., 2014. When
855 phenology matters: Age - size truncation alters population response to trophic mismatch. Proc.
856 R. Soc. B Biol. Sci. 281, 1–7. <https://doi.org/10.1098/rspb.2014.0938>

857 Palkovacs, E.P., Kinnison, M.T., Correa, C., Dalton, C.M., Hendry, A.P., 2012. Fates beyond traits:
858 Ecological consequences of human-induced trait change. Evol. Appl. 5, 183–191.
859 <https://doi.org/10.1111/j.1752-4571.2011.00212.x>

860 Palkovacs, E.P., Moritsch, M.M., Contolini, G.M., Pelletier, F., 2018. Ecology of harvest-driven
861 trait changes and implications for ecosystem management. Front. Ecol. Environ. 16, 20–28.
862 <https://doi.org/10.1002/fee.1743>

863 Pettorelli, N., Coulson, T., Durant, S.M., Gaillard, J.M., 2011. Predation, individual variability and
864 vertebrate population dynamics. Oecologia 167, 305–314. [https://doi.org/10.1007/s00442-011-](https://doi.org/10.1007/s00442-011-2069-y)
865 2069-y

866 Philips, B.L., Brown, G.P., Shine, R., 2010. Life-history evolution in range-shifting populations.
867 Ecology 91, 1617–1627. <https://doi.org/https://doi.org/10.1890/09-0910.1>

868 R Core Team 2019. R: A language and environment for statistical computing. R Foundation for
 869 Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

870 Reznick, D. A., Bryga, H. and Endler, J.A., 1990. Experimentally induced life-history evolution in a
 871 natural population. Nat. Publ. Gr. 346, 357–359. <https://doi.org/10.1038/346357a0>

872 Roff, D.A., 2002. Life history evolution. Sinauer, Sunderland, MA.

873 Roman, J., Estes, J.A., Morissette, L., Smith, C., Costa, D., McCarthy, J., Nation, J., Nicol, S.,
 874 Pershing, A., Smetacek, V., 2014. Whales as marine ecosystem engineers. Front. Ecol.
 875 Environ. 12, 377–385. <https://doi.org/10.1890/130220>

876 Rosenzweig, M.. L., MacArthur, R.H., 1963. Graphical Representation and Stability Conditions of
 877 Predator-Prey Interactions. Am. Nat. 97, 209–223.

878 Saavedra, C., 2017. Multispecies population modelling of the common dolphin (*Delphinus delphis*),
 879 the bottlenose dolphin (*Tursiops truncatus*) and the southern stock of European hake
 880 (*Merluccius merluccius*), in Atlantic waters of the Iberian Peninsula. University of Vigo,
 881 Spain.

882 Santos, M.B., Fernández, R., López, A., Martínez, J.A., Pierce, G.J., 2007. Variability in the diet of
 883 bottlenose dolphin, *Tursiops truncatus*, in Galician waters, north-western Spain, 1990-2005. J.
 884 Mar. Biol. Assoc. United Kingdom 87, 231–241. <https://doi.org/10.1017/S0025315407055233>

885 Santos, M.B., Saavedra, C., Pierce, G.J., 2014. Quantifying the predation on sardine and hake by
 886 cetaceans in the Atlantic waters of the Iberian Peninsula. Deep. Res. Part II Top. Stud.
 887 Oceanogr. 106, 232–244. <https://doi.org/10.1016/j.dsr2.2013.09.040>

888 Scharf, F.S., Juanes, F., Rountree, R.A., 2000. Predator size - Prey size relationships of marine fish
 889 predators: Interspecific variation and effects of ontogeny and body size on trophic-niche
 890 breadth. Mar. Ecol. Prog. Ser. 208, 229–248. <https://doi.org/10.3354/meps208229>

891 Sion, L., Zupa, W., Calculli, C., Garofalo, G., Hidalgo, M., 2019. Spatial distribution pattern of
 892 European hake, *Merluccius merluccius* (Pisces : *Merlucciidae*), in the Mediterranean Sea. Sci.

893 Mar. 83, 000–000. <https://doi.org/10.3989/scimar.04988.12A>

894 Shuter, B.J., Giacomini, H.C., de Kerckhove, D., Vascotto, K., 2015. Fish life history dynamics:
 895 shifts in prey size structure evoke shifts in predator maturation traits. *Can. J. Fish. Aquat. Sci.*
 896 73, 693–708. <https://doi.org/10.1139/cjfas-2015-0190>

897 Spitz, J., Ridoux, V., Trites, A.W., Laran, S., Authier, M., 2018. Prey consumption by cetaceans
 898 reveals the importance of energy-rich food webs in the Bay of Biscay. *Prog. Oceanogr.* 166,
 899 148–158. <https://doi.org/10.1016/j.pocean.2017.09.013>

900 Swain, D.P., Sinclair, A.F., Hanson, J.M., 2007. Evolutionary response to size-selective mortality in
 901 an exploited fish population. *Proc. R. Soc. B Biol. Sci.* 274, 1015–1022.
 902 <https://doi.org/10.1098/rspb.2006.0275>

903 Waples, R.S., Audzijonyte, A., 2016. Fishery-induced evolution provides insights into adaptive
 904 responses of marine species to climate change. *Front. Ecol. Environ.* 14, 217–224.
 905 <https://doi.org/10.1002/fee.1264>

906 Wood, Z.T., Palkovacs, E.P., Kinnison, M.T., 2018. Eco-evolutionary Feedbacks from Non-target
 907 Species Influence Harvest Yield and Sustainability. *Sci. Rep.* 8, 6389.
 908 <https://doi.org/10.1038/s41598-018-24555-0>

909 Yamamichi, M., Miner, B.E., 2015. Indirect evolutionary rescue: Prey adapts, predator avoids
 910 extinction. *Evol. Appl.* 8, 787–795. <https://doi.org/10.1111/eva.12295>

911 Zhou, S., Smith, A.D.M., Punt, A.E., Richardson, A.J., Gibbs, M., Fulton, E.A., Pascoe, S.,
 912 Bulman, C., Bayliss, P., Sainsbury, K., 2010. Ecosystem-based fisheries management requires
 913 a change to the selective fishing philosophy. *Proc. Natl. Acad. Sci.* 107, 9485–9489.
 914 <https://doi.org/10.1073/pnas.0912771107>

915

916 E – Sources:

917 Wikipedia Commons: https://commons.wikimedia.org/wiki/File:Merluccius_merluccius_mapa.svg;

918 Accessed on 16.12.2019
919 European Commission, Fisheries:
920 https://ec.europa.eu/fisheries/marine_species/wild_species/hake_en;

921 Accessed on 16.12.2019

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936 **Tables**

937 **Table 1.** Set values of model parameters (M = natural mortality; M_{rc} = natural mortality due to cost
938 of reproduction; N_i = prey initial population size; CC = prey carrying capacity; p_i = predator initial
939 population density; a = predator attack rate; e = assimilation efficiency; d = predator death rate; h =
940 predator handling time; F = fishing intensity; a, b = fishing selectivity parameters; L_{50} = length at
941 50% retention).

942

Set model parameters	Prey initial values	Parameters	Values	Sources	Predation	Parameters	Values	Sources
		M	0.1	Estimated through model parameterization		e	0.90	Lockyer, 2007
		M_{rc}	0.1			d	0.06	Estimated through model parameterization
		N_i	4000			h	0.5	
		CC	6000			F	0.15	
	Predation	p_i	0.3	Fishing	a , b	-10.85; 0.411	Estimated through model parameterization	
		a	0.07		L_{50}	24.6 cm		ICES, 2019

Figure captions

Figure 1. The European hake and its native distribution range in European, African, and Asian seas (Wikipedia Commons, user Misigon; ec.europe.eu).

Figure 2. Visual presentation of an individual growth trajectory and density-dependent growth (plot A) as well as a stepwise introduction of the size-selective removals by predation (plot B) and fisheries (plot C). In all plots fish body length is plotted against the simulation time steps (T) with

the curved line depicting the asymptotic growth trajectory (full line denoting the trajectory with evolution and dashed without evolution). In plot A), each individual trajectory is defined by length at maturity (L_{mat}) and maximum theoretical length (L_{∞}). If the hake population density is high, the time left for the individual to grow (i.e., the interval between T and $T+1$) is reduced by 50% (adapted from Kuparinen et al., 2012). Plot B) demonstrates the introduction of predation, which is present throughout the entire simulation time (denoted by the green background), selecting only individuals between 21 and 51 cm of length (denoted by dotted horizontal lines). In plot C), with predation present, fishing is introduced through a narrower time period in the simulation (denoted by the red background), but targeting a range of hake length with a 50% retention at the length of 26.4 cm (L_{50} ; red dot-dashed horizontal line). The lower limit for the hake size range is set by the age of the individual (> 2) denoted in the red dotted horizontal line, which approximately corresponds to 22–23 cm of length. However, it is important to note that the actual retention by the fishing gear is described by logistic, asymptotic curve.

Figure 3. Scheme of eight tested scenarios denoted by S1–S8, as follows: S1) non-evolutionary scenario without predation, a sex-specific difference in asymptotic length (L_{∞}), as well as without evolution (reference scenarios); S2) evolutionary scenario without predation and a sex-specific difference in L_{∞} ; S3) non-evolutionary scenario without predation; S4) evolutionary scenario without predation; S5) non-evolutionary scenario with FRII predation; S6) evolutionary scenario with FRII predation; S7) non-evolutionary scenario with FRIII predation; and S8) evolutionary scenario with FRIII predation.

Figure 4. Graphs showing the average values of individual hake growth parameters, asymptotic length (a) and the intrinsic growth rate (b). The parameters are plotted against the simulation time on the x-axis, where vertical dashed lines delineate the fishing period of 30 time steps from the pre-fishing period (up to 399 time steps) and recovery period (431–600). Non-evolutionary and

988 evolutionary scenarios are denoted by dotted and bold lines, respectively. Scenarios without
 989 predation are depicted in grey, FRII in green, and FRIII in red. Scenarios without predation and a
 990 sex-specific difference in L_{∞} (reference scenarios) are presented in black.

991

992 **Figure 5.** Average values of hake individual parameters showing the age (a) and size at maturity
 993 (b), and genotypic diversity (c) with respect to each scenario. Plots a) and b) are plotted against
 994 simulation time steps where the fishing period is denoted with vertical, dashed lines (400–430). In
 995 plot c), the cumulative number (N) of juveniles is plotted against the genotype values. In all plots,
 996 the dotted lines denote the non-evolutionary scenarios, whereas the bold lines denote the
 997 evolutionary scenarios. All reference scenarios are depicted in black color, while no-predation, FRII
 998 and FRIII scenarios are depicted in grey, green and red colors, respectively.

999

1000 **Figure 6.** Hake demographic structure based on each scenario. The x-axis denotes the scenarios as:
 1001 S1) non-evolutionary scenario without predation and a sex-specific difference in L_{∞} (reference
 1002 scenario without evolution); S2) evolutionary scenario without predation and a sex-specific
 1003 difference in L_{∞} (reference scenario with evolution); S3) non-evolutionary scenario without
 1004 predation; S4) evolutionary scenario without predation; S5) non-evolutionary scenario with FRII
 1005 predation; S6) evolutionary scenario with FRII predation; S7) non-evolutionary scenario with FRIII
 1006 predation, and S8) evolutionary scenario with FRIII predation. Each bar per scenario depicts the
 1007 average size at age as well as each age category ranging from 0–12 years. Size at L_0 (first size bar)
 1008 is the same for all scenarios and set to 3.9 cm.

1009

1010 **Figure 7.** Graphs presenting the average values of hake demographic parameters for each scenario,
 1011 as follows: a) recruitment; b) population biomass with respect to carrying capacity (BM to CC); and
 1012 c) hake female to male density ratio. Simulation time steps on the x-axis are divided into pre-fishing

1013 (0–399), fishing (400–430), and recovery periods (431–600) by vertical dashed lines. Scenarios
1014 differ according to the presence or absence of evolution as bold or dotted lines, while a black color
1015 depicts reference scenarios, a grey color depicts scenarios without predation, green with FRII, and
1016 red with FRIII predation scenarios.

1017

1018 **Figure 8.** Hake and bottlenose dolphin population dynamics under fishing pressure with respect to
1019 each scenario. Average values of the following population parameters are presented: a) caught hake
1020 biomass; b) hake and bottlenose dolphin density ratio; and c) bottlenose dolphin density. Simulation
1021 time steps (x-axis) are divided by vertical dashed lines into pre-fishing (0–399), fishing (400–430),
1022 and recovery periods (431–600), although plot a) only shows the fishing period. Dotted and bold
1023 lines depict non-evolutionary and evolutionary scenarios, respectively, where a grey color denotes
1024 scenarios without predation, green FRII, and red FRIII predation scenarios. Reference scenarios are
1025 denoted in black only in plot a).

1026

1027

1028

1029

1030

1031 **Figures**

1032

1033

1034 **Figure 1.** The European hake and its native distribution range in European, African, and Asian seas
1035 (Wikipedia Commons, user Misigon; ec.europe.eu).

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

Figure 2. Visual presentation of an individual growth trajectory and density-dependent growth (plot A) as well as a stepwise introduction of the size-selective removals by predation (plot B) and fisheries (plot C). In all plots fish body length is plotted against the simulation time steps (T) with the curved line depicting the asymptotic growth trajectory (full line denoting the trajectory with evolution and dashed without evolution). In plot A), each individual trajectory is defined by length at maturity (L_{mat}) and maximum theoretical length (L_{∞}). If the hake population density is high, the time left for the individual to grow (i.e., the interval between T and $T+1$) is reduced by 50% (adapted from Kuparinen et al., 2012). Plot B) demonstrates the introduction of predation, which is present throughout the entire simulation time (denoted by the green background), selecting only individuals between 21 and 51 cm of length (denoted by dotted horizontal lines). In plot C), with predation present, fishing is introduced through a narrower time period in the simulation (denoted

by the red background), but targeting a range of hake length with a 50% retention at the length of 26.4 cm (L_{50} ; red dot-dashed horizontal line). The lower limit for the hake size range is set by the age of the individual (> 2) denoted in the red dotted horizontal line, which approximately corresponds to 22–23 cm of length. However, it is important to note that the actual retention by the fishing gear is described by logistic, asymptotic curve.

Figure 3. Scheme of eight tested scenarios denoted by S1–S8, as follows: S1) non-evolutionary scenario without predation, a sex-specific difference in asymptotic length (L_{∞}), as well as without evolution (reference scenarios); S2) evolutionary scenario without predation and a sex-specific difference in L_{∞} ; S3) non-evolutionary scenario without predation; S4) evolutionary scenario without predation; S5) non-evolutionary scenario with FRII predation; S6) evolutionary scenario with FRII predation; S7) non-evolutionary scenario with FRIII predation; and S8) evolutionary scenario with FRIII predation.

Figure 4. Graphs showing the average values of individual hake growth parameters, asymptotic length (a) and the intrinsic growth rate (b). The parameters are plotted against the simulation time on the x-axis, where vertical dashed lines delineate the fishing period of 30 time steps from the pre-fishing period (up to 399 time steps) and recovery period (431–600). Non-evolutionary and evolutionary scenarios are denoted by dotted and bold lines, respectively. Scenarios without predation are depicted in grey, FRII in green, and FRIII in red. Scenarios without predation and a sex-specific difference in L_{∞} (reference scenarios) are presented in black.

1106
1107
1108
1109
1110
1111
1112

1113 **Figure 5.** Average values of hake individual parameters showing the age (a) and size at maturity
 1114 (b), and genotypic diversity (c) with respect to each scenario. Plots a) and b) are plotted against
 1115 simulation time steps where the fishing period is denoted with vertical, dashed lines (400–430). In
 1116 plot c), the cumulative number (N) of juveniles is plotted against the genotype values. In all plots,
 1117 the dotted lines denote the non-evolutionary scenarios, whereas the bold lines denote the
 1118 evolutionary scenarios. All reference scenarios are depicted in black color, while no-predation, FRII
 1119 and FRIII scenarios are depicted in grey, green and red colors, respectively.

Figure 6. Hake demographic structure based on each scenario. The x-axis denotes the scenarios as: S1) non-evolutionary scenario without predation and a sex-specific difference in L_{∞} (reference scenario without evolution); S2) evolutionary scenario without predation and a sex-specific difference in L_{∞} (reference scenario with evolution); S3) non-evolutionary scenario without predation; S4) evolutionary scenario without predation; S5) non-evolutionary scenario with FRII predation; S6) evolutionary scenario with FRII predation; S7) non-evolutionary scenario with FRIII predation, and S8) evolutionary scenario with FRIII predation. Each bar per scenario depicts the average size at age as well as each age category ranging from 0–12 years. Size at L_0 (first size bar) is the same for all scenarios and set to 3.9 cm.

1170 **Figure 8.** Hake and bottlenose dolphin
 1171 population dynamics under fishing
 1172 pressure with respect to each scenario.
 1173 Average values of the following
 1174 population parameters are presented: a)
 1175 caught hake biomass; b) hake and
 1176 bottlenose dolphin density ratio; and c)
 1177 bottlenose dolphin density. Simulation
 1178 time steps (x-axis) are divided by

vertical dashed lines into pre-fishing (0–399), fishing (400–430), and recovery periods (431–600), although plot a) only shows the fishing period. Dotted and bold lines depict non-evolutionary and evolutionary scenarios, respectively, where a grey color denotes scenarios without predation, green FRII, and red FRIII predation scenarios. Reference scenarios are denoted in black only in plot a).