<https://helda.helsinki.fi>

Upregulation of CRISP-3 and kallikrein in stallion seminal plasma is associated with poor tolerance of cooled storage

Kareskoski, Anna Maria

2020-04

Kareskoski , A M , Palviainen , M , Johannisson , A & Katila , T 2020 , ' Upregulation of CRISP-3 and kallikrein in stallion seminal plasma is associated with poor tolerance of cooled storage ' , *Reproduction in Domestic Animals* , vol. 55 , no. 4 , pp. 496-502 . <https://doi.org/10.1111/rda.13643>

<http://hdl.handle.net/10138/324963>

<https://doi.org/10.1111/rda.13643>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

**Upregulation of CRISP-3 and kallikrein in stallion seminal plasma is associated
with poor tolerance of cooled storage**

Short title: Stallion seminal plasma proteins

Anna Maria Kareskoski^a, Mari Palviainen^b, Anders Johannisson^c, Terttu Katila^a

^aUniversity of Helsinki, Faculty of Veterinary Medicine, Department of Production
Animal Medicine, Paroninkuja 20, 04920 Saarentaus, Finland

^bUniversity of Helsinki, Faculty of Veterinary Medicine, Department of Equine and
Small Animal Medicine, Central Laboratory, P.O. Box 57, 00014 Helsinki

^cSwedish University of Agricultural Sciences, Faculty of Veterinary Medicine and
Animal Science, Box 7054, 75007 Uppsala, Sweden

Corresponding author:

Maria Kareskoski; e-mail: maria.kareskoski@helsinki.fi, address: University of
Helsinki, Faculty of Veterinary Medicine, Department of Production Animal
Medicine, Paroninkuja 20, 04920 Saarentaus, Finland

Contents

For unknown reasons, stallion fertility and sperm longevity during cooled storage of semen vary markedly between individuals. Spermatozoa from individual stallions react differently to the presence, or the removal, of seminal plasma (SP). The aim was to evaluate differences in protein content in stallion seminal plasma with either a positive or a negative effect on sperm chromatin integrity during storage. Stallion semen samples from different ejaculate fractions were stored at 5°C for 24h. Sperm survival was assessed after storage using a sperm chromatin structure assay. Protein expression in SP with either positive or negative effects on sperm survival during storage was studied using two-dimensional differential gel electrophoresis and liquid chromatography - mass spectrometry. Lower sperm chromatin integrity was associated with upregulation of the proteins kallikrein, CRISP-3, and HSP-1, while higher chromatin integrity was associated with upregulation of TIMP-2. In the sperm-rich fractions, kallikrein and CRISP-3 differed significantly between SP samples with differing effects on sperm chromatin integrity. In the sperm-poor fractions, TIMP-2 and HSP-1 differed significantly between the two SP groups. Differences in the seminal plasma proteome are associated with sperm longevity during cooled storage.

Keywords:

seminal plasma protein, spermatozoa, sperm chromatin integrity, sperm storage, stallion

Introduction

Stallion fertility and sperm longevity during cooled storage of semen vary markedly between individuals, but the reasons for this are largely unknown. The composition of

stallion seminal plasma (SP) varies between stallions and different fractions of the ejaculate (Kareskoski *et al.* 2010a, Kareskoski *et al.* 2010b). The ejaculate fractions differ in sperm concentration and semen volume, with the first few jets, the sperm-rich fractions, containing the main portion of spermatozoa. The last sperm-poor fractions consist of secretions from the seminal vesicles with low sperm concentration (Tischner *et al.* 1974, Kosiniak 1975, Varner *et al.* 1987, Magistrini *et al.* 1995).

Reducing the proportion of SP to less than 10-20% for cooled storage of semen is beneficial for sperm survival during storage (Jasko *et al.* 1991, Pruitt *et al.* 1993, Braun *et al.* 1994, Todd *et al.* 2001, Alghamdi *et al.* 2002, Love *et al.* 2005), but the effects of SP on sperm survival can vary between the fractions of the ejaculate. Most studies have reported higher sperm motility and quality in sperm-rich fractions as compared to the sperm-poor fractions (Varner *et al.* 1987, Sieme *et al.* 2004, Kareskoski *et al.* 2011).

Variation between stallions in the effects of SP on spermatozoa is evident. It has been reported that spermatozoa from individual stallions react differently to the presence, or the removal, of SP (Padilla & Foote 1991, Jasko *et al.* 1992, Akcay *et al.* 2006).

Exchange studies, where spermatozoa from one stallion were stored in another stallion's SP, have also been carried out. Seminal plasma from some stallions seem to have a less negative, or even a beneficial, effect on sperm survival during storage (Aurich *et al.* 1996). It has been recommended that stallions producing semen with poor tolerance to cooling and storage should be evaluated for the effects of their SP on semen quality, and the semen processing technique modified accordingly (Brinsko *et al.* 2000, Love *et al.* 2005).

Stallion SP contains several different types of proteins. The main SP proteins are proteins carrying fibronectin type II (Fn-2) modules, cysteine-rich secretory proteins (CRISPs), and spermadhesins. The suggested functions of SP proteins include involvement in several essential steps preceding fertilization, such as regulating capacitation, establishment of the oviductal sperm reservoir, modulation of the uterine immune response, sperm transport in the female genital tract, and gamete interaction and fusion (reviewed by Töpfer-Petersen *et al.* 2005).

The aim of this study was to evaluate protein expression in stallion SP with either positive or negative effects on sperm survival during storage. Two-dimensional differential gel electrophoresis and liquid chromatography - mass spectrometry was used for proteomic analysis and protein identification.

Materials and methods

Animals

Semen was collected during one breeding season from 43 stallions (ages 4 to 23 years) residing at nine stud farms in Finland. Twenty stallions were Standardbred trotters (ages 4 - 23 years), 17 Finnhorses (ages 8 - 23 years), four warmblood riding horses (ages 9 - 19 years) and two were ponies (ages 8 and 9 years). The breeding history and current use of the stallions was variable, with the stallions serving from 2 to 150 (median 14) mares during the study season. Most of the stallions (33 stallions) were used for collection of both fresh and transported semen, and four stallions were also used for natural breeding. The eight stallions that were selected for proteomic

analyses were aged 10 to 23 years, and were used for commercial breeding regularly during the year of the study. Ethical approval was not required for this study according to the Finnish Act on the Protection of Animals Used for Scientific or Educational Purposes (497/2013).

Semen collection

The ejaculatory jets were collected as three to four fractions using one of the three types of artificial vaginas (AV): an open-ended artificial vagina equipped with a collection bag attached to a funnel (Krakow AV; Tischner et al. 1974) a Missouri AV with a collection bag (Kareskoski *et al.* 2011) or a computer-controlled fractioning phantom with an integrated AV (Equidame phantom, Haico Oy, Loimaa, Finland; Lindeberg et al. 1999), depending on stallion preference and stud farm. To provide comparable samples, fractions with the highest and lowest sperm concentrations within each ejaculate were included in the statistical analyses. The mean concentrations for the fractions were 300.0×10^6 and 54.0×10^6 sperm/ml, respectively, with no significant differences in sperm concentration in fractions collected with the different methods. Of the stallions selected for proteomic analyses, all were collected using the open-ended AV, except for stallion E, which was collected with the modified closed AV.

Semen processing and storage

After semen collection, any gel was removed if present, and the volume of each fraction measured. The sperm concentration in each fraction was determined using a Bürker counting chamber. For the sperm chromatin structure assay (SCSA), a sample of $2 - 10 \times 10^6$ spermatozoa from each fraction of raw semen was pipetted into

120 cryovials and TNE-buffer (9.48 g Tris-HCl, 52.6 NaCl, 2.23 g disodium-EDTA, aqua
121 ster. ad 600 mL, pH 7.4) was added (ad 1.5 mL in each vial). The SCSA samples were
122 placed in liquid nitrogen vapor (3cm above the liquid surface) for 10min, and then
123 plunged into liquid nitrogen.

124
125 Each fraction was divided into two parts: one-half was centrifuged for the preparation
126 of SP (SP samples), and the other half was processed for cooled storage (semen
127 samples). A portion (10%) of the volume of each fraction was combined to form a
128 sample representing the whole ejaculate (WE). For cooled storage, semen samples
129 were extended in a semen:extender ratio of 1:1 using skim milk extender (Kenney *et*
130 *al.* 1975). Each fraction of the semen samples was divided into two centrifuge tubes
131 and centrifuged at 500 x g (10 min). After removal of the major part (about 5% was
132 left) of the supernatant (i.e. extender and SP), the sperm pellet was extended in either:
133 a) skim milk extender only (these samples stored without SP were named SP0
134 samples), or b) a combination of supernatant and skim milk extender (these samples
135 stored with SP were named SP1). The final sperm concentration was 50×10^6
136 sperm/ml. The final SP to extender ratio in the SP1 samples was 1:2. The semen
137 samples were stored in 1-mL vials at 5°C for 24h. After the 24-h cooled storage, $5 \times$
138 10^6 spermatozoa from each semen sample were pipetted into cryovials for SCSA, and
139 TNE-buffer was added (ad 1.5 mL in each vial). The samples were placed in liquid
140 nitrogen vapor (3cm above the liquid surface) for 10min and plunged into liquid
141 nitrogen.

142
143 The SP samples were centrifuged at 4000 x g (15 min), and the supernatant was
144 filtered using 0.45- μ m filters (Millex-HV, Millipore, Billerica, MA, USA). Protease

inhibitor (Trasylol, 10000 Kallikrein Inactivator Units (KIU)/mL, Bayer Schering Pharma, Berlin, Germany) was added to each sample (500 KIU/mL of SP) to decrease proteolytic activity on enzymes and other proteins. The SP samples were stored frozen in 1-mL aliquots in -75°C until analyzed.

Sperm chromatin structure assay

Sperm chromatin stability of sperm from both the raw and stored semen samples was measured as the susceptibility of sperm DNA to denaturation using the sperm chromatin structure assay (SCSA) as described by Evenson *et al.* (1999). Acridine orange was added to each thawed sample after exposure to an acid detergent solution that induces partial DNA denaturation *in situ*. Flow cytometry was then used to measure either red (abnormal sperm with denatured DNA) or green (normal sperm) fluorescence). Sperm DNA fragmentation index (DFI) was assessed by calculating the amount of red fluorescence divided by the total (red plus green) fluorescence, indicating the amount of denatured sperm DNA over the total DNA in each sperm cell.

Selection of samples for proteomic analyses according to sperm survival during cooled storage with or without seminal plasma

The samples of eight stallions were chosen for the proteomic analyses based on the effect of SP on sperm chromatin integrity during storage, represented here by the difference in DFI (diff-DFI) between semen samples stored with SP and the samples stored without SP (diff-DFI = SP1 – SP0). The samples with the lowest and the highest diff-DFI (-5.9 to +0.8% and +10.6 to +13.5%, respectively) of all the stallions were selected, and the samples were grouped into two groups according to the diff-

DFI. The effect of SP on sperm chromatin integrity was negative in the samples with high diff-DFI (group HIGH DIFF), and the effect of SP was positive or neutral in the samples with low diff-DFI (group LOW DIFF). The diff-DFI values varied significantly between the groups according to Student's t-test ($p=0.008$) (Table 1).

Sample preparation for proteomic analysis and CyDye labelling

Seminal plasma samples were prepared to proteomic analysis by precipitation with 2D Clean Up-kit and solubilized in 50 μ l of labeling buffer (7 M Urea, 2 M Thiourea, 4 % CHAPS, 30 mM Tris). The concentration of proteins in the samples was measured by using 2D Quant Kit according to manufacturer's instructions. The samples were then labeled with Cy2, Cy3 and Cy5 dyes (CyDye DIGE Fluor minimal dyes) according to Ettan two-dimensional difference gel electrophoresis (DIGE) protocol. Briefly, 50 μ g of protein from each sample were labeled with 400 pmol of the Cy3 and Cy5 dyes. Equal amount aliquots of each sample were combined to establish internal standard labeled with Cy2 dye. The labeling reaction was incubated 30 minutes on ice in the dark and halted by adding of 1 mM lysine to the reaction following 10 minutes incubation as earlier. All materials and equipment by GE Healthcare, Piscataway, NJ, USA.

Two-dimensional difference gel electrophoresis

The labeled proteins were analyzed by two-dimensional difference gel electrophoresis (2D-DIGE) as described earlier (Ünlü *et al.* 1997). An immobilized pH gradient (IPG) strip (24 cm, pH 3-10, nonlinear, GE Healthcare, Piscataway, NJ, USA) was used for isoelectric focusing. IPG strips were loaded with 150 μ g of protein in total by using cup-loading method. Isoelectric focusing was performed using IPGPhor (GE

Healthcare, Piscataway, NJ, USA) at 20°C as follows: 3h at 150 V, 3h at 300 V, then linear ramping to 10 000 V and 10 000 V for 50 000 Vh with maximum current per strip being 75 µA. After focusing the isoelectric strips were prepared for the second dimension gels by incubation in I (50 mM Tris pH 8.8, 6 M urea, 30% glycerol, 2% sodium dodecyl sulphate (SDS), 0.2% bromophenol blue, with added 10mg/ml dithiothreitol (DTT)) equilibrium buffer solution for 15 minutes. This was followed by equilibration in a II (50 mM Tris pH 8.8, 6 M urea, 30% glycerol, 2% SDS, 0.2% bromophenol blue, supplemented with 25mg/ml iodoacetamide) buffer solution for another 15 minutes. The prepared IPG strips were then placed on 12% sodium dodecyl sulphate polyacrylamide gels (SDS-PAGE) and sealed with overlay agarose (Bio-Rad, Hercules, CA, USA). Electrophoresis was initiated at 50 V for 30 minutes, which was followed by 400 V for 3 hours. The gels were scanned between low-fluorescence glass plates using an FLA-5100 laser scanner (Fujifilm, Tokyo, Japan) at wavelengths 473 (for Cy2), 532 (for Cy3) and 635 nm (Cy5) using voltages of 420, 410 and 400 V. After scanning the gels were stained by PlusOne silver staining kit (GE Healthcare, Piscataway, NJ, USA) without aldehyde.

Image analysis and data processing

The gel images were analyzed and statistically assessed using DeCyder 7.0 software (GE Healthcare, Piscataway, NJ, USA). First the gels were automatically analyzed using the batch processor function to normalize the Cy2, Cy3 and Cy5 images from each gel. Spot volumes were calculated and compared to Cy2 volumes (internal standard) to correct the inter-gel variations. In the biological variation module, the Cy2 images of all gels in groups were matched and the spot volumes compared. Approximately 1000 separate spots were detected on each gel. Protein spots

demonstrating a minimum of 1.5-fold difference in average spot volume ratios between groups using Student's *t* test (*p*-value less than 0.05) were chosen for identification.

Protein identification

Protein spots of interest were manually excised from the gel and digested in-gel using trypsin (Trypsin Gold, Promega, Madison, WI, USA) as described earlier (Shevchenko *et al.* 1996; Jensen *et al.* 1998). The peptides were extracted, vacuum dried (Vacufuge Plus vacuum concentrator, Eppendorf, Hamburg, Germany), centrifuged (Eppendorf microcentrifuge, Eppendorf, Hamburg, Germany) and re-dissolved in 20 µl 5% formic acid. Digested peptides were de-salted using C18 Zip-Tips (Millipore, Burlington, MA, USA) according to the manufacturer's instructions. The resulting peptides were identified by fragment ion analysis with LC-MS/MS using Ultimate 3000 nano-LC system (Dionex, Sunnyvale, CA, USA) and QSTAR Elite hybrid quadrupole TOF mass spectrometer (Applied Biosystems/MDS Sciex, Foster City, CA, USA) with nano-ESI ionization. The identification of proteins was performed using the local Mascot version 2.2 (Matrix Science, London, UK) against the in-house database with the following criteria: one missed cleavage allowed, carbamidomethyl modification of cysteine as a fixed modification, and oxidation of methionine as a variable modification.

Results

Approximately 1000 different proteins were expressed in the 2D-DIGE analysis. Only four proteins were significantly (*p*<0.05) differentially expressed in SP of the different storage groups (SP0 and SP1). The identified proteins were tissue inhibitor

of metalloproteinase-2 (TIMP-2), horse seminal protein-1 (HSP-1), kallikrein, and cysteine-rich secretory protein-3 (CRISP-3).

In the sperm-rich fractions, kallikrein and CRISP-3 were upregulated in SP samples with high diff-DFI (i.e. a low rate of sperm survival during storage with SP). In the sperm-poor fractions, TIMP-2 was upregulated in SP samples with low diff-DFI (i.e. a high rate of sperm survival during storage), and HSP-1 was upregulated in samples with high diff-DFI (Table 2).

Discussion

Poor sperm survival during cooled storage in the presence of SP was associated with upregulation of the SP proteins kallikrein, CRISP-3, and HSP-1, while better sperm survival was associated with upregulation of TIMP-2. Sperm survival was evaluated using a chromatin structure assay, which has been shown to be correlated with sperm survival and fertility in stallions (Love *et al.* 2005, Johannisson *et al.* 2018). The effects of SP on sperm motility can differ from those on sperm chromatin integrity, and SP can damage chromatin markedly while motility is maintained (Morrell *et al.* 2012). In this study, the stallions were categorized according to diff-DFI alone, because of the correlation between fertility and sperm chromatin integrity (Love 2005, Morrell *et al.* 2008).

Increased levels of CRISP-3 were associated with high diff-DFI, which reflects poor sperm survival during storage when SP is present. This is not necessarily directly linked to decreased fertility, which is a multifactorial trait. Most studies on CRISP-3 in stallion semen associate upregulation of the protein with increased fertility,

270 whereas in our study upregulation was associated with decreased sperm survival
271 during storage. Based on genomic studies, CRISP-3 is associated with stallion fertility
272 and semen quality traits (Hamann *et al.* 2007, Novak *et al.* 2010, and Gottschalk *et al.*
273 2016). In raw semen, high levels of CRISP-3 in SP result in higher sperm motility and
274 improved semen quality index (Restrepo *et al.* 2019), but in our study, higher levels
275 of CRISP-3 decreased sperm longevity during storage.

276
277 Proteins that are correlated to fertility in raw or frozen semen may not be associated
278 with sperm longevity during cooled storage. During storage, proteolytic changes in SP
279 may alter the protein composition, as SP contains several different proteases and
280 protease inhibitors (Laflamme & Wolfner 2013). At present, the magnitude and
281 significance of these proteolytic alterations are not known. The relationship between
282 SP proteins and sperm survival during cooled storage has not previously been studied,
283 except for our earlier study on SP proteins and cooled semen storage for 24h where
284 we found no significant correlation between the level of HSP-1 and sperm motility
285 parameters (Kareskoski *et al.* 2011). The protein HSP-1 has been shown to protect
286 other proteins against thermal and chemical stress conditions (Sankhala *et al.* 2012),
287 and to prevent lipid peroxidation under oxidative stress (Kumar & Swamy 2016). In
288 our study, the levels of HSP-1 itself were higher in samples with poor sperm survival.

289
290 The mechanisms behind the association of these proteins with semen quality,
291 freezability and fertility are largely unknown. The CRISP-3 protein is an androgen-
292 regulated protein expressed only in postpubertal horses (Fedorka *et al.* 2017).
293 Associations between this protein and semen quality traits are not necessarily direct
294 and causal, but rather indirect correlations of androgen-dependent traits. A possible

role of CRISP-3 in the regulation of oxidative pathways in semen has been proposed (Restrepo *et al.* 2019), based on the high content of cysteine residues in CRISP-3 (Udby *et al.* 2005) and the contribution of these residues to oxidative reactions (Requejo *et al.* 2010). In human semen, seminal vesicular fluid makes SP a zinc-binding medium after ejaculation. This can deplete spermatozoa of zinc and affect sperm chromatin decondensation (Björndahl & Kvist 2010), and CRISP proteins could also enhance the appearance of DNA lesions in sperm (Rodrigues *et al.* 2016). This could explain the association between CRISP-3 and DNA injuries shown in our study.

Kallikrein was associated with high diff-DFI and DNA injuries during cooled storage. It is a protease similar to prostate-specific antigen (PSA) produced by the prostate gland in humans. Kallikrein and HSP-1 have earlier been shown to be negatively correlated to freezability (Jobim *et al.* 2011) and pregnancy rate, especially at low sperm concentrations and low dilution rates, when the amount of SP is high in relation to the number of spermatozoa. It was suggested that these proteins bind to spermatozoa and reduce their ability to fertilize the ovum (Novak *et al.* 2010). In our study, the negative effects of kallikrein on sperm chromatin integrity were found in the sperm-rich fraction of the ejaculate.

Both matrix metalloproteinases (MMPs) and their inhibitors (TIMPs) are found in SP (Baumgart *et al.* 2002). In our study, less sperm DNA damage during storage was associated with higher levels of TIMP-2. Similarly, high levels of TIMP-2 are associated with low levels of sperm DNA fragmentation in men (Belardin *et al.* 2019) and high fertility in bulls (McCauley *et al.* 2001, Viana *et al.* 2018).

320

321 In the sperm-rich fractions, kallikrein and CRISP-3 differed significantly between
322 HIGH DIFF and LOW DIFF stallions. In the sperm-poor fractions, TIMP-2 and HSP-
323 1 differed significantly between these groups. The composition of these fractions
324 differs to some extent, but in our earlier study, there were no significant differences in
325 the amount of HSP-1 or CRISP-3 between fractions (Kareskoski *et al.* 2010b).
326 However, both sperm chromatin integrity (Björndahl & Kvist 2010) and the function
327 of some SP proteins, such as kallikrein and CRISP-3 (Carvalho *et al.* 2002,
328 Schambony *et al.* 1998), are Zn-dependent, and hence the levels of Zn may affect the
329 observed fraction differences. The levels of Zn in stallion ejaculates are correlated to
330 sperm concentration (Pesch *et al.* 2006), which suggests that the seminal vesicles,
331 contributing mostly to the sperm-poor fractions of the ejaculate (Magistrini *et al.*
332 1995), are not the main source of Zn in stallion SP. In primates, Zn is mostly derived
333 from the prostate gland, and is therefore found in the sperm-rich fractions (Srivastava
334 *et al.* 1984). The Zn-regulated proteins kallikrein and CRISP-3 were significant for
335 sperm chromatin integrity in only the sperm-rich fraction in our study.

336

337 **Conclusions**

338 Poor sperm survival during cooled storage when SP was present was associated with
339 upregulation of the SP proteins kallikrein, CRISP-3, and HSP-1, while better sperm
340 survival was associated with upregulation of TIMP-2.

341

342 **Conflict of interest statement**

343 None to declare.

344

Data availability statement

The data that support the findings of this study are available upon reasonable request from the corresponding author. The data are not publicly available due to privacy restrictions, e.g. their containing information that could compromise the privacy of research participants.

References

- Akcay, E., Reilas, T., Andersson, M., & Katila, T. (2006). Effect of seminal plasma fractions on sperm survival after cooled storage. *Journal of Veterinary Medicine*, 53, 481-485. *doi: 10.1111/j.1439-0442.2006.00882.x*
- Alghamdi, A. S., Troedsson, M. H., Xue, J. L., & Crabo, B. G. (2002). Effect of seminal plasma concentration and various extenders on postthaw motility and glass wool-Sephadex filtration of cryopreserved stallion semen. *American Journal of Veterinary Research*, 63, 880-885. *doi: 10.2460/ajvr.2002.63.880*
- Aurich, J.E., Kühne, A., Hoppe, H., & Aurich, C. (1996). Seminal plasma affects membrane integrity and motility of equine spermatozoa after cryopreservation. *Theriogenology*, 46, 791-797. *doi: 10.1016/s0093-691x(96)00237-3*
- Baumgart, E., Lenk, S. V., Loening, S. A., & Jung, K. (2002). Quantitative differences in matrix metalloproteinase (MMP)-2, but not in MMP-9, tissue inhibitor of metalloproteinase (TIMP)-1 or TIMP-2, in seminal plasma of normozoospermic and azoospermic patients. *Human Reproduction*, 17, 2919-2923. *doi: 10.1093/humrep/17.11.2919*

370

371 Belardin, L. B., Antoniassi, M. P., Camargo, M., Intasqui, P., Fraietta, R., & Bertolla,
372 R. P. (2019). Semen levels of matrix metalloproteinase (MMP) and tissue inhibitor of
373 metalloproteinases (TIMP) protein families members in men with high and low sperm
374 DNA fragmentation. *Scientific Reports*, 29, 903. *doi:* 10.1038/s41598-018-37122-4

375

376 Björndahl, L., & Kvist, U. (2010). Human sperm chromatin stabilization: a proposed
377 model including zinc bridges. *Molecular Human Reproduction*, 16, 23-29.
378 *doi:* 10.1093/molehr/gap099

379

380 Braun, J., Sakai, M., Hochi, S., & Oguri, N. (1994). Preservation of ejaculated and
381 epididymal stallion spermatozoa by cooling and freezing. *Theriogenology*, 41, 809-
382 818. *doi:* 10.1016/0093-691x(94)90497-7

383

384 Brinsko, S. P., Crockett, E. C., & Squires, E. L. (2000). Effect of centrifugation and
385 partial removal of seminal plasma on equine spermatozoal motility after cooling and
386 storage. *Theriogenology*, 54, 129-136. *doi:* 10.1016/S0093-691X(00)00331-9

387

388 Doty, A., Buhi, W. C., Benson, S., Scoggin, K. E., Pozor, M., Macpherson, M., Mutz,
389 M., & Troedsson, M. H. T. (2011). Equine CRISP3 modulates interaction between
390 spermatozoa and polymorphonuclear neutrophils. *Biology of Reproduction*, 85, 157-
391 164. *doi:* 10.1095/biolreprod.110.084491

392

393 Evenson, D. P., Jost, L. K., Marshall, D., Zinaman, M. J., Clegg, E., Purvis, K., de
394 Angelis, P., & Claussen, O. P. (1999). Utility of the sperm chromatin structure assay

395 as a diagnostic and prognostic tool in the human fertility clinic. *Human Reproduction*,
396 14, 1039-49. *doi:* 10.1093/humrep/14.4.1039

397

398 Fedorka, C. E., Scoggin, K. E., Squires, E. L., Ball, B. A., & Troedsson, M. H. T.
399 (2017). Expression and localization of cysteine-rich secretory protein-3 (CRISP-3) in
400 the prepubertal and postpubertal male horse. *Theriogenology*, 87, 187-192.
401 *doi:* 10.1016/j.theriogenology.2016.08.027

402

403 Gottschalk, M., Metzger, J., Martinsson, G., Sieme, H., & Distl, O. (2016). Genome-
404 wide association study for semen quality traits in German Warmblood stallions.
405 *Animal Reproduction Science*, 171, 81-86. *doi:* 10.1016/j.anireprosci.2016.06.002

406

407 Hamann, H., Jude, R., Sieme, H., Mertens, U., Töpfer-Petersen, E., Distl, O., & Leeb,
408 T. (2007). A polymorphism within the equine CRISP3 gene is associated with stallion
409 fertility in Hanoverian warmblood horses. *Animal Genetics*, 38, 259-264.
410 *doi:* 10.1111/j.1365-2052.2007.01594.x

411

412 Jasko, D. J., Moran, D. M., Farlin, M. E., & Squires, E. L. (1991). Effect of seminal
413 plasma dilution or removal on spermatozoa motion characteristics of cooled semen.
414 *Theriogenology*, 35, 1059-1067.

415

416 Jasko, D. J., Hathaway, V. L., Schaltenbrand, V. L., Simper, W. D., & Squires, E. L.
417 (1992). Effect of seminal plasma and egg yolk on motion characteristics of cooled
418 stallion spermatozoa. *Theriogenology*, 37, 1241-1252.

419

420 Jensen, O. N., Larsen, M. R., & Roepstorff, P. (1998). Mass spectrometric
421 identification and micro characterization of proteins from electrophoretic gels:
422 strategies and applications. *Proteins: Structure, Function, and Bioinformatics*, 33, 74–
423 89.

424

425 Jobim, M. I., Trein, C., Zirkler, H., Gregory, R. M., Sieme, H., & Mattos, R. C.
426 (2011). Two-dimensional polyacrylamide gel electrophoresis of equine seminal
427 plasma proteins and their relation with semen freezability. *Theriogenology*, 76, 765-
428 71. *doi: 10.1016/j.theriogenology.2011.04.010*

429

430 Johannisson, A., Figueiredo, M. I., Al-Kass, Z., & Morrell, J. M. (2018).
431 Simultaneous evaluation of superoxide content and mitochondrial membrane potential
432 in stallion semen samples provides additional information about sperm quality.
433 *Animal Reproduction Science*, 192, 290-297. *doi: 10.1016/j.anireprosci.2018.03.030*

434

435 Kareskoski, A. M., Reilas, T., Sankari, S., Andersson, M., Güvenc, K., & Katila, T.
436 (2010a). Alkaline and acid phosphatase, β -glucuronidase, and electrolyte levels in
437 fractionated stallion ejaculates. *Reproduction in Domestic Animals*, 45, e369-e374.
438 *doi: 10.1111/j.1439-0531.2009.01579.x*

439

440 Kareskoski, A. M., Rivera del Alamo, M. M., Güvenc, K., Reilas, T., Calvete, J. J.,
441 Rodriguez-Martinez, H., Andersson, M., & Katila, T. (2010b). Protein composition of
442 seminal plasma in fractionated stallion ejaculates. *Reproduction in Domestic Animals*,
443 46, e79-e84. *doi: 10.1111/j.1439-0531.2011.01789.x*

444

445 Kareskoski, A. M., Sankari, S., Johannisson, A., Kindahl, H., Andersson, M., &
 446 Katila, T. (2011). The association of the presence of seminal plasma and its
 447 components with stallion fertility and sperm longevity in fractionated ejaculates.
 448 *Reproduction in Domestic Animals*, 46, 1073-1081. *doi*: 10.1111/j.1439-
 449 0531.2011.01789.x
 450
 451 Kenney, R. M., Bergman R. V., Cooper, W. L., & Morse, G. W. (1975). Minimal
 452 contamination techniques for breeding mares: techniques and preliminary findings.
 453 *Proceedings of the 21st Annual Convention of the American Association of Equine*
 454 *Practitioners*, 327-336.
 455
 456 Kosiniak, K. (1975). Characteristics of the successive jets of ejaculated semen of
 457 stallions. *Journal of Reproduction and Fertility Supplement*, 23, 59-61.
 458
 459 Kumar, C. S., & Swamy, M. J. (2016). HSP-1/2, a major horse seminal plasma
 460 protein, acts as a chaperone against oxidative stress. *Biochemical and Biophysical*
 461 *Research Communications*, 473, 1058-1063. *doi*: 10.1016/j.bbrc.2016.04.015
 462
 463 Laflamme, B. A., & Wolfner, M. F. (2013). Identification and function of proteolysis
 464 regulators in seminal fluid. *Molecular Reproduction and Development*, 80, 80-101.
 465 *doi*: 10.1002/mrd.22130
 466
 467 Lindeberg, H., Karjalainen, H., Koskinen, E., Katila, T. (1999). Quality of stallion
 468 semen obtained by a new semen collection phantom (Equidame) versus a Missouri

469 artificial vagina. *Theriogenology*, 51, 1157-1173. *doi:* 10.1016/S0093-
470 691X(99)80019-3
471
472 Love, C. C., Brinsko, S. P., Rigby, S. L., Thompson, J. A., Blanchard, T. L., &
473 Varner, D. D. (2005). Relationship of seminal plasma level and extender type to
474 sperm motility and DNA integrity. *Theriogenology*, 63, 1584-1591.
475 *doi:* 10.1016/j.theriogenology.2004.05.030
476
477 Love, C. C. (2005). The sperm chromatin structure assay: a review of clinical
478 applications. *Animal Reproduction Science*, 89, 39-45.
479 *doi:*10.1016/j.anireprosci.2005.06.019
480
481 Magistrini, M., Lindeberg, H., Koskinen, E., Beau, P., & Seguin, F. (2000).
482 Biophysical and ¹H magnetic resonance spectroscopy characteristics of fractionated
483 stallion ejaculates. *Journal of Reproduction and Fertility Supplement*, 56, 101-110.
484
485 McCauley, T. C., Zhang, H. M., Bellin, M. E., & Ax, R. L. (2001). Identification of a
486 heparin-binding protein in bovine seminal fluid as tissue inhibitor of
487 metalloproteinases-2. *Molecular Reproduction and Development*, 58, 336-341.
488 *doi:* 10.1002/1098-2795(200103)58:3<336::AID-MRD12>3.0.CO;2-Z
489
490 Morrell, J. M., Johannisson, A., Dalin, A. M., Hammar, L., Sandebert, T., Rodriguez-
491 Martinez, H. (2008). Sperm morphology and chromatin integrity in Swedish
492 warmblood stallions and their relationship to pregnancy rates. *Acta Veterinaria*
493 *Scandinavica*, 50, 2-8. *doi:* 10.1186/1751-0147-50-2.

494 Morrell, J. M., Pihl, J., Dalin, A. M., Johannisson, A. (2012). Restoration of seminal
495 plasma to stallion spermatozoa selected by colloid centrifugation increases sperm
496 progressive motility but is detrimental to chromatin integrity. *Theriogenology*, 78,
497 345-52. doi: 10.1016/j.theriogenology.2012.02.009.

498

499 Novak, S., Smith, T. A., Paradis, F., Burwash, L., Dyck, M. K., Foxcroft, G. R., &
500 Dixon, W. T. (2010). Biomarkers of in vivo fertility in sperm and seminal plasma of
501 fertile stallions. *Theriogenology*, 74, 956-67.
502 doi:10.1016/j.theriogenology.2010.04.025

503

504 Padilla, A. W., & Foote, R. H. (1991). Extender and centrifugation effects on the
505 motility patterns of slow-cooled stallion spermatozoa. *Journal of Animal Science*, 69,
506 3308-3313. doi: 10.2527/1991.6983308x

507

508 Pesch, S., Bergmann, M., & Bostedt, H. (2006). Determination of some enzymes and
509 macro- and microelements in stallion seminal plasma and their correlations to semen
510 quality. *Theriogenology*, 66, 307-313. doi: 10.1016/j.theriogenology.2005.11.015

511

512 Pruitt, J. A., Arns, M. J., & Pool, K. C. (1993). Seminal plasma influences recovery of
513 equine spermatozoa following in vitro culture (37 °C) and cold-storage (5 °C).
514 *Theriogenology*, 39, 291.

515

516 Requejo, R., Hurd, T. R., Costa, N. J., & Murphy, M. P. (2010). Cysteine residues
517 exposed on protein surfaces are the dominant intramitochondrial thiol and may protect

518 against oxidative damage. *FEBS Journal*, 277, 1465-1480. *doi*: 10.1111/j.1742-

519 4658.2010.07576.x

520

521 Restrepo, G., Rojano, B., & Usuga, A. (2019). Relationship of cysteine-rich secretory

522 protein-3 gene and protein with semen quality in stallions. *Reproduction in Domestic*

523 *Animals*, 54, 39-45. *doi*: 10.1111/rda.13309

524

525 Rodrigues, M. F., Trentin, J. M., Boligon de Araujo, Centeno, L. A. M., Schenatto, R.

526 O., Pessoa, G. A., Neves, A. P., & Rubin, M. I. B. (2016). Seminal plasma: Effect on

527 motility, membrane functionality, and spermatid chromatin dispersion of equine

528 sperm treated with N-acetyl-L-cysteine at 5°C. *Acta Scientiae Veterinariae*, 44, 1371.

529

530 Sankhala, R. S., Kumar, C. S., Singh, B. P., Arangasamy, M. J., & Swamy, M. J.

531 (2012). HSP-1/2, a major protein of equine seminal plasma, exhibits chaperone-like

532 activity. *Biochemical and Biophysical Research Communications*, 427, 18-23.

533 *doi*: 10.1016/j.bbrc.2012.08.120

534

535 Shimokawa, K., Katayama, M., Matsuda, Y., Takahashi, H., Hara, I., Sato, H., &

536 Kaneko, S. (2002). Matrix metalloproteinase (MMP)-2 and MMP-9 activities in

537 human seminal plasma. *Molecular Human Reproduction*, 8, 32-36.

538 *doi*: 10.1093/molehr/8.1.32

539

540 Shevchenko, A., Wilm, M., Vorm, O., & Mann, M. (1996). Mass spectrometric

541 sequencing of proteins from silver-stained polyacrylamide gels. *Analytical Chemistry*

542 68, 850–858. *doi*: 10.1021/ac950914h

543

544 Sieme, H., Katila, T., & Klug, E. (2004). Effect of semen collection practices on
545 sperm characteristics before and after storage and on fertility of stallions.

546 *Theriogenology*, 61, 769-784. *doi*: 10.1016/s0093-691x(03)00251-6

547

548 Srivastava, A., Chowdhury, A. R., & Setty, B. S. (1984). Zinc content in the
549 epididymis, vas deferens, prostate, and seminal vesicles of juvenile rhesus monkey
550 (*Macaca mulatta*): effect of androgen and estrogen. *Prostate*, 5, 153-8.

551 *doi*: 10.1002/pros.2990050205

552

553 Tischner, M., Kosiniak, K., & Bielanski, W. (1974). Analysis of pattern of ejaculation
554 in stallions. *Journal of Reproduction and Fertility*, 41, 329-335.

555 *doi*: 10.1530/jrf.0.0410329

556

557 Todd, P., Arns M. J., Chenoweth, P., & Schultz, B. (2001). Influence of seminal
558 plasma and processing on cold-stored stallion spermatozoa. *Animal Reproduction*
559 *Science*, 68, 335-336.

560

561 Troedsson, M. H. T., Doty, A., Macpherson, M. L., Connor, M. C., Verstegen, J. P.,
562 Pozor, M. A., Buhi, W. C. (2010). CRISP-3 in equine seminal plasma is involved in
563 selective uterine sperm transport. *Animal Reproduction Science*, 121, 192-193.

564

565 Töpfer-Petersen, E., Ekhlasi-Hundrieser, M., Kirchhoff, C., Leeb, T., & Sieme, H.
566 (2005). The role of stallion seminal plasma proteins in fertilisation. *Animal*

567 *Reproduction Science*, 89, 159-170. *doi*: 10.1016/j.anireprosci.2005.06.018

568

569 Udby, L., Calafat, J., Sørensen, O.E., Borregaard, N., & Kjeldsen, L. (2002).

570 Identification of human cysteine-rich secretory protein 3 (CRISP-3) as a matrix

571 protein in a subset of peroxidase-negative granules of neutrophils and in the granules

572 of eosinophils. *Journal of Leukocyte Biology*, 72, 462-9. *doi*:10.1189/jlb.72.3.462

573

574 Udby, L., Bjartell, A., Malm, J., Egesten, A., Lundwall, K., Cowland, J., & Kjeldsen,

575 I. (2005). Characterization and localization of cysteine-rich secretory protein 3

576 (CRISP-3) in the human male reproductive tract. *Journal of Andrology*, 26, 333-342.

577 *doi*: 10.2164/jandrol.04132

578

579 Ünlü, M., Morgan, M. E., & Minden, J. S. (1997). Difference gel electrophoresis: a

580 single gel method for detecting changes in protein extracts. *Electrophoresis*, 18, 2071-

581 7. *doi*: 10.1002/elps.1150181133

582

583 Usuga, A., Rojano, B., & Restrepo, G. (2018). Association of the cysteine-rich

584 secretory protein-3 (CRISP-3) and some of its polymorphisms with the quality of

585 cryopreserved stallion semen. *Reproduction, Fertility and Development*, 30, 563-

586 569. *doi*: 10.1071/RD17044

587

588 Varner, D. D., Blanchard, T. L., Love, C. C., Garcia, M. C., & Kenney, R. M. (1987).

589 Effects of semen fractionation and dilution ratio on equine spermatozoa motility

590 parameters. *Theriogenology*, 28, 709-723. *doi*: 10.1016/0093-691x(87)90288-3

591

592

593 Viana, A. G. A., Martins, A. M. A., Pontes, A. H., Fontes, W., Castro, M. S., Ricart,
594 C. A. O., Sousa, M. V., Kaya, A., Topper, E., Memili, E., & Moura, A. A. (2018).

595 Proteomic landscape of seminal plasma associated with dairy bull fertility. Scientific

596 Reports, 8, 16323. *doi:* 10.1038/s41598-018-34152-w

597

598

599

600

Table 1. DNA fragmentation index (DFI) in semen samples stored at 5°C for 24h with or without seminal plasma (SP). Difference in DFI = (DFI when stored with SP) – (DFI when stored without SP).

Stallion	DFI (%) before storage	DFI (%) after storage		
		Stored without SP	Stored with SP	Difference in DFI
A	10.3	11.8	5.8	-5.9
B	22.7	29.7	28.5	-1.2
C	11.1	13.5	13.5	-0.01
D	17.6	25.0	25.8	0,8
E	17.3	21.6	32.2	10.6
F	22.4	30.6	41.6	11.0
G	11.1	7.6	19.5	11.9
H	35.8	33.1	46.5	13.5

606 Table 2. Seminal plasma proteins with significant (p=0.05) upregulation in samples with a low rate of sperm survival during cooled storage with
607 seminal plasma (evaluated as changes in sperm chromatin integrity), compared with samples with a high rate of sperm survival during cooled
608 storage. (CRISP-3 = cysteine-rich secretory protein-3; TIMP-2 = tissue inhibitor of metalloproteinase-2; HSP-1 = horse seminal protein-1; pI =
609 isoelectric point).

Gel ID	Identified protein	Accession number	pI	Molecular weight (kDa)	Sequence coverage (%)	MASCOT score	Upregulation in
Seminal plasma in sperm-rich fractions							
6	Kallikrein	18250639	5.4	21.4	13.1	71	Low sperm survival
5	CRISP-3	3023562	7.4	27.0	21.2	111	Low sperm survival
Seminal plasma in sperm-poor fractions							
2	TIMP-2	135853	7.4	24.0	7.3	60	High sperm survival
5	HSP-1	1168021	8.2	14.0	42.9	112	Low sperm survival