<https://helda.helsinki.fi>

Sharpening species boundaries in the *Micarea prasina* group, with a new circumscription of the type species *M. prasina*

Launis, Annina

2019-07-04

Launis , A , Malicek , J , Svensson , M , Tsurykau , A , Serusiaux , E & Myllys , L 2019 , '
Sharpening species boundaries in the *Micarea prasina* group, with a new circumscription of
the type species *M. prasina* ' , *Mycologia* , vol. 111 , no. 4 , pp. 574-592 . <https://doi.org/10.1080/00275514.2019.1603044>

<http://hdl.handle.net/10138/326192>

<https://doi.org/10.1080/00275514.2019.1603044>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Sharpening species boundaries in the *Micarea prasina* group with notes on the type species *M. prasina*

a Annina Launis, b Jiří Malíček, c Måns Svensson, d Andrei Tsurykau, e Emmanuël Sérusiaux & a Leena Myllys

a Botany Unit, Finnish Museum of Natural History, P.O. Box 7, FI-00014 University of Helsinki, Finland; b The Czech Academy of Sciences, Institute of Botany, Zámek 1, 252 43 Průhonice, Czech Republic; c Museum of Evolution, Uppsala University, Villavägen 9, 752 36 Uppsala, Sweden; d Department of Biology, F. Skorina Gomel State University, Sovetskaya str. 104, 246019 Gomel, Belarus; Department of Ecology, Botany and Nature Protection, Institute of Natural Sciences, Samara National Research University, Moskovskoye shosse 34, 443086 Samara, Russia; e Evolution and Conservation Biology Unit, InBios research center, University of Liège, Sart Tilman B22, B-4000 Liège, Belgium.

CONTACT: Annina Launis, annina.launis@helsinki.fi (2020 onwards: annina.kantelinen@helsinki.fi)

ABSTRACT

Micarea is a lichenized genus in the family Pilocarpaceae (Ascomycota). We studied the phylogeny and reassessed the current taxonomy of the *M. prasina* group. We focused especially on the taxonomic questions concerning the type species *M. prasina* and, furthermore, challenges concerning type specimens that are too old for successful DNA barcoding and molecular studies. The phylogeny was reconstructed using nuc rDNA internal transcribed spacer region (ITS1-5.8S-ITS2), mito rDNA small subunit (mtSSU) and replication licensing factor *MCM7* gene from 31 species. Fifty-six new sequences were generated. The data was analyzed using maximum parsimony and maximum likelihood methods. The results revealed four undescribed, well-supported lineages. Three lineages represent new species described here as *M. fallax*, *M. flavoleprosa* and *M. pusilla*. In addition, our results support the recognition of *M. melanobola* as a distinct species. *Micarea fallax* is characterized by a vivid to olive green thallus composed of aggregated granules and whitish or brownish apothecia sometimes with greyish tinge (Sedifolia-grey pigment). *Micarea flavoleprosa* has a thick, wide-spreading yellowish-, whitish to olive-green sorediate thallus and lacks the Sedifolia-grey pigmentation. The species is mostly anamorphic, developing apothecia rarely. *Micarea melanobola* is characterized by a pale to dark vivid green granular thallus and darkly pigmented apothecia (Sedifolia-grey). *Micarea pusilla* is characterized by a whitish-green to olive-green thinly granular or membranous thallus, numerous and very small whitish apothecia lacking the Sedifolia-grey pigment, and by the production of methoxymicareic acid. *Micarea fallax*, *M. flavoleprosa* and *M. melanobola* produce micareic acid. The reliability of crystalline granules as a character for species delimitation was investigated and was highly informative for linking the old type specimen of *M. prasina* to fresh material.

KEY WORDS:

crystalline granules; diversity; dead wood; epitypes; lichens; molecular phylogenetics; new species; species delimitation; taxonomy; type species.

INTRODUCTION

Micarea Fr. is a lichen genus in the family Pilocarpaceae, comprising ca. 100 species (Andersen & Ekman 2005; Kirk *et al.* 2008; Coppins 2009). All species are crustose in growth form and occur across a wide range of habitats. In many cases, traditional methods have proven insufficient for species delimitation because there are only a small number of phenotypic characters and/or there are difficulties in their interpretation (Coppins 1983; Czarnota 2007; Czarnota & Guzew-Krzemińska 2010). Largely based on molecular methods, the understanding of species boundaries and species diversity has improved (Czarnota & Guzew-Krzemińska 2010; van den Boom & Ertz 2014; Guzew-Krzemińska *et al.* 2016; van den Boom *et al.* 2017). Recent molecular phylogenies show, for instance, that the morphological concept of *Micarea* is paraphyletic (Andersen & Ekman 2005; Sérusiaux *et al.* 2010), even after the introduction of a new genus *Brianaria* S. Ekman & M. Svensson for the *M. sylvicola* group (Ekman & Svensson 2014). However as explained below, before a reliable genus-level phylogenetic reconstruction can be proposed, several taxonomic issues concerning the type species, *M. prasina* Fr., should be addressed (e.g. Czarnota & Guzew-Krzemińska 2010; Launis *et al.* 2019). Also, species delimitation within the *M. prasina* group needs further studies (Andersen & Ekman 2005; Sérusiaux *et al.* 2010; Schmull *et al.* 2011).

The *M. prasina* group is monophyletic within *Micarea*, characterized by a “micareoid” photobiont (a coccoid green alga with cells 4–7.5 µm diam.), immarginate small apothecia, a hyaline hypothecium, branched paraphyses and an ascus of the *Micarea*-type with a K/I+ blue amyloid tholus and a more lightly staining axial body often with a darkly stained lining (Hafellner 1984; Czarnota 2007; Ekman *et al.* 2008). Many species form effuse thalli composed of goniocysts and produce Sedifolia-grey pigment (K+ violet, C+ violet), which is typically present in the pycnidia and in the upper layer of the hymenium of the apothecia (Coppins 1983; Czarnota & Guzew-Krzemińska 2010). The group originally included *M. prasina*, *M. hedlundii* Coppins, *M. levicula* (Nyl.) Coppins and with some uncertainty also *M. misella* (Nyl.) Hedl., *M. melanobola* (Nyl.) Coppins and *M. synotheoides* (Nyl.) Coppins (Coppins 1983, but see also Hedlund 1892). Since, delimitation of the group was examined using molecular markers and several new species were described (e.g. Launis *et al.* 2019 and the references therein).

Challenges in species delimitation within the *M. prasina* group were discussed by Coppins (1983), Czarnota (2007), Czarnota & Guzew-Krzemińska (2010) and Launis *et al.* (2019). The type species, *M.*

prasina, is notorious for its phenotypic variability. Coppins (1983) treated *M. prasina* in a broad sense with high morphological variability and three chemical races. Later, *M. prasina* was shown to be non-monophyletic and two morphologically and chemically distinct lineages were recognized as species, namely *M. subviridescens* (Nyl.) Hedl. producing prasinic acid and *M. micrococca* (Körb.) Gams ex Coppins producing methoxymicareic acid (Coppins 2002 and 2009). Furthermore, three distinct lineages were discovered within *M. micrococca* and named *M. byssacea* (Th. Fr.) Czarnta, Guzow-Krzemińska & Coppins, *M. micrococca* s. str. (Körb.) Gams ex Coppins and a third species (Czarnta & Guzow-Krzemińska 2010) that was later described as *M. czarntae* Launis, van den Boom, Sérus. & Myllys (Launis *et al.* 2019). Following the reintroduction of these species, *M. prasina* in a strict sense was primarily characterized by having a goniocystoid thallus combined with the presence of micareic acid (e.g. Czarnta 2007).

Changes in species delimitation raise the need to reevaluate the synonymies. Coppins (1983) placed several names as synonyms under his concept of *M. prasina*, many of them earlier treated as forms or variations by Hedlund (1892). These synonyms were reorganized based on their secondary chemistry, i.e. methoxymicareic, micareic or prasinic acid, and either moving as synonyms for recircumscribed *M. byssacea* and *M. micrococca* (Czarnta & Guzow-Krzemińska 2010), remaining as synonyms of *M. prasina* (Czarnta 2007) or reintroduced as new species (Coppins 2002, Hafellner & Türk 2016). During this process, one synonym of *M. prasina* sensu Coppins, i.e. *Lecidea declivitatum*, was left untreated and its taxonomic rank needs clarification. In addition, the taxonomic rank of specimens with black apothecia, earlier regarded as *M. melanobola* (Nyl.) Coppins, have remained unclear (Czarnta 2007). Historically, this taxon was treated as a species (Nylander 1867, Coppins 1983), as a form of *M. prasina* (Hedlund 1892) or as a synonym of *M. prasina* (Vezda & V. Wirth 1976, Czarnta 2007). Confusion also arose from the infraspecific genetic variation between European and North American specimens of *M. prasina* (Czarnta & Guzow-Krzemińska 2010).

Because of the small number of distinct phenotypic traits and challenges in molecular studies of old lichen specimens (Sohrabi *et al.* 2010), we continue to investigate the value of crystalline granules as a character for species delimitation in *Micareia*. These granules were detected previously in the *M. byssacea* and *M. micrococca* complexes in sections of apothecia and thallus examined in polarized light (Launis *et al.* 2019). The presence, distribution, size and solubility of such granules are considered important characters for the

identification of some genera of crustose lichen species (e.g. Brodo 1984; Spribille *et al.* 2011), but their significance in many other groups, including *Micarea*, is still poorly understood (Orange *et al.* 2010).

In this study, we investigate the species diversity within the *M. prasina* group and disentangle the taxonomy concerning the type species *M. prasina*. We use phenotypic characters and sequence data from three loci (ITS, mtSSU and *MCM7*). All relevant type specimens synonymized under *M. prasina* by Czarnota (2007) were re-examined. The need for investigating species boundaries with molecular markers is evident following studies demonstrating that the species-delimiting phenotypic features are subtle in this group (Czarnota & Guzow-Krzemińska 2010; Launis *et al.* 2019).

MATERIALS AND METHODS

The phylogeny of the *Micarea prasina* group was studied using 31 species (Table 1). Fresh specimens were collected from Belarus, the Czech Republic, Finland, the Netherlands, Scotland, Sweden and Maine, U.S.A. during 2010–2015. Type material of related *Micarea* species from the herbaria G, H, HBG and UPS was studied for comparison (Thiers B. Index Herbariorum). Additional material was studied from personal herbaria of F. Berger and J. Malíček.

DNA extraction, polymerase chain reaction and DNA sequencing. —DNA was extracted from 1–3 apothecia of max. three years ago stored specimens. For most specimens DNA was extracted using DNeasy® Blood & Tissue kit (QIAGEN) following the protocol described by Myllys *et al.* (2011). PCR reactions were prepared using PuReTaq Ready-To-Go PCR beads (GE Healthcare). Each 25 µL reaction volume contained 19 µL dH₂O, 1 µL of each primer (10 µM) and 4 µL extracted DNA.

For the ITS region, PCR was run under the following conditions: initial denaturation for 5 min at 95 C followed by five cycles of 30 s at 95 C (denaturation), 30 s at 58 C (annealing), and 1 min at 72 C (extension); for the remaining 40 cycles the annealing temperature was decreased to 56 C and the PCR program ended with a final extension for 7 min at 72 C. Primers ITS1-LM (Myllys *et al.* 1999) and ITS4 (White *et al.* 1990) were used both for PCR amplification and sequencing of ITS.

For the mtSSU gene, PCR was run under the following conditions: initial denaturation for 10 min at 95 C followed by six cycles of 1 min at 95 C (denaturation), 1 min at 62 C (annealing), and 105 s at 72 C (extension); for the remaining 35 cycles the annealing temperature was decreased to 56 C; the PCR profile

ended with a final extension of 10 min at 72 C. Primers mrSSU1 and mrSSU3R (Zoller *et al.* 1999) were used both for PCR amplification and sequencing.

For the *MCM7* gene, PCR was run under two different conditions depending on the primers selected: For the first, an initial denaturation for 10 min at 94 C was followed by 38 cycles of 45 s at 94 C (denaturation), 50 s at 55 C (annealing), and 1 min at 72 C (extension), with the PCR profile ending with a final extension for 5 min at 72 C. Primers *MCM7_AL1r* and *MCM7_AL2f* (Launis *et al.* 2019) were used both for PCR amplification and sequencing. The second protocol used an initial denaturation for 10 min at 94 C, followed by 38 cycles of 45 s at 94 C (denaturation), 50 s at 56 C (annealing), and 1 min at 72 C (extension); the PCR program ended with a final extension for 5 min at 72 C. Primers *x.Mcm7.f* (Leavitt *et al.* 2011) and *Mcm7.1348R* (Schmitt *et al.* 2009) were used both for PCR amplification and sequencing. PCR products were cleaned and sequenced by Macrogen Inc., South Korea (www.macrogen.com).

Phylogenetic analyses.— Thirty-nine ITS sequences, 82 mtSSU sequences and 49 *MCM7* sequences, were aligned separately with MUSCLE 3.8.31 (Edgar 2004) using EMBL-EBI's freely available webservice (<http://www.ebi.ac.uk/Tools/msa/muscle/>). The single gene trees, reconstructed by using the same methods and algorithms as subjected to the concatenated dataset described below, showed no strongly supported conflicts following the approach of Kauff & Lutzoni (2002) (with threshold bootstrap values $\geq 75\%$). The three datasets were combined into a concatenated matrix in MacClade 4.08 (Maddison & Maddison 2005). *Micarea peliocarpa* (Anzi) Coppins & R. Sant. was used as outgroup for the *M. prasina* group. *Micarea misella*, *M. adnata*, *M. elachista*, *M. globulosella*, *M. pycnidiophora*, *M. synotheoides* and *M. stipitata* were included to examine the monophyly of the ingroup *sensu* van den Boom *et al.* (2017). Portions of the alignment with ambiguous positions that might not have been homologous were manually excluded. The concatenated dataset, including 83 terminals was subjected to maximum parsimony analysis (MP) as implemented in TNT 1.1 (Goloboff *et al.* 2008) and to maximum likelihood analysis (ML) using RAxML 8.1.15 (Stamatakis 2014) on the CSC-IT Center for Science server (<http://www.csc.fi/home>). The MP analysis was performed using “traditional search” with random addition of sequences with 100 replicates and the TBR branch swapping algorithm. Ten trees were saved for each replicate and gaps were treated as missing data. Node support was estimated by bootstrapping (Felsenstein 1985) with 1000 replicates. Bootstrap values $> 75\%$ were considered significant. For the ML analysis, the combined dataset was assigned to seven partitions: ITS1, 5.8S, ITS2,

mtSSU and each of three codon positions of *MCM7*. The hypervariable region at the end of the mtSSU was removed from the analyses (characters 649–804 in the alignment). An independent GTR+G model was used for each subset and branch lengths were assumed to be proportional across subsets. Node support was estimated with 1000 bootstrap replicates using the rapid bootstrap algorithm. The alignments are deposited in TreeBASE (study no. 24165, <http://purl.org/phylo/treebase/phyloids/study/TB2:S24165>).

Morphology and chemistry.— Hand-cut apothecial sections and squashed thallus preparations were examined with a dissecting or compound microscope. Ascospores and other anatomical details were studied and measured in water, or in 10% potassium hydroxide (K) if features were otherwise unseparated. Measurements are given in a format of minimum and maximum measure. Rare minimum or maximum measures of spores are marked in parenthesis. Chemical spot tests were performed under a compound microscope using sodium hypochlorite (C) and (K) (Orange *et al.* 2010). Pigments were defined following the system of Coppins (1983), Meyer and Printzen (2000) and Czarnota (2007). Specimens were further studied using thin-layer chromatography (solvent C) following the methods of Culberson and Kristinsson (1970) and Orange *et al.* (2010). The crystalline granules were investigated by using a compound microscope with polarization filters. Crystalline granules were studied from sequenced specimens and from the type specimen of *M. prasina*.

RESULTS

Altogether 56 new sequences were generated for the analyses and 114 were obtained from GenBank. Our multiloci dataset and the constructed phylogeny (Fig. 1) included 31 OTUs, 170 sequences and 1728 characters, of which 969 were parsimony informative. Since the topologies of the ML and MP analyses did not have any strongly supported conflicts, only the tree obtained from the ML analysis is shown (Fig. 1).

The *Micarea prasina* group is monophyletic, though remains unsupported. It includes 23 species. *Micarea tomentosa* and a new lineage discovered in this study, *M. pusilla* sp. nov. Launis, Malíček & Myllys (represented by 13 samples from Czech Republic, Finland and Russia), form a sister group and are resolved as basal species in the *M. prasina* group. The remaining species are divided into two strongly supported clades. The first clade consists of species of the *M. prasina* complex. In the second clade *M. hedlundii* and *M. xanthonica* are resolved as basal species. *Micarea byssacea* and *M. micrococca* complexes discussed in Launis *et al.* (2019) form a strongly supported group and are both recovered as monophyletic.

The *M. prasina* complex includes *M. herbarum* Brand, Coppins, Sérus. & van den Boom, *M. meridionalis* van den Boom, Brand, Coppins & Sérus., *M. nowakii* Czarnota & Coppins, *M. prasina*, *M. soralifera* Guz.-Krzemiń., Czarnota, Łubek & Kukwa and *M. subviridescens*, and in addition, the here described new species *M. fallax* Launis & Myllys (including 12 specimens from Czech Republic, Belarus, Finland, Scotland and Sweden) and *M. flavoleprosa* Launis, Malíček & Sérus. (4 specimens from Czech Republic and France). The results also support the distinction of *M. melanobola* as a species-level taxon (11 specimens from Finland). *Micareia prasina* splits into three well supported lineages (referred to as *M. prasina* 1, 2 and 3, see Fig. 1).

To examine the identity of the three *M. prasina* lineages, the specimens were compared to the lectotype specimen of Fries published in 1825. Based on careful morphological and anatomical investigation, including studies of the crystalline granules discussed below, we were able to determine that *M. prasina* 3 is conspecific with to the holotype. Because of its age, we did not attempt to generate DNA sequence data from the holotype and were thus unable to confirm the identity of lineage 3 by molecular methods.

Small crystalline granules, soluble in K, were studied in polarized light, and are shown in detail in Fig 3. The type specimen of *M. prasina* had crystalline granules in the epihymenium of the apothecial section; the thallus itself was not studied because of the high value of the holotype specimen. The granules were also studied in apothecial sections of *M. prasina* 1, 2 and 3. In *M. prasina* 3, the granules detected in the epihymenium were similar to those in the holotype. In *M. prasina* 1 and 2, the crystalline granules were mostly detected in the lower half of the hymenia and also in the thalli. *Micareia fallax* and *M. melanobola* produced granules in the hymenium of the apothecia at least sometimes, and always in the thallus. No granules were detected in apothecia of *M. flavoleprosa* and *M. pusilla*, but polarization was detected in the thallus of the first.

Our results show that the species complexes differ in their secondary metabolites. Species in the *M. prasina* complex, including the three new lineages, produce micareic acid, except *M. subviridescens*, which produces prasinic acid, and *M. herbarum* with no secondary metabolites detected. Species in the *M. byssacea* and *M. micrococca* complexes, on the other hand, produce methoxymicareic acid.

TAXONOMY

Micarea fallax Launis & Myllys sp. nov.

FIGS. 2 A & B

MycoBank No.: MB 830205

Typification: FINLAND, VARSINAIS-SUOMI: Karkkila, Myllypuro, mixed-forest between lake Vahermanjärvi and lake Tarkeelanjärvi, near river Myllypuro, on bark of *Pinus sylvestris*, in shaded and moist microhabitat, WGS84 Lat: 60° 33,18022', Lon: 23° 59,67047', 10.Sept.2011, A. Launis 109115 (**holotype** H). GenBank: mtSSU: MK454757.

Etymology: *fallax* (Latin), meaning deceptive, referring to the challenges in identifying this species with morphological characters.

Diagnosis: Thallus vivid green, pale olive green to dark olive green, granular, warted or membranous; apothecia numerous, cream white, pale brownish, honey brown to medium brown, sometimes with pale greyish tinge because of the Sedifolia-grey pigment, usually hemispherical, up to 0.4 (–0.5) mm diam.; ascospores oblong-ellipsoidal or obovoid, 0–1 septate, 8–11 × 3–4 µm; produces micareic acid.

Description: Thallus effuse, vivid green or pale olive green to dark olive green, granular, composed of goniocysts 20–40 µm diam., often coalescing to form larger granules or a ± thick almost continuous and cracked thallus, if less developed warted-granular, small-areolate or membranous and ± shiny. Photobiont micareoid, algal cells 4.5–7 µm. Apothecia numerous, 0.2–0.4 (–0.5) mm diam., cream white, pale brownish, honey brown to medium brown, sometimes with pale greyish tinge, and then K+ violet and C+ violet because of the Sedifolia-grey pigment, usually hemispherical or subglobose, sometimes adnate or convex and semi-immersed in the thallus, simple or tuberculate. Hypothecium hyaline, sometimes yellowish. Hymenium hyaline, ca 35–45 µm high. Epihymenium hyaline, pale grey or pale brown. Paraphyses numerous, branched, not or only scarcely wider at apices, 0.5–1 (–1.5) µm wide. Asci clavate, *Micarea*-type, 28–38 × 10–13 µm. Ascospores 8 per ascus, oblong-ellipsoidal or obovoid, 0–1 septate, 8–11 × 3–4 µm. Pycnidia of two types, rare or numerous depending on a specimen, whitish, K– and C–. Mesopycnidia small and usually inconspicuous, 50–100 µm wide, sessile or mostly semi-immersed in surrounding gonicysts, globose or barrel-like, sometimes with extruding white conidial mass. Mesoconidia cylindrical or cylindrical-fusiform, (4–) 4.5–5.5 (–6) × 1–1.5 µm. Micropycnidia small and inconspicuous, immersed in surrounding gonicysts, globose,

40–80 µm wide. Microconidia straight or sometimes curved, bacilliform or narrowly fusiform, $5.5\text{--}7.5 \times 1$ µm. Crystals often visible in the hymenium and in the thallus using polarized light, soluble in K.

Chemistry: Micareic acid.

Habitat and distribution: *Micareia fallax* is widely distributed in Europe. Specimens have so far been collected from Belarus, Czech Republic, Finland, Scotland and Sweden. The collections are from several tree species (*Pinus sylvestris*, *Abies alba*, *Quercus* spp., *Picea abies* and *Alnus* spp.) where *M. fallax* was found to occur on bark and decaying wood. The specimens are from mature managed and old-growth forests.

Additional specimens examined: BELARUS. GOMEL REGION: Gomel district, Kalinino forest, 1 km NE of Tereshkovichi village, pine forest, on *Pinus sylvestris*, 52°15'55''N, 30°58'51''E, 2011, *Tsurykau 001c4* (H). CZECH REPUBLIC. EASTERN MORAVIA: Vsetín, Beskydy Protected Landscape Area, Velké Karlovice – Razula National Nature Reserve, old-growth beech-silver fir forest, on bark of *Abies alba*, ca 49°21'38''N, 18°22'48''E, alt. 700–750 m., 2013, *Malíček 6127 & Vondrák* (Hb. Malíček). SOUTHERN BOHEMIA: Prachatice, Šumava Protected landscape Area, Kubova Hut' – Milešický prales Nature Reserve, old-growth beech-spruce forest, on fallen decaying wood, 48°59'05''N, 13°50'20''E, alt. 1090–1120 m., 2018, *Malíček 11821 & Palice* (Hb. Malíček). CENTRAL BOHEMIA: Beroun, Křivoklátsko Protected Landscape Area, Kublov coniferous forest 0.5 km W of Velké Čihátko Hill, on stump of *Picea abies*, 49°56'48''N, 13°54'35''E, alt. 500 m., 2018, *Malíček 11992 & Vondrák* (Hb. Malíček). FINLAND. KAINUU: Sotkamo, Rommakkovaara Nature Reserve, old-growth forest dominated by *Picea abies* and *Betula* spp., on bark of fallen decaying (early stage) *Picea abies*, WGS84 Lat: 63° 53,41124', Lon: 28° 29,15969', 2012, *Launis 27122, Myllys & Kuusinen* (H). Pohjois-Karjala, Lieksa, Koli National Park, E slope of Koli, old natural forest, on wood of fallen decaying (early stage) *Picea abies*, WGS84 Lat: 63° 6,19996', Lon: 29° 48,83939', 2013, *Launis 59132* (H). UUSIMAA: Kirkkonummi, near Nuuksio Nature Reserve, recently protected *Picea abies* – dominated forest, on bark of fallen decaying (mid-stage) *Picea abies*, WGS84 Lat: 60° 17,09232', Lon: 28° 44,86853', 2013, *Launis 1710132 & Myllys* (H). UUSIMAA: Tuusula, Korso, shaded and dense *Picea abies* dominated mixed managed forest, on wood of fallen decaying (early stage) *Picea abies*, WGS84 Lat: 60° 21,26638', Lon: 25° 1,93227', 2013, *Launis 1010138* (H). Ibid. on wood of fallen decaying (late stage) *Picea abies*, WGS84 Lat: 6° 3,22812', Lon: 26° 1,13417', 2013, *Launis 1010139* (H). —SCOTLAND. EAST LOTHIAN (vc 82): Humbie, Church wood, on wood of old *Quercus*, British National Grid NT 46287, BNG

63808, 2014, *Launis 171143 & Coppins* (H). SWEDEN. SMÅLAND: Västra Ed parish, 1.7 km ESE of Forsby manor house, about 200 m E of Ingelsberg, on hard wood on stump of *Pinus sylvestris* in mixed coniferous swamp forest, 58°06'N, 16°58'E, SWEREF99: 6436288 593161, 2011, *Svensson 2398* (H).

Other type specimens examined:

GERMANY: Schleswig-Holstein, Kreis Pinneberg, Gehege Voßloch Barmstedt, an *Fraxinus excelsior*, 1939, Erichsen (holotype of *Lecidea abdita*, HBG).

Notes: *Micarea fallax* is characterized by a vivid to olive green thallus that is composed of aggregated granules, small-areolate, warted or membranous. The less developed forms of thalli are found especially when the species grows on dead wood. Apothecia are usually cream-white, pale brownish or pale greyish, rarely with darker shades of brown in parts. Morphologically *M. fallax* resembles *M. prasina* s. str. and *M. laeta*, a new species in the *M. byssacea* complex described by Launis *et al.* (2019).

The main morphological distinguishing character separating *M. fallax* from its close relative *M. prasina* s. str. is the crystalline granules: *M. prasina* forms crystals in the epihymenium whereas *M. fallax* produces granules only in the hymenium. Less distinctive differences occur: *M. prasina* usually forms a well-developed granular thallus and prefers decaying wood as a substrate, whereas *M. fallax* lacks a well-developed thallus if growing on decaying wood and is more common on bark. Also, the production of the Sedifolia-grey pigment is quite rare in *M. fallax* and the ascospore size is smaller.

The most distinctive differences between *M. fallax* and *M. laeta* are in their secondary metabolites: *M. fallax* produces micareic acid, whereas *M. laeta* produces methoxymicareic acid. The species also have morphological differences, especially in the apothecia. *Micarea fallax* usually has smaller apothecia that are hemispherical rather than adnate, and quite often brownish and/or greyish pigmented. Also, the thallus color in *M. fallax* is often paler and the structure is more aggregated. Despite the differences listed above, some specimens are very difficult to distinguishing without thin layer chromatography (TLC) or DNA analysis. The difficulties in the identification occur especially with young, less developed specimens of *M. laeta*, because the apothecia size tends to be the same as in *M. fallax*. Furthermore, the species have similar spore sizes and both species produce crystalline granules in the hymenium.

Our results show that *M. fallax* is sister to *M. melanobola*. However, morphologically these two species clearly differ: *M. fallax* has pale to brownish or greyish apothecia, whereas *M. melanobola* always has dark gray to blackish apothecia.

Micarea fallax was also compared to the type specimens of names synonymized under *M. prasina*. *Micarea prasiniza* Nyl. from 1874, resembles *M. fallax* and is probably a close relative. However, the type of *M. prasiniza* has bigger, subglobose apothecia that are homogeneously dark brown (K– and C–). It has granular thalli that are less aggregated than in *M. fallax* and they are never clearly warted, areolate or membranous. Both taxa produce micareic acid. Crystalline granules were not detected in the apothecia of *M. prasiniza*, but the investigated sample size was small. Furthermore, *Lecidea abdita* Erichsen, described in 1939, has some similarity with *M. fallax*, but the ascospores are larger at $(7-12-14(-17) \times 3(-3.5) \mu\text{m}$. The secondary metabolite profile of the holotype of *L. abdita* is unknown.

Micarea flavoleprosa Launis, Malíček & Sérus. sp. nov.

FIG: 2 C

MycoBank No.: MB 830206

Typification: CZECH REPUBLIC. NORTHERN MORAVIA: Jeseník, Jeseníky Protected Landscape Area, Bělá pod Pradědem – N part of Vysoký vodopád Nature Reserve, valley of Studený p. brook, on stump, 50°07′06″N, 17°12′02″E, alt. 900 m, 13.Jul.2012, J. Malíček 5098 (**holotype** PRA). GenBank: mtSSU MK454756, MCM7 MK456615.

Etymology: The name refers to the sordidously granular and farinose thallus that is often yellowish green in colour.

Diagnosis: Thallus yellowish-, whitish- to olive green, bright when fresh, minutely granular or farinose; apothecia rare, cream-white, adnate to hemispherical, up to 0.6 mm diam.; ascospores oblong-ellipsoidal or obovoid, 0–2 septate, $(10-12-16 \times 4-6 \mu\text{m}$; apothecia lack Sedifolia-grey pigment; produces micareic acid.

Description: Thallus effuse and wide-spreading, uneven, thick, yellowish green, whitish green to olive green, bright yellow green in fresh specimens, dark olive green in thin parts and paler yellowish to whitish green in thicker parts, granular or farinose, composed of minute soredia of small goniocysts 12–20 μm diam., which often coalesce to form larger granules up to 65 μm , K–, C–. Photobiont micareoid, algal cells 4.5–7 μm .

Apothecia rarely developed (found on one sample), if present few, 0.4–0.6 mm diam., cream-white, adnate to hemispherical, K[–] and C[–] in cross section (Sedifolia-grey pigment absent). Hypothecium hyaline. Hymenium hyaline, ca 40–55 µm high. Epihymenium hyaline. Paraphyses numerous, branched, 0.5–1.5 µm wide. Asci clavate, *Micarea*-type. Ascospores 8 per ascus, oblong-ellipsoidal or obovoid, 0–2 septate, (10–) 12–16 × 4–6 µm. Pycnidia of two types, rare or numerous, whitish, K[–] and C[–]. Mesopycnidia small and inconspicuous, 50–90 µm wide, usually immersed in surrounding goniocysts, barrel-like or globose, sometimes with extruding white conidial mass. Mesoconidia cylindrical or cylindrical-fusiform, 4–5.5 × 1.2–1.5 µm. Micropycnidia 40–100 µm wide, globose, immersed in surrounding gonicysts or distinctly sessile with gaping ostiole. Microconidia straight or sometimes curved, bacilliform or narrowly fusiform, 5.5–9 × 1 µm. Crystals (studied in polarized light) not detected in the apothecia, but polarization in the thallus. Owing to the rarity of apothecia the investigated sample size was small.

Chemistry: Micareic acid, with traces of 2-4 related compounds.

Habitat and distribution: *Micarea flavoleprosa* is so far known from two localities in Austria, three in the Czech Republic (Northern Moravia and Southern Bohemia) and one locality in France. Collections are from beech dominated managed and old-growth forests, where it often covers large patches on stumps of decaying wood.

Additional specimens examined: AUSTRIA. OBERÖSTERREICH: Koberausserwald, am Schwarzmoosbach, Plenterwald 200 m WSW Einfahrt Schottergrube, on Totholz *Picea*, 48°04'18"N, 13°20'38"E, alt. 640 m, 2018, *Berger* 32900 (Hb. Berger). Ibid., Hinterstoder, Dietltal, Schluchtwald am Dietlbach, on Totholz, 47°40'10"N, 14°06'04"E, 2018, alt. 655 m, *Berger* 33573 (Hb. Berger). CZECH REPUBLIC. SOUTHERN BOHEMIA: Prachatice, Šumava National Park, Nová Pec old-growth spruce-beech forest above road 1.5 km NNE of top of Smrčina Mt., on stump, 48° 45'05"N, 13° 55'40"E, alt. 1200 m., 2012, *Maliček* 4699, *Bouda & Syrovátková* (Hb. Maliček). Ibid., Kubova Hut' – Milešický prales Nature Reserve, old-growth beech-spruce forest, on fallen decaying wood, 48°59'05"N, 13°50'20"E, alt. 1090–1120 m., 2018, *Maliček* 11823 & *Palice* (Hb. Maliček). FRANCE. VOSGES: Dépt. Haut-Rhin, Hohneck, Frankenthal nature preserve, under the Falimont pass, avalanche corridor with coppice of *Sorbus aucuparia*, with *Fagus* and *Acer pseudoplatanus*, on dead tree in scree, 48°02.39' N 7°01.07'E, 1100 m alt., 2005, E. Sérusiaux s.n. (LG).

Notes: *Micarea flavoleprosa* is characterized by the thick yellowish-, whitish to olive-green thallus, rarity of apothecia, and the production of micareic acid. Morphologically, and by occupying decaying wood as a growing substrate, the species resembles especially *M. viridileprosa*, *M. soralifera*, and sterile forms of *M. prasina*. Our phylogenetic analyses show that *M. flavoleprosa* is a close relative of the North American and central European lineages of *M. prasina* s. lato (*M. prasina* 1 and 2 in Fig. 1).

Micarea viridileprosa mainly differs from *M. flavoleprosa* by the production of gyrophoric acid, detectable by TLC and C+ red reaction in thallus and apothecia, and by developing a thinner, brighter and clearer green thallus. Apothecia are usually absent in both species, but if present, the ascospores are shorter and narrower in *M. viridileprosa* ($8\text{--}14 \times 2.5\text{--}4 \mu\text{m}$) than in *M. flavoleprosa* ($10\text{--}12\text{--}16 \times 4\text{--}6 \mu\text{m}$).

Micarea soralifera and *M. flavoleprosa* both produce micareic acid as a secondary metabolite and usually lack apothecia. However, the species differ morphologically rather clearly: *M. soralifera* develops a distinctly thinner thallus and well-delimited greyish-green soralia; these soralia react K+ violet and C+ violet in exposed parts because of presence of Sedifolia-grey.

Micarea prasina, on the other hand, differs from *M. flavoleprosa* by developing a thinner, more evenly colored thallus that is rarely minutely granular, farinose or yellowish-green. Also, *M. prasina* usually forms apothecia. Both species produce micareic acid.

Micarea flavoleprosa is so far known only from decaying wood in old-growth forests. In areas of forest management this has become a rare and diminishing substrate. Therefore, its special ecological requirements should be noted in conservation planning and red list evaluations.

Micarea melanobola (Nyl.) Coppins

FIGS: 2 D & E

MycoBank: MB110358]

≡ *Lecidea melanobola* Nyl. in *Flora*, Jena **50**: 371. 1867.

≡ *Catillaria melanobola* (Nyl.) Vain. in *Acta Soc. Fauna Fl. fenn.* **57** (2): 465. 1934.

≡ *Lecidea erysiboides* * [ssp.] *L. melanobola* (Nyl.) Nyl. in Hue in *Revue Bot. Courrensan* **5**: 103. 1886.

≡ *Catillaria prasiniza* β *byssacea* f. *melanobola* (Nyl.) Blomb. & Forss., *Enum. Pl. Scand.*: 91. 1880.

≡ *M. prasina* f. *melanobola* Hedl. in *Bih. K. Svenska VetenskAkad. Handl.* III, **18** (3): 87. 1892.

= *Lecidea prasinella* Müll. Arg., Switzerland, Valais, *Bovernier*, on *Larix*, *Flora*, Regensburg 55: 484. 1872 (holotype G G00292613 and isotypes G G00292614, G00292615, G00292616).

Typification: FINLAND. TAVASTIA AUSTRALIS: Kuhmois (Kuhmoinen), ad abietem (= *Picea abies*), 1866, *Norrlin* (lectotype H-NYL 21614, H9510482; isolectotypes: H-NYL p.m. 4504, H9510483, H9504133).

Description: Thallus effuse, pale to dark vivid green, sometimes olivaceous, granular, composed of goniocysts 14–35 µm diam., often coalescing to form larger granules, or if less developed warted-areolate; in darker parts of the thallus the goniocysts sometimes surrounded by olivaceous K± violet pigment (Sedifolia-grey). Photobiont micareoid, rounded, algal cells 4.5–7 µm. Apothecia numerous, 0.15–0.4 mm, hemispherical to subglobose, simple or tuberculate, dark grey or blackish, sometimes dark brownish, K+ violet and C+ violet in cross section because of Sedifolia-grey pigment. Hypothecium hyaline. Hymenium hyaline, ca 35–55 µm high. Epihymenium grey, dark grey, blackish grey, sometimes brownish grey or even greenish grey, K+ (intense) violet and C+ violet. Paraphyses numerous, richly branched, 0.5–1 µm wide. Asci clavate, *Micarea*-type 30–35 × 9–11 µm. Ascospores 8 per ascus, oblong-ellipsoidal or obovoid, 0–1 septate, 7–11 × 2.5–3.5 (–4) µm. Pycnidia of two types, rare or numerous depending on a specimen, mostly pigmented, grey to blackish grey, K+ violet and C+ violet. Mesopycnidia usually numerous, 30–90 µm wide, often immersed between goniocysts, globose or barrel-like with gaping ostiole often extruding white conidial mass. Mesoconidia cylindrical or cylindrical-fusiform, 3.5–4.5 × 1–1.5 µm. Micropycnidia small and inconspicuous, 40–90 µm wide, immersed or rarely sessile, globose. Microconidia straight or sometimes curved, bacilliform or narrowly fusiform, 5–7 (–7.5) × 0.5–1 µm. Crystals (studied in polarized light) detected in sections of the apothecia (hymenium), and polarization also in the thallus. Soluble in K. The character sometimes varies within a single individual thallus: some apothecia produce crystalline granules and some do not. The granules often form vertical streaks in the hymenium.

Chemistry: Micareic acid.

Habitat and distribution: Found on wood and bark of *Picea abies* and *Pinus sylvestris* in boreal forests. Usually in rather shaded and moist habitats e.g. on northern side of tree trunks and/or near ground. Known so far from southern and central Finland and from one old locality in Switzerland. In addition, Czarnota (2007)

mentioned one possible collection from Estonia, but no sequence is available of the specimen for comparison and it has not been seen by the authors. Collections are from old-growth and managed forests.

Additional specimens examined: FINLAND. POHJOIS-KARJALA: Lieksa, Koli National Park, E slope of Koli, old natural forest, on wood of fallen decaying (early-stage) *Picea abies*, WGS84 Lat: 63° 6,16967', Lon: 29° 48,89951', 2013, *Launis* 79133 (H). KAINUU: Sotkamo, Rommakkovaara Nature Reserve, old-growth forest dominated by *Picea abies* and *Betula* spp., on bark of fallen decaying (early-stage) *Picea abies*, WGS84 Lat: 63° 53,41124', Lon: 28° 29,15969', 2012, *Launis* 27123, *Myllys & Kuusinen* (H). UUSIMAA: Vantaa, Herukkapuro Nature Reserve, old-growth forest, dominated by *Picea abies* and *Betula* spp., near a river, on bark of fallen decaying (mid-stage) *Picea abies*, WGS84 Lat: 60° 19,28983', Lon: 24° 45,94960', 2013, *Launis* 11014 (H). ETELÄ-HÄME: Muurame, Kettumäki, mixed old-growth forest dominated by *Betula* spp., *Picea abies* and *Pinus sylvestris*, on standing decaying *P. abies*, on bark, WGS84 Lat: 62° 11,10836', Lon: 16° 31,73796', 2014, *Launis* 49141 (H). POHJOIS-HÄME: Rautalampi, Kalajanvuori, mixed old-growth forest dominated by *Betula* spp., *Picea abies*, *Pinus sylvestris* and *Populus tremula*, on standing decaying *P. sylvestris*, on bark, WGS84 Lat: 62° 34,67921', Lon: 26° 40,75558', 2015, *Launis* 116152 (H). ETELÄ-HÄME: Padasjoki, Vesijako, Strict Nature Reserve, mixed old-growth forest dominated by *Betula* spp., *Picea abies* and *Pinus sylvestris*, on standing decaying *P. sylvestris*, on bark, WGS84 Lat: 61° 21,02120', Lon: 25° 6,11605', 2015, *Launis* 56151 (H). ETELÄ-HÄME: Muurame, Kirkkokangas, mixed old-growth forest dominated by *Betula* spp., *Picea abies* and *Pinus sylvestris*, on standing decaying *P. abies*, on bark, WGS84 Lat: 62° 13,76004', Lon: 25° 34,42501', 2015, *Launis* 39151 (H). POHJOIS-HÄME: Jyväskylä, Toivakka, mixed managed forest dominated by *Betula* spp., *Picea abies*, *Pinus sylvestris* and *Populus tremula*, on standing decaying *P. abies*, on bark, WGS84 Lat: 62° 4,86551', Lon: 26° 10,30759', 2015, *Launis* 286152 (H). POHJOIS-HÄME: Saarijärvi, Pyhä-Häkki National Park, mixed old-growth forest dominated by *Betula* spp., *Picea abies* and *Pinus sylvestris*, on standing decaying *P. abies*, on bark, WGS84 Lat: 62° 51,06825', Lon: 25° 29,14815', 2015, *Launis* 266151 (H). ETELÄ-SAVO: Joutsa, Leivonmäki National Park, mixed old-growth forest dominated by *Betula* spp., *Picea abies* and *Pinus sylvestris*, on standing decaying *P. abies*, on bark, WGS84 Lat: 61° 56,72980', Lon: 26° 0,92040', 2015, *Launis* 166151 (H).

Notes: *Micarea melanobola* was described by Nylander (1867). Later, however, it was considered a form of *M. prasina* by Hedlund (1892) and a synonym of *M. prasina* by Vězda & V. Wirth (1976). Coppins

(1983), on the other hand, treated *M. melanobola* as a species-level taxon. However, it was once again treated as a synonym of *M. prasina* by Czarnota (2007) with an emphasis on the need for molecular studies.

Micarea melanobola is characterized by its darkly pigmented apothecia (high amount of Sedifolia-grey pigment). The pigment is strongly visible even in specimens collected from shaded habitats. The principal morphological characters separating *M. melanobola* from its close relative *M. prasina* involve the degree of pigmentation, size of the ascospores and location of the crystalline granules. *Micarea prasina* produces crystalline granules in the epihymenium, whereas *M. melanobola* produces granules in the hymenium. Also, *M. prasina* prefers decaying wood as growing substrate whereas *M. melanobola* usually occurs on bark.

Micarea melanobola also resembles *M. herbarum* and *M. nowakii*, both belonging to the *M. prasina* complex. These three species have dark grey or blackish apothecia because of the high amount of Sedifolia-grey pigment. *Micarea nowakii* and *M. melanobola* sometimes have a similar finely warted thallus and also both produce micareic acid. However, *M. nowakii* has smaller ascospores of $6-8 (-8.5) \times 2-3 \mu\text{m}$ and emergent or even shortly stalked pycnidia often extruding a white mass of conidia. *Micarea nowakii* also often develops a thicker clearly warted and areolate thallus. Contrary to *M. melanobola* and *M. nowakii*, *M. herbarum* develops a very thin thallus and no secondary metabolites have been detected by TLC. The three species differ in their ecological preferences as well: *M. melanobola* prefers rather shaded and moist niches of dead and living trees, *M. herbarum* occurs on soft decaying wood and on dead moist stems of herbaceous plants, and *M. nowakii* occupies decaying hard wood and is usually found in localities exposed to light such as clearings in managed forests.

Micarea melanobola was also compared to the type specimens synonymized under *M. prasina* and / or type specimens containing micareic acid. We found that *M. melanobola* is morphologically identical with *Lecidea prasinella* Müll., a species described in 1872. Previously, *L. prasinella* was treated with some hesitation as a synonym of *M. micrococca* (Czarnota 2007), but based on its darkly pigmented epihymenium (Sedifolia-grey pigment) the taxon is not referable to the new circumscription of *M. micrococca* s. str. (Czarnota & Guzow-Krzemińska 2010). Also, *L. prasinella* likely contains micareic acid, instead of methoxymicareic acid found in *M. micrococca*. TLC was not applied to the holotype of *L. prasinella*, but an isotype contains micareic acid (TLC conducted by Pape & Clerc in 2013). We propose *L. prasinella* as a new synonym of *M. melanobola*.

Micarea prasina Fr. s. str. Syst. orb. veg. 257. 1825.

FIG: 2 F

MycoBank: MB119388

≡ *Biatora prasina* Fr., Stirp. agri femsion.: 36. 1825., nom. illeg. (Art. 63). – *Biatora prasina* (Fr.)

Trevisan, Linnea **28**: 288. 1856. – *Catillaria prasina* (Fr.) Th. Fr., Lich. Scand. **2**: 572. 1874.

= *Lecidea prasiniza* Nyl., Flora **57**: 312. 1874. – *Micarea prasina* f. *byssacea* subf. *prasiniza* (Nyl.) Th. Fr. in Hedl., Bih. Kongl. Svenska Vetensk. –Akad. Handl. III, **18**(3): 77. 1892. Type: Fennia, Tavastia australis, Padasjoki: Nyystölä, lepän kuorella (= ad corticem *Alni*), 1872, *E. Lang* 160 (lectotype H-NYL 21604, H9510484) [see notes in Czarnota 2007: 153].

= *Lecidea sordidescens* Nyl., Flora **57**: 312. 1874. – *Catillaria prasina* var. *byssacea* f. *sordidescens* (Nyl.) Blomb. & Forss., Enum. Pl. Scand.: 91. 1880. – *Micarea prasina* f. *byssacea* subf. *sordidescens* (Nyl.) Th. Fr. in Hedl., Bih. Kongl. Svenska Vetensk. –Akad. Handl. III, **18**(3): 77. 1892. – *Lecidea byssacea* var. *sordidescens* (Nyl.) Vain., Term. Füz. **22**: 320. 1899. – *Catillaria prasina* var. *sordidescens* (Nyl.) Lettau, Hedwigia **52**: 136. 1912. – *Micarea prasina* var. *sordidescens* (Nyl.) Brodo, Bull. New York State Mus. Sci. Serv. **410**: 152. 1968. Type: Switzerland, Zürich, Hepp, *Flecht. Eur. No.* 278 (isolectotype H-NYL 21632; isolectotype WRS�; lectotype E, n.v.).

= ?*Catillaria micrococca* f. *glebulosa* Erichsen, Ann. Mycol. **36**: 139. 1938. Type: Germany (holotype HBG).

Typification: SWEDEN. SMÅLAND: Femsjö, on wood, *E. M. Fries* (lectotype UPS L-167451)

Description: Thallus effuse, bright green to olive green, granular or softly isidious in appearance, composed of goniocysts 12–40 µm diam., often coalescing to form larger granules, or if less developed warted-granular. Photobiont micareoid, rounded, algal cells 4.5–7 µm. Apothecia often numerous, sometimes few, 0.2–0.5(–0.8) mm, hemispherical to sometimes subglobose, simple or tuberculate, creamy white, pale grey to partly dark grey, sometimes brownish, K± violet and C± violet in cross section because of Sedifolia-grey pigment. Variable in size and colour even in the same collection. Hypothecium hyaline. Hymenium hyaline, ca 30–60 µm high. Epihymenium hyaline or pale grey to darkish grey, sometimes brownish, K+ violet and C+ violet if Sedifolia-grey pigment is present. Paraphyses numerous, richly branched, 0.5–1 (–1.5) µm wide, only slightly widened above. Asci clavate, *Micarea*-type 25–55 × 8–12 µm. Ascospores 8 per ascus, oblong-

ellipsoidal or obovoid, 0–1 septate, $8\text{--}12\text{--}(14) \times 3\text{--}4.5\text{--}(5) \mu\text{m}$. Pycnidia of two types often present, sessile or immersed in surrounding gonocysts, whitish to gray, $K\pm$ violet and $C\pm$ violet. Mesopycnidia usually numerous, $50\text{--}150 \mu\text{m}$ wide, often immersed between gonocysts, sometimes emergent, globose or barrel-like with wide ostiole. Mesoconidia cylindrical or cylindrical-fusiform, $4\text{--}5.5\text{--}(6) \times 1\text{--}1.5 \mu\text{m}$. Micropycnidia small and often inconspicuous, $30\text{--}60 \mu\text{m}$ wide, immersed or sometimes sessile with wide ostiole, globose. Microconidia straight or sometimes curved, bacilliform or narrowly fusiform, $5\text{--}8\text{--}(9) \times 0.5\text{--}1 \mu\text{m}$. Crystals detected in sections of apothecia in epihymenium and sometimes also hymenium when studied in polarized light, polarization also visible in the thallus. Soluble in K.

Chemistry: Micareic acid.

Habitat and distribution: This is a common forest species in Fennoscandia and probably also in central Europe. Our study also includes reliably identified specimens from Maine, USA. The species usually occurs on medium to soft wood of stumps and logs of coniferous trees, and rarely also on bark (one specimen). The species is mainly encountered in natural forest habitats.

Additional specimens studied: AUSTRIA. Steiermark (=Styria): 9 km WNW of Bruck an der Mur, Floning-Zug, Kotzgraben 4 km NW of St. Dionysen, end of a valley floor, NW below a farmstead called Brandner, *Picea* –forest, on strongly decayed stump, $47^{\circ}26'N$, $15^{\circ}09'40''E$, MTB 8556/4, 880 m alt., 1990, *J. Hafellner* 43132 & *A. Hafellner* (H). CANADA. NEW BRUNSWICK: Charlotte Co.: Lepreau Falls Provincial Park, next to waterfalls parking area, in moist conifer forest by the falls, $45.1686^{\circ}N$, $66.4699^{\circ}W$, 2014, *Ahti* 74382 & *Clayden* (H). FINLAND. UUSIMAA: Kirkkonummi, Meiko Nature Reserve, *Vaccinium myrtillus* – *Picea abies* forest type, on wood of decaying *Picea abies*, WGS84 Lat: $60^{\circ} 7,74525'$, Lon: $15^{\circ} 22,23422'$, 2010, *Launis* 265101 & *Myllys* (H). ETELÄ-HÄME: Hämeenlinna, Evo, Kotinen Nature Reserve, SW of lake Valkea-Kotinen, on wood of a dead standing *Pinus sylvestris*, 2010, WGS84 Lat: $61^{\circ} 14,53326'$, Lon: $25^{\circ} 3,63680'$, *Launis* 229101 (H); Ibid., Lat: $61^{\circ} 14,49954'$, Lon: $25^{\circ} 3,69032'$, *Launis* 229106 (H). ETELÄ-SAVO: Joensuu, Ahveninen, Sorsasalo, old forest nature reserve, SE from the lake Lyly, on wood of decayed conifer stump against rock wall, WGS84 Lat: $62^{\circ} 20,73748'$, Lon: $27^{\circ} 40,11872'$, 2010, *Launis* 199105 (H). POHJOIS-KARJALA: Lieksa, Koli National Park, E slope of Koli, *Picea abies* -dominated old-growth forest, on wood of fallen *P. abies*, WGS84 Lat: $63^{\circ} 5,32176'$, Lon: $29^{\circ} 49,83050'$, 2013, *Launis* 59131 (H); Ibid., Lat: $63^{\circ} 5,30002'$, Lon: $29^{\circ} 49,94729'$ *Launis* 89131 (H); Ibid., on bark remnants of fallen *P. abies*, WGS84 Lat: 63°

5,30002', Lon: 29° 49,94729'*Launis* 89135 (H). U.S.A. MAINE: Washington Co.: Town of Steuben, Dyer Neck, Eagle Hill, Eagle Hill Institute, red trail, costal mixed forest dominated by *Picea rubens*, *Abies*, *Betula* and *Acer rubrum*, on wood of dead standing conifer, near ground, 44.45955 °N, 67.93161 °W, 2012, *Launis* 76122 (H). Town of Beals, Great Wass Island Reserve, open *Pinus banksiana* forest with heathy understory of *Kalmia*, *Ledum*, *Rhododendron canadense* etc., on wood of decaying horizontal conifer, 44.48099 °N, 67.59472 °W, 2012, *Launis* 66127 (H). Town of Cutler, Cutler Public Reserve Land, coast trail 0–1.5 miles between ME 191 and coast, humid mixed conifer (*Abies*, *Picea*, *Thuja*) and hardwood (*Acer*, *Sorbus*, *Betula*) forest, on wood of dead standing *Abies* sp., near ground, 44 41 54 °N, 67 0928 °W, 2012, *Launis* 136121 (H); Ibid., near seashore, on wood of dead standing *Abies balsamea* (on several stumps), near ground, 44 41 38 °N, 67 0930 °W, 2012, *Launis* 136123, 136124, 136129, 1361210, 1361211, 1361212 (H).

Notes: Although *M. prasina* was described in 1825, it was insufficiently known because of taxonomic uncertainties and changes in its circumscription. Our results suggest that even lately, as characterized by the presence of goniocystoid thallus and micareic acid (Czarnota 2007), the species has been misunderstood and actually included several taxa, such as *M. fallax*, *M. flavoleprosa* and *M. melanobola*.

Micarea prasina s.str. is characterized by a granular or softly isidious thallus, crystalline granules in the epihymenium and a preference for a dead wood as a substrate. The principal characters separating *M. prasina* from its close relatives *M. fallax* and *M. melanobola* involve ecological preferences and morphology: the latter two usually occur on bark instead of dead wood and always produce crystalline granules in the hymenium instead of the epihymenium. Furthermore, *M. fallax* produces somewhat smaller ascospores $8\text{--}11 \times 3\text{--}4 \mu\text{m}$ than *M. prasina*, where they are $8\text{--}12\text{--}(14) \times 3\text{--}4\text{--}(5) \mu\text{m}$, and the structure of its thallus is never well-developed and granular when encountered on dead wood. Production of Sedifolia-grey pigment is also rarer in *M. fallax*. *Micarea prasina* s.str. also resembles *M. melanobola*, but these two species can be separated by the quantity of pigmentation in apothecia, which is always very dark in *M. melanobola*, and ascospore sizes, which are $7\text{--}11 \times 2.5\text{--}3.5\text{--}(4) \mu\text{m}$ in *M. melanobola*. All three species produce micareic acid.

Micarea prasina s.str. also resembles three sterile species in the *M. prasina* group: *M. flavoleprosa*, *M. soralifera* and *M. viridileprosa*. *Micarea prasina* s.str. is usually fertile, but if sterile it can be distinguished by secondary metabolite profiles and/or characters of the thalli. Compared to *M. flavoleprosa*, *M. prasina* s.str. develops a thinner, more evenly colored thallus that is rarely as minutely granular, farinose or yellowish-green.

Micarea soralifera develops a thin thallus and well-delimited greyish-green soralia. These soralia have K+ violet and C+ violet reactions in exposed parts because of presence of Sedifolia-grey. *Micarea viridileprosa*, on the other hand, produces gyrophoric acid instead of micareic acid and is characterized by bright and clear green thallus.

Our results (Fig. 1) show that *M. prasina* s.l. includes three lineages: *M. prasina* 1, 2, and s. str. (earlier referred to as 3), of which the first two represent putative new species. The North American *M. prasina* 1 is characterized by a coralloid thallus that is pale green to whitish green in color. Compared to *M. prasina* s. str., the apothecia are adnate rather than hemispherical and lack Sedifolia-grey pigment. The European collections of *M. prasina* 2 display a rather similar morphology than *M. prasina* 1 except for the apothecia, which are somewhat smaller. Both taxa exhibit crystalline granules in the hymenium instead of only in the epihymenium. Because of the small number of collections of these two species (one for *M. prasina* 1, two for *M. prasina* 2) no taxonomic changes are proposed in this study.

Future studies should also clarify the taxonomic rank of two synonyms previously placed under *M. prasina* sensu Coppins (1983). During the process of new species delimitations based mainly on their secondary metabolites (Coppins 2002, Czarnota 2007, Czarnota & Guzow-Krzemińska 2010), *Lecidea declivitatum* (prasinic acid) was left untreated. The taxonomic rank of *M. polytrichi* Poelt & Döbbeler (secondary metabolites unknown) also varies in the current literature (e.g. Hafellner & Türk 2016, Nimis et al. 2018). This bryophilous taxon was described in 1975 (Poelt & Döbbeler), then regarded as a synonym of *M. prasina* (Coppins 1983), and lately reintroduced as a species (Hafellner & Türk 2016).

Micarea pusilla Launis, Malíček & Myllys sp. nov.

FIGS: 2 G & H

MycoBank: MB 830207

Typification: FINLAND. UUSIMAA: Tuusula, near Korso, shaded and dense *Picea abies* dominated managed forest, on wood of fallen decaying (early-stage) *Picea abies*, WGS84 Lat: 60° 21,26638', Lon: 25° 1,93227', 10.Oct.2013, A. Launis 101035 (**holotype** H). Genbank: mtSSU MK454753, MCM7 MK456612.

Etymology: The name refers to the very small size of the species and its apothecia.

Diagnosis: Thallus granular, warted-granular or membranous; apothecia whitish, crowded, very small, up to 0.2 mm in diam; ascospores oblong-ellipsoidal or obovoid, 0–1 septate, $7\text{--}9\text{ (–}9.5\text{)} \times 2\text{--}3\text{ }\mu\text{m}$; apothecia lack Sedifolia-grey pigment; produces methoxymicareic acid.

Description: Thallus effuse but rather small, whitish-green to olive-green, usually inconspicuous, membranous, or if well developed thinly granular or warted-granular, composed of small goniocysts $20\text{--}35\text{ }\mu\text{m}$ diam. that are often aggregated, K– and C–. Photobiont micareoid, algal cells $4.5\text{--}7\text{ }\mu\text{m}$. Apothecia numerous and crowded, very small $(0.07\text{--}) 0.1\text{--}0.15\text{ (–}0.2\text{)}\text{ mm}$, white or cream white, convex to hemispherical, simple, K– and C– in cross section (Sedifolia-grey pigment absent). Hypothecium hyaline. Hymenium hyaline, ca $30\text{--}35\text{ }\mu\text{m}$ high. Epihymenium hyaline. Paraphyses branched, $0.5\text{--}1$ wide. Asci clavate, *Micarea*-type, $16\text{--}25 \times 7\text{--}8\text{ }\mu\text{m}$. Ascospores 8 per ascus, oblong-ellipsoidal or obovoid, 0–1 septate, $7\text{--}9\text{ (–}9.5\text{)} \times 2\text{--}3\text{ }\mu\text{m}$. Pycnidia of two types, whitish, K– and C–. Mesopycnidia sessile or mostly immersed in surrounding goniocysts, small and usually inconspicuous, $30\text{--}70\text{ }\mu\text{m}$ wide. Mesoconidia fusiform, narrowly-ellipsoidal or sometimes cylindrical, $4\text{--}4.5\text{ (–}5\text{)} \times 1\text{--}1.5\text{ }\mu\text{m}$. Micropycnidia mostly immersed in surrounding goniocysts, small and inconspicuous, $30\text{--}50\text{ }\mu\text{m}$ wide. Microconidia bacilliform or narrowly fusiform: $(5\text{--}) 5.5\text{--}7\text{ (–}7.5\text{)} \times 1\text{ }\mu\text{m}$. Crystals not detected in sections of the apothecia or of thallus studied in polarized light.

Chemistry: Methoxymicareic acid.

Habitat and distribution: Usually on decaying wood of e.g. *Picea abies*. Sometimes on bark and then found especially at bases of branches. Samples are collected from mixed, beech or coniferous –dominated shaded old-growth or managed forests. Known from the Czech Republic, southern Finland and Russia (Caucasus and Dagestan). From the Czech Republic, it has already been published under a provisional name *Micarea inconspicua* nom. ined. (Vondrák *et al.* 2016, Malíček *et al.* 2017). This is probably a common species on dead wood, at least in eastern and northern Europe, but the small size and inconspicuousness of the species has left it overlooked.

Additional specimens examined: CZECH REPUBLIC: Český les Mts., Tachov, Rozvadov, old-growth mixed montane forest in protected area Diana, on log, 49.632021N, 12.5796478E, alt 510 m., 2016, Vondrák 14632 (PRA). Ibid., central part of the reserve, on fallen trunk, 49°37'55''N, 12°34'46''E, alt. 515 m., 2016, Malíček 9590, Kocourková, Palice, Vondrák (Hb Malíček). Ibid., managed beech forest at settlement Diana, on log, 49.6130N, 12.5937E, alt. 530 m., 2016, Vondrák 14634 (PRA). Ibid., Domažlice, Vranov, old ash-

maple forest on hill with ruin of Starý Herštýn, SE-S-SW slope, on stump, 49°28'17''N, 12°42'50''E, alt. 830–870 m, 2016, *Vondrák 14633* (PRA). SOUTHERN BOHEMIA: Prachatice, Šumava National Park, Volary–České Žleby, managed beech forest with spruce and sycamores intermixed on top of Mt Spáleníště, on fallen tree, 48°52'42''N, 13°47'24''E, alt. 940 m., 2016, *Maliček 9903, Palice & Vondrák* (Hb Maliček). SOUTHERN BOHEMIA: Český Krumlov, Novohradské hory Mts, Pohorská Ves–Žofín, Pivonické skály Nature Reserve, managed beech forest with intermixed spruces, 130 years old, NW-facing slope, on fallen tree, 48°40'01''N, 14°43'02''E, alt. 825 m., 2016, *Maliček 9636, Kocourková, Palice, Vondrák* (Hb Maliček). PRAEBOHEMICUM: Náměšt nad Oslavou, Březník, ruin Lamberk in Oslava river valley, on log, 49.1662897N, 16.1677097E, alt. 300–320 m., 2016, *Vondrák 16643* (PRA). CENTRAL BOHEMIA: Příbram, Brdy Hills, Jince managed coniferous forest 0.3 km NW of Velcí pond, on stump of *Picea abies*, 49°45'44''N, 13°56'21''E, alt. 580 m., 2018, *Maliček 12017 & Vondrák* (Hb Maliček). FINLAND. UUSIMAA: Tuusula, near Korso, shaded and dense *Picea abies* dominated managed forest, on bark of dead standing *Picea abies*, at base of a branch, WGS84Lat: 60° 21,26638', Lon: 25° 1,93227', 2013, *Launis 1010136* (H); Ibid., on bark of fallen decaying *Picea abies*, 2013, *Launis 1010137* (H). RUSSIA. REPUBLIC OF DAGESTAN: Magaramkentsky district, State nature Sanctuary Samurski, on log, 41.865059N, 48.503064E, 2015, *Ismailov, Urbanavichus, Vondrák 14668* (PRA). Southwestern Russia–Caucasus Mts, Caucasian Biosphere Reserve, Guzeripl primeval mixed forest (*Abies, Fagus, Carpinus, Quercus, Rhododendron ponticum*) on siliceous ridge 1 km SE of village, on decaying wood, 43°59'12''N, 40°08'29''E, alt. 940 m., 2016, *Maliček 10449, Palice, Vondrák, Urbanavichus* (Hb Maliček).

Notes: Small, numerous and crowded whitish apothecia are characteristic for *M. pusilla*. The species resembles especially *M. micrococca*, but the apothecia and ascospores are smaller and the thallus usually olive green and warted-granular or membranous. *Micarea micrococca* is characterized by a thallus of bright green aggregated granules. *Micarea pseudomicrococca* Launis & Myllys is a close relative of *M. micrococca* and morphologically similar; however it develops larger apothecia up to 0.4 mm diam., longer ascospores up to 14 µm and wider paraphyses up to 2 µm. *Micarea pusilla* can also be confused with some species in the *M. prasina* complex. Specimens of *M. fallax* growing on wood particularly resembles *M. pusilla*. The main distinguishing characters are secondary metabolites, with *M. fallax* producing micareic acid. Also the ascospore widths of *M. pusilla*, at 2–3 µm, and *M. fallax* at 3–4 µm, differ, and *M. fallax* produce Sedifolia-

grey at least occasionally. Despite the similarities between *M. pusilla* and the species of the *M. micrococca* and *M. prasina* complexes, our phylogenetic studies show that *M. pusilla* resolves outside these groups and is a sister to *M. tomentosa*. Morphologically these two species clearly differ: *M. tomentosa* develops emergent tomentose pycnidia, whereas such pycnidia have never been encountered in *M. pusilla*.

Key to the described species in the *Micarea prasina* group in Europe

1. Thallus containing micareic acid, prasinic acid, xanthonenes or no compounds detectable by thin layer chromatography 2

1'. Thallus containing methoxymicareic or gyrophoric acid 13

2. Thallus containing micareic acid, xanthonenes, or no secondary metabolites, usually on decaying wood or herbs, rarely on bark 3

2'. Thallus containing prasinic acid, usually on soil *M. subviridescens*

3. No secondary metabolites 4

3'. Thallus containing micareic acid, thiophanic acid or xanthonenes 6

4. Stalked tomentose pycnidia always present, apothecia pale to pale brown 5

4'. Pycnidia immersed, apothecia darkly pigmented *M. herbarum*

5. Pycnidia numerous, brownish with white tomentum, walls K+ violet and C+ violet (Sedifolia grey), thallus dull green or olive green, containing orange-brown droplets reacting K+ violet (Intrusa yellow) *M. hedlundii*

5'. Pycnidia brownish with whitish tomentum at least in lower parts, walls K– or if sometimes greyish K± violet (Sedifolia grey), thallus bright green, no orange-brown droplets in the thallus *M. tomentosa*

6. Apothecia usually absent, or if present few 7

6'. Apothecia present and usually abundant 9

7. Forms well-delimited soralia or scattered to confluent clusters of proliferating granules 8
- 7'. Thallus minutely and soresidiously granular or farinose, thick, yellowish, whitish to olive green, bright green when fresh, K– and C– *M. flavoleprosa*
8. Soralia/clusters of proliferating granules pale yellow-green, K– and C+ persistent orange, contains xanthones *M. xanthonica*
- 8'. Soralia greyish-green, usually K+ violet and C+ violet (Sedifolia grey), no xanthones, micareic acid *M. soralifera*
9. Thallus granular, sometimes warted if thin, but never distinctly areolate 10
- 9'. Thallus distinctly areolate and partly granular *M. meridionalis*
10. Apothecia always dark grey to blackish, epihymenium K+ violet, C+ violet (Sedifolia grey), even if growing in shade 11
- 10'. Apothecia brownish or pale to medium grey, usually K+ violet and C+ violet (Sedifolia grey), sometimes partly dark grey if in well-lit habitat 12
11. On decaying hard wood in habitats exposed to light, ascospores $6-8 (-8.5) \times 2-3 \mu\text{m}$, pycnidia numerous especially when apothecia are rare, emergent to stalked *M. nowakii*
- 11'. Usually on bark in shaded and moist habitats, ascospores $7-11 \times 2.5-3.5 (-4) \mu\text{m}$, pycnidia immersed *M. melanobola*
12. Usually on decaying wood, thallus softly granular and medium thick, crystalline granules always in the epihymenium and sometimes also in the hymenium *M. prasina* s.str.

12'. Usually on bark, if on decaying wood thallus thin or membranous, crystalline granules in the hymenium and not in the epihymenium *M. fallax*

13. Thallus and apothecia containing methoxymicareic acid, apothecia present and abundant 14

13'. Thallus and apothecia containing gyrophoric acid (KC+ red), thallus $\pm$ leprose, bright green, apothecia usually absent or rarely few *M. viridileprosa*

14. Apothecia up to 0.6(–0.7) mm wide, often adnate (*M. byssacea* complex) 15

14'. Apothecia up to 0.4 mm wide, rarely adnate 17

15. Thallus minutely granular, olive green, apothecia usually at least partly greyish, K+ and C+ violet (Sedifolia grey) *M. byssacea*

15'. Thallus granular or partly areolate, vivid green, olive green, pale olive green, whitish green or sometimes partly bright green, apothecia cream white to brownish (K– and C–) 16

16. Thallus at least partly small-areolate, apothecia cream white, ascospores 2–3 μ m wide *M. microareolata*

16'. Thallus granular and/or continuous crust, apothecia cream white or brownish, ascospores 3–4 μ m wide *M. laeta*

17. Apothecia 0.2–0.4 mm wide, cream white or greyish and then K+violet, C+ violet (Sedifolia grey) (*M. micrococca* complex) 18

17'. Apothecia very small up to 0.15 (–0.2) mm diam., numerous and crowded, always cream white, K–, C–, thallus thinly granular or membranous *M. pusilla*

18. Thallus warted, cracked to continuous crust without crystalline granules, olive green, apothecia greyish tinged, K+ and C+ violet (Sedifolia grey), ascospores 7–10 $\times$ 2.0–3.5 μ m, paraphyses up to 1.5 μ m wide *M. czarnotae*

18'. Thallus granular, bright to pale olive green with crystalline granules, apothecia cream white (K–, C–), ascospores up to 16 µm long, paraphyses up to 1.5 or up to 2–3 µm wide 19

19. Thallus bright green, ascospores 3–4.5µm wide, one type of paraphyses up to 1.5 µm wide *M. micrococca*

19'. Thallus pale olive green or sometimes partly bright green, ascospores 2–3 µm wide, two types of paraphyses up to 2 µm wide (apices up to 3 µm wide) *M. pseudomicrococca*

DISCUSSION

In this study, we explored species relationships and diversity within the *Micarea prasina* group, focusing on the *M. prasina* complex. Our multilocus phylogeny agrees with the previous single- and multilocus phylogenies of this group (Czarnota & Guzow-Krzemińska 2010; Guzow-Krzemińska *et al.* 2016; van den Boom *et al.* 2017; Launis *et al.* 2019). It should be noted that the *M. prasina* group remains unsupported if treated in a strict sense (van den Boom *et al.* 2017) or unresolved if *M. misella* is included (see Czarnota & Guzow-Krzemińska 2010) (see Fig. 1). Clearly, more gene regions are needed to examine the delimitation of the group. At the species level, the clades are strongly supported. Mainly based on new collections, we discovered three previously unrecognized, well-supported lineages. These new lineages are supported by unique molecular and morphological traits, and also secondary metabolite profiles.

Our major goal was to disentangle the taxonomy of the type species *M. prasina*. The species is morphologically variable (Coppins 1983, Czarnota 2007) and infraspecific genetic variation between European and North American specimens have was (e.g. Czarnota & Guzow-Krzemińska 2010). Our results based on three-loci phylogenies show that *M. prasina* divide into three separate lineages referred here as *M. prasina* 1, 2 and 3 (Fig. 1). Because the original type specimen of *M. prasina* was collected in 1825, and hence is too old for successful DNA amplification, detailed morphological and anatomical studies were performed including microscopic studies of crystalline granules in sections of the apothecia for connecting the correct taxon with the original type. Interestingly, the holotype of *M. prasina* had crystalline granules only in the epihymenium of the apothecial section, a unique feature among the studied species of *Micarea*. Based on these features, and considering ecological and distributional data of the collections, we were able to confirm *M. prasina* 3 as *M. prasina* s. str. We consider the crystalline granules in the epihymenium a significant link in connecting the

specimens. This result also agrees with the earlier morphological and ecological notes of Czarnota & Guzow-Krzemińska (2010). *Micarea prasina* groups 1 and 2, on the other hand, represent putative new taxa, and are discussed in detail under the description of *M. prasina* s.str.

Our results show that the presence and distribution of the crystalline granules may vary within species and between species in the *M. prasina* group. Most of the investigated species, including the new species *M. fallax* and *M. melanobola*, produce granules at least sometimes. However, granules were not detected in the apothecia of *M. flavoleprosa* and *M. pusilla*, although granules were visible in the thallus of the first species. The results for *M. flavoleprosa* should be treated cautiously, because the species is mostly anamorphic and therefore the investigated number of apothecia was very low. It is still unclear what chemical compound or compounds these crystals actually are. Hymenial crystals that dissolve in K have previously been found in *M. neostipitata* Coppins & P. May, and were thought to be lobaric acid (Coppins & May 2001).

Because of the taxonomic uncertainties concerning the *M. prasina* complex, especially the type species *M. prasina* s. str., we discussed the need for an epitype. According to the nomenclatural code “an epitype may be designed when the type material associated with a validly published name is demonstrably ambiguous and cannot be reliably identified for purposes of the precise application of the name of a taxon” (Turland *et al.* 2018). The original holotype of *M. prasina* is in good condition despite its age. Issues concerning the challenges in sequencing old type specimens are quite often used as an argument for designating an epitype (Ariyawansa *et al.* 2014). In the era of DNA barcoding and molecular species identification, epitypification is certainly a practical approach. However, at least for now, the absence of a sequence from a holo- or lectotype is not sufficient reason for creating an epitype. In the case of *M. prasina*, we successfully connected fresh material with the original well preserved type specimen by investigating morphology and the crystalline granules. Therefore, creating an epitype was considered unnecessary.

Previous studies (Coppins 2002; Czarnota 2007; Czarnota & Guzow-Krzemińska 2010; Guzow-Krzemińska *et al.* 2016; van den Boom *et al.* 2017; Launis *et al.* 2019) and our own TLC experiments indicate that secondary metabolites correlate with the phylogenetic relationships in the *M. prasina* group. This correlation appears quite clearly between species complexes. Species in the closely related *M. byssacea* and *M. micrococca* complexes produce methoxymicareic acid whereas species in the *M. prasina* complex produce micareic acid, with two exceptions: *M. subviridescens* produces prasinic acid and no secondary metabolites

were detected in *M. herbarum*. Our results indicate correspondence between growth substrate and secondary chemistry as well: e.g. methoxymicareic acid seems to be common in species that mainly colonize bark. On the other hand, species mostly found on dead wood (e. g. *M. flavoleprosa*, *M. hedlundii*, *M. misella*, *M. nowakii* and *M. tomentosa*) never produce methoxymicareic acid, instead micareic acid or no secondary metabolites at all.

Even in relatively well-studied areas of Europe, species diversity is still rather poorly known. DNA sequencing is often essential for delimiting species, but other characters are still needed especially when DNA methods are unavailable or specimens are too old for sequencing. For example, ascospore size was previously regarded too variable and of little diagnostic value within *M. prasina* s. lato (Coppins 1983), but our new species descriptions show that ascospore width is an important character for species delimitation (see also Launis et al. 2019). However, it is not always easy to determine where one species begins and the other ends, especially in the studies of small crustose lichens such as *Micarea*. For example, what is now known as *M. byssacea*, *M. micrococca* and *M. prasina* complexes, were previously – with some hesitation – interpreted as phenotypic variation within only one species, *M. prasina* (Coppins 1983). Also ecological research and conservation planning rely on our understanding of species and species boundaries. For example *M. flavoleprosa*, is known only from decaying wood, which is a rare and diminishing substrate in managed forests. *Micarea fallax* and *M. melanobola* are, on the other hand, probably quite common. Observations such as these are essential for reliable conservation planning and red list evaluations.

ACKNOWLEDGMENTS

The authors wish to thank emeritus curator Arto Kurtto and emeritus professor Teuvo Ahti for their valuable help with nomenclatural issues. Franz Berger and Jan Vondrák kindly provided their collections for this study. Dr. Juan Carlos Zamora Senoret gave valuable technical help with TreeBASE submission. Also, we are grateful to two anonymous reviewers for their insightful comments and suggestions that improved the manuscript. Financial support for this study was provided by the Societas pro Fauna et Flora Fennica (a personal grant for the first author), and from the research project “Conservation of wood-inhabiting Ascomycetes in changing forest landscapes” (grant number 7000T-YTB079), as part of the research program

on deficiently known and threatened forest species (PUTTE) financed by the Ministry of the Environment.

Work of Dr. Malíček was supported by the long-term research development project RVO 67985939.

Figure 1. Phylogenetic relationships of *Micarea fallax* sp. nov., *M. flavoleprosa* sp. nov., *M. pusilla* sp. nov., and *M. melanobola* (shown in bold). A maximum likelihood phylogram obtained from the RAxML analysis based on the combined ITS, mtSSU and *MCM7* dataset. Branches supported with bootstrap values $\geq 75\%$ in both analyses (RAxML and TNT) are indicated in bold. Bootstrap values $\geq 75\%$ only supported in maximum likelihood analysis are shown above nodes.

Figure 2. Morphological and anatomical features. A–B. *Micarea fallax* sp. nov. (Launis 1710132, H) A apothecial section; B habitus. C. *Micarea flavoleprosa* sp. nov. (holotype, H) habitus. D–E. *Micarea melanobola* (Launis 27123, H) D apothecial section; E habitus. F. *Micarea prasina* s.str. (Launis 229106, H) habitus. G–H. *Micarea pusilla* sp. nov. (holotype, H) G apothecial section; H habitus. Scales: B, C, E, F, H = 1 mm; A, D, G = 100 μ m.

Figure 3. Crystalline granules detected in apothecial sections in polarized light. A. *Micarea fallax* sp. nov. (Launis 1710132, H). B. *Micarea melanobola* (Launis 27123, H). C. *Micarea prasina* 1 (Tønsberg 30856, BG). D. *Micarea prasina* 2 (Sérusiaux, DNA 3437, LG). E. *Micarea prasina* s. str. (Launis 229106, H).

Table 1. List of specimens used for phylogenetic analyses. New species and new sequences generated for the current study are in bold.

Specimens	Locality	Collector, collection number, DNA sample number (where appropriate), herbarium	GenBank accession number		
			ITS	mtSSU	Mcm7
<i>Micarea peliocarpa</i>	USA, Maine	Launis 66123, DNA A324, (H)	MG521544	MG707741	MG692505
<i>M. adnata</i>	Norway	Andersen 48 (BG)	—	AY567751	—
<i>M. byssacea</i>	Finland	Launis 289103, DNA A98, (H)	MG521562	MG707768	MG692527
<i>M. byssacea</i>	Finland	Launis 289102, DNA A97, (H)	MG521563	MG707769	MG692528
<i>M. byssacea</i>	Finland	Launis 66128, DNA A96, (H)	MG521564	MG707770	MG692529
<i>M. czarnotae</i>	Poland	Czarnota 3179 (GPN)	—	EF453674	—
<i>M. czarnotae</i>	Poland	Czarnota 4059 (GPN)	—	EF453663	—
<i>M. czarnotae</i>	Finland	Launis 109111, DNA A604, (H)	—	MG707759	—
<i>M. czarnotae</i>	Finland	Launis 1010133, DNA A455, (H)	MG521557	MG707760	MG692517
<i>M. czarnotae</i>	Belgium	P. van den Boom 50312, DNA 3712, (LG)	—	MG707761	—
<i>M. elachista</i>	Finland	Launis 67113, DNA A340, (H)	MG521548	MG707745	—
<i>M. fallax</i>	Finland	Launis 109115, DNA A605, (H)	—	MK454757	—
<i>M. fallax</i>	Czech Republic	Malíček 6127, DNA A608	—	MK454758	MK456616
<i>M. fallax</i>	Finland	Launis 59132, DNA A559, (H)	MK454942	MK454759	MK456617
<i>M. fallax</i>	Belarus	Tsurykau 001c4, DNA A397, (H)	MK454943	MK454760	MK456618
<i>M. fallax</i>	Scotland	Launis 171143, DNA A646, (H)	—	MK454761	MK456619
<i>M. fallax</i>	Sweden	Svensson 2398, DNA MSv2398, (H)	—	MK454762	—
<i>M. fallax</i>	Finland	Launis 27122, DNA A440, (H)	—	MK454763	—
<i>M. fallax</i>	Finland	Launis 1710132, DNA A718, (H)	—	MK454764	—
<i>M. fallax</i>	Finland	Launis 1010138, DNA A461, (H)	MK454944	MK454765	MK456620
<i>M. fallax</i>	Finland	Launis 1010139, DNA A453, (H)	MK454945	MK454766	MK456621
<i>M. flavoleprosa</i>	France	Sérusiaux s.n., DNA 3841, (LG)	—	MK454754	MK456613
<i>M. flavoleprosa</i>	Czech Republic	Malíček 5098, DNA A616, (PRA)	—	MK454756	MK456615
<i>M. flavoleprosa</i>	Czech Republic	Malíček 4699, DNA A614	—	MK454755	MK456614
<i>M. globulosella</i>	Finland	Launis 67112, DNA A240, (H)	MG521546	MG707743	MG692507
<i>M. globulosella</i>	Finland	Launis 67114, DNA A243, (H)	MG521547	MG707744	MG692508
<i>M. hedlundii</i>	Finland	Launis 67119, DNA A254, (H)	MG521551	MG707749	MG692512
<i>M. herbarum</i>	Netherlands	Brand 63193 (LG)	—	KX459350	—
<i>M. herbarum</i>	Netherlands	P. & G. van den Boom 52575 (LG)	—	KX459349	MG692513
<i>M. laeta</i>	Finland	Launis 59153, DNA A825, (H)	MG521565	MG707771	MG692530
<i>M. laeta</i>	Finland	Launis 49151, DNA A819, (H)	MG521566	MG707772	MG692531
<i>M. laeta</i>	Finland	Launis 1010134, DNA A478, (H)	MG521569	MG707779	MG692539
<i>M. laeta</i>	Finland	Launis 59154, DNA A824, (H)	MG521567	MG707773	MG692532
<i>M. melanobola</i>	Finland	Launis 49141, DNA A808, (H)	—	MK454767	MK456622
<i>M. melanobola</i>	Finland	Launis 116152, DNA A791, (H)	—	MK454768	MK456623
<i>M. melanobola</i>	Finland	Launis 79133, DNA A633, (H)	—	MK454769	MK456624
<i>M. melanobola</i>	Finland	Launis 27123, DNA A437, (H)	MK454946	MK454770	MK456625
<i>M. melanobola</i>	Finland	Launis 56151, DNA A788, (H)	—	—	MK456626
<i>M. melanobola</i>	Finland	Launis 39151, DNA A817, (H)	MK454947	MK454771	MK456627
<i>M. melanobola</i>	Finland	Launis 286152, DNA A813, (H)	MK454948	MK454772	MK456628
<i>M. melanobola</i>	Finland	Launis 266151, DNA A818, (H)	MK454949	MK454773	MK456629
<i>M. melanobola</i>	Finland	Launis 11014, DNA A424, (H)	MK454950	MK454774	MK456630
<i>M. melanobola</i>	Finland	Launis 166151, DNA A809, (H)	—	—	MK456631
<i>M. meridionalis</i>	Portugal	van den Boom hb., DNA 4279 (LG)	—	KX459354	—
<i>M. meridionalis</i>	Portugal	van den Boom hb., DNA 4281 (LG)	—	KX459355	—
<i>M. microareolata</i>	Sweden	Launis 148132, DNA A394 (H)	MG521559	MG707763	MG692519
<i>M. microareolata</i>	Finland	Launis 59152, DNA A826 (H)	MG521560	MG707764	MG692520

<i>M. microareolata</i>	Finland	Pykälä 47787, DNA A797 (H)	—	MG707765	MG692522
<i>M. microareolata</i>	Finland	Launis 59133, DNA A565 (H)	MG521561	MG707766	MG692523
<i>M. microareolata</i>	Finland	Launis 89133, DNA A629 (H)	—	MG707767	MG692524
<i>M. micrococca</i>	Finland	Launis 299101, DNA A100 (H)	MG521552	MG707753	MG692514
<i>M. micrococca</i>	USA, Maine	Launis 146127, DNA A320 (H)	MG521553	MG707754	MG692515
<i>M. misella</i>	Finland	Launis 108111, DNA A264 (H)	MG521545	MG707742	MG692506
<i>M. nowakii</i>	Finland	Launis 245131, DNA A684 (H)	—	MG707751	—
<i>M. nowakii</i>	Poland	Czarnota & Guzow-Krzemińska 4181 (GPN)	—	EF453688	—
<i>M. prasina s.str.</i>	Finland	Launis 265101, DNA A92 (H)	MG521549	MG707747	MG692510
<i>M. prasina s.str.</i>	Finland	Launis 199105, DNA A93 (H)	MG521550	MG707748	MG692511
<i>M. prasina s.str.</i>	USA, Maine	Launis 1361212, DNA A327 (H)	MK517714	MK517718	MK520933
<i>M. prasina s.str.</i>	Scotland	Launis 171144, DNA A644 (H)	MK517713	MK517717	MK520932
<i>M. prasina s.str.</i>	Poland	Czarnota 4319 (GPN)	—	EF453679	—
<i>M. prasina s.str.</i>	Poland	Czarnota 4489 (GPN)	—	EF453678	—
<i>M. prasina s.str.</i>	Poland	Czarnota 3913 (GPN)	—	EF453675	—
<i>M. prasina 2</i>	France	Sérusiaux, DNA 3437 (LG)	—	KX459362	—
<i>M. prasina 2</i>	Belgium	Sérusiaux, DNA 3609 (LG)	—	KX459363	—
<i>M. prasina 1</i>	USA, Tennessee	Tønberg 30856 (BG)	—	AY756452	—
<i>M. pseudomicrococca</i>	Finland	Launis 59151, DNA A811 (H)	MG521554	MG707755	—
<i>M. pseudomicrococca</i>	Finland	Launis 89132, DNA A599 (H)	MG521555	MG707756	—
<i>M. pseudomicrococca</i>	Finland	Launis 258131, DNA A603 (H)	—	MG707757	—
<i>M. pseudomicrococca</i>	Scotland	Launis 171141, DNA A645 (H)	MG521556	MG707758	MG692516
<i>M. pusilla</i>	Finland	Launis 1010137, DNA A460 (H)	MK454941	MK454752	MK456611
<i>M. pusilla</i>	Finland	Launis 101035, DNA A464 (H)	—	MK454753	MK456612
<i>M. pusilla</i>	Finland	Launis 1010136, DNA A470 (H)	MK454940	MK454751	MK456610
<i>M. pycnidiphora</i>	USA	Tønberg 30881 (BG)	—	AY567754	—
<i>M. soralifera</i>	Poland	Kukwa 13001 (GPN)	KT119887	KT119886	—
<i>M. soralifera</i>	Finland	Launis 1710131, DNA A714 (H)	—	MG707746	MG692509
<i>M. sp. lineage A</i>	Scotland	Launis 171142, DNA A648 (H)	MG521571	MG707782	MG692542
<i>M. stipitata</i>	USA	Ekman s.n.	—	AY567753	—
<i>M. subviridescens</i>	Scotland	Czarnota 3599 (GPN)	—	EF453666	—
<i>M. synotheoides</i>	Norway	Andersen 47 (BG)	—	AY567756	—
<i>M. tomentosa</i>	Finland	Launis 11013, DNA A773 (H)	—	MG707750	—
<i>M. tomentosa</i>	Poland	Czarnota 3949 (GPN)	—	EF453686	—
<i>M. viridileprosa</i>	Poland	Czarnota 3436 (GPN)	—	EF453671	—
<i>M. viridileprosa</i>	Poland	Czarnota 3869 (GPN)	—	EF453673	—
<i>M. xanthonica</i>	USA	Tønberg 25674 (BG)	—	AY756454	—

LITERATURE CITED

- Andersen H, Ekman S. 2005. Disintegration of the *Micarea* (lichenized *Ascomycota*): a molecular phylogeny based on mitochondrial rDNA sequences. *Mycological Research* 109: 21–30.
- Ariyawansa HA, Hawksworth DL, Hyde KD, Jones EBG, Maharachchikumbura SSN, Manamgoda DS, Thambugala KM, Udayanga D, Camporesi E, Daranagama A, Jayawardena R, Liu J-K, McKenzie EHC, Phookamsak R, Senanayake IC, Shivas RG, Tian Q, Xu J-C. 2014. Epitypification and neotypification: guidelines with appropriate and inappropriate examples. *Fungal Diversity* 69: 57–91.
- Brodo IM. 1984. The North American species of the *Lecanora subfusca* group. *Nova Hedwigia* 79: 63–185.
- Coppins BJ. 1983. A taxonomic study of the lichen genus *Micarea* in Europe. *Bulletin of the British Museum (Natural History)*, Botany series 11: 17–214.
- Coppins BJ, May PF. 2001. *Micarea neostipitata*, a new species with pale stipitate pycnidia from eastern North America. *Lichenologist* 33: 487–490.
- Coppins BJ. 2002. Checklist of lichens of Great Britain and Ireland 86. 2002 London: British Lichen Society.
- Coppins BJ. 2009. *Micarea*. In: Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW, Wolseley PA eds. *The Lichens of Great Britain and Ireland*: London: British Lichen Society. p. 583–606.
- Culberson CF, Kristinsson HD. 1970. A standardized method for the identification of lichen products. *Journal of Chromatography A* 46: 85–93.
- Czarnota P, Guzow-Krzemińska B. 2010. A phylogenetic study of the *Micarea prasina* group shows that *Micarea micrococca* includes three distinct lineages. *Lichenologist* 42: 7–21.

- Czarnota P. 2007. The lichen genus *Micarea* (Lecanorales, Ascomycota) in Poland. Polish Botanical Studies 23: 190 p.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic acids research 32: 1792–1797.
- Ekman S, Andersen H, Wedin M. 2008. The Limitations of Ancestral State Reconstruction and the Evolution of the Ascus in the Lecanorales (Lichenized Ascomycota). Systematic Biology 57(1):141-156.
- Ekman S, Svensson M. 2014. *Brianaria* (Psoraceae), a new genus to accommodate the *Micarea sylvicola* group. Lichenologist 46: 285–294.
- Erichsen CFE. 1939. Mitt. Inst. Allg. Bot. Hamburg 10: 407.
- Felsenstein J. 1985. Phylogenies and the comparative method. American Naturalist 125: 1–15.
- Goloboff P, Farris J, Nixon K. 2008. TNT, a free program for phylogenetic analysis. Cladistics 24: 774–786.
- Guzow- Krzemińska B, Czarnota P, Łubek A, Kukwa M. 2016. *Micarea soralifera* sp. nov., a new sorediate species in the *M. prasina* group. Lichenologist 48: 161–169.
- Hafellner J. 1984. Studien in Richtung einer natürlicheren Gliederung der Sammelfamilien Lecanoraceae und Lecideaceae. Beiheft zur Nova Hedwigia 79: 241–371.
- Hafellner J, Türk R. 2016. Die lichenisierten Pilze Österreichs – eine neue Checkliste der bisher nachgewiesenen Taxa mit Angaben zu Verbreitung und Substratökologie. [The lichenized fungi of Austria – a new checklist of the taxa so far recorded, with data to distribution and substrate ecology.] Stapfia 104/1: 98.

- Hedlund JT. 1892. Kritische Bemerkungen über einige Arten der Flechtengattungen *Lecanora* (Ach.), *Lecidea* (Ach.) und *Micarea* (Fr.). Bihang till Kongliga Svenska Vetenskaps-Akademiens Handlingar III 18: 1–104.
- Kauff F, Lutzoni F. 2002. Phylogeny of the *Gyalectales* and *Ostropales* (Ascomycota, Fungi): among and within order relationships based on nuclear ribosomal RNA small and large subunits. *Molecular Phylogenetics and Evolution* 25: 138–156.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Ainsworth & Bisby's Dictionary of the Fungi. 10th edition ed. CABI Publishing, UK.
- Launis A, Pykälä J, van den Boom P, Sérusiaux E, Myllys L. 2019. Four new epiphytic species in the *Micarea prasina* group from Europe. *The Lichenologist* 51(1): 7–25.
- Leavitt SD, Johnson L, Goward T, St. Clair L. 2011. Species delimitation in taxonomically difficult lichen-forming fungi: an example from morphologically and chemically diverse *Xanthoparmelia* (Parmeliaceae) in North America. *Molecular Phylogenetics and Evolution* 60: 317–32.
- Maddison DR, Maddison WP. 2005. MacClade 4: Analysis of Phylogeny and Character Evolution. Sunderland, Massachusetts: Sinauer Associates.
- Malíček J, Berger F, Bouda F, Cezanne R, Eichler M, Halda JP, Langbehn T, Palice Z, Šoun J, Uhlík P & Vondrák J 2017. Lišejníky zaznamenané během bryologicko-lichenologického setkání v Mohelně na Třebíčsku na jaře 2016 [Lichens recorded during the Bryological and Lichenological meeting in Mohelno (Třebíč region, southwestern Moravia) in spring 2016]. *Bryonora* 60: 24–45.
- Meyer B, Printzen C. 2000. Proposal for a standardized nomenclature and characterization of insoluble lichen pigments. *Lichenologist* 32: 571–583.

- Myllys, L., Lohtander, K., Källersjö, M., Tehler, A. 1999: Sequence insertion and ITS data provide congruent information in *Rocella canariensis* and *R. tuberculata* (Arthoniales, Euascomycetes) phylogeny. *Molecular Phylogenetics and Evolution* 12: 295-309.
- Myllys L, Velmala S, Holien H, Halonen P, Wang LS, Goward T. 2011. Phylogeny of the genus *Bryoria*. *Lichenologist* 43: 617–638.
- Nimis PL, Hafellner J, Roux C, Clerc P, Mayrhofer H, Martellos S, Bilovitz PO. 2018. The lichens of the Alps – an annotated checklist. *MycKeys* 31: 1-634. <https://doi.org/10.3897/mycokeys.31.23568>
- Nylander W. 1867. Flora oder Allgemeine botanische Zeitung herausgegeben von der Königl. bayer. botanischen Gesellschaft in Regensburg. p. 371.
- Nylander W. 1874. *Lecidea prasiniza*. *Flora* 57: 312.
- Orange A, James PW, White FJ. 2001. Microchemical methods for the identification of lichens. *British Lichen Society* p. 44–45.
- Poelt J, Döbbeler P. 1975: Über moosparasitische Arten der Flechtengattung *Micarea* und *Vezdaea*. - *Bot. Jahrb. Syst.* 96(1-4): 328–352.
- Schmitt I., Crespo A, Divakar P, Fankhauser J, Herman-Sackett E, Nelsen M. 2009. New primers for single-copy protein-coding genes for fungal systematics. *Persoonia* 23: 35–40.
- Schmull M, Miadlikowska J, Pelzer M, Stocker-Wörgötter E, Hofstetter V, Fraker E, Hodkinson B, Reeb V, Kukwa M, Lumbsch HT. 2011. Phylogenetic affiliations of members of the heterogeneous lichen-forming fungi of the genus *Lecidea* sensu Zahlbruckner (Lecanoromycetes, Ascomycota). *Mycologia* 103: 983–1003.

Sérusiaux E, Brand AM, Motiejūnaitė J, Orange A, Coppins BJ. 2010. *Lecidea doliiformis* belongs to *Micarea*, *Catillaria alba* to *Biatora* and *Biatora ligni-mollis* occurs in western Europe. *Bryologist* 113: 333–344.

Sohrabi M, Myllys L, Stenroos S. 2010. Successful DNA sequencing of a 75 year-old herbarium specimen of *Aspicilia aschabadensis* (J. Steiner) Mereschk. *Lichenologist* 43: 626–628.

Spribille T, Klung B, Mayerhofer H. 2011. A phylogenetic analysis of the boreal lichen *Mycoblastus sanguinarius* (Mycoblastaceae, lichenized Ascomycota) reveals cryptic clades correlated with fatty acid profiles. *Molecular Phylogenetics and Evolution* 59: 603–614.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313.

Thiers B. Continuously updated. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/>.

Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber WH, Li DZ, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ & Smith GF (eds.). 2018. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Vegetabile* 159. Glashütten: Koeltz Botanical Books. DOI <https://doi.org/10.12705/Code.2018>

van den Boom PPG, Brand AM, Coppins BJ, Sérusiaux E. 2017. Two new species in the *Micarea prasina* group from Western Europe. *Lichenologist* 49: 13–25.

Vězda A, Wirth V. 1976. Zur Taxonomie der Flechtengattung *Micarea* Fr. em. Hedl. *Folia Geobot. Phytotax.* 11: 93–102.

Vondrák J, Malíček J, Palice Z, Coppins B, Kukwa M, Czarnota P, Sanderson N & Acton A 2016. Methods for obtaining more complete species lists in surveys of lichen biodiversity. *Nordic Journal of Botany* 34: 619–626.

White TJ, Bruns T, Lee S, Taylor JW. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. *PCR protocols: A guide to the methods and applications*. New York, NY: Academic Press. p. 315–322.

Zoller S, Scheidegger C, Sperisen C. 1999. PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming Ascomycetes. *Lichenologist* 31: 511–516.