<https://helda.helsinki.fi>

New insights from XRF core scanning data into boreal lake
ontogeny during the Eemian (Marine Isotope Stage 5e) at Sokli,
northeast Finland

Kylander, Malin E.

2018-01

Kylander , M E , Pliikk , A , Rydberg , J , Löwemark , L , Salonen , J S ,
Fernandez-Fernandez , M & Helmens , K 2018 , ' New insights from XRF core scanning data
into boreal lake ontogeny during the Eemian (Marine Isotope Stage 5e) at Sokli, northeast
Finland ' , Quaternary Research , vol. 89 , no. 1 , pp. 352-364 . <https://doi.org/10.1017/qua.2017.84>

<http://hdl.handle.net/10138/326392>
<https://doi.org/10.1017/qua.2017.84>

---

cc_by_nc_nd
acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **New Insights from XRF Core Scanning Data into Boreal Lake**
2 **Ontogeny During the Eemian (MIS 5e) at Sokli, NE Finland**

3 Malin E. Kylander^{1,2}, Anna Pliikk^{2,3}, Johan Rydberg⁴, Ludvig Löwemark⁵, J. Sakari
4 Salonen⁶, María Fernández-Fernández⁷, and Karin Helmens^{2,3}

5 ¹*Department of Geological Sciences, Stockholm University, SE-10691 Stockholm, Sweden Tel: +46*
6 *(0)8-674 7898; Fax: +46 (0)8-674-7897*

7 ²*The Bolin Centre for Climate Research, Stockholm University, SE-10691 Stockholm, Sweden*

8 ³*Department of Physical Geography, Stockholm University, SE-10691 Stockholm, Sweden*

9 ⁴*Department of Ecology and Environmental Sciences, Umeå University, SE-901 87 Umeå, Sweden*

10 ⁵*Department of Geosciences, National Taiwan University, 106 17, Taipei, Taiwan*

11 ⁶*Department of Geosciences and Geography, FI-00014 University of Helsinki, Finland*

12 ⁷*Instituto de Investigaciones Agrobiológicas de Galicia, CSIC, Apartado 122, E-15780 Santiago de*
13 *Compostela, Spain*

14 *corresponding author: malin.kylander@geo.su.se*

15 **Key words:** Eemian, XRF core scanning, geochemistry, lake sediment, boreal, ontogeny

17 Biological proxies from the Sokli Eemian (MIS 5e) paleolake sequence from NE
18 Finland have previously shown that, unlike many postglacial records from boreal
19 sites, the lake becomes increasingly eutrophic over time. Here, principal
20 components (PC) were extracted from a high resolution multi-element XRF core
21 scanning dataset to describe minerogenic input from the wider catchment (PC1),
22 the input of S, Fe, Mn, and Ca-rich detrital material from the surrounding Sokli
23 Carbonatite Massif (PC2), and chemical weathering (PC3). Minerogenic inputs to
24 the lake were elevated early in the record and during two abrupt cooling events
25 when soils and vegetation in the catchment were poor. Chemical weathering in
26 the catchment generally increased over time, coinciding with higher air
27 temperatures, catchment productivity, and the presence of acidic conifer species.
28 Abiotic edaphic processes play a key role in lake ontogeny at this site stemming
29 from the base cation- and nutrient-rich bedrock, which supports lake alkalinity
30 and productivity. The climate history at this site, and its integrated effects on the
31 lake system, appear to override development processes and alters its long-term
32 trajectory.

33

34   **Introduction**

35   The last interglacial, the Eemian (roughly equivalent to Marine Isotope Stage
36   (MIS) 5e, ca. 130-115 ka ago), was a time when certain regions of the world
37   experienced a warmer-than-present climate. While studies from temperate
38   central Europe suggest that the Eemian was climatically rather stable,
39   experiencing only minor short-term variability (Cheddadi et al., 1998; Kühl et al.,
40   2007; Rioual et al., 2007; Bińka et al., 2011), records from northern Europe and
41   the North Atlantic region have registered pronounced Eemian climate changes,
42   seen as cooling and aridity pulses (Cortijo et al., 1994; Seidenkrantz et al., 1995;
43   Fronval and Jansen, 1996; Björck et al., 2000; Bauch et al., 2011; Irvalı et al.,
44   2012; Galaasen et al., 2014). Due to glacial erosion at high latitudes, Eemian
45   sequences are rare at boreal sites in northern Europe. An exception to this is the
46   thick and continuous Eemian lake sediment sequence from Sokli, NE Finland.
47   This sedimentation basin was formed through the action of ice and water
48   working along a fracture zone bisecting the Sokli Carbonatite Massif (SCM), a
49   carbonatite-alkaline intrusion in the crystalline Precambrian Shield (Vartiainen,
50   1980).

51   Palynological analysis first proposed an Eemian age for the diatom gyttja
52   deposit (Ilvonen, 1973) found between 25 and 16 m of depth at Sokli. The
53   bracketing glaciofluvial sands and gravel (28.5 and 25 m) and fluvial sands (16 to
54   13 m) were later dated by thermoluminescence (TL) and infrared stimulated
55   luminescence (IRSL) to >ca. 110 ka and <ca. 150-180 ka, respectively (Helmens
56   et al., 2000; 2007). Additional optically stimulated luminescence (OSL) dating of
57   the upper fluvial sands revealed an age >ca. 95 ka (Alexanderson et al., 2008).

While an absolute chronology for this sequence is unavailable, previous studies have divided the lake development into five main phases: glacio-lacustrine, early-, early mid-, late mid-, and late-Eemian (Plikk et al., 2016). Superimposed on these phases are two abrupt climate change events (Helmens et al., 2015; Plikk et al., 2016). The earlier event reveals a cooling similar to that seen in the North Atlantic Ocean (Bauch et al., 2011; Irvali et al., 2012; Galaasen et al., 2014). Occurring after full interglacial conditions were reached, the “Tunturi event” saw summer temperatures drop 2-4°C over a time frame of ca. 500-1000 years. Similarly, the later event implies cooler and drier conditions at Sokli (Helmens et al., 2015; Plikk et al., 2016).

Reconstructing past paleoenvironmental change using postglacial lake sediments must consider a lake’s natural development, or *ontogeny*, as well as the external climate signal we are often trying to capture. Studies in boreal landscapes on late glacial/Holocene and recent timescales suggest a common pattern towards natural oligotrophication over time. After ice retreat the catchment is stabilized through the gradual build-up of soils and colonization by vegetation. In boreal regions this vegetation typically sees the transition from *Betula* to more acidic *Pinus* and *Picea* conifer species (Donner, 1995). The growing catchment vegetation also retains more nutrients, making them unavailable for lake biota. Soluble minerals in the soils and exposed bedrock surfaces (e.g., apatite, biotite, hornblende) are weathered and base cations are consumed, leading to the loss of alkalinity in the lake water. As soil organic matter (SOM) and dissolved organic carbon (DOC) in the catchment increase there is an influx of increasingly acidic water to the lake. These effects can be further enhanced by the inhibition of groundwater recharge by hardened soil

horizons; this reduces contact with the bedrock and soils, and therefore leaching of base cations, and increases flow through humic-rich soils (Ford, 1990; Pienitz et al., 1999; Engstrom et al., 2000; Fritz et al., 2004; Boyle, 2007; Fritz and Anderson, 2013; Law et al., 2015).

In contrast to the pattern commonly seen at boreal sites, previous studies based on biological proxies (diatoms, pollen, biogenic silica (BSi), total organic carbon (TOC), chironomids) show that the lake at Sokli becomes increasingly eutrophic over time (Helmens et al., 2015; Pliikk et al., 2016). Within the literature, there is a discussion as to the relative importance of abiotic and biotic processes to lake development. Most studies have however, been restricted to lakes in base cation-poor and mixed terrains (e.g., Whitehead et al., 1989; Ford, 1990; Pienitz et al., 1999; Engstrom et al., 2000; Fritz et al., 2004; Law et al., 2015) or modelling exercises (Boyle, 2007). Sokli differs from a typical boreal lake in that it sits in the base cation-rich SCM, providing us with an alternative geological, and therefore abiotic, setting in which to examine postglacial boreal lake ontogeny. The elemental chemistry of the sediments is controlled by the interplay between the geology, hydrology, and biology of the lake and its catchment and the external climate changes. We use Principal Component Analysis (PCA) and supporting ratio and correlative tools (e.g., Kylander et al., 2011; Ljung et al., 2015) from a high-resolution XRF core scanning dataset to reveal detail about ontogeny relevant processes, such as soil development/erosion, nutrient and base cation supply, and productivity. We then place these processes into context by comparing with previously published proxy data and attempt to understand what factors control the unusual lake ontogeny at Sokli.

108 **Materials and Methods**

109 ***Sampling Site***

110 Sokli (67°48' N, 29°18'E), surface elevation 220 m a.s.l., is situated in the SCM
111 located in NE Finland (Figure 1A). The present day climate around Sokli is cold
112 boreal with a mean annual temperature of 1°C and annual precipitation averages
113 between 500-550 mm yr⁻¹ (Drebs et al., 2002). The vegetation is northern boreal
114 forest and mires are widespread in the area. Soils in the region are generally
115 composed of acidic podzol soils.

116 The carbonate-rich magma intrusion of the SCM penetrated through the
117 crystalline Precambrian Shield about 350 Ma ago. Today, the deeply eroded
118 carbonatite intrusion forms a circular depression, ca. 5 km in diameter,
119 surrounded by a hilly ring of fenites (up to 320 m a.s.l.)(Figure 1B). Dispersed
120 residual phosphorous deposits can be found across the carbonatite bedrock
121 (Talvitie et al., 1981). The deeper part of the depression is now infilled with
122 sedimentary deposits, the thickest of which are found in a string of hollows
123 extending along a NE-SW trending fracture valley that crosses the SCM. The
124 sequence studied here is from the Sokli 2010/4 borehole which was retrieved by
125 percussion drilling in March 2010 by the Geological Survey of Finland.

126 ***Sample Analysis***

127 After retrieval all sediment cores were described in the laboratory. Prior to any
128 sub-sampling, the sediments were scanned at the SLAM Lab at the Department of
129 Geological Sciences, Stockholm University, using an ITRAX XRF Core Scanner
130 from Cox Analytical Systems (Gothenburg, Sweden). Analyses were made using a
131 Mo tube set at 30 kV and 45 mA with a step size of 2.5 mm and a dwell time of 60

s. Based on analytical performance (counting statistics above background), reliable data was acquired for Si, S, K, Ca, Ti, Mn, Fe, Rb, Sr, and Zr. All data presented here are normalized to the (incoherent+coherent) scattering to remove various instrumental effects and then smoothed using a 5-point running mean to capture the main shifts. Full analytical details of BSi, TOC, siliceous microfossil, and pollen analyses can be found in Plikk et al. (2016).

Statistical Methods

Data interpretation was aided by PCA, which was made using JMP 9.0.0 software in correlation mode using a Varimax rotation. Before analysis all data were converted to Z-scores calculated as $(X_i - X_{avg})/X_{std}$, where X_i is the normalized elemental peak areas and X_{avg} and X_{std} are the series average and standard deviation, respectively, of the variable X_i . This was done to avoid scaling effects and to obtain average-centred distributions (Eriksson et al., 1999). All principal components (PCs) with Eigenvalues above 1 were extracted. In order to reduce the amount of data manipulation when making correlations between data of different resolution, a single XRF core scanning data point was selected for matching TOC, TIC, and BSi sample depths.

Results

Stratigraphy

The sediment between 25 and 16 m is mainly comprised of diatom gyttja, which is divided into thirteen units based on the stratigraphy and sediment properties. Due to the open core head during drilling several cores were compressed but this has been corrected for in the stratigraphic column presented here. Note that

155 this compression may have affected the visibility of some laminations, namely in
156 Units C, F, G, and H. Details of the compressed sections are given in Plikk et al.
157 (2016).

158   The lowermost Unit A (25.00-23.41 m) consists of partly laminated
159 brown gyttja with thin yellowish laminations (Figures 2-5). In Unit B (23.41-
160 23.33 m) laminations stop and the diatom gyttja becomes distinctly more silty.
161 The beginning of Unit C (23.33-21.50 m) is composed of silty diatom gyttja.
162 Laminations disappear in the diatom gyttja between 22.75-22.25 m. Partly
163 laminated brown gyttja is found above this layer which is replaced again by a
164 homogenous diatom gyttja. Unit D (21.50-21.00 m) consists of dark brown gyttja
165 with thin yellowish laminae and 1 cm thick dark bands every 5-10 cm. In the
166 upper part of the unit the thin laminae become slightly distorted (21.20-21.00
167 m). In Unit E (21.00-20.73 m) the diatom gyttja is vaguely laminated and slightly
168 reddish. The sediment was disturbed during coring between 20.73-20.50 m and
169 is not considered. Unit F (20.50-19.86 m) has light brown gyttja with less
170 distinct and more sporadic laminations. With the transition to Unit G (19.86-
171 19.50 m) the laminations disappear. Instead, this unit consists of darker brown
172 homogeneous gyttja. Unit H (19.50-17.41 m) consist of brown homogeneous
173 gyttja although dark laminations were observed during coring. In Unit I (17.41-
174 17.33 m), the gyttja once again becomes silty. In Unit J (17.33-17.07 m) the
175 homogeneous brown gyttja is spotted with light yellow granules. Unit K (17.07-
176 16.75 m) shows a gradual transition into a light grey-reddish spotted diatom
177 gyttja. Unit L (16.75-16.58 m) is the last unit consisting of brown homogeneous
178 diatom gyttja. Unit M (16.58-16.00 m) consists of ca. 10 cm thick layers of brown
179 gyttja with small yellowish spots intercalated with five 2-3 cm thick layers of

180 homogeneous sand centred on 16.57, 16.47, 16.37, 16.30, and 16.23 m. In the
181 uppermost 20 cm the gyttja is sandy.

182 ***XRF Core Scanning Data***

183 In a paleoclimatic context, it is the relative changes in the elemental XRF core
184 scanning profiles, rather than the absolute concentrations, that are of interest.
185 XRF core scanning analyses can be negatively affected by large variations in the
186 matrix (Chawchai et al., 2016). At Sokli the studied sediments are mainly
187 composed of gyttja, which reduces the importance of matrix effects on the data.
188 With the aim of reconstructing paleoenvironmental change at Sokli, the acquired
189 data were used in a PCA. The raw data are presented in the Supplementary
190 Information and can be downloaded from www.bolin.su.se/data/xxx.

191 ***PCA, Total Organic Carbon, and Elemental Ratios***

192 Three main PC explain 81% of the variance in the data (Table 1). The first
193 component (PC1), explaining 42% of the data, is associated with K, Ti, Rb, Zr, Si,
194 and Ca (Table 2). PC2 explains 26% of the variance in the data and is associated
195 with S, Fe, Ca, and Mn. PC3 explains 13% of the variance in the data linked to the
196 behaviour of Sr and Ca.

197 The plotting of PC factors scores by depth indicates when the processes
198 described by the PC are more intense (having more positive values) or less
199 intense (having more negative values). PC1 factor scores are slightly elevated at
200 the base of Unit A and then decrease towards the middle of the unit, fluctuating
201 around a value of zero (Figure 2). PC1 factors scores peak at the end of the unit
202 at 23.55 m. A second peak is observed at the transition to Unit B at 23.41 m in the

203 siltier layer. In Unit C factor scores decrease slightly from the base but remain
204 just below zero up into Unit H. High positive values are observed in Unit I, again
205 coinciding with a siltier layer. Peaks in PC1 are observed in Unit M matching with
206 visible sand layers. Although the TOC data is much lower resolution (n=102 vs.
207 n=2404), PC1 and TOC are significantly negatively correlated ($r=-0.57$, $n=102$,
208 $p=0.001$),

209   For PC2 the highest and largest amplitude changes in factor scores are
210 found overprinted on an overall decreasing trend in Unit A (Figure 3). This is
211 capped by a large peak in Unit B. From Unit C through to Unit H factor scores are
212 slightly negative with occasional positive incursions and the frequency of the
213 shifts generally tightens moving up the profile. The lowest PC2 factor scores of
214 the profile are seen at the boundary of Units J and K while values increase up into
215 Unit L. Peaks observed in Unit M are slightly out of phase with the sand layers.
216 PC2 is significantly negatively correlated with inferred values of BSi from FTIR-
217 measurements ($r=-0.57$; $n=802$; $p\text{-value} < 0.001$).

218   In general PC3 factor scores are negative in Unit A (Figure 4). PC3 factor
219 scores peak in Unit B while Unit C initially has the lowest factor scores of the
220 profile. From Unit D through to Unit H there is a long-term trend towards
221 increasingly positive values. At the top of Unit H positive values are reached and
222 more prominent peaks are observed. PC3 factor scores increase in Unit I and
223 then decrease in Units J and K; this trend is reversed in Unit L. In Unit M peaks in
224 PC3 occur for the most part at the same time as the sand layers. PC3 is positively
225 correlated to inferred values of total inorganic carbon (TIC) from FTIR-
226 measurements ($r=0.58$; $n=851$; $p\text{-value} < 0.001$; Rosén et al. 2010).

227 Si/Ti increases from the base of Unit A but has some of the lowest values
228 seen in the profile at the transition into Unit B (Figure 4). From this unit upwards
229 Si/Ti gradually increases peaking at the top of Unit E. An apparent decrease in
230 Si/Ti occurs in Unit F followed by a gradual increase until approximately halfway
231 through Unit H. From this point Si/Ti starts a gradual long-term decline that
232 continues to the top of the sequence and is punctuated by low values in Units I
233 and J. Si/Ti and BSi values are significantly correlated ($r=0.67$, $n=802$, $p<0.001$).

234 **Discussion**

235 The Sokli Eemian record has been divided into five main lake phases (L1-L5)
236 based on a number of biotic proxies (Plikk et al., 2016). Additionally, two abrupt
237 climate change events, E1 (the Tunturi Event) and E2, are distinguished
238 (Helmens et al., 2015). Note that L1 corresponds to the initial phase with glacial
239 lake sedimentation and is not included in this study. We first establish what
240 processes control the sediment geochemistry and then place these into context
241 by comparing with previously published proxy data. Finally, we relate changes in
242 our record to key factors controlling lake ontogeny in order to understand why
243 Sokli differs from other boreal sites by becoming increasingly eutrophic.

244 ***Identifying the Processes: Interpretation of the PCA and Si/Ti Ratios***

245 The purpose of PCA is to reduce the dimensions of a multivariate data set and
246 explain as much variability as possible with a few components (Reimann et al.,
247 2008), thus enabling the identification of variables (i.e., chemical elements in this
248 case) whose behaviour is similar and, thereby, likely controlled by the same
249 process. Three PC were extracted and based on the elemental groupings we are

able to reconstruct changes in detrital input from outside the SCM (PC1)(Figure 2), detrital input from within the SCM (PC2)(Figure 3), and chemical weathering (PC3)(Figure 4). Si/Ti is used as an indicator of lake productivity (Figure 4).

PC1 is represented by Zr, Ti, K, Rb, Ca, and Si (Table 1), which are all part of the crystal structure of common minerals (e.g., silicates, feldspars, micas, titanium oxides, zircon, calcite, apatite) and are therefore transported as detrital material to the lake (Deer et al., 1992). This interpretation is supported by the stratigraphy, where high PC1 scores coincide with visible silt and sand layers (Figure 2). Based on our interpretation of PC2 given below, PC1 is thought to be dominated by materials coming from outside the SCM or possibly, till deposited within the SCM, with periods of higher/lower input indicated by positive/negative factors scores. TOC, which is negatively correlated to PC1, is similarly linked to allogenic inputs from the catchment. When TOC is high, the catchment is more productive with more vegetation available to hold materials in place and reduce erosion, which results in negative PC1 scores.

PC2 is driven by variations in S, Fe, Mn, and Ca (Table 1). The carbonatite rocks of the SCM have relatively high Ca concentrations (6-34%) while some of the later stages of magmatic carbonatite formation resulted in rocks with up to 5% S, as well as higher than average Fe and Mn concentrations (Vartiainen, 1980). The elements associated with PC2 can be transported to the lake as particulate material bound to minerals, in dissolved form, or bound to organic matter (Holmer and Storkholm, 2001). Peaks in PC2 coincide with laminations in the sediments, suggesting a relation to anoxic conditions (Figure 3)(c.f. Pliikk et al., 2016). However, even though Fe, S, Mn, and Ca can all precipitate in the water column after being transported in solution to the lake, they experience very

different behaviour when it comes to redox conditions (Davison, 1993). If changes in redox conditions and the preservation of precipitated Fe- and Mn-hydroxides and oxyhydroxides were the process driving PC2, it would be expected that Fe and Mn would show opposite patterns, with stronger anoxia giving high Fe and low Mn and vice versa (Wetzel, 2001). Since both elements load positively on PC2 it is unlikely that anoxia is controlling their accumulation in the sediment. Additionally, even if the SCM contains carbonate minerals, the sediment itself contains very low amounts of carbonates (<0.5 % based on FTIR modelled TIC; Rosén et al., 2010) and hence, Ca is most likely associated with crystalline mineral particles (see below regarding carbonates in association to PC3). Thus increases/decreases in PC2 factor scores describe greater/lesser inputs of non-carbonaceous detrital material originating from SCM rock types rich in mainly S and Fe, but also Ca and Mn (e.g., phoscorites, micas, late stage magmatic carbonatites; Vartiainen, 1980). This said, there have likely been variations in the redox conditions in the lake during the Eemian, and periods preserving laminations most likely had more anoxic conditions as compared to periods without laminations (cf. Plikk et al., 2016).

PC3 is associated with Sr and Ca (Table 1). Strontium is hosted in Ca bearing minerals such as plagioclase, amphibole, pyroxene, and carbonate minerals (Cohen, 2003). It can enter the system detritally or as Sr^{2+} chemically weathered from the soils and rocks of the catchment. Strontium-rich veins of dolomite carbonatite are only found near the centre of the SCM complex and the surrounding fenites (Lee et al., 2006). The preferential erosion of these veins to produce Sr- and Ca-rich detrital material seems unrealistic and if this did occur, it would be integrated into PC2. This indicates that Sr and Ca are being washed

into the lake in solution. Once in the water column Sr and Ca are either co-precipitated and/or absorbed onto the surface of organic matter or Fe-Mn oxides, eventually sinking to the bottom of the lake and being buried in the sediments (Cohen, 2003). Given the fact that Fe and Mn do not load on PC3, it is most likely that soluble Ca and Sr are delivered to the sediment through precipitation, although the carbonate concentrations of the sediment are low, and absorption to organic matter. That carbonate precipitation plays a role in Sr and Ca variations and PC3 is verified by the positive correlation between TIC and PC3. Zeng et al. (2013) showed that non-residual Sr has a strong positive correlation with chemical weathering and can be used as a proxy for this process. The pattern of chemical weathering itself is controlled by a number of often related, time-variable factors including physical erosion and the presence of fresh material, temperature, source mineralogy, precipitation, runoff, soil cover, organic matter, and vegetation (White and Blum, 1995; Oliva et al., 2003; Riebe et al., 2004; West et al., 2005; Mavris et al., 2010). As such we stress that PC3 is the integrated expression of several factors related to chemical weathering and the input of soluble Sr and Ca to the lake with increased/decreased chemical weathering signalled by higher/lower factor scores (Figure 4).

It should be kept in mind that the PCs model the integrated behaviour of several elements based on the variability observed in the entire profile. Therefore some short term events observed in the PC factor scores may not capture the true behaviour of an element at all times. An inverse correlation should be observed between $1/\text{Sr}$ and PC3 in those samples where Sr is the dominant control on the behaviour of PC3 (Figure 5). This is found to be the case except during periods with higher detrital input (i.e., E1, E2, and sand layers in

Unit M). Thus, during periods with higher minerogenic flux (positive PC1 scores) we interpret our chemical weathering proxy with caution.

Bearing in mind the fundamental theory of PCA, the individual PC should also not correlate when considering the whole profile. This does not exclude the possibility however, that during a given lake phase the PC cannot show some co-variation, as the processes described by the PC may be diminished or enhanced at the same time (Figure 6, Table 3). PC1 and PC2 correlation little (with the exception of the negative correlation in L3) in the different lake phases. This demonstrates that the importance of sources outside the SCM (PC1) and within the SCM (PC2) change over time. This could be a result of changes in (i) vegetation and linked soil erosion; (ii) hydrological pathways; (iii) the size of the lake; and (iv) energy of the transporting waters. While we do not have grain size analyses, PC1 increases during the visibly siltier and sandier layers, which would evoke a coarser grain size for those minerals associated with PC1. In the majority of lake phases PC2 and PC3 are correlated to some degree, which reflects the fact that they are both dominated by sources from within the SCM. PC3 correlates with PC1 and PC2 in E1 and E2. As seen in the plots of 1/Sr vs PC3, the influx of coarse-grained material acts to overprint the chemical weathering signal during these two events.

Silicon can be hosted in both mineral and biological matrices and can play a dual role in lake systems. In contrast, Ti is hosted by mineral phases. Due to this, increases in the Si/Ti ratio can either indicate increases in biological productivity (i.e., more biogenic Si in the sediment) or, as we are dealing with a ratio, decreased input of Ti to the system. As seen in records elsewhere (e.g., Brown et al., 2007; Tanaka et al., 2007; Minyuk et al., 2014), Si/Ti and BSi curves

are correlated despite the fact that Si/Ti represents two processes (biological activity and minerogenic input)(Figure 4). In the most obvious cases where the records differ (e.g., 22.20-22.58 m, 21.05-20.98 m, 19.30-19.05 m) we can assume that it is the increased input of Ti and/or a change in the mineral matter character that is driving the mismatch (Chawchai et al., 2016). We recognize that the BSi curve is more representative of within lake biological productivity but compliment this data with Si/Ti ratios which are higher resolution.

Lake Development

Lake Phase 2 (L2), Early Eemian lake phase (25.0 to 23.45 m)

PC1 and PC2 factor scores suggest that relatively higher minerogenic inputs occur during L2a as compared to L2b (Figure 2 and 3). In that the area was recently deglaciated, this decreasing minerogenic input signals the stabilization of the catchment in the form of soil development (Kylander et al., 2013). This is confirmed by the colonization of the area by open subarctic birch, *Pinus*, and *Juniperus* (L2a) and its gradual replacement by interglacial boreal forest with *Picea* (L2b). The ratio of Arboreal to Non-Arboreal Pollen (AP/NAP) indicates increasing forest cover density from L2a to L2b (Figure 2)(Helmens et al., 2015).

Much of L2 is characterized by laminations, reflecting a regularly anoxic hypolimnion and a stably stratified lake (Renberg, 1981). The diatom assemblages present at this time describe short mixing periods and potentially strong seasonality with long, cold winters and short, warm summers (Plikk et al., 2016). Of special note in L2 is the presence of *Cyclotella radiosa* (Grun.) Lemmermann and *Cyclotella michiganiana* Skvortz (Figure 3), an oligo-mesotrophic taxa that favours deep, warm waters and stable summer

stratification (Stoermer, 1993; Wolin, 1996; Whitlock et al., 2012). The diatom community also indicates that nutrient levels were initially low (L2a) but gradually rose creating a more eutrophic system (L2b) (Plikk et al., 2016) with BSi and Si/Ti values similarly showing increasing productivity (Figure 3). Soil development, as evoked by the increases in TOC and AP/NAP (Figure 2), and the leaching of base cations from the unconsolidated glacial materials and the surrounding bedrock after deglaciation could explain the moderate, but still variable behaviour of PC3 during this lake phase (Figure 4); this would help raise lake nutrient levels.

During L2c, minerogenic inputs start to increase again which, given the decrease in AP/NAP, can be linked to a vegetation shift (Figure 2 and 3). The lake flora imply increasingly cold and dynamic conditions and a slight decrease in trophic status (Plikk et al., 2016). Si/Ti values decrease marginally during L2c, likely as a result of the increased minerogenic inputs and cooler conditions (Figure 3). BSi has perhaps a small, but unremarkable, decrease. A very slight decrease in PC3 and chemical weathering occurs which could have contributed to the change in trophic status (Figure 4).

Event 1 (E1), Tunturi Event (23.45 to 22.55 m)

The Tunturi Event is clearly seen in the Sokli record, signalled by the siltier layer at the beginning of E1a. PC1 and PC2 show a double peak just prior to, and after, the boundary to Unit B (Figure 2 and 3). The increase in erosional input is in response to reduced catchment productivity (i.e., TOC), the replacement of the soil protecting mixed boreal forest by an open subarctic birch-dominated woodland (i.e., decrease in AP/NAP) and a temperature drop of 2-4°C (Helmens

et al., 2015). The instability of the soil cover is confirmed by the increase in fossil pollen of *Juniperus* and *Hippophaë rhamnoides*, which can cope with such conditions. Aquatic proxy indicators evidence prolonged ice cover as verified by minor increases in *Fragilaria* (mainly *Pseudostaurosira brevistriata* (Grun.) Williams & Round *Pseudostaurosira pseudoconstruens* Marciniak and *Staurosira venter* (Grun.) Williams & Round)(Lotter and Bigler, 2000), lowered lake levels and drier conditions, intense mixing and an earlier breakdown of thermal summer stratification (Pliikk et al., 2016). The significant increase in minerogenic input during E1a overprints PC3 and the chemical weathering signal (Figure 4 and 5) and affects biological signals either directly by reducing productivity through increased turbidity or indirectly by diluting BSi in the sediment (Figure 3).

PC1 and PC2 factor scores recover by E1b and slowly decrease in E1c indicating recovery of the catchment soils (Figure 2 and 3); a mirror change is observed in TOC and AP/NAP ratios. Increasing biological productivity within the lake is argued by changes in the Si/Ti and BSi data (Figure 3). Laminated sediments return but the stratification, and perhaps the seasonality, is not as extreme as that seen in L2 since diatoms indicating an extended mixing period length are present (e.g., *Aulacoseira ambigua* (Grun.) Simonsen and *Stephanodiscus medius* Håkansson)(Figure 3). PC3 is low and stable after E1a with reduced chemical weathering in the catchment, possibly as an effect of the inferred dry conditions of the time (Figure 4)(Pliikk et al., 2016).

In E1c laminations disappear indicating there was mixing of bottom waters. Productivity continues to increase during E1c (Figure 3). In examining the elemental data this sub-event appears rather uneventful in terms of detrital

input and chemical weathering. This is curious as the biological data from both within the lake (i.e., diatoms and chironomids) and the catchment (i.e., pollen) show that E1c is a period of minor cooling (Helmens et al., 2015). It is possible that the hydrological conditions did not vary much between E1b and E1c (all affecting the processes described by the elemental data) while the temperature did.

Lake Phase 3, Early mid-Eemian lake phase (22.55 to 20.85 m)

The factor scores for PC1 and PC2 generally remain just below zero during L3 (Figure 2 and 3). The observed stability is expected considering the return to interglacial conditions, with July temperatures an estimated 3°C warmer than present day, and the presence of arboreal-rich vegetation (i.e., increase in AP/NAP) (Helmens et al., 2015). There are again visible laminations in L3 indicating oxygen deficient conditions in the hypolimnion. The increased productivity as inferred by Si/Ti and BSi may have been enhanced by chemical weathering in the catchment (increase in PC3, Figure 4) and subsequent nutrient addition to the lake, which sees meso-eutrophic and eutrophic species inhabiting the lake (e.g., *Stephanodiscus* spp. and *A. ambigua*)(Figure 3). Additionally, these species are likely favoured by an enhanced internal loading of nutrients (especially P) during this phase, as Fe-bound P that is released and accumulated in the anoxic hypolimnion during the summer stratification is circulated back into the water column during the mixing periods (Bradbury and Dieterich-Rurup, 1993; Wetzel, 2001; Plikk et al., 2016).

Lake Phase 4 (L4), Late mid-Eemian (20.85 to 17.43 m)

During L4 the continued high interglacial temperatures that began in L3 are maintained. It is during L4 that visible laminations disappear, signifying a significant change in the lake system. PC1 and PC2 indicate that minerogenic material input decreases slightly at the start of L4a but thereafter remains rather steady during L4 (Figure 2 and 3). This would imply that the catchment is stable, which is supported by continued relatively high AP/NAP ratio (Helmens et al., 2015).

In L4 the long-term lake infilling that started after deglaciation appears to cross a key depth threshold. There are several indications of lower lake levels, increased mixing, and reduced bottom water anoxia. For example, there is change from weak laminations (Unit F) to no visible laminations (Unit G). The aquatic assemblages increase in benthic diatoms and littoral green algae evoking lake shallowing (Pliikk et al., 2016), e.g. *Staurosira construens* v. *binodis* (Ehrenb.) Hamilton that prefers shallow, macrophyte-rich, eutrophic lakes (Figure 2)(Bradbury and Winter, 1976). The increase in *A. ambigua*, a species often found in shallow lakes and near-shore areas of deep lakes and which requires high levels of turbulence and light to prosper (Bradbury and Dieterich-Rurup, 1993), also validates the idea that the lake in L4 is shallow and well mixed (Figure 3). This lake phase also sees the highest values in the productivity indicators Si/Ti and BSi, particularly in L4a.

L4 is also distinct in that PC3 shows gradually increasing positive values over several meters of accumulation (Figure 4). At this time acidophilous diatoms such as *Aulacoseira alpigena* (Grun.) Krammer and *Tabellaria flocculosa* (Roth.) Kütz. appear in low amounts (Van Dam et al., 1994), indicating a slight

decrease in the pH of the lake waters (Figure 4). The on-going development of soils would increase the production of SOM and DOC, promoting acid conditions (Pienitz et al., 1999; Engstrom et al., 2000; Fritz et al., 2004). Within the catchment there is an increase in coniferous tree species like *Picea* in the mixed boreal forest present; this leads to increased DOC transport from the catchment soils as compared to deciduous vegetation (Van Nevel, 2013). The increase in pollen of several floating leaved macrophytes such as *Nuphar* spp. and unidentified Nymphaeaceae, especially in L4b, attests to decreasing transparency associated with increasing DOC in the lake (Plikk et al., 2016). The increasing chemical weathering however, appears to offset significant acidification as there is continued presence of species found in near-shore areas of eutrophic lakes (e.g., *S. parvus*, *Fragilaria capucina* v. *mesolepta* (Rabenh.) Rabenh. and *Cyclotella choctawhatcheeana* Theriot, Stoermer & Håkansson comb. Nov.) (Plikk et al., 2016).

Event 2 (E2)(17.43 to 17.05 m)

E2 clearly mimics E1 and evidences another significant climate perturbation with cool conditions (Helmens et al., 2015) and a sharp influx of minerogenic material early in E2 (i.e., PC1 and PC3 increase) (Figure 2 and 4). Decreases in TOC (Figure 2), Si/Ti, and BSi (Figure 3) may signal a decrease in biological productivity in both the lake and the catchment but dilution by minerogenic influx is also again possible. The main difference between E2 and E1 lays in the fact that during the latter event the lake is much shallower due to progressive infilling. The increase in diatom species that thrive in subaerial conditions and rivers as well as an increase in *Sphagnum* spores and unidentified terrestrial

plant remains suggests decreased lake levels (Figure 2)(Valiranta, 2006; Plikk et al., 2016).

After this seemingly abrupt event the system seems to recover again with PC1, PC3, TOC, BSi, and Si/Ti going back to pre-event levels indicating more productive conditions and moderate chemical weathering (Figure 2, 3, and 4). The sediment at this time is characterized by light yellow spots which might be related to subaerial exposure or bottom freezing, and planktonic diatoms almost completely disappear (Plikk et al., 2016).

Lake Phase 5 (17.05-16.00 m)

Early in this lake phase, PC1 and PC2 indicate that minerogenic inputs are rather constant in L5a despite low AP/NAP ratios (Figure 2 and 3). Likewise, PC3 is low at this time (Figure 4). This may be related to infilling and the expansion of *Sphagnum* in the area, which acts to filter incoming sediment when lake levels are low. TOC levels also increase suggesting a vegetated catchment. While diatoms indicate increased lake levels compared to E2, the presence of diatoms common to acidic (e.g., *A. alpigena*) and shallow telmatic conditions (e.g., *S. construens* v. *binodis*) corroborates this continued infilling (Plikk et al., 2016) (Figure 2 and 4). Productivity is initially high but starts to decline leading up to the start of L5b (Figure 3). In these conditions there appears to be available oxygen and high mixing, as shown by a peak in *A. ambigua*.

In L5b the stratigraphy, PC1, PC2, and PC3 have a distinct pattern representing an unstable environment with the in-wash of silt and sand in flood type events (Figure 2). The general instability of L5b is verified by the increased presence of the diatoms *S. venter* and *S. pinnata*, which can tolerate rapidly

changing environments and are often found in shallow waters with sandy substrates (Haworth, 1976; Jones and Birks, 2004). *Sphagnum* grows in importance as infilling proceeds (Figure 2). The diatom flora move towards those species found in subarctic/arctic lakes (e.g., *Staurosirella lapponica* (Grun.) Williams & Round, *S. venter*, *S. pinnata*) and *Betula* becomes more abundant in the pollen data, indicating a move towards cool conditions (Pliikk et al., 2016).

Postglacial Boreal Lake Ontogeny

Lakes in northern boreal regions are commonly shallow, extremely dilute, clear, and oligotrophic (e.g., for Fennoscandia and the Kola Peninsula; Korhola and Weckström, 2004). Paleoclimate records reconstructed from late glacial and Holocene lake sediments in these regions integrate both climate as well as natural lake ontogeny, which often see the long-term postglacial oligotrophication of the lake (Ford, 1990; Pienitz et al., 1999; Engstrom et al., 2000; Fritz et al., 2004; Boyle, 2007; Fritz and Anderson, 2013; Law et al., 2015). As is common in the early phase of postglacial lake evolution elsewhere (e.g., Korhola and Weckström, 2004; Kylander et al., 2011), Sokli sees an initial period of high minerogenic input that is curbed by the development of catchment soils and colonizing vegetation. In L2a oligo-mesotrophic diatom species are already present and, with the exception of during E1 and E2, Sokli follows a trajectory towards meso-eutrophic conditions, a productive lake, and gradual infilling of the basin. Clearly, Sokli follows a path of long-term eutrophication rather than oligotrophication.

A number of abiotic and biotic processes are thought to drive postglacial oligotrophication of lakes. The high alkalinity and productivity observed within

542 the first few millennia after deglaciation are linked to (i) primary succession and
543 species immigration and (ii) leaching of cations from soils and bedrock (Fritz and
544 Anderson, 2013). At Sokli there are shrubs (i.e., low AP/NAP), *Betula*, and *Alnus*
545 present after deglaciation followed by the arrival of *Pinus* and *Picea* and an
546 increasing proportion of trees. This vegetation succession is rather typical of
547 postglacial landscapes in Fennoscandinavia (Donner, 1995) and does not seem to
548 explain why Sokli becomes increasingly eutrophic over time. The early presence
549 of *Alnus* at Sokli could be important given its role in N fixation and supply of
550 nutrients to the catchment (Fritz et al., 2004; Engstrom and Fritz, 2006). *Alnus* is
551 continuously present throughout the Eemian record at Sokli but this should be
552 interpreted with caution, given the fact that *Alnus* is an abundant pollen
553 producer (Bradshaw, 1981) which can be overrepresented in pollen data when
554 locally present (Tinsley and Smith, 1974). Catchment and soil weathering,
555 particularly within the base cation- and P-rich SCM (Talvitie et al., 1981), would
556 maintain both the alkalinity and nutrient supply to the lake, rather than it being
557 exhausted as seen in typical granitic and gneissic landscapes (Fritz and
558 Anderson, 2013). Several studies suggest that this climate independent, abiotic
559 edaphic process is a more important control on lake ontogeny than biological
560 processes (Boyle, 2007; Law et al., 2015) and the lake evolution at Sokli would
561 seem to lend support to this. The vegetative succession and the (iii)
562 accumulation of SOM and increased DOC input was also interrupted at Sokli
563 during E1 and E2 where conditions are glacial-like with high minerogenic inputs,
564 low temperature, increased *Betula*, *Juniper*, and shrubs. This could potentially
565 alter the trajectory of the lake with the external climate signals overriding lake
566 ontogeny processes. Finally, (iv) hardpan formation and the expansion of

wetlands can inhibit leaching of base cations and the maintenance of alkalinity (Engstrom et al., 2000; Fritz and Anderson, 2013). While we are not able to say much in regards to ground water input, *Sphagnum* percentages indicate the development of wetlands only late in the Sokli sequence.

Conclusions

The long-term oligotrophication normally seen in postglacial boreal lakes is not observed at Sokli and reasons for this were explored using the elemental geochemistry of the lake sediments. Lake sediment geochemistry is controlled by a number of integrated factors including the geological setting, lake and catchment development, and climate variability. Three main processes controlling the sediment geochemistry at Sokli were identified: (i) detrital input from sources dominantly beyond the SCM; (ii) detrital input from sources dominantly within the SCM; and (iii) chemical weathering. In general, the lake evolution was characterized by stable minerogenic inputs with varying importance of sources from both outside and inside the SCM over time. These inputs contributed to the long-term infilling of the lake. Chemical weathering increased over the span of the Eemian record at Sokli. The SCM provided an abundant and continual supply of base cations and nutrients, which helped to maintain a relatively alkaline and increasingly eutrophic lake. We note the highest levels of chemical weathering occurred during L3 and L4 when temperatures were higher than present day; there was increased presence of acidic *Picea*; and slightly higher catchment productivity (TOC). Trends in the processes described above were however, interrupted during both E1 and E2, which saw the influx of significant amounts of detrital material, overwhelming

the geochemical signals. At Sokli it appears that the forcing from abiotic edaphic process and climate override the typical postglacial oligotrophication normally seen in boreal lakes.

Acknowledgements

We gratefully acknowledge the Geological Survey of Finland for field and coring assistance. This study was funded by the Swedish Nuclear Fuel and Waste Management Company (SKB) and the Bolin Centre for Climate Research at Stockholm University. Additional funding was provided by Stiftelsen Längmanska kulturfonden, Academy of Finland (project no. 278692) and the Finnish Cultural Foundation. MEK dedicates this article to her father and thanks all the staff at the Virginia Commonwealth University who helped him through a successful open-heart surgery during the resubmission of this article.

References

- Alexanderson, H., Eskola, K.O., Helmens, K.F., 2008. Optical dating of a late Quaternary sediment sequence from Sokli, northern Finland. *Geochronometria* 32: 51–59.
- Battarbee, R.W., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L., Juggins, S., 2003. Diatoms, in: Smol, J.P., Birks, H.J.B., Last, W.M. (Eds.), *Tracking Environmental Change Using Lake Sediments: Terrestrial, Algal and Silicious Indicators, Developments in Paleoenvironmental Research*. Kluwer Academic Publishers, Dordrecht, pp. 155–202.
- Bauch, H.A., Kandiano, E.S., Helmke, J., Andersen, N., Rosell-Mele, A., Erlenkeuser, H., 2011. Climatic bisection of the last interglacial warm period in the Polar North Atlantic. *Quaternary Science Reviews* 30: 1813–1818.
- Bińka, K., Nitychoruk, J., Dzierżek, J., 2011. Climate stability during the Eemian – new pollen evidence from the Nidzica site, northern Poland. *Boreas* 40: 342–350.
- Björck, S., Noe-Nygaard, N., Wolin, J., Houmark-Nielsen, M., Jørgen Hansen, H., Snowball, I., 2000. Eemian Lake development, hydrology and climate: a multi-stratigraphic study of the Hollerup site in Denmark. *Quaternary Science Reviews* 19: 509–536.
- Boyle, J.F., 2007. Loss of apatite caused irreversible early-Holocene lake

acidification. *The Holocene* 17:543-547.

Bradbury, J.P., Dieterich-Rurup, K.V., 1993. Holocene diatom paleolimnology of Elk Lake, Minnesota. In: Bradbury, J.P., Dean, W.E. (Eds.), Elk Lake, Minnesota: Evidence for Rapid Climate Change in the North-central United States. Geological Society of America Special Paper, Boulder, Colorado, pp. 215-237.

Bradbury, J.P., Winter, T.C., 1976. Areal Distribution and Stratigraphy of Diatoms in the Sediments of Lake Sallie, Minnesota. *Ecology* 57, 1005-1014.

Bradshaw, R.H.W., 1981. Modern pollen-representation factors for woods in south-east England. *Journal of Ecology* 69: 45-70.

Brown, E.T., Johnson, T.C., Scholz, C.A., Cohen, A.S., King, J.W., 2007. Abrupt change in tropical African climate linked to the bipolar seesaw over the past 55,000 years. *Geophysical Research Letters* 34. doi:10.1029/2007GL031240

Chawchai, S., Kylander, M.E., Chabangborn, A., 2016. Testing commonly used X-ray fluorescence core scanning-based proxies for organic-rich lake sediments and peat. *Boreas* 45:180-189.

Cheddadi, R., Mamakowa, K., Guiot, J., de Beaulieu, J.-L., Reille, M., Andrieu, V., Granoszewski, W., Peyron, O., 1998. Was the climate of the Eemian stable? A quantitative climate reconstruction from seven European pollen records. *Palaeogeography, Palaeoclimatology and Palaeoecology* 143: 73-85.

Cohen, A.S., 2003. *Paleolimnology: The History and Evolution of Lake Systems*. Oxford University Press, Oxford.

Cortijo, E., Duplessy, J.C., Labeyrie, L., Leclair, H., Duprat, J., Van Wearing, T., 1994. Eemian cooling in the Norwegian Sea and North Atlantic ocean preceding continental ice-sheet growth. *Nature* 372: 446-449.

Davison W., 1993. Iron and Manganese in Lakes. *Earth-Science Reviews* 34: 119-163.

Deer, W.A., Howie, R.A., Zussman, J., 1992. *An introduction to the rock-forming minerals*. Harlow: Longman Scientific & Technical.

Donner, J. (1995) *The Quaternary History of Scandinavia*. Cambridge University Press, Cambridge, UK.

Drehs, A., Nordlund, A., Karlsson, P., Helminen, J., Rissanen, P., 2002. Climatological statistics of Finland 1971-2000. Finish Meteorological Institute, Helsinki.

Engstrom, D.R., Fritz, S.C., 2006. Coupling between primary terrestrial succession and the trophic development of lakes at Glacier Bay, Alaska. *Journal of Paleolimnology* 35: 873-880.

Engstrom, D.R., Fritz, S.C., Almendinger, J.E., Juggins, S., 2000. Chemical and biological trends during lake evolution in recently deglaciated terrain. *Nature* 408: 161-166.

Eriksson, L., Johansson, E., Kettaneh-Wodl, N., Wold, S., 1999. *Introduction to Multi- and Mega-Variate Data Analysis using Projection Methods (PCA & PLS)*. Umetrics AB, Umeå.

Ford, M.S.J., 1990. A 10,000-year history of natural ecosystem acidification. *Ecological Monographs* 60: 57-89.

Fritz, S.C., Anderson, N.J., 2013. The relative influences of climate and catchment processes on Holocene lake development in glaciated regions. *Journal of Paleolimnology* 49: 349-362.

Fritz, S.C., Engstrom, D.R., Juggins, S., 2004. Patterns of early lake evolution in

672 boreal landscapes: a comparison of stratigraphic inferences with a modern
673 chronosequence in Glacier Bay, Alaska. *The Holocene* 14: 828–840.

674 Fronval, T., Jansen, E., 1996. Rapid changes in ocean circulation and heat flux in
675 the Nordic seas during the Last Interglacial period. *Nature* 383: 745–844.

676 Galaasen, E.V., Ninnemann, U.S., Irval, N., Kleiven, H.F., Rosenthal, Y., Kissel, C.,
677 Hodell, D.A., 2014. Rapid Reductions in North Atlantic Deep Water During the
678 Peak of the Last Interglacial Period. *Science* 343: 1129–1132.

679 Haworth, E.Y., 1976. Two late-glacial (late Devensian) diatom assemblage
680 profiles from northern Scotland. *New Phytologist* 77: 227–256.

681 Helmens, K.F., Bos, J.A.A., Engels, S., Van Meerbeeck, C.J., Bohncke, S.J.P., Renssen,
682 H., Heiri, O., Brooks, S.J., Seppä, H., Birks, H.J.B., Wohlfarth, B., 2007. Present-
683 day temperatures in northern Scandinavia during the last glaciation. *Geology*
684 35: 987–990.

685 Helmens, K.F., Räsänen, M.E., Johansson, P.W., 2000. The last interglacial-glacial
686 cycle in NE Fennoscandia: a nearly continuous record from Sokli (Finnish
687 Lapland). *Quaternary Science Reviews* 19: 1605–1623.

688 Helmens, K.F., Salonen, J.S., Pliik, A., Engels, S., Välranta, M., Kylander, M.,
689 Brendryen, J., Renssen, H., 2015. Major cooling intersecting peak Eemian
690 Interglacial warmth in northern Europe. *Quaternary Science Reviews* 122:
691 293–299.

692 Holmer, M., Storkholm, P., 2001. Sulphate reduction and sulphur cycling in lake
693 sediments: a review. *Freshwater Biology* 46: 431–451.

694 Ilvonen, E., 1973. Eem - kerrostuma savukosken soklilla. *Julkaisija: Suomen*
695 *Geologinen Seura* 25: 81–84.

696 Irvali, N., Ninnemann, U.S., Galaasen, E.V., Rosenthal, Y., Kroon, D., Oppo, D.W.,
697 Kleiven, H.F., Darling, K.F., Kissel, C., 2012. Rapid switches in subpolar North
698 Atlantic hydrography and climate during the Last Interglacial (MIS 5e).
699 *Paleoceanography* 27: PA2207.

700 Jones, V., Birks, H., 2004. Lake-sediment records of recent environmental change
701 on Svalbard: results of diatom analysis. *Journal of Paleolimnology* 31: 445–
702 466.

703 Jones, B.F., Bowser, C.J., 1978. The mineralogy and related chemistry of lake
704 sediments. In Lerman, A. (Ed.), *Lakes: Chemistry, Geology, Physics*. Springer
705 New York, pp. 179–235.

706 Jørgensen, B.B., 1990. The sulfur cycle of freshwater sediments: Role of
707 thiosulfate. *Limnology and Oceanography* 35: 1329–1342.

708 Korhola, A., Weckström, J., 2004. Paleolimnological studies in Arctic
709 Fennoscandia and the Kola Peninsula (Russia). In Pienitz, R., Douglas,
710 M.S.V., Smol, J. (Eds). *Developments in Paleoenvironmental Research*, Springer
711 Academic, pp. 381–418.

712 Köhl, N., Litt, T., Schölzel, C., Hense, A., 2007. Eemian and Early Weichselian
713 temperature and precipitation variability in northern Germany. *Quaternary*
714 *Science Reviews* 26: 3311–3317.

715 Kylander, M.E., Ampel, L., Wohlfarth, B., 2011. High-resolution X-ray fluorescence
716 core scanning analysis of Les Echets (France) sedimentary sequence: new
717 insights from chemical proxies. *Journal of Quaternary Science* 26: 109–117.

718 Kylander, M.E., Klaminder, J., Wohlfarth, B., Löwemark, L., 2013. Geochemical
719 responses to paleoclimatic changes in southern Sweden since the late glacial:
720 the Hässeldala Port lake sediment record. *Journal of Paleolimnology* 50: 57–

721 70.

722 Law, A.C., Anderson, N.J., McGowan, S., 2015. Spatial and temporal variability of

723 lake ontogeny in south-western Greenland. *Quaternary Science Reviews* 126:

724 1-16.

725 Lee, M.J., Lee, J.I., Garcia, D., Moutte, J., Williams, C.T., Wall, F., Kim, Y., 2006.

726 Pyrochlore chemistry from the Sokli phoscorite-carbonatite complex,

727 Finland: Implications for the genesis of phoscorite and carbonatite

728 association. *Geochemical Journal* 40: 1-13.

729 Lee, M.J., Lee, J.I., Moutte, J., 2005. Compositional variation of Fe-Ti oxides from

730 the Sokli complex, northeastern Finland. *Geosciences Journal* 9: 1-13.

731 Ljung, K., Holmgren, S., Kylander, M., Sjolte, J., Van der Putten, N., Kageyama, M.,

732 Porter, C.T., Björck, S., 2015. The last termination in the central South

733 Atlantic. *Quaternary Science Reviews* 123: 193-214.

734 Lotter, A.F., Bigler, C., 2000. Do diatoms in the Swiss Alps reflect the length of ice-

735 cover? *Aquatic Sciences* 62: 125-141.

736 Mavris, C., Egli, M., Plötze, M., Blum, J.D., Mirabella, A., Giaccari, D., Haeberli, W.,

737 2010. Initial stages of weathering and soil formation in the Morteratsch

738 proglacial area (Upper Engadine, Switzerland). *Geoderma* 155: 359-371.

739 Minyuk, P.S., Borkhodoev, V.Y., Wennrich, V., 2014. Inorganic geochemistry data

740 from Lake El'gygytgyn sediments: marine isotope stages 6-11. *Climate of the*

741 *Past* 10: 467-485.

742 Oliva, P., Viers, J., Dupré, B., 2003. Chemical weathering in granitic environments.

743 *Chemical Geology* 202: 225-256.

744 Pienitz, R., Smol, J.P., MacDonald, G.M., 1999. Paleolimnological Reconstruction of

745 Holocene Climatic Trends from Two Boreal Treeline Lakes, Northwest

746 Territories, Canada. *Arctic, Antarctic, and Alpine Research* 31: 82-93.

747 Pliikk, A., Helmens, K.F., Fernandez-Fernandez, M., Kylander, M.E., Löwemark, L.,

748 Risberg, J., Salonen, J.S., Väiliranta, M., Weckström, J., 2016. Development of an

749 Eemian (MIS 5e) Interglacial palaeolake at Sokli (N Finland) inferred using

750 multiple proxies. *Palaeogeography, Palaeoclimatology, Palaeoecology* 463:

751 11-26.

752 Reimann, C., Filzmoser, P., Garret, R., Dutter, R., 2008. Statistical data analysis

753 explained: Applied environmental statistics. John Wiley & Sons Ltd.

754 Renberg, I., 1981. Formation, structure and visual appearance of iron-rich,

755 varved lake sediments. *Proceedings - International Association of Theoretical*

756 *and Applied Limnology* 21: 94-101.

757 Riebe, C.S., Kirchner, J.W., Finkel, R.C., 2004. Erosional and climatic effects on

758 long-term chemical weathering rates in granitic landscapes spanning diverse

759 climate regimes. *Earth and Planetary Science Letters* 224: 547-562.

760 Rioual, P., Andrieu-Ponel, V., de Beaulieu, J.L., Reille, M., Svobodova, H., Battarbee,

761 R.W., 2007. Diatom responses to limnological and climatic changes at Ribains

762 Maar (French Massif Central) during the Eemian and Early Würm.

763 *Quaternary Science Reviews* 26: 1557-1609.

764 Rosén, P., Vogel, H., Cunningham, L., Reuss, N., Conley, D., Persson, P., 2010.

765 Fourier transform infrared spectroscopy, a new method for rapid

766 determination of total organic and inorganic carbon and biogenic silica

767 concentration in lake sediments. *Journal of Paleolimnology* 43: 247-259.

768 Seidenkrantz, M.S., Kristensen, P., Knudsen, K.L., 1995. Marine evidence for

769 climatic instability during the last interglacial in shelf records from

- northwest Europe. *Journal of Quaternary Science* 10: 77–82.
- Shchukarev, A., Gälman, V., Rydberg, J., Sjöberg, S., Renberg, I., 2008. Speciation of iron and sulphur in seasonal layers of varved lake sediment: an XPS study. *Surface and Interface Analysis* 40: 354–357.
- Stoermer, E. 1993. Evaluating diatom succession: some peculiarities of the Great Lakes case. *Journal of Paleolimnology* 8: 71–83.
- Talvitie, J., Lehmuspelto, P., Vuotovesi, T., 1981. Airborne Thermal Surveying of the Ground in Sokli, Finland. Geological Survey of Finland.
- Tanaka, K., Akagawa, F., Yamamoto, K., Tani, Y., Kawabe, I., Kawai, T., 2007. Rare earth element geochemistry of Lake Baikal sediment: its implication for geochemical response to climate change during the Last Glacial/Interglacial transition. *Quaternary Science Reviews* 26: 1362–1368.
- Tinsley, H., Smith, R.T., 1974. Surface pollen studies across a woodland/heath transition and their application to the interpretation of pollen diagrams. *New Phytologist* 73: 547–565.
- Van Dam, H., Mertens, A., Sinkeldam, J., 1994. A coded checklist and ecological indicator values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic Ecology* 28: 117–133.
- Van Nevel, L., Mertens, J., De Schrijver, A., Baeten, L., De Neve, S., Tack, F.M., Meers, E., Verheyen, K., 2013. Forest floor leachate fluxes under six different tree species on a metal contaminated site. *Science of the Total Environment* 447: 99–107.
- Vartiainen, H., 1980. The Petrography, mineralogy and petrochemistry of the Sokli carbonatite massif, northern Finland. Geological Survey of Finland.
- Väliiranta, M., 2006. Terrestrial plant macrofossil records; possible indicators of past lake-level fluctuations in north-eastern European Russia and Finnish Lapland. *Acta Palaeobotanica* 46: 235–243.
- West, A., Galy, A., Bickle, M., 2005. Tectonic and climatic controls on silicate weathering. *Earth and Planetary Science Letters* 235: 211–228.
- Wetzel, R.G., 2001. Limnology. Academic Press.
- White, A.F., Blum, A.E., 1995. Effects of climate on chemical weathering in watersheds. *Geochimica et Cosmochimica Acta* 59: 1729–1747.
- Whitehead, D.R., Charles, D.F., Jackson, S.T., Smol, J.P., Engstrom, D.R., 1989. The developmental history of Adirondack (NY) lakes. *Journal of Paleolimnology* 2: 185–206.
- Whitlock, C., Dean, W., Fritz, S., Stevens, L., Stone, J., Power, M., Rosenbaum, J., Pierce, K. and Bracht-Flyr, B. 2012. Holocene seasonal variability inferred from multiple proxy records from Crevice Lake, Yellowstone National Park, USA. *Palaeogeography, Palaeoclimatology, Palaeoecology* 331: 90–103.
- Wittkop, C., Teranes, J., Lubenow, B., Dean, W.E., 2014 Carbon- and oxygen-stable isotopic signatures of methanogenesis, temperature, and water column stratification in Holocene siderite varves. *Chemical Geology* 389: 153–166.
- Wolin, J. 1996. Late Holocene lake-level and lake development signals in Lower Herring Lake, Michigan. *Journal of Paleolimnology* 15: 19–45.
- Zeng, Y., Chen, J., Xiao, J., Qi, L., 2013. Non-residual Sr of the sediments in Daihai Lake as a good indicator of chemical weathering. *Quaternary Research* 79: 284–291.

818 **Table 1.** Variance explained by each factor

Factor	Variance	%	Cumulative %
1	4,19	41,91	41,91
2	2,61	26,15	68,06
3	1,33	13,30	81,37

819 **Table 2.** Rotated factor loadings. Shaded data are significant at the α 0.05 level.

Element	Factor 1	Factor 2	Factor 3
K	0,94	0,25	0,00
Ti	0,94	0,26	0,03
Rb	0,93	0,17	-0,09
Zr	0,83	-0,28	0,19
Si	0,71	0,16	0,05
Fe	0,28	0,80	-0,37
S	0,09	0,93	0,06
Mn	0,02	0,63	-0,29
Ca	0,54	0,65	0,43
Sr	0,07	-0,21	0,93

820

821 **Table 3.** Correlation coefficients between the PC during different lake phases and events

Lake Phase/Event		PC2	PC3
L5	PC1	0.44	0.03
	PC2		0.68
E2	PC1	0.64	0.81
	PC2		0.80
L4	PC1	0.39	-0.53
	PC2		0.74
L3	PC1	-0.67	-0.24
	PC2		0.67
E1	PC1	0.23	0.62
	PC2		0.68
L2	PC1	-0.24	-0.50
	PC2		0.15

822

823 **Figure Captions**

- 824 1. Location of the study site in northern Finland (A). The Sokli Eemian paleolake
825 sits in the Sokli Carbonatite Massif (SCM), a carbonatite-alkaline complex
826 surrounded by crystalline Precambrian Shield. The work presented here is
827 from Borehole 2 (B).
- 828 2. Depth profiles of PC1, TOC, *S. construens* v. *binodis*, AP/NAP ratios, and
829 *Sphagnum* pollen percentages. PC1 is representative of minerogenic inputs
830 from sources outside the SCM while TOC tracks catchment productivity. *S.*
831 *construens* v. *binodis* is used here as an indicator of lake infilling while
832 AP/NAP tells us about the relative importance of trees versus other species
833 where trees require thicker soils and milder conditions to grow. The presence
834 of *Sphagnum* is also linked to infilling of the lake. The bracketing glaciofluvial
835 sands and gravel (28.5 and 25 m) and fluvial sands (16 to 13 m) have been
836 dated to >ca. 95 and 110 ka and <ca. 150-180 ka, respectively (Helmens et al.,
837 2000; 2007; Alexanderson et al., 2008).
- 838 3. Depth profiles of PC2, *C. radiosa*, *C. michiganiana*, *A. ambigua*, *S. medius*, Si/Ti,
839 and BSi. PC2 represents inputs from S, Fe, Mn, and Ca-rich sources within the
840 SCM. Both *C. radiosa* and *C. michiganiana* are species that favour deep, warm
841 waters with stable summer stratification. *A. ambigua* and *S. medius* require an
842 extended mixing period length and eutrophic and productive conditions. Si/Ti
843 and BSi are indicators of in-lake productivity.
- 844 4. Depth profiles of PC3, *T. flocculosa*, *A. alpigena*, *Picea* pollen percentages, and
845 *Picea* stomata. PC3 represents the integrated signal of those processes
846 controlling chemical weathering. Both *T. flocculosa* and *A. alpigena* are
847 acidophilic species and are used as indicators of pH trends in the lake. *Picea*

848 pollen and stomata are used to highlight the increase presence of these
849 acidifying species.

850 5. Depth profiles of PC1, PC2, and PC3 show that the PC co-vary during certain
851 phases of lake development.

852 6. Plot of $1/Sr$ vs. PC3 where a negative correlation indicates that Sr is the main
853 control on PC3 variability.

854 **Supplementary Information**

855 Raw data elemental profiles by depth given in peak areas.

856
857

Kylander et al.-Figure 2

858
859

Kylander et al.-Figure 3

860
861

Kylander et al.-Figure 4

862
863

Kylander et al.-Figure 5

864

Kylander et al.-Figure 6

865