The circadian clock at the intercept of sleep and pain

Palada, Vinko

2020-05

Palada , V , Gilron , I , Canlon , B , Svensson , C & Kalso , E 2020 , ' The circadian clock at the intercept of sleep and pain ' , Pain , vol. 161 , no. 5 , pp. 894-900 . <https://doi.org/10.1097/j.pain.0000000000001786>

<http://hdl.handle.net/10138/328478>

<https://doi.org/10.1097/j.pain.0000000000001786>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

The circadian clock at the intercept of sleep and pain

Vinko Paladà¹, Ian Gilron², Barbara Canlon¹, Camilla I. Svensson¹, Eija Kalso^{1,3,*}

¹Department of Physiology and Pharmacology, Karolinska Institute, Stockholm, Sweden

²Department of Anesthesiology and Perioperative Medicine, Queen's University, Kingston, Canada

³Pain Clinic, Department of Anesthesiology, Intensive Care, and Pain Medicine, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

Number of pages: 29

Number of figures: 2

*Author of correspondence:

Name: Eija Kalso, MD, PhD

Address: Pain Clinic, Helsinki University Hospital, Haartmaninkatu 2A (PB 140), 00029 HUS, Helsinki, Finland

Telephone number: +358-9-47175885

Fax number: +358-9-4717641

Email address: eija.kalso@helsinki.fi

Keywords: Circadian; gene; pain; neuropathym; Sleep

1. Introduction

It is estimated that 50-88% of the patients with chronic pain suffer from inadequate sleep.⁹⁴ Not only does pain disturb sleep, but insomnia symptoms often precede the onset of pain in patients with e.g. fibromyalgia,⁷⁴ musculoskeletal pain⁷⁹ and headache.⁸⁰ Insomnia also leads to increased pain intensity.¹⁰⁵ Recent studies suggest that both sleep and pain are regulated by the circadian clock and circadian genes have been associated with nociception and an altered sleep-wake cycle.^{38,93} Most of the current knowledge regarding circadian regulation of pain comes from animal models; thus, there is an urgent need to better understand the effect of circadian rhythms in the development and maintenance of chronic pain states. Therefore, in this topical review we discuss the association between the circadian clock, sleep and pain focusing on circadian regulation of neuropathic and inflammatory pain, headache and migraine. We will also address the effect of circadian regulation on pain pharmacology and analgesia.

1.1. Bidirectional relationship between sleep and pain

The relationship between sleep and pain is bidirectional since insufficient sleep can precede and maintain pain while pain can impair one's ability to initiate and maintain sleep.⁶² Experimental studies show that acute sleep deprivation leads to reduced pain thresholds^{55,58,77,92,98}, increased pain sensitivity^{33,95}, and a female-specific reduction in descending pain inhibition.^{33,95} A recent fMRI study revealed that acute sleep deprivation in healthy adults caused increased pain reactivity in primary somatosensory cortex and reduced activity in insula and striatum— brain regions involved in descending pain inhibition.¹⁰² In addition to a direct effect on pain, sleep deprivation has been found to promote depression and anxiety, which are frequent pain

comorbidities.^{5,6,57,66,75} However, some studies also report that sleep deprivation can have anti-depressant effects.¹⁶

1.2. Circadian clock at the interplay between sleep and pain

Emerging data suggest that both sleep^{4,24,56,84,88,101,107,108} and pain^{20,23,31,62,109,113} can be regulated by the circadian clock on the physiological and molecular level (Figure 1). Master regulators of the mammalian circadian clock are the responsive neurons in the suprachiasmatic nucleus (SCN) of the hypothalamus that respond to light stimuli from retina⁴⁸ and synchronize the clocks in peripheral tissues through the autonomic nervous system and secretion of hormones and neurotransmitters (melatonin, cortisol, noradrenaline and dopamine) involved in regulation of sleep-wake cycle and nociception.⁵¹ The SCN neurons suppress synthesis of melatonin from the pineal gland during daytime whereas pineal melatonin production is elevated during the night to induce sleep.¹⁹ Disturbed melatonin secretion has been suggested to contribute to insomnia, particularly in the elderly,⁶⁴ and melatonin administration was shown to improve the sleep quality³⁴ and to exert analgesic effects.¹⁰⁶ Similarly, secretion of cortisol from the adrenal cortex shows circadian oscillations. The levels of cortisol increase steadily during sleep progression to promote wakefulness and to initiate gluconeogenesis resulting in peak cortisol levels in the early morning.³⁹ Long-term stress has been associated with elevated cortisol levels in the evening, and a sleep onset, which is linked to insomnia and pain chronification.⁷⁸ It is important to note that diurnal fluctuation in the cortisol levels can also occur due to psychological distress^{28,69} and physical activity¹⁰ may influence the melatonin levels which might indirectly affect the diurnal rhythmicity in pain intensity independently of the circadian input from the SCN. Diurnal rhythmicity due to psychological distress, physical activity and yet unknown mechanisms independent of circadian regulation, might also explain

the variable levels of coupling between the circadian clock and pain in different pain cohorts as reported in patients with migraine.^{31,83} Additionally, it has been shown that SCN neurons indirectly project to noradrenergic neurons in the locus coeruleus (LC)^{8,44} as well as to ventral tegmental area (VTA) containing dopaminergic neurons.⁶⁷ Levels of noradrenaline in the blood circulation are reduced during sleep in humans compared to wakefulness,⁷¹ which is in line with higher discharge rates of noradrenergic LC neurons during wakefulness compared to sleep.⁹ Also, a recent study on mice showed that inhibiting VTA dopaminergic neurons suppresses wakefulness to promote sleep while optogenetic stimulation induced long-term wakefulness and suppressed sleep.³² Besides regulation of the sleep-wake cycle, both noradrenaline and dopamine are key modulators of descending pain inhibition pathway.¹¹ On the molecular level, circadian regulation of sleep and pain is achieved through the feedback loops that promote the rhythmicity in gene expression and endogenous periodicity of circa 24h.⁵⁴ The main regulation loops of the circadian genes are described in Figure 2.

2. Circadian clock and sleep

The sleep-wake cycle can be divided into wakefulness, non-rapid eye movement (NREM), and rapid eye movement (REM) sleep. The electroencephalogram (EEG) can distinguish between REM sleep and wakefulness characterized by fast, desynchronized oscillations and NREM sleep with synchronized cortical oscillations.⁸¹ Several single nucleotide polymorphisms (SNPs) in the core circadian genes CLOCK, BMAL1, CRY1, PER2, and PER3 have been associated with inter-individual differences in sleep time, sleep duration, as well as with sleep disorders.^{4,56,60,84,101}

Double Cry1/2^{-/-} knockout mice display hypersomnolence and recurrent episodes of prolonged sleep, associated with increased duration of NREM sleep in the EEG and followed by changes in the expression of circadian genes in the cerebellum.¹⁰⁷ Changes in the cortical expression of

the mouse clock genes *CLOCK*, *BMAL1*, *CRY2* and *NPAS2* due to sleep deprivation were found to correlate with differences in EEG patterns between different mice strains indicating that circadian gene expression might drive the EEG differences among these strains.¹⁰⁸ The EEG measurements on *BMAL1* knockout macaque monkeys indicated reduced REM sleep duration during the evening⁸⁸ which is opposite to increased sleep duration in *Bmal1* knockout mice⁸¹ indicating the differences in sleep regulation between diurnal and nocturnal species. A recent study in healthy adults showed that the *PER2* SNP, rs6753456, shapes sleep and EEG patterns in humans evidenced by lower EEG power density during NREM sleep and lower slow-wave activity in the early part of the sleep.

3. Circadian clock rhythmicity and pain

Pain intensity is characterized by diurnal fluctuations both in clinical conditions and experimental pain.^{20,42} Circadian clock rhythmicity plays an important role in regulating the rhythmic expression of the voltage-gated calcium channel subunit α_1 ⁵⁹ and NMDA glutamate receptor subunits on nociceptive neurons in dorsal ganglions (DRGs).¹¹³ The circadian genes *CLOCK* and *BMAL1* also regulate the rhythmic expression of *TAC1* gene in DRG neurons which encode for the neuropeptide substance P.¹¹³ Besides DRGs, circadian rhythm was also associated with rhythmic expression of opioid receptors in periaqueductal gray, part of brainstem important for descending pain inhibition⁹⁹ and with expression of the NMDA receptor subunit NR2B in the dorsal horn of spinal cord.¹⁰⁹

3.1. Circadian regulation of neuropathic pain

Patients with neuropathic pain in conditions such as diabetic neuropathy and postherpetic neuralgia, usually experience the most severe pain in the evening hours.^{40, 82} The morning-

evening pain intensity difference has been observed to be more pronounced in females and in diabetic neuropathy compared with postherpetic neuralgia suggesting that various biological factors affect the circadian control of pain.⁴¹ Altered circadian expression of opioid and melatonin receptors was reported in the spared nerve injury (SNI) mouse model of neuropathic pain. Gene expression of melatonin MT1A and MT1B receptors in hypothalamus was significantly higher in SNI mice at 2:00 and lower at 8:00 time point compared with sham controls.⁸¹ Spinal dorsal horn expression of circadian clock genes PER1, PER2, and CRY1 was suppressed in the lumbar dorsal horn 7 days after sciatic nerve ligation (PSNL) in mice, which was followed by increased mechanical hypersensitivity.⁷³ These studies suggest that altered expression of clock genes in sensory neurons contribute to neuropathic pain following peripheral nerve injury. Considering that neuroinflammation also associates with neuropathic pain, it is important to notice that microglia and inflammatory cytokines TNF α , IL-1 β , and IL-6 show strong rhythmic expression in CNS.^{36,76}

3.2. Inflammatory pain and circadian clock

Patients with inflammatory pain conditions such as rheumatoid arthritis (RA) have the most severe joint pain and stiffness in the morning.^{13,14} While pain in osteoarthritis (OA) is primarily associated with abnormal mechanical loading of joints,³⁷ some OA patients also have an inflammatory pain component. It has been reported that pain in OA patients has higher intensity in the morning and early afternoon compared with the evening hours when the level of physical activity is reduced.^{2,12,14} This temporal pattern is apparent also in acute inflammatory pain conditions,¹⁸ and interestingly, this is distinctly different from the pattern in neuropathic pain conditions where pain generally gets worse in the evening. The inflammatory pain pattern in RA patients is associated with circadian rhythmicity in the serum levels of the pro-inflammatory

cytokine IL-6 with the highest concentrations in the early hours of the day (7-10 AM).⁷ The overnight increase in the level of pro-inflammatory cytokines is partially due to reduced release of cortisol in RA patients.^{25,26} Furthermore, addition of pro-inflammatory cytokines IL-1 or TNF- α to cultured synovial fibroblasts from RA patients delayed the peak of circadian BMAL1 expression by up to 6 hours,⁴⁹ while the protein level of PER2 was reduced in synoviocytes upon lipopolysaccharide (LPS)-induced inflammation.⁶³ Finally, a recent study on partial medial meniscectomy model of OA pain showed that activation of ERK-ERB with the agonist SR9009 leads to reduced BMAL1 expression, decreased mechanical hyperalgesia, and severe cartilage degeneration.²⁸ These studies suggest that pain in RA and OA is under circadian regulation.

3.3. Role of circadian clock in migraine and headache

Migraine has a 24h pattern which is characterized by frequent morning onset although migraine patients have reported headache episodes during the whole day.^{31,83,96} Interestingly, forebrain specific knockdown of the calcium voltage-gated channel Ca_v2.1 in a mouse model of migraine led to impaired circadian rhythmicity as the mice were running both during the active phase at night and during the resting phase during daytime.⁶⁸ Patients with cluster headache (CH) usually experience the most severe pain in the middle of the night^{90,97} and both cortisol and melatonin levels are dysregulated in patients with CH.^{20,104} In a recent study, the CLOCK gene SNP rs1264507 was significantly associated with CH with elevated gene expression of CLOCK in primary fibroblasts from CH patients.³⁶

4. Circadian regulation of analgesia and pharmacology of analgesics

So far, most studies of circadian effects on pain medication have focused on opioid-induced analgesia. The circadian pattern of analgesic efficacy was observed for morphine injections in mice, with the highest efficacy and fastest recovery from tolerance observed during the dark, compared with the light phase. This is consistent with the 24h rhythm of μ -opioid receptor expression.¹¹⁰ Mice with deletion mutation in *Per2* gene showed reduced morphine-induced tolerance and withdrawal.⁸⁷ Interestingly, a recent study reported that mice presenting opioid-induced hyperalgesia after chronic morphine exposure had overexpression of circadian core genes such as *Per2* and *Per3* in pain regulating trigeminal ganglia and nucleus accumbens compared with controls.¹¹⁴ Rhythmicity in analgesic efficacy of opioids was detected in healthy human volunteers with peak efficacy of fentanyl reported at 17:30 whereas the efficacy of dihydrocodeine and tramadol was highest at 20:00 compared with efficacy in the morning.^{17,52} Diurnal variation was previously described for opioid requirements after gastric surgery with morning doses of morphine 15% higher compared with the evening dosage and with the peak analgesic efficacy at 09:00.⁴⁷ This time-dependent analgesic efficacy was also observed for the $\alpha_2\delta_1$ ligands gabapentin and pregabalin. The i.p. injection of gabapentin alleviated tactile allodynia in the PSNL mouse model of neuropathic pain at 5:00 compared with 17:00 which correlated with the circadian oscillation of the $\alpha_2\delta_1$ protein level in DRGs.⁵⁹ Additionally, i.p. administration of the NMDA-receptor antagonist ketamine to rats during the resting phase reduced secretion of melatonin from the pineal gland and locomotor activity up to 3 days after anesthesia⁷² and was shown to reduce *Per2* oscillation in NG108-15 neuronal cells up to 12h after ketamine administration.¹⁵

In addition to the pharmacodynamic effects, the circadian rhythms may have effects on the pharmacokinetics of analgesics. Peak concentrations ($C_{\max}$) of morphine and morphine-6-glucuronide (M6G), an active metabolite of morphine, in plasma from patients with cancer pain were detected at 18:00 compared with 14:00 and 10:00 time points.⁴⁶ The synthetic opioid analgesic pethidine showed peak $C_{\max}$ and area under the drug concentration-time curve (AUC) values in serum during the dark phase (21:00) compared with the light phase (9:00) whereas the total serum clearance had the opposite trend in Balb/C mice.¹¹² The $C_{\max}$ and AUC values of the opioid analgesic bucinnazine were significantly increased at 9:00 compared with 21:00 in serum from i.p. injected Sprague Dawley rats.¹¹¹ Furthermore, mice with pain due to streptozotocin-induced diabetic polyneuropathy showed the highest analgesic efficacy when oral pregabalin was administered at the early dark phase which was associated with increased absorption due to circadian oscillations in the intestinal expression of organic cation transporter Octn1.¹

It is becoming more apparent that the timing of analgesic administration may be important for improved efficacy. A well-studied example of the chrono-release drug is prednisone which is a glucocorticoid anti-inflammatory drug for RA. It has been reported that administration of prednisone with modified release (at 2am or 4am) to RA patients is more effective in reducing morning stiffness and/or joint pain compared with immediate-release standard glucocorticoids taken in the morning.^{21,22,27} Similarly, oral administration of sustained release indomethacin, a nonsteroidal anti-inflammatory drug, to patients with osteoarthritis showed a different efficacy based on the timing of administration with evening dosing being more effective in patients with predominantly nocturnal or morning pain whereas morning or noon dosing was more effective in patients with afternoon or evening pain.⁶⁵ When considering chronotherapy for different drugs, it is important to consider not only the half-life of the compound but also whether the target is

rhythmic. Interestingly, the targets of many commonly used drugs are controlled by circadian genes making the timing of administration important¹¹⁵, especially when the half-life of the drug is short. These drugs include e.g. rituximab for rheumatoid arthritis (targeting circadian genes *Fcgr2b*, *Ms4a1*, *Fcgr3*), lidocaine patch for pain (targeting circadian genes *Slc22a5*, *Cyp2b10*, *Egfr*, *Abcb1a*), and eszopiclone for insomnia (targeting circadian genes *Ptgs1*, *Tspo*, *Gabra3*).¹¹⁵

5. Future prospects

Emerging evidence suggests a vicious cycle between sleep and pain which are both regulated by the circadian clock. Whereas the association of genetic variants in circadian genes with sleep regulation and sleep disorders is well characterized, currently there are no genome-wide association studies (GWAS) performed on pain cohorts to identify associations between circadian SNPs and pain with only recent report of association between variable number tandem repeat variant *PER3*^{5/5} and reduced descending pain modulation during day in healthy subjects.²³

Studies on mouse models investigating the connection between sleep and pain should consider that most of the inbred mouse lines are melatonin deficient except for C3H/He and CBA strains.^{45,86} Data from patients and experimental models demonstrate a circadian effect on the efficacy and dosage of analgesics, which suggests that time-targeted analgesic administration should be considered to optimize efficacy. Future pharmacological studies should investigate the potential of small-molecule inhibitors for core circadian genes to fine-tune the circadian clock and to improve pharmacokinetics of currently available analgesics.⁵⁰ Circadian cycle enhancing molecules (CEMs), which can regulate circadian oscillation in gene expression, present novel therapeutic targets for dysregulated circadian clocks in sleep and pain.⁴³

Finally, since the majority of current preclinical pain research is conducted on nocturnal animals such as rodents which are active during the night and sleep during the day, it is crucial to consider the circadian effect on pain and analgesia to successfully translate the findings from nocturnal animal models to patients with diurnal variation in the sleep-wake cycle.

6. Acknowledgments.

VP has no conflicts of interest and has received support from IBSA Foundation, Ulla and Gustaf af Uggle Foundation, KI Rheumatology Foundation and Lars Hiertas Minne Foundation.

IG has received support from Biogen, Adynxx, TARIS Biomedical, AstraZeneca, Pfizer, and Johnson and Johnson and has received support from the Canadian Institutes of Health Research, Physicians' Services Incorporated Foundation, CIHR SPOR Chronic Pain Network, and Queen's University.

BC has no conflicts of interest and has received support from Communication Disorders of the National Institutes of Health R21DC013172 and 1R56DC016415-01, the Swedish Medical Research Council K2014-99X-22478-01-3, Karolinska Institute, Tysta Skolan, and from the Office of the Assistant Secretary of Defense for Health Affairs, through the Neurosensory and Rehabilitation under Award No. W81XWH-16-1-0032.

CS has no conflicts of interest and has received support from Swedish Research Council, Knut and Alice Wallenberg Foundation and Leif Lundblad family.

EK has no conflicts of interest but has received payment for participation in the advisory board of Orion Pharma, Grünenthal and Pierre Fabre.

Opinions, interpretations, conclusions and recommendations are those of the authors and are not influenced by funding sources.

7. References

- [1] Akamine T, Koyanagi S, Kusunose N, Hashimoto H, Taniguchi M, Matsunaga N, Ohdo S. Dosing time-dependent changes in the analgesic effect of pregabalin on diabetic neuropathy in mice. *J Pharmacol Exp Ther* 2015;354:65-72.
- [2] Allen KD, Coffman CJ, Golightly YM, Stechuchak KM, Keefe FJ. Daily pain variations among patients with hand, hip, and knee osteoarthritis. *Osteoarthritis Cartilage* 2009;17:1275-82.
- [3] Alexandre C, Latremoliere A, Ferreira A, Miracca G, Yamamoto M, Scammell TE, Woolf CJ. Decreased alertness due to sleep loss increases pain sensitivity in mice. *Nat Med* 2017;23:768-74.
- [4] Allebrandt KV, Teder-Laving M, Akyol M, Pichler I, Müller-Myhsok B, Pramstaller P, Merrow M, Meitinger T, Metspalu A, Roenneberg T. CLOCK gene variants associate with sleep duration in two independent populations. *Biol Psychiatry* 2010;67:1040-7.
- [5] Alvaro PK, Roberts RM, Harris JK. A Systematic Review Assessing Bidirectionality between Sleep Disturbances, Anxiety, and Depression. *Sleep* 2013;36:1059-68.
- [6] Anyan J, Verwey M, Amir S. Individual differences in circadian locomotor parameters correlate with anxiety- and depression-like behavior. *PLoS One* 2017;12:e0181375.

- [7] Arvidson NG, Gudbjornsson B, Elfman L, Ryden AC, Totterman TH, Hallgren R. Circadian rhythm of serum interleukin-6 in rheumatoid arthritis. *Annals of the Rheumatic Diseases* 1994;53:521-24.
- [8] Aston-Jones G, Chen S, Zhu Y, Oshinsky ML. A neural circuit for circadian regulation of arousal. *Nat Neurosci* 2001;4:732-8.
- [9] Aston-Jones G, Bloom FE. Activity of norepinephrine-containing locus coeruleus neurons in behaving rats anticipates fluctuations in the sleep-waking cycle. *J Neurosci* 1981;1:876-86.
- [10] Atkinson G, Drust B, Reilly T, Waterhouse J. The relevance of melatonin to sports medicine and science. *Sports Med* 2003; 33:809-831.
- [11] Bannister K, Dickenson AH. What do monoamines do in pain modulation? *Curr Opin Support Palliat Care* 2016;10:143-8.
- [12] Bellamy N, Sothorn RB, Campbell J. Rhythmic variations in pain perception in osteoarthritis of the knee. *J Rheumatol* 1990;17:364–372.
- [13] Bellamy N, Sothorn RB, Campbell J, Buchanan WW. Circadian rhythm in pain, stiffness, and manual dexterity in rheumatoid arthritis: relation between discomfort and disability. *Ann Rheum Dis* 1991;50:243–248.
- [14] Bellamy N, Sothorn RB, Campbell J, Buchanan WW. Rhythmic variations in pain, stiffness, and manual dexterity in hand osteoarthritis. *Ann Rheum Dis* 2002;61:1075–1080.
- [15] Bellet MM, Vawter MP, Bunney BG, Bunney WE, Sassone-Corsi P. Ketamine influences CLOCK:BMAL1 function leading to altered circadian gene expression. *PLoS One* 2011;6:e23982.

- [16] Boland EM, Rao H, Dinges DF, Smith RV, Goel N, Detre JA, Basner M, Sheline YI, Thase ME, Gehrman PR. Meta-analysis of the antidepressant effects of acute sleep deprivation. *J Clin Psychiatry* 2017;78:e1020–e1034.
- [17] Boom M, Grefkens J, van Dorp E, Olofsen E, Lourenssen G, Aarts L, Dahan A, Sarton E. Opioid chronopharmacology: influence of timing of infusion on fentanyl's analgesic efficacy in healthy human volunteers. *J Pain Res* 2010;3:183-90.
- [18] Boscariol R, Gilron I, Orr E. Chronobiological characteristics of postoperative pain: diurnal variation of both static and dynamic pain and effects of analgesic therapy. *Can J Anaesth* 2007;54:696-704.
- [19] Brzezinski A. Melatonin in humans. *N Engl J Med* 1997;336:186-95.
- [20] Burish MJ, Chen Z, Yoo SH. Emerging relevance of circadian rhythms in headaches and neuropathic pain. *Acta Physiol (Oxf)* 2019;225:e13161.
- [21] Buttgereit F, Doering G, Schaeffler A, Witte S, Sierakowski S, Gromnica-Ihle E, Jeka S, Krueger K, Szechinski J, Alten R. Efficacy of modified-release versus standard prednisone to reduce duration of morning stiffness of the joints in rheumatoid arthritis (CAPRA-1): a double-blind, randomised controlled trial. *Lancet* 2008;371:205-14.
- [22] Buttgereit F, Kent JD, Holt RJ, Grahn AY, Rice P, Alten R, Yazici Y. Improvement Thresholds for Morning Stiffness Duration in Patients Receiving Delayed- Versus Immediate-Release Prednisone for Rheumatoid Arthritis. *Bull Hosp Jt Dis* (2013) 2015; 73:168-77.
- [23] Carvalho F, Pedrazzoli M, Gasparin A, Dos Santos F, Zortea M, Souza A, da Silva Lucena Torres I, Fregni F, Caumo W. PER3 variable number tandem repeat (VNTR) polymorphism

modulates the circadian variation of the descending pain modulatory system in healthy subjects. *Sci Rep* 2019;9:9363.

[24] Chang AM, Bjornnes AC, Aeschbach D, Buxton OM, Gooley JJ, Anderson C, Van Reen E, Cain SW, Czeisler CA, Duffy JF, Lockley SW, Shea SA, Scheer FA, Saxena R. Circadian gene variants influence sleep and the sleep electroencephalogram in humans. *Chronobiol Int* 2016;33:561-73.

[25] Crofford LJ, Kalogeras KT, Mastorakos G, Magiakou MA, Wells J, Kanik KS, Gold PW, Chrousos GP, Wilder RL. Circadian relationships between interleukin (IL)-6 and hypothalamic-pituitary-adrenal axis hormones: failure of IL-6 to cause sustained hypercortisolism in patients with early untreated rheumatoid arthritis. *J Clin Endocrinol Metab* 1997;82:1279-83.

[26] Cutolo M, Foppiani L, Minuto F. Hypothalamic-pituitary-adrenal axis impairment in the pathogenesis of rheumatoid arthritis and polymyalgia rheumatica. *J Endocrinol Invest* 2002; 25:19-23.

[27] Cutolo M, Iaccarino L, Doria A, Govoni M, Sulli A, Marcassa C. Efficacy of the switch to modified-release prednisone in rheumatoid arthritis patients treated with standard glucocorticoids. *Clin Exp Rheumatol* 2013; 31:498-505.

[28] Daly M, Delaney L, Doran PP, Maclachlan M. The role of awakening cortisol and psychological distress in diurnal variations in affect: a day reconstruction study. *Emotion* 2011; 11:524-532.

- [29] Das V, Kc R, Li X, Varma D, Qiu S, Kroin JS, Forsyth CB, Keshavarzian A, van Wijnen AJ, Park TJ, Stein GS, O-Sullivan I, Burris TP, Im HJ. Pharmacological targeting of the mammalian clock reveals a novel analgesic for osteoarthritis-induced pain. *Gene* 2018; 655:1-12.
- [30] DeBruyne JP, Weaver DR, Reppert SM. CLOCK and NPAS2 have overlapping roles in the suprachiasmatic circadian clock. *Nat Neurosci* 2007; 10:543-5.
- [31] de Tommaso M, Delussi M. Circadian rhythms of migraine attacks in episodic and chronic patients: a cross sectional study in a headache center population. *BMC Neurol* 2018; 18:94.
- [32] Eban-Rothschild A, Rothschild G, Giardino WJ, Jones JR, de Lecea L. VTA dopaminergic neurons regulate ethologically relevant sleep-wake behaviors. *Nat Neurosci* 2016; 19:1356-66.
- [33] Eichhorn N, Treede RD, Schuh-Hofer S. The Role of Sex in Sleep Deprivation Related Changes of Nociception and Conditioned Pain Modulation. *Neuroscience* 2018; 387:191-200.
- [34] Ferracioli-Oda E, Qawasmi A, Bloch MH. Meta-analysis: melatonin for the treatment of primary sleep disorders. *PLoS One* 2013; 8:e63773.
- [35] Fonken LK, Frank MG, Kitt MM, Barrientos RM, Watkins LR, Maier SF. Microglia inflammatory responses are controlled by an intrinsic circadian clock. *Brain, Behavior, and Immunity* 2015; 45:171-1.
- [36] Fourier C, Ran C, Zinnegger M, Johansson AS, Sjöstrand C, Waldenlind E, Steinberg A, Belin AC. A genetic CLOCK variant associated with cluster headache causing increased mRNA levels. *Cephalalgia* 2018; 38:496-502.
- [37] Fu K, Robbins SR, McDougall JJ. Osteoarthritis: the genesis of pain. *Rheumatology (Oxford)* 2018;57(suppl_4):iv43–iv50.

- [38] Franken P. A role for clock genes in sleep homeostasis. *Curr Opin Neurobiol* 2013; 23:864-72.
- [39] Gamble KL, Berry R, Frank SJ, Young ME. Circadian clock control of endocrine factors. *Nat Rev Endocrinol* 2014; 10:466-75.
- [40] Gilron I, Bailey JM, Tu D, Holden RR, Weaver DF, Houlden RL. Morphine, gabapentin, or their combination for neuropathic pain. *New England Journal of Medicine* 2005; 352:1324-34.
- [41] Gilron I, Bailey JM, Vandekerckhof EG. Chronobiological characteristics of neuropathic pain: clinical predictors of diurnal pain rhythmicity. *Clin J Pain* 2013; 29:755-9.
- [42] Gilron I, Ghasemlou N. Chronobiology of chronic pain: focus on diurnal rhythmicity of neuropathic pain. *Curr Opin Support Palliat Care* 2014; 8:429-36.
- [43] Gloston GF, Yoo SH, Chen ZJ. Clock-Enhancing Small Molecules and Potential Applications in Chronic Diseases and Aging. *Front Neurol* 2017; 8:100.
- [44] González MM, Aston-Jones G. Circadian regulation of arousal: role of the noradrenergic locus coeruleus system and light exposure. *Sleep* 2006; 29:1327-36.
- [45] Goto M, Oshima I, Tomita T, Ebihara S. Melatonin content of the pineal gland in different mouse strains. *J Pineal Res.* 1989; 7:195–204.
- [46] Gourlay GK, Plummer JL, Cherry DA. Chronopharmacokinetic variability in plasma morphine concentrations following oral doses of morphine solution. *Pain* 1995; 61:375-81.

- [47] Graves DA, Batenhorst RL, Bennett RL, Wettstein JG, Griffen WO, Wright BD, Foster TS. Morphine requirements using patient-controlled analgesia: influence of diurnal variation and morbid obesity. *Clin Pharm* 1983;2:49-53.
- [48] Gu C, Ramkisoensing A, Liu Z, Meijer JH, Rohling JH. The proportion of light-responsive neurons determines the limit cycle properties of the suprachiasmatic nucleus. *J Biol Rhythms* 2014; 29:16-27.
- [49] Haas S, Straub RH. Disruption of rhythms of molecular clocks in primary synovial fibroblasts of patients with osteoarthritis and rheumatoid arthritis, role of IL-1 β /TNF. *Arthritis Res Ther* 2012; 14:R1222012.
- [50] He B, Chen Z. Molecular Targets for Small-Molecule Modulators of Circadian Clocks. *Curr Drug Metab* 2016; 17:503-12.
- [51] Herzog ED, Hermansteyne T, Smyllie NJ, Hastings MH. Regulating the suprachiasmatic nucleus (SCN) circadian clockwork: interplay between cell-autonomous and circuit-level mechanisms. *Cold Spring Harb Perspect Biol* 2017;9:a027706.
- [52] Hummel T, Kraetsch HG, Lötsch J, Hepper M, Liefhold J, Kobal G. Analgesic effects of dihydrocodeine and tramadol when administered either in the morning or evening. *Chronobiol Int* 1995;12:62-72.
- [53] Iacovides S, George K, Kamerman P, Baker FC. Sleep fragmentation hypersensitizes healthy young women to deep and superficial experimental pain. *J Pain* 2017;18:844-54.
- [54] Jagannath A, Taylor L, Wakaf Z, Vasudevan SR, Foster RG. The genetics of circadian rhythms, sleep and health. *Hum Mol Genet* 2017;26:R128-R138.

- [55] Krause AJ, Prather AA, Wager TD, Lindquist MA, Walker MP. The pain of sleep loss: A brain characterization in humans. *J Neurosci* 2019;2408-18.
- [56] Kripke DF, Kline LE, Nievergelt CM, Murray SS, Shadan FF, Dawson A, Poceta JS, Cronin J, Jamil SM, Tranah GJ, Loving RT, Grizas AP, Hahn EK. Genetic variants associated with sleep disorders. *Sleep Med* 2015; 16:217-24.
- [57] Kumaran Satyanarayanan S, Chien YC, Pei-Chen Chang J, Huang SY, Guu TW, Su H, Su KP. Melatonergic agonist regulates circadian clock genes and peripheral inflammatory and neuroplasticity markers in patients with depression and anxiety. *Brain Behav Immun* 2019; S0889-1591(19)30099-6.
- [58] Kundermann B, Sernal J, Huber MT, Krieg JC, Lautenbacher S. Sleep deprivation affects thermal pain thresholds but not somatosensory thresholds in healthy volunteers. *Psychosom Med* 2004;66:932-7.
- [59] Kusunose N, Koyanagi S, Hamamura K, Matsunaga N, Yoshida M, Uchida T, Tsuda M, Inoue K, Ohdo S. Molecular basis for the dosing time-dependency of anti-allodynic effects of gabapentin in a mouse model of neuropathic pain. *Mol Pain* 2010; 6:83.
- [60] Lane JM, Vlasac I, Anderson SG, Kyle SD, Dixon WG, Bechtold DA, Gill S, Little MA, Luik A, Loudon A, Emsley R, Scheer FA, Lawlor DA, Redline S, Ray DW, Rutter MK, Saxena R. Genome-wide association analysis identifies novel loci for chronotype in 100,420 individuals from the UK Biobank. *Nat Commun* 2016; 7:10889.

- [61] Laposky A, Easton A, Dugovic C, Walisser J, Bradfield C, Turek F. Deletion of the mammalian circadian clock gene BMAL1/Mop3 alters baseline sleep architecture and the response to sleep deprivation. *Sleep* 2005; 28:395-409.
- [62] Lautenbacher S, Kundermann B, Krieg JC. Sleep deprivation and pain perception. *Sleep Med Rev* 2006; 10:357-69.
- [63] Lee H, Nah SS, Chang SH, Kim HK, Kwon JT, Lee S, Cho IH, Lee SW, Kim YO, Hong SJ, Kim HJ. PER2 is downregulated by the LPS-induced inflammatory response in synoviocytes in rheumatoid arthritis and is implicated in disease susceptibility. *Mol Med Rep* 2017; 16:422-28.
- [64] Leger D, Laudon M, Zisapel N. Nocturnal 6-sulfatoxymelatonin excretion in insomnia and its relation to the response to melatonin replacement therapy. *Am J Med*; 116:91-5.
- [65] Levi F, Le Louarn C, Reinberg A. Timing optimizes sustained-release indomethacin treatment of osteoarthritis. *Clin Pharmacol Ther* 1985; 37:77-84.
- [66] Liberman AR, Kwon SB, Vu HT, Filipowicz A, Ay A, Ingram KK. Circadian Clock Model Supports Molecular Link Between PER3 and Human Anxiety. *Sci Rep* 2017;7:9893.
- [67] Luo AH, Aston-Jones G. Circuit projection from suprachiasmatic nucleus to ventral tegmental area: a novel circadian output pathway. *Eur J Neurosci* 2009;29:748-60.
- [68] Mallmann RT, Elgueta C, Sleman F, Castonguay J, Wilmes T, van den Maagdenberg A, Klugbauer N. Ablation of Ca(V)2.1 voltage-gated Ca²⁺ channels in mouse forebrain generates multiple cognitive impairments. *PLoS One* 2013;8:e78598.

- [69] Marchand A, Durand P, Juster RP, Lupien SJ. Workers' psychological distress, depression, and burnout symptoms: associations with diurnal cortisol profiles. *Scand J Work Environ Health* 2014;40:305-314.
- [70] May ME, Harvey MT, Valdovinos MG, Kline RH 4th, Wiley RG, Kennedy CH. Nociceptor and age specific effects of REM sleep deprivation induced hyperalgesia. *Behav Brain Res* 2005;159:89-94.
- [71] Meerlo P, Sgoifo A, Suchecki D. Restricted and disrupted sleep: effects on autonomic function, neuroendocrine stress systems and stress responsivity. *Sleep Med Rev* 2008; 12:197-210.
- [72] Mihara T, Kikuchi T, Kamiya Y, Koga M, Uchimoto K, Kurahashi K, Goto T. Day or night administration of ketamine and pentobarbital differentially affect circadian rhythms of pineal melatonin secretion and locomotor activity in rats. *Anesth Analg* 2012; 115:805-13.
- [73] Morioka N, Saeki M, Sugimoto T, Higuchi T, Zhang FF, Nakamura Y, Hisaoka-Nakashima K, Nakata Y. Downregulation of the spinal dorsal horn clock gene *Per1* expression leads to mechanical hypersensitivity via c-jun N-terminal kinase and CCL2 production in mice. *Mol Cell Neurosci* 2016; 72:72-83.
- [74] Mork PJ, Nilsen TI. Sleep problems and risk of fibromyalgia: Longitudinal data on an adult female population in Norway. *Arthritis Rheum* 2012; 64:281-4.
- [75] Nagy AD, Iwamoto A, Kawai M, Goda R, Matsuo H, Otsuka T, Nagasawa M, Furuse M, Yasuo S. Melatonin adjusts the expression pattern of clock genes in the suprachiasmatic nucleus

and induces antidepressant-like effect in a mouse model of seasonal affective disorder. *Chronobiol Int* 2015; 32:447-57.

[76] Narasimamurthy R, Hatori M, Nayak SK, Liu F, Panda S, Verma IM. Circadian clock protein cryptochrome regulates the expression of proinflammatory cytokines. *PNAS* 2012; 109:12662-7.

[77] Nascimento DC, Andersen ML, Hipólido DC, Nobrega JN, Tufik S. Pain hypersensitivity induced by paradoxical sleep deprivation is not due to altered binding to brain mu-opioid receptors. *Behav Brain Res* 2007; 178:216-20.

[78] Nees F, Löffler M, Usai K, Flor H. Hypothalamic-pituitary-adrenal axis feedback sensitivity in different states of back pain. *Psychoneuroendocrinology* 2019; 101:60-66.

[79] Nitter AK, Pripp AH, Forseth K. Are sleep problems and non-specific health complaints risk factors for chronic pain? A prospective population-based study with 17 year follow-up. *Scandinavian Journal of Pain* 2012; 3:210-7.

[80] Odegard SS, Sand T, Engstrom M, Stovner LJ, Zwart JA, Hagen K. The Long-Term Effect of Insomnia on Primary Headaches: A Prospective Population-Based Cohort Study (HUNT-2 and HUNT-3) Headache: The Journal of Head and Face Pain 2011; 51:570-80.

[81] Odo M, Koh K, Takada T, Yamashita A, Narita M, Kuzumaki N, Ikegami D, Sakai H, Iseki M, Inada E, Narita M. Changes in circadian rhythm for mRNA expression of melatonin 1A and 1B receptors in the hypothalamus under a neuropathic pain-like state. *Synapse* 2014; 68:153-8.

- [82] Odrich M, Bailey JM, Cahill CM, Gilron I. Chronobiological characteristics of painful diabetic neuropathy and postherpetic neuralgia: Diurnal pain variation and effects of analgesic therapy. *Pain* 2006; 120:207-12.
- [83] Park JW, Cho SJ, Park SG, Chu MK. Circadian variations in the clinical presentation of headaches among migraineurs: A study using a smartphone headache diary. *Chronobiol Int* 2018; 35:546-554.
- [84] Parsons MJ, Lester KJ, Barclay NL, Archer SN, Nolan PM, Eley TC, Gregory AM. Polymorphisms in the circadian expressed genes *PER3* and *ARNTL2* are associated with diurnal preference and $GN\beta 3$ with sleep measures. *Sleep Med* 2015; 16:217-24.
- [85] Partch CL, Green CB, Takahashi JS. Molecular architecture of the mammalian circadian clock. *Trends Cell Biol* 2014; 24:90-9.
- [86] Peirson SN, Brown LA, Potheary CA, Benson LA, Fisk AS. Light and the laboratory mouse. *J Neurosci Methods* 2018; 15;300:26-36.
- [87] Perreau-Lenz S, Sanchis-Segura C, Leonardi-Essmann F, Schneider M, Spanagel R. Development of morphine-induced tolerance and withdrawal: involvement of the clock gene *mPer2*. *Eur Neuropsychopharmacol* 2010; 20:509-17.
- [88] Qiu P, Jiang J, Liu Z, Cai Y, Huang T, Wang Y, Liu Q, Nie Y, Liu F, Cheng J, Li Q, Tang YC, Poo M, Sun Q, Chang HC. *BMAL1* knockout macaque monkeys display reduced sleep and psychiatric disorders. *National Science Review* 2019; 1:87-100.
- [89] Roehrs TA, Harris E, Randall S, Roth T. Pain sensitivity and recovery from mild chronic sleep loss. *Sleep* 2007; 30:494-505.

- [90] Rozen TD, Fishman RS. Cluster headache in the United States of America: demographics, clinical characteristics, triggers, suicidality, and personal burden. *Headache* 2012; 52:99-113.
- [91] Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE. Sleep state switching. *Neuron* 2010; 68:1023-42.
- [92] Schrimpf M, Liegl G, Boeckle M, Leitner A, Geisler P, Pich C. The effect of sleep deprivation on pain perception in healthy subjects: a meta-analysis. *Sleep Med* 2015; 16:1313-20.
- [93] Segal JP, Tresidder KA, Bhatt C, Gilron I, Ghasemlou N. Circadian control of pain and neuroinflammation. *J Neurosci Res* 2018; 96:1002-20.
- [94] Smith MT, Perlis ML, Smith MS, Giles DE, Carmody TP. Sleep quality and presleep arousal in chronic pain. *Journal of Behavioral Medicine* 2000; 23:1-13.
- [95] Smith MT, Edwards RR, McCann UD, Haythornthwaite JA. The effects of sleep deprivation on pain inhibition and spontaneous pain in women. *Sleep* 2007; 30:494-505.
- [96] Smolensky MH, Portaluppi F, Manfredini R, Hermida RC, Tiseo R, Sackett-Lundeen LL, Haus EL. Diurnal and twenty-four hour patterning of human diseases: acute and chronic common and uncommon medical conditions. *Sleep Med Rev* 2015; 21:12-22.
- [97] Steinberg A, Fourier C, Ran C, Waldenlind E, Sjöstrand C, Belin AC. Cluster headache - clinical pattern and a new severity scale in a Swedish cohort. *Cephalalgia* 2018; 38:1286-95.
- [98] Sutton BC, Opp MR. Sleep fragmentation exacerbates mechanical hypersensitivity and alters subsequent sleep-wake behavior in a mouse model of musculoskeletal sensitization. *Sleep* 2014; 37:515-24.

- [99] Takada T, Yamashita A, Date A, Yanase M, Suhara Y, Hamada A, Sakai H, Ikegami D, Iseki M, Inada E, Narita M. Changes in the circadian rhythm of mRNA expression for μ -opioid receptors in the periaqueductal gray under a neuropathic pain-like state. *Synapse* 2013; 67:216-23.
- [100] Takahashi JS. Transcriptional architecture of the mammalian circadian clock. *Nat Rev Genet* 2017; 18:164-79.
- [101] Toh KL, Jones CR, He Y, Eide EJ, Hinz WA, Virshup DM, Ptáček LJ, Fu YH. An hPer2 phosphorylation site mutation in familial advanced sleep phase syndrome. *Science* 2001; 291:1040-3.
- [102] Tracey I, Mantyh PW. The cerebral signature for pain perception and its modulation. *Neuron* 2007; 55:377-91.
- [103] Vgontzas AN, Bixler EO, Lin HM, Prolo P, Mastorakos G, Vela-Bueno A, Kales A, Chrousos GP. Chronic insomnia is associated with nyctohemeral activation of the hypothalamic-pituitary-adrenal axis: clinical implications. *J Clin Endocrinol Metab* 2001; 86:3787-94.
- [104] Waldenlind E, Gustafsson SA, Ekblom K, Wetterberg L. Circadian secretion of cortisol and melatonin in cluster headache during active cluster periods and remission. *J Neurol Neurosurg Psychiatry*. 1987; 50:207-213.
- [105] Wei Y, Blanken TF, Van Someren EJW. Insomnia Really Hurts: Effect of a Bad Night's Sleep on Pain Increases With Insomnia Severity. *Front Psychiatry* 2018;9:377.

- [106] Wilhelmsen M, Amirian I, Reiter RJ, Rosenberg J, Gögenur I. Analgesic effects of melatonin: a review of current evidence from experimental and clinical studies. *J Pineal Res* 2011; 51:270-277.
- [107] Wisor JP, O'Hara BF, Terao A, Selby CP, Kilduff TS, Sancar A, Edgar DM, Franken P. A role for cryptochromes in sleep regulation. *BMC Neurosci* 2002; 3:20.
- [108] Wisor JP, Pasumarthi RK, Gerashchenko D, Thompson CL, Pathak S, Sancar A, Franken P, Lein ES, Kilduff TS. Sleep deprivation effects on circadian clock gene expression in the cerebral cortex parallel electroencephalographic differences among mouse strains. *J Neurosci* 2008; 28:7193-201.
- [109] Xia T, Cui Y, Qian Y, Chu S, Song J, Gu X, Ma Z. Regulation of the NR2B-CREB-CRTC1 signaling pathway contributes to circadian pain in murine model of chronic constriction injury. *Anesthesia and Analgesia* 2016; 122:542-52.
- [110] Yoshida M, Ohdo S, Takane H, Tomiyoshi Y, Matsuo A, Yukawa E, Higuchi S. Chronopharmacology of analgesic effect and its tolerance induced by morphine in mice. *J Pharmacol Exp Ther* 2003; 305:1200-5.
- [111] Yu Z, Li X, Feng C, Lei K, He W, Zhang C, Liu D. Circadian variations in the pharmacokinetics of bucinnazine in rats. *Biological Rhythm Research* 2019. In Press.
- [112] Zhang C, Yu Z, Li X, Xu Y, Liu D. Chronopharmacodynamics and chronopharmacokinetics of pethidine in mice. *PLoS One* 2014; 9:e102054.

- [113] Zhang J, Li H, Teng H, Zhang T, Luo Y, Zhao M, Li YQ, Sun ZS. Regulation of peripheral clock to oscillation of substance P contributes to circadian inflammatory pain. *Anesthesiology* 2012; 117:149-60.
- [114] Zhang P, Moye LS, Southey BR, Dripps I, Sweedler JV, Pradhan A, Rodriguez-Zas SL. Opioid-Induced Hyperalgesia Is Associated with Dysregulation of Circadian Rhythm and Adaptive Immune Pathways in the Mouse Trigeminal Ganglia and Nucleus Accumbens. *Mol Neurobiol* 2019; In Press.
- [115] Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB. A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci USA* 2014; 111:16219-24.

FIGURE LEGENDS

Figure 1. **Circadian clock at the interplay between sleep and pain.** Light responsive neurons in the suprachiasmatic nucleus to light stimuli from retina are master regulators of circadian clock¹⁰⁰ which synchronize the peripheral clocks in peripheral tissues through the autonomic nervous system and secretion of hormones, cytokines and neurotransmitters involved in regulation of both sleep and pain.^{35,76,93} Furthermore, the core circadian genes promote stable oscillations in gene expression for some genes involved in regulation of sleep and pain.⁵⁴ The relationship between sleep and pain is bidirectional since insufficient sleep can precede and maintain pain while pain can impair our ability to initiate and maintain sleep.⁶²

Figure 2. The molecular clock machinery. The core mammalian circadian genes are two transcriptional factors, circadian locomotor output cycles kaput (CLOCK) and brain and muscle aryl hydrocarbon receptor nuclear translocator-like 1 (BMAL1), which – in the form of CLOCK/BMAL1 heterodimer – promote the rhythmic transcription of repressor proteins Period (Per1/2/3) and Cryptochrome (Cry1/2).^{30,100} The PER and CRY proteins drive the negative feedback loop by forming a complex which translocates into the nucleus during the evening to repress the transcription of CLOCK/BMAL1 and therefore its own transcription.⁸⁵ In addition to regulation of Per and Cry transcription, CLOCK/BMAL1 activates the nuclear receptors retinoid-related orphan receptor alpha (ROR α ; also known as NR1F1) and REV-ERB α (also known as NR1D1) which compete with each other to bind a promoter of Bmal1. The ROR α is an activator, whereas REV-ERB α is a repressor, of Bmal1 transcription which contributes to cyclic expression of the Bmal1 gene.⁵⁴

Light input from the retina through the retinohypothalamic tract

Master circadian clock

Light responsive neurons in suprachiasmatic nucleus

Circadian genes

Secretion of hormones, cytokines and neurotransmitters

Sleep

Pain

↑ Insufficient sleep

↑ Pain sensitivity
↑ Pain intensity
↓ Descending pain inhibition