<https://helda.helsinki.fi>

Staff Training Interventions to Improve End-of-Life Care of Nursing Home Residents: A Systematic Review

Lamppu, Pauli J

2021-02

Lamppu , P J & Pitkälä , K 2021 , ' Staff Training Interventions to Improve End-of-Life Care of Nursing Home Residents: A Systematic Review ' , Journal of the American Medical Directors Association , vol. 22 , no. 2 , pp. 268-278 . <https://doi.org/10.1016/j.jamda.2020.09.011>

<http://hdl.handle.net/10138/335660>

<https://doi.org/10.1016/j.jamda.2020.09.011>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Staff training interventions to improve end-of-life care of nursing home residents: A systematic review

Pauli Lamppu^{a,b} M.D., Kaisu H. Pitkala^{a,c} M.D., PhD

^a Department of General Practice and Primary Health Care, University of Helsinki

^b Department of Social Services and Health Care, Geriatric Clinic, Helsinki Hospital, Helsinki, Finland

^c Unit of Primary Health Care, Helsinki University Hospital, Helsinki, Finland

Correspondence to:

Pauli Lamppu, M.D.

pauli.lamppu@helsinki.fi

Department of General Practice and Primary Health Care

University of Helsinki

P.O. Box 20

00014 University of Helsinki

Finland

Running title: Staff training effects on residents' end-of-life care

Words: 224 (abstract) 3163 (main text), 3 tables, 1 figure, 72 references

Key words: advance care planning, education, end-of-life care, nursing home, palliative care, systematic review

Funding: This study was supported by Helsinki University Hospital and with scholarships from the Uulo Arhio Foundation, the Finnish Medical Foundation, and the Foundation of General Practitioners in Finland. The funding sources had no role in the design and conduct of the study, in the collection, analysis, and interpretation of the data, or in the preparation, review, and approval of the manuscript.

29 *Summary:* 16 randomized controlled trials examined the effect of staff training on residents' quality
30 of life, end-of-life care, and hospitalizations. These interventions had minor effects, and further
31 research is needed to guide policies.

32

33 **Staff training interventions to improve end-of-life care of nursing home residents:**

34 **A systematic review**

35 **Abstract**

36 **Objectives:** The aim was to review evidence from all randomized controlled trials (RCTs) using
37 palliative care education or staff training as an intervention to improve nursing home residents'
38 quality of life (QOL) or quality of dying (QOD) or to reduce burdensome hospitalizations.

39 **Design:** A systematic review with a narrative summary.

40 **Setting and participants:** Residents in nursing homes and other long-term care facilities

41 **Methods:** We searched MEDLINE, CINAHL, PsycINFO, the Cochrane Library, Scopus, and Google
42 Scholar, references of known articles, previous reviews, and recent volumes of key journals. RCTs
43 were included in the review. Methodological quality was assessed.

44 **Results:** The search yielded 932 articles after removing the duplicates. Of them, 16 cluster RCTs
45 fulfilled inclusion criteria for analysis. There was a great variety in the interventions with respect to
46 learning methods, intensity, complexity, and length of staff training. Most interventions featured other
47 elements besides staff training. In the six high-quality trials only one showed a reduction in
48 hospitalizations, whereas among six moderate-quality trials two suggested a reduction in
49 hospitalizations. None of the high-quality trials showed effects on residents' QOL or QOD. Staff
50 reported an improved QOD in one moderate-quality trial.

51 **Conclusion and implications:** Irrespective of the means of staff training, there were surprisingly few
52 effects of education on residents' QOL, QOD, or burdensome hospitalizations. Further studies are
53 needed to explore the reasons behind these findings.

54

55 **Introduction**

56 Long-term care facilities take care of a large majority of older people at the end of their life.^{1,2}
57 Multimorbidity and functional and cognitive decline are the major challenges in their day-to-day care,
58 whereas the approaching end of life gets less attention.³ High-turnover nursing home staff have large
59 educational needs.^{1,2} The need for quality end-of-life care in this setting has been widely recognized
60 with various development projects to improve residents' end-of-life care.⁴

61 Successful end-of-life care in nursing homes requires the staff to have a certain set of practical skills.⁵
62 Active advance care planning (ACP) to guide actions in different scenarios is important.⁶ In addition,
63 skills to communicate with the residents and their proxies⁷ and the ability to deliver good symptom
64 care are required.^{1,2} Previous research has suggested that hospitalizations of nursing home residents
65 may lead to many adverse effects such as delirium, infections, functional decline, falls, pressure
66 ulcers, and decreased comfort without survival benefit.^{8,9} Nursing home staff should therefore have
67 the necessary skills and attitudes to prevent all avoidable hospitalizations and organize the required
68 treatments in place without hospitalization.

69 Previous systematic reviews focusing on improvements in palliative care have been derived from
70 various perspectives.^{10–13} Training has an obvious potential to introduce and reinforce practical skills
71 and change in operating culture.¹³ Many of the studies examining training interventions evaluate
72 outcomes related to change in staff knowledge or other staff-level outcomes.¹⁴ However, there seems
73 to be a paucity of evidence on how this gained knowledge would transfer into clinically relevant
74 resident-level outcomes.¹⁵ Previous studies suggest that resident's quality of life (QOL), quality of
75 dying (QOD), symptom management, proxy satisfaction, and the likelihood of burdensome
76 hospitalizations are important resident-level measures to define successful end-of-life care in this
77 population.^{11,16,17}

78 In this systematic review, we wanted to gather the evidence from randomized controlled trials (RCTs)
79 examining the effect of an intervention featuring an element of nursing home staff training on the
80 resident-level outcomes QOL, QOD, symptom management, proxy satisfaction, or residents' risk of
81 hospital use.

82

83 **Methods**

84 The protocol for this systematic review was devised prior to the final searches and review process.
85 The protocol was published and can be found in PROSPERO (ID 160497).

86

87 *Study eligibility criteria*

88 We aimed for a clinically comprehensive yield with various types of interventions and aims. The
89 following criteria had to be met:

- 90 - Study was an RCT or cluster RCT.
- 91 - Study patients resided in a nursing home or comparable long-term care facility.
- 92 - Intervention included structured education or on-site training in end-of-life related topics for
93 a large proportion of staff (e.g., lectures, seminars, videos, facilitated small group discussions,
94 case conferences, role plays, clinical coaching, benchmarking, and feedback). Staff taking
95 care of a comparison group of patients received no training, continued usual care, or were
96 provided with only attention in some way.
- 97 - Outcome: resident-related outcomes such as QOL, QOD, hospitalizations or use of hospital
98 days, or proxy satisfaction with care.
- 99 - Abstract and full text in English.

100

101 *Search strategy*

102 The primary search was performed on March 3, 2020 and repeated before submitting for publication
 103 in June 2020. We searched MEDLINE, PsycINFO, CINAHL, Scopus, Cochrane Library, and Google
 104 Scholar using the following terms adapted to each database: (nursing home OR assisted living OR
 105 long-term care OR residential care OR care home) AND (education OR quality improvement OR
 106 training OR train-the-trainer) AND (end-of-life care OR palliative care OR terminal care OR quality
 107 of dying OR comfort OR hospitalization OR quality of life OR health-related quality of life OR
 108 satisfaction with care OR advance care planning) AND (randomized controlled trial OR cluster
 109 randomized OR RANDOMI*).

110 We also hand-searched reference lists of relevant previous systematic reviews^{5,7,10,11,11–14,18–23}, known
 111 key articles, and recent volumes of key journals. See Figure 1 for the study selection flowchart. When
 112 our search identified a trial protocol meeting our criteria, we additionally searched for an additional
 113 article presenting the results.

114 *Methodological quality*

115 Both authors (P.L. and K.H.P.) independently assessed the methodological quality of the included
 116 articles according to 10 predetermined quality criteria. The list of criteria was composed from a past
 117 version of the Cochrane Handbook²⁴, a Delphi list²⁵, and criteria developed by the Evidence-based
 118 Medicine Working Group.²⁶ For all the criteria, see Table 1. Each criterion was worth one point and
 119 each criterion and study were discussed until consensus was reached. A total of 8–10 points was
 120 considered good methodological quality, a score of 5–7 moderate quality, and a score < 5 low quality.

121 *Path from theory to resident-related effects*

122 We explored each trial in terms of what kind of pedagogical methods it used in training the staff and
 123 which learning theories it implemented as the basis for intervention. We also noted the length and

intensity of the training and the staff training attrition rate. The complexity of the entire intervention and the role of training in its entirety was assessed.

The outcome measures we required for inclusion were patient-related.^{15,16} Residents' QOL, QOD, and symptom burden were considered such outcomes. In addition, acute hospitalizations or hospital days were included as patient-related outcomes. We also explored resident and proxy satisfaction with care. These patient-related outcomes were chosen by the authors from previous literature as being the most appropriate.^{2,15-17}

131

132 **Results**

133 *Study selection*

The search yielded 1046 studies. After screening the titles of the studies, we first removed duplicates (N = 114) and then those not fulfilling our criteria (N = 730). We screened abstracts of all the remaining 202 studies for eligibility. Of them, 140 did not fulfill our inclusion criteria. Then all full texts of 62 articles were assessed in detail for eligibility criteria and discussed between authors until consensus on inclusion was reached. Overall, 16 studies met the inclusion criteria. A systematic literature search yielded 15 studies and one additional study was identified from a previous systematic review.¹⁰ Reasons for exclusion were mainly related to intervention or outcome not fulfilling our inclusion criteria (see Figure 1).

142 *Characteristics of studies*

Table 2 provides an overview of the included studies. All studies had a cluster randomized trial design and were conducted in nursing homes. One study⁵³ employed a stepped-wedge design, while the rest were conventional parallel-group trials. The number of participants ranged from 72 to 23 478 per study and altogether the trials reported data on 40 712 participants. The participants' mean age was

high (mean age range from 82 to 90) and participants were predominantly female. Of the studies, four included only patients with dementia.^{30,35,38,40} and the oldest study included an atypical nursing home population with mainly cancer patients.²⁷ In five studies^{35,37,40,45,51} the final analyses included only participants who died during the follow-up.

The education intervention varied greatly between the studies. Although articles described their interventions, in most studies the description was not detailed enough to plan a valid replication of education. None of the trials presented any learning theory to justify their type of education. Furthermore, they rarely discussed how well the staff had complied with education or adopted new competences. A wide variety of learning methods was used in these studies. Few studies used active involvement of staff in incorporating the new approach of palliative care into their units.^{30,32,45,49} It seems that lectures or webinars were implied in nine studies^{30,32,33,37,38,42,44,45,47}, although in several studies it was not clearly stated how the education was delivered in sessions for a large number of professionals. Seminars or active small group discussions were mentioned in five studies^{27,31,33,47,49} and case-based learning in three studies.^{33,35,53} Coaching as a form of supportive assistance and facilitation was used in 10 studies.^{31–33,35,37,42,44,45,51,53} Role plays for communication training were used in three studies.^{27,47,49} Feedback information about care outcomes was featured in four studies.^{32,33,37,40} From the available information we judged that nine studies employed some form of train-the-trainer methodology where the study teams trained a group of “facilitators” or “facility champions” who in turn distributed the intervention to the facilities.^{30,31,35,37,42,44,47,49,51} Two studies trained also nursing home residents’ proxies using videos.^{31,38} Training was often used as part of a complex intervention featuring also, for example, tool kits or guides for managing various situations in end-of-life care^{32,37,42}, discussion aids^{35,38}, or informing proxies via video aids.^{31,38} Some interventions even featured major organizational changes such as the formation of new units³⁰ or systematic multicomponent reviews⁴⁷. The most common other components were organization of ACP sessions or facilitating their formation in other ways^{31,33,35,37,38,47,49,51}, medication reviews⁴⁷, and

172 organizing meaningful activities^{30,47}.

173 The intensity of training also varied greatly. Whereas some interventions lasted only hours or
 174 days^{27,38,53}, others included intensive weekly seminars or other inputs for up to several months
 175 ^{33,45,47,51}; some even consisted of up to 18–24 months of continuous implementation^{35,37}.

176 *Methodological quality*

177 Of the 16 studies, six were considered high quality whereas six were considered moderate quality and
 178 four low quality. The most common methodological problems were not using blinded assessments,
 179 not using intention-to-treat analysis when assessing the outcomes, and not describing the dropouts.
 180 Several studies analyzed only deceased participants and did not discuss others who had been
 181 recruited. Furthermore, randomization was only briefly mentioned in several studies and the method
 182 was not described. In addition, many studies gave very little information on the baseline
 183 characteristics of their participants.

184 *Effects of training interventions*

185 *Quality of life*

186 Of the included studies, five included QOL as an outcome.^{27,35,42,45,47} In the included trials, QOL (or
 187 health-related QOL (HRQOL)) was assessed with depression scaling, presence of pain, and with five
 188 different validated measures: QUALIDEM (a dementia-specific QOL assessment instrument)⁵⁶,
 189 QUALID (Quality of Life in Late-Stage Dementia scale)⁵⁷, EQ-5D (EuroQol 5D) and its subscale the
 190 EQ-VAS (EuroQol Visual Analogue Scale)⁵⁸, and the Health Utilities Index (HUI)⁵⁹. Two studies^{35,47}
 191 used nurses to assess QOL, one⁴² used a research assistant, and in two^{27,45} the outcomes were resident
 192 reported. None of the good- and moderate-quality studies^{35,42,45,47} reporting QOL produced clear
 193 benefits. Arendts et al.⁴² reported two QOL measures of which the HUI instrument in the intervention
 194 group showed less lowering of HRQOL.

195

196

197 *Quality of dying, symptom management, and proxy satisfaction with care*

198 QOD, symptom management, or proxy satisfaction was assessed in eight studies.^{30,31,35,38,40,44,51,53}

199 End-Of-Life in Dementia (EOLD) scales include three different QOD measurement scales.⁶⁰ Two of

200 these scales – Symptom Management (SM-EOLD) or Comfort Assessment in Dying (CAD-EOLD)

201 – were used in five studies.^{35,38,40,44,51} Only one low-quality trial⁴⁴ showed effects in these scales,

202 while four higher-quality trials did not. One moderate-⁵³ and one high⁵¹-quality trial used Quality of

203 dying in long-term care (QOD-LTC)⁶¹ and Quality of Death and Dying (QODD)⁶² scales and both

204 showed modest but statistically significant intervention effects.

205 Satisfaction with care, rated by proxy or staff member after death using SWC-EOLD, was used as a

206 proxy for QOD. It was assessed in four studies, none of these producing effects.^{36,38,40,51} The Family

207 Perception of Care Score (FPCS) evaluates similar aspects of care such as quality of communication

208 with proxies and preparatory tasks for death and palliative care.⁶³ It was assessed in one low-quality

209 trial with intervention showing benefits.⁴⁴ One study³¹ assessed satisfaction with care in living

210 residents using the Relative and Patient Satisfaction Indexes⁶⁴ without intervention effects in this

211 outcome.

212 The Discomfort Scale for Dementia of the Alzheimer's Type (DS-DAT)⁶⁵ was used to evaluate

213 symptoms in a single low-quality study with positive intervention effects.³⁰

214 *Hospital use*

215 Hospitalizations were measured in 10 trials.^{31–33,35,37,38,42,45,49,53} Of the high-quality trials, three

216 showed a decreased number of hospitalizations in the intervention group compared with the control

217 group.^{31,38,53} The largest high-quality trial with a pragmatic design did not have any effect on

218 hospitalizations.³⁷ Only two trials measured hospital days as an outcome measure.^{33,54} The high-
219 quality trial suggested no effect of intervention³³, whereas the moderate-quality trial⁵⁴ showed a
220 reduction in hospital days.

221 *Characteristics of effective training interventions*

222 Table 3 summarizes the characteristics and findings of high- and moderate-quality trials in order to
223 explore what kind of pedagogical interventions and resident target group are needed for effective
224 intervention.

225 Overall, few trials had effectiveness. There seem to be no particular characteristics in the residents,
226 pedagogical methods, or length or intensity of training related to these effective trials. The effects
227 related to hospital use were seen in trials which provided education to proxies in addition to staff
228 ^{31,38}. In the third trial showing lower hospital use an external palliative care nurse rounded with the
229 ward staff discussing selected patients and offering expert consultations when needed.⁵⁴ In addition,
230 in the same trial staff reported better QOD among their residents.⁵³

231

232 **Discussion**

233 Our systematic review identified 16 cluster randomized trials examining various educational
234 interventions aiming to improve end-of-life care in nursing homes: improvements in residents' QOL
235 and QOD, or reducing their burdensome hospitalizations. The effects of interventions were
236 surprisingly mild, with only one high-quality trial and two moderate-quality trials showing reductions
237 in burdensome hospitalizations. None of the six high-quality trials showed improvements in
238 residents' QOL or QOD, whereas in one moderate-quality trial the staff reported improved QOD.
239 There has been a high number of participants in these educational trials. The learning methods,
240 intensity, and length have varied greatly.

241 The quality of the included studies is quite good. The high number of participants allows enough
242 statistical power to evaluate also categorical outcomes such as hospitalizations. For example, about
243 1000 participants randomized into two groups suggests > 80% power to detect 30% of differences
244 when mean hospitalizations per participant is 0.7 and SD 1.5. In addition, measures used in studies
245 on QOL and QOD have been well-validated. Thus, the lack of effectiveness is not due to problems
246 in statistical power, insensitive outcome measures, or inappropriate statistical methods. The
247 pedagogical approach also used various means. Thus, if education is an effective intervention to
248 improve nursing home residents' end-of-life care, this systematic review of 16 trials should have
249 shown more positive findings. The problem of ineffective complex interventions in primary care
250 cRCTs has been discussed in a previous study. This systematic review called for studies with
251 alternative designs but also discussed that cRCTs generally give more trustworthy results than
252 observational studies.⁶⁶

253 In Table 3 summarizing the effectiveness and elements of training interventions in high- and
254 moderate-quality trials, two of the trials providing reductions in burdensome hospitalizations had
255 significant intervention elements of proxy education. The residents' proxies received video training
256 to enhance ACP decision-making³⁸ or proxies were trained to complete advance directives.³¹ In
257 addition, one trial showed a decrease in hospitalizations and hospital days.⁵⁴ In this trial, external
258 palliative care nurses were integrated into nursing home teams to enhance ACP. They discussed
259 selected residents on monthly visits and offered external expert consultations when needed.
260 Furthermore, only one moderate-quality trial suggested some staff-reported effects on QOD.⁵³ In this
261 study, the staff assessed the effects of their own work, leading to a risk of bias. Thus, it seems to be
262 difficult to change the culture and attitudes of staff to improve end-of-life care at resident-level
263 outcomes.

264

265 A wide variety of training interventions has been used in these trials. They included behavioristic
266 interventions such as lectures and also activating learning methods such as case-based learning or
267 role play. However, only very small effects were seen in resident outcomes. None of the trials have
268 justified their training with respect to learning theories or an adult pedagogical approach. However,
269 many trials included possibilities for reflection and integration of new competencies into everyday
270 practice. These have been highlighted as an effective means to enhance adult learning.⁶⁷⁻⁷⁰ In end-of-
271 life care training it should be essential to tackle professionals' own feelings, for example related to
272 fear of death, in order to change their attitudes and culture.²⁸ None of the trials described whether the
273 learning needs or especially the motivation of staff had been assessed before the training. Motivation
274 was found to be an important feature determining the outcome in the INTERACT project.⁷¹ The
275 popular train-the-trainer approach might be efficient in distributing interventions and to engage
276 facilities by making the interventions originate from "inside" the facility, with the ability to adjust the
277 intervention to suit the practices in each facility. On the other hand, nurses or nurse practitioners who
278 were most often recruited to be facilitators did training without any apparent pedagogical training and
279 will be more susceptible to pressure from other staff members or managers than an outside expert
280 might be.

281 Implementing training elements into practice is a critical element and tools to do so were used in
282 several trials. Several trials evaluated the use of the new skills or practices after initial training –
283 giving feedback to trainees or even re-training. However, this effort did not translate into better
284 resident outcomes. It is impossible to determine whether low effectiveness is due to training or
285 implementation. It is important to assess the fidelity of new practices and skills also in future studies
286 to determine how well the changes are implemented.

287

288 The two largest included trials were both complex interventions. The early pilot INTERACT project
289 showed significant reductions in burdensome hospitalizations⁷² but later more rigorous trials did not

290 manage to show effects.³⁷ Also, the large European multicenter collaboration PACE, with its well
291 outlined and long-lasting intervention, failed to show an effect on end-of-life care.⁵¹ A previous
292 systematic review noted that interventions having narrow aims and low complexity were more likely
293 to be beneficial in changing care practices than those with ambitious aims and high complexity.¹³
294 Ambitious and complex interventions may be hindered by common barriers such as high staff
295 turnover, a high training attrition rate, low organizational support, a lack of resources or commitment,
296 or opposing cultures and attitudes.

297

298 *Strengths and limitations*

299 Previous systematic reviews noted that staff training may influence relevant skills and has been
300 mostly well received by staff.^{13,14} However, to our knowledge this is the first systematic review to
301 assess the effects of staff training on resident-related end-of-life care outcomes. We managed to find
302 a large number of studies with robust RCT design and with a clinically oriented research question.
303 Most of our included studies were not included in previous related reviews; thus most of the
304 contribution is novel. Our findings suggest that influencing resident-related outcomes is hard, and
305 positive changes in staff-related outcomes may often fail to translate into resident-level outcomes.

306 The included studies presented varying perspectives and interventions. The interventions were too
307 diverse to perform a meta-analysis for example on hospitalizations. We wanted to focus on training
308 but most of the studies had complex interventions where training played a varying but often limited
309 role, thus making the assessment of the training effects challenging. Our evaluation of methodological
310 quality seems to give a useful numerical and categorical evaluation of the different methodologies
311 but has not been optimized to evaluate clustered trials.

312 **Conclusions and implications**

313 In this systematic review, we explored evidence of the effect of cluster RCTs using staff training
314 intervention on the resident-level outcomes QOL, QOD, symptom management, proxy satisfaction,
315 and residents' risk of hospital use. The effects were very mild and controversial. Our findings may
316 guide future research. We showed that the pedagogical and theoretical backgrounds of training
317 interventions were rarely clarified. It might be beneficial to investigate how to best motivate cultural
318 change and ownership of development in end-of-life care to be continuous processes within facilities.

319 *Conflicts of interest:*

320 We wish to confirm that the authors have no conflicts of interest associated with this publication.

321

322

323 **Figure 1. Study selection flowchart**

324

References

1. Morley JE, Caplan G, Cesari M, et al. International Survey of Nursing Home Research Priorities. *J Am Med Dir Assoc*. 2014;15(5):309–312. doi:10.1016/j.jamda.2014.03.003
2. Rolland Y, Aquino J-P, Andrieu S, et al. IAGG Workshop: Identification of the main domains for quality of care and clinical research in nursing homes. *J Nutr Health Aging*. 2011;15(5):410-424. doi:10.1007/s12603-011-0091-8
3. Briggs R, Robinson S, Martin F, O'Neill D. Standards of medical care for nursing home residents in Europe. *Eur Geriatr Med*. 2012;3(6):365-367. doi:10.1016/j.eurger.2012.07.455
4. Ouslander JG, Lamb G, Tappen R, et al. Interventions to Reduce Hospitalizations from Nursing Homes: Evaluation of the INTERACT II Collaborative Quality Improvement Project. *J Am Geriatr Soc*. 2011;59(4):745–753. doi:10.1111/j.1532-5415.2011.03333.x
5. Laging B, Ford R, Bauer M, Nay R. A meta-synthesis of factors influencing nursing home staff decisions to transfer residents to hospital. *J Adv Nurs*. 2015;71(10):2224–2236. doi:10.1111/jan.12652 [doi]
6. Flo E, Husebo BS, Bruusgaard P, et al. A review of the implementation and research strategies of advance care planning in nursing homes. *BMC Geriatr*. 2016;16:4. doi:10.1186/s12877-016-0179-4 [doi]
7. Gonella S, Basso I, Dimonte V, et al. Association Between End-of-Life Conversations in Nursing Homes and End-of-Life Care Outcomes: A Systematic Review and Meta-analysis. *J Am Med Dir Assoc*. 2019;20(3):249-261. doi:10.1016/j.jamda.2018.10.001
8. Dwyer R, Gabbe B, Stoelwinder JU, Lowthian J. A systematic review of outcomes following emergency transfer to hospital for residents of aged care facilities. *Age Ageing*. 2014;43(6):759-766. doi:10.1093/ageing/afu117
9. Lemoyne SE, Herbots HH, De Blick D, Remmen R, Monsieurs KG, Van Bogaert P. Appropriateness of transferring nursing home residents to emergency departments: a systematic review. *BMC Geriatr*. 2019;19(1):17. doi:10.1186/s12877-019-1028-z
10. Hall S, Kolliakou A, Petkova H, Froggatt K, Higginson IJ. Interventions for improving palliative care for older people living in nursing care homes. *Cochrane Database Syst Rev*. 2011;2011(3). doi:10.1002/14651858.CD007132.pub2
11. Martin RS, Hayes B, Gregorevic K, Lim WK. The Effects of Advance Care Planning Interventions on Nursing Home Residents: A Systematic Review. *J Am Med Dir Assoc*. 2016;17(4):284-293. doi:10.1016/j.jamda.2015.12.017
12. Murphy E, Froggatt K, Connolly S, et al. Palliative care interventions in advanced dementia. *Cochrane Database Syst Rev*. 2016;2016(12). doi:10.1002/14651858.CD011513.pub2
13. Low L-F, Fletcher J, Goodenough B, et al. A Systematic Review of Interventions to Change Staff Care Practices in Order to Improve Resident Outcomes in Nursing Homes. *PLOS ONE*. 2015;10(11):e0140711. doi:10.1371/journal.pone.0140711
14. Anstey S, Powell T, Coles B, Hale R, Gould D. Education and training to enhance end-of-life care for nursing home staff: a systematic literature review. *BMJ Support Palliat Care*. 2016;6(3):353–361. doi:10.1136/bmjspcare-2015-000956

15. Kearns T, Cornally N, Molloy W. Patient reported outcome measures of quality of end-of-life care: A systematic review. *Maturitas*. 2017;96:16–25. doi:10.1016/j.maturitas.2016.11.004
16. Kavalieratos D, Corbelli J, Zhang D, et al. Association Between Palliative Care and Patient and Caregiver Outcomes: A Systematic Review and Meta-analysis. *Jama*. 2016;316(20):2104–2114. doi:10.1001/jama.2016.16840 [doi]
17. van Soest-Poortvliet MC, van der Steen JT, Zimmerman S, et al. Selecting the best instruments to measure quality of end-of-life care and quality of dying in long term care. *J Am Med Dir Assoc*. 2013;14(3):179-186. doi:10.1016/j.jamda.2012.09.019
18. Konetzka RT, Spector W, Limcangco MR. Reducing Hospitalizations From Long-Term Care Settings. *Med Care Res Rev*. 2008;65(1):40-66. doi:10.1177/1077558707307569
19. Robinson L, Dickinson C, Rousseau N, et al. A systematic review of the effectiveness of advance care planning interventions for people with cognitive impairment and dementia. *Age Ageing*. 2012;41(2):263–269. doi:10.1093/ageing/afr148
20. Houben CH, Spruit MA, Wouters EF, Janssen DJ. A randomised controlled trial on the efficacy of advance care planning on the quality of end-of-life care and communication in patients with COPD: the research protocol. *BMJ Open*. 2014;4(1):4465. doi:10.1136/bmjopen-2013-004465 [doi]
21. Phillips JL, West PA, Davidson PatriciaM, Agar M. Does case conferencing for people with advanced dementia living in nursing homes improve care outcomes: Evidence from an integrative review? *Int J Nurs Stud*. 2013;50(8):1122-1135. doi:10.1016/j.ijnurstu.2012.11.001
22. Graverholt B, Forsetlund L, Jamtvedt G. Reducing hospital admissions from nursing homes: a systematic review. *BMC Health Serv Res*. 2014;14:36. doi:10.1186/1472-6963-14-36
23. Chung HO, Oczkowski SJ, Hanvey L, Mbuagbaw L, You JJ. Educational interventions to train healthcare professionals in end-of-life communication: a systematic review and meta-analysis. *BMC Med Educ*. 2016;16:x. doi:10.1186/s12909-016-0653-x [doi]
24. Higgins J, Altman D, Sterne J, eds. Chapter 8: Assessing risk of bias in included studies. In: *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (Updated March 2011)*. Vol 2011. ; 2011. <http://www.handbook.cochrane.org>
25. Verhagen AP, de Vet HC, de Bie RA, et al. The Delphi list: a criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *J Clin Epidemiol*. 1998;51(12):1235-1241. doi:10.1016/s0895-4356(98)00131-0
26. Guyatt GH, Sackett DL, Cook DJ. Users' guides to the medical literature. II. How to use an article about therapy or prevention. A. Are the results of the study valid? Evidence-Based Medicine Working Group. *JAMA*. 1993;270(21):2598-2601. doi:10.1001/jama.270.21.2598
27. Linn MW, Linn BS, Stein S, Stein EM. Effect of nursing home staff training on quality of patient survival. *J Aging*. 1989;28(4):305-315.
28. Linn MW, Linn BS, Stein S. Impact on nursing home staff of training about death and dying. *JAMA*. 1983;250(17):2332-2335.
29. Doherty K, Stein S, Linn MW. An in-Service Guide for Death Education. *Gerontol Geriatr Educ*. 1981;2(3):191-198. doi:10.1300/J021v02n03_05

30. Kovach CR, Wilson SA, Noonan PE. The effects of hospice interventions on behaviors, discomfort, and physical complications of end stage dementia nursing home residents. *Am J Alzheimers Dis.* 1996;11(4):7-15. doi:10.1177/153331759601100402
31. Molloy DW, Guyatt GH, Russo R, et al. Systematic Implementation of an Advance Directive Program in Nursing Homes: A Randomized Controlled Trial. *JAMA.* 2000;283(11):1437–1444. doi:10.1001/jama.283.11.1437
32. Boyd M, Armstrong D, Parker J, et al. Do Gerontology Nurse Specialists Make a Difference in Hospitalization of Long-Term Care Residents? Results of a Randomized Comparison Trial. *J Am Geriatr Soc.* 2014;62(10):1962-1967. doi:10.1111/jgs.13022
33. Connolly MJ, Boyd MNP, Broad JBBA, et al. The Aged Residential Care Healthcare Utilization Study (ARCHUS): A Multidisciplinary, Cluster Randomized Controlled Trial Designed to Reduce Acute Avoidable Hospitalizations From Long-Term Care Facilities. *J Am Med Dir Assoc.* 2015;16(1):49–55. doi:10.1016/j.jamda.2014.07.008
34. Foster SJ, Boyd M, Broad JB, et al. Aged Residential Care Health Utilisation Study (ARCHUS): a randomised controlled trial to reduce acute hospitalisations from residential aged care. *BMC Geriatr.* 2012;12(1):54. doi:10.1186/1471-2318-12-54
35. Agar M, Luckett T, Luscombe G, et al. Effects of facilitated family case conferencing for advanced dementia: A cluster randomised clinical trial. *PLOS ONE.* 2017;12(8):e0181020. doi:10.1371/journal.pone.0181020
36. Agar M, Beattie E, Luckett T, et al. Pragmatic cluster randomised controlled trial of facilitated family case conferencing compared with usual care for improving end of life care and outcomes in nursing home residents with advanced dementia and their families: the IDEAL study protocol. *BMC Palliat Care.* 2015;14:63.
37. Kane RL, Huckfeldt P, Tappen R, et al. Effects of an Intervention to Reduce Hospitalizations From Nursing Homes. *JAMA Intern Med.* 2017;177(9):1257. doi:10.1001/jamainternmed.2017.2657
38. Hanson LC, Zimmerman S, Song M-K, et al. Effect of the Goals of Care Intervention for Advanced Dementia. *JAMA Intern Med.* 2017;177(1):24-31. doi:10.1001/jamainternmed.2016.7031
39. Einterz SF, Gilliam R, Lin FC, McBride JM, Hanson LC. Development and Testing of a Decision Aid on Goals of Care for Advanced Dementia. *J Am Med Dir Assoc.* 2014;15(4):251-255. doi:10.1016/j.jamda.2013.11.020
40. Boogaard JA, de Vet HC, van Soest-Poortvliet MC, Anema JR, Achterberg WP, van der Steen JT. Effects of two feedback interventions on end-of-life outcomes in nursing home residents with dementia: A cluster-randomized controlled three-armed trial. *Palliat Med.* 2018;32(3):693-702. doi:10.1177/0269216317750071
41. Boogaard JA, van Soest-Poortvliet MC, Anema JR, et al. Feedback on end-of-life care in dementia: the study protocol of the FOLLOW-up project. *BMC Palliat Care.* 2013;12(1):29. doi:10.1186/1472-684X-12-29
42. Arendts G, Deans P, O'Brien K, et al. A clinical trial of nurse practitioner care in residential aged care facilities. *Arch Gerontol Geriatr.* 2018;77:129-132. doi:10.1016/j.archger.2018.05.001

43. Arendts G, Etherton-Beer C, Howard K, et al. Nurse led care coordination: Trial protocol and development of a best practice resource guide for a cluster controlled clinical trial in Australian aged care facilities. *Arch Gerontol Geriatr*. 2014;58(1):15-19. doi:10.1016/j.archger.2013.07.004
44. Verreault R, Arcand M, Misson L, et al. Quasi-experimental evaluation of a multifaceted intervention to improve quality of end-of-life care and quality of dying for patients with advanced dementia in long-term care institutions. *Palliat Med*. 2018;32(3):613-621. doi:10.1177/0269216317719588
45. Temkin-Greener H, Mukamel DB, Ladd H, et al. Impact of Nursing Home Palliative Care Teams on End-of-Life Outcomes. *Med Care*. 2018;56(1):11–18. doi:10.1097/MLR.0000000000000835
46. Temkin-Greener H, Ladwig S, Ye Z, Norton SA, Mukamel DB. Improving palliative care through teamwork (IMPACTT) in nursing homes: Study design and baseline findings. *Contemp Clin Trials*. 2017;56:1-8. doi:10.1016/j.cct.2017.01.011
47. Husebø BS, Ballard C, Aarsland D, et al. The Effect of a Multicomponent Intervention on Quality of Life in Residents of Nursing Homes: A Randomized Controlled Trial (COSMOS). *J Am Med Dir Assoc*. 2019;20(3):330-339. doi:10.1016/j.jamda.2018.11.006
48. Husebo BS, Flo E, Aarsland D, et al. COSMOS—improving the quality of life in nursing home patients: protocol for an effectiveness-implementation cluster randomized clinical hybrid trial. *Implement Sci*. 2015;10(1):131. doi:10.1186/s13012-015-0310-5
49. Saevar Reid TJL, Thoresen L, Gjerberg E, Lillemoen L, Pedersen R. Improved patient participation through advance care planning in nursing homes-A cluster randomized clinical trial. *Patient Educ Couns*. 2019;102(12):2183-2191. doi:10.1016/j.pec.2019.06.001
50. Saevar Reid TJL, Lillemoen L, Thoresen L, Forde R, Gjerberg E, Pedersen R. Implementing advance care planning in nursing homes - study protocol of a cluster-randomized clinical trial. *BMC Geriatr*. 2018;18(1):180. doi:10.1186/s12877-018-0869-1
51. Van den Block L, Honinx E, Pivodic L, et al. Evaluation of a Palliative Care Program for Nursing Homes in 7 Countries: The PACE Cluster-Randomized Clinical Trial. *JAMA Intern Med*. Published online November 11, 2019;1-10. doi:10.1001/jamainternmed.2019.5349
52. Smets T, Onwuteaka-Philipsen BBD, Miranda R, et al. Integrating palliative care in long-term care facilities across Europe (PACE): protocol of a cluster randomized controlled trial of the 'PACE Steps to Success' intervention in seven countries. *BMC Palliat Care*. 2018;17. doi:10.1186/s12904-018-0297-1
53. Liu W, Koerner J, Lam L, et al. Improved Quality of Death and Dying in Care Homes: A Palliative Care Stepped Wedge Randomized Control Trial in Australia. *J Am Geriatr Soc*. Published online November 4, 2019;jgs.16192. doi:10.1111/jgs.16192
54. Forbat L, Liu W-M, Koerner J, et al. Reducing time in acute hospitals: A stepped-wedge randomised control trial of a specialist palliative care intervention in residential care homes. *Palliat Med*. Published online January 2, 2020;0269216319891077. doi:10.1177/0269216319891077
55. Forbat L, Chapman M, Lovell C, Liu W-M, Johnston N. Improving specialist palliative care in residential care for older people: a checklist to guide practice. *BMJ Support Palliat Care*. 2018;8(3):347-353. doi:10.1136/bmjspcare-2017-001332

56. Ettema TP, Dröes R-M, de Lange J, Mellenbergh GJ, Ribbe MW. QUALIDEM: development and evaluation of a dementia specific quality of life instrument. Scalability, reliability and internal structure. *Int J Geriatr Psychiatry*. 2007;22(6):549-556. doi:10.1002/gps.1713
57. Weiner MF, Martin-Cook K, Svetlik DA, Saine K, Foster B, Fontaine CS. The quality of life in late-stage dementia (QUALID) scale. *J Am Med Dir Assoc*. 2000;1(3):114-116.
58. Naglie G, Tomlinson G, Tansey C, et al. Utility-based Quality of Life measures in Alzheimer's disease. *Qual Life Res Int J Qual Life Asp Treat Care Rehabil*. 2006;15(4):631-643. doi:10.1007/s11136-005-4364-8
59. Horsman J, Furlong W, Feeny D, Torrance G. The Health Utilities Index (HUI®): concepts, measurement properties and applications. *Health Qual Life Outcomes*. 2003;1:54. doi:10.1186/1477-7525-1-54
60. Volicer L, Hurley AC, Blasi ZV. Scales for evaluation of End-of-Life Care in Dementia. *Alzheimer Dis Assoc Disord*. 2001;15(4):194–200. doi:10.1097/00002093-200110000-00005
61. Munn JC, Zimmerman S, Hanson LC, et al. Measuring the Quality of Dying in Long-Term Care. *J Am Geriatr Soc*. 55(9):1371-1379.
62. Downey L, Curtis JR, Lafferty WE, Herting JR, Engelberg RA. The Quality of Dying and Death (QODD) Questionnaire. *J Pain Symptom Manage*. 2010;39(1):9. doi:10.1016/j.jpainsymman.2009.05.012
63. Vohra JU, Brazil K, Hanna S, Abelson J. Family Perceptions of End-of-Life Care in long-term care facilities. *J Palliat Care*. 2004;20(4):297–302.
64. Guyatt GH, Mitchell A, Molloy DW, Capretta R, Horsman J, Griffith L. Measuring patient and relative satisfaction with level or aggressiveness of care and involvement in care decisions in the context of life threatening illness. *J Clin Epidemiol*. 1995;48(10):1215-1224. doi:10.1016/0895-4356(95)00024-X
65. Hurley AC, Volicer BJ, Hanrahan PA, Houde S, Volicer L. Assessment of discomfort in advanced Alzheimer patients. *Res Nurs Health*. 1992;15(5):369-377. doi:10.1002/nur.4770150506
66. Siebenhofer A, Paulitsch MA, Pregartner G, et al. Cluster-randomized controlled trials evaluating complex interventions in general practices are mostly ineffective: a systematic review. *J Clin Epidemiol*. 2018;94:85-96. doi:10.1016/j.jclinepi.2017.10.010
67. Knowles MS. *The Modern Practice of Adult Education : From Pedagogy to Andragogy*. Association Press; 1980.
68. Kolb DA. *Experiential Learning : Experience as the Source of Learning and Development*. Prentice-Hall; 1984.
69. Dolmans DHJM, Grave WD, Wolfhagen IHAP, van der Vleuten CPM. Problem-based learning: future challenges for educational practice and research. *Med Educ*. 2005;39(7):732–741. doi:10.1111/j.1365-2929.2005.02205.x
70. Schön DA. *Educating the Reflective Practitioner: Toward a New Design for Teaching and Learning in the Professions*. Jossey-Bass; 1987:xvii, 355.
71. Huckfeldt PJ, Kane RL, Yang Z, et al. Degree of Implementation of the Interventions to Reduce Acute Care Transfers (INTERACT) Quality Improvement Program Associated with Number of Hospitalizations. *J Am Geriatr Soc*. 2018;66(9):1830-1837. doi:10.1111/jgs.15476

72. Ouslander JG, Perloe M, Givens JH, Kluge L, Rutland T, Lamb G. Reducing Potentially Avoidable Hospitalizations of Nursing Home Residents: Results of a Pilot Quality Improvement Project. *J Am Med Dir Assoc*. 2009;10(9):644–652. doi:10.1016/j.jamda.2009.07.001

Figure 2. Study selection flowchart

Table 1. Evaluation of quality criteria fulfillment in randomized controlled trials examining effects of training on nursing home residents' end-of-life care

*Criteria: (1) The inclusion and exclusion criteria are satisfactorily described. (2) Groups are comparable at baseline. (3) The study has sufficient statistical power to detect an effect ($N \geq 25/\text{group}$) or an adequate power calculation is presented. (4) The randomization method is valid (a computerized randomization program or a separate randomization center) and adequately described. (5) The intervention is adequately described. (6) The measurements and outcome measures are valid and well defined. (7) The group allocation is blinded when assessing the outcomes. (8) The dropouts are described and the analyses take them into account. (9) Intention-to-treat analysis is applied. (10) Appropriate statistical analyses are used (a comparison is made in relation to outcome variables between the groups).

Study	Criteria*										Total
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	
Linn et al. 1989 ²⁷	+	–	+	–	?	+	+	–	–	–	3
Kovach et al. 1996 ³⁰	+	+/-	+	+/-	+	+	–	–	–	–	4
Molloy et al. 2000 ³¹	+	+/-	+	+/-	+	+	–	+	+	+	7
Boyd et al. 2014 ³²	+/-	+/-	+	+	+	+	–	–	–	+	4
Connolly et al. 2015 ³³	+	+/-	+	+	+	+	+	+	+	+	9
Agar et al. 2017 ³⁵	+	+	+	+	+	+	+	+/-	+/-	+	8
Hanson et al. 2017 ³⁸	+	+	+	+	+	+	+	+	+	+	10
Kane et al. 2017 ³⁷	+	+	+	+	+	+	+/-	+/-	+	+	8
Arendts et al. 2018 ⁴²	+	+	+	+/-	+	+	+	+/-	?	+	7
Boogaard et al. 2018 ⁴⁰	+	+	+	+/-	+	+	–	–	–	+/-	5
Temkin-Greener et al. 2018 ⁴⁵	+/-	+/-	+	+	+	+	?	+	?	+	7
Verreault et al. 2018 ⁴⁴	+	+/-	+	–	+	+	–	–	–	–	4
Husebø et al. 2019 ⁴⁷	+	+	+	+	+	+	+	+	+	+	10
Sævareid et al. 2019 ⁴⁹	+	+	+	+/-	+	+/-	+/-	+	+	+	7
Van den Block et al. 2019 ⁵¹	+	+	+	+	+	+	+/-	+	+/-	+	8
Liu et al. 2020 ⁵³	+	+	+	+	+	+	–	+/-	–	+	7

High quality 8–10, moderate quality 5–7, low quality 0–4

Table 2. Characteristics of studies

Study, year, country	Design	Participants, age, mean age, females (F) %	Intervention	Control	Outcomes (primary outcome as stated by original study)	Comments	Methodological quality
Linn et al., 1989, USA ²⁷⁻²⁹	cRCT / matched control	10 NHs, N=306, 2/3 cancer diagnosis, age NA, F=NA	Training covered feelings about death, lack of preparation to deal with dying, palliative vs curative care, common fears, and feelings of dying patients and their families.	Usual care	QOL: Patients in trained NHs had less depression (POMS) and greater satisfaction with care. Follow-up 3 mo. Primary outcome not defined.	Only 201 patients were analyzed. Staff turnover was 50%/year.	Low 4
Kovach et al., 1996, USA ³⁰	RCT	3 LTCFs, N=72, severe dementia, mean age 88y, F=81%	End-stage dementia units were formed. More homelike environment, more activities, and encouragement to treat behavioral symptoms with painkillers. Use of hospice nurse as case manager. Staff had one-day training for hospice.	Usual care	Behavioral symptoms (BEHAVE-AD) did not differ between the groups post-test. QOD: There was a significant difference in discomfort (DS-DAT) between the groups post-test. No difference in physical complications. Primary outcome not defined.	Only 2-mo follow-up. No information on BEHAVE-AD or DS-DAT at baseline. No blinding.	Low 4
Molloy et al., 2000, Canada "Let me decide" ³¹	cRCT	Six NHs, N=1292, 79-85y, F=72%	Implementation of "Let me decide" AD program. Three registered nurses from I NHs received 2d training. They trained staff, residents, proxies, and hospital staff about the program in several sessions and helped residents and proxies to complete ADs.	Usual care	HSP: Hospitalizations per resident were lower in 18-mo follow-up (0.27 vs 0.48; p<0.001). Total health care costs were lower in the intervention homes (CAD 3490 vs CAD 5239, p=0.01). SWC: Change in satisfaction with care or mortality did not differ between the two groups. Satisfaction with health care utilization (primary outcome) did not differ.	In service use all residents were included. Unclear how nonconsenting participants were otherwise handled in analyses.	Moderate 7
Boyd et al., 2014, New Zealand, "RACIP" ³²	cRCT	54 RACFs with 2553 beds, about 85y, F=NA	I NHs were trained and coached to avoid hospitalizations, 5 to 6 standardized educational sessions in each facility each year, clinical support and care coordination for high-risk residents.	Usual care	HSP: Hospitalization rates increased in both groups, in the intervention group significantly less. Hospitalizations for medical (non-surgical) causes showed the greatest difference between the groups. Primary outcome not defined.	Unclear how comprehensively residents' data could be used.	Low 4
Connolly et al., 2015, New Zealand, "ARCHUS" ^{3,34}	cRCT	36 RACFs, N=1998, 96% >65y; F=70%	Three 1h monthly meetings with cases discussed. Gerontology education including e.g., ACP, nutrition/hydration, end-stage dementia care, communication with families.	Usual care	HSP: All hospitalizations (0.68 vs 0.67), avoidable hospitalizations (primary outcome) (0.57 vs 0.53), and days spent in hospital (4.18 vs 4.41) were similar in I and C. Time to first hospital admission, time to death and mortality were similar.	All residents were included if the facility participated.	High 9

Study, year, country	Design	Participants, age, mean age, females (F) %	Intervention	Control	Outcomes (primary outcome as stated by original study)	Comments	Methodological quality
Agar et al., 2017, Australia “IDEAL” ^{35,36}	cRCT	20 NHs, N=286, all had advanced dementia, mean age 85y, F=60%	Facilitated case conferencing with families to promote ACP. RN was trained in each unit to organize case conference sessions. Staff were trained in person-centered palliative care and to develop ACP.	Usual care	In I NHs symptoms were better documented. QOD: No group differences in family-(primary outcome) or nurse-rated QOD (SM-EOLD, CAD-EOLD, SWC-EOLD) QOL: Nurse-rated QUALID did not differ. HSP: No difference.	Only 76% of deceased residents had evaluations using EOLD scale.	High 8
Hanson et al., 2017, USA “Goals of Care” ^{38,39}	cRCT	22 NHs, N=302, advanced dementia, mean age 86–87y, F=82%	Goals of care intervention (video and decision aid printed material) to patients’ surrogate decision makers to improve palliative care in people with advanced dementia. 1h training for staff in communication and discussion guide.	Usual care	Family decision makers reported better quality of communication scores (primary outcome) overall and regarding EOL care in I. Treatment consistent with preferences (primary outcome) did not differ. HSP: Hospital transfers were significantly lower in I than in C. QOD: No difference in SWC-EOLD or SM-EOLD scores or survival.	Intervention was mainly targeted to family decision makers to prepare for ACP. Only 1h staff training.	High 10
Kane et al., 2017, USA “INTERACT” ³⁷	cRCT	85 NHs, N=23 478, mean age 80–82y, F=66–67%	Each I NH selected “champion” and “co-champion” to facilitate INTERACT training and implementation of tool kit (>30 tools) to address the key factors leading to avoidable hospital admissions and ED visits among NH residents.	1. pure control 2. attention control	HSP: No reductions in hospitalization rates or ED visits without admission (primary outcomes). Intervention NHs exhibited a reduction in potentially avoidable hospitalizations (primary outcome), but this effect was not robust to a Bonferroni correction for multiple comparisons.	Well-designed long-term quality improvement project with very large number of NHs and participants.	High 8
Arendts et al., 2018, Australia ^{42,43}	cRCT	6 RACFs, N=200, mean age 89y, F=76%	Nurse practitioners acted as care managers for residents. They coordinated care of chronic and acute conditions and arranged acute care if needed. They promoted use of “best practice resource folder” and organized staff training.	Usual care	HSP: No significant difference in ED transfers (primary outcome) QOL measured by EQ-5D: no significant difference between the groups QOL by HUI: I group decreased less in QOL during follow-up.	Nurse practitioner had major role and education minor role in intervention.	Moderate 7
Boogaard et al., 2018, the Netherlands ^{40,41}	cRCT	18 NHs, N=423, mean age 86y, F=69%	SWC-EOLD and CAD-EOLD scale family ratings on quality of EOL care and QOD were used to give feedback to NH staff: I1: generic feedback, I2: patient-specific feedback	Usual care	QOD: Generic strategy resulted in lower quality of EOL care (primary outcome) compared with C group. Patient-specific strategy did not affect the quality of EOL care but increased comfort in unadjusted analyses.	High number of dropouts. Mild intervention. The primary outcome of trial was the same as feedback tool.	Moderate 5

Study, year, country	Design	Participants, age, mean age, females (F) %	Intervention	Control	Outcomes (primary outcome as stated by original study)	Comments	Methodological quality
Temkin-Greener et al., 2018 USA, "IMPACTT" ^{45,46}	cRCT	25 NHs, N=5830 decedents, mean age 86y, F=74%	Team development and palliative care training (14 sessions available in both). Online EOL care materials available. Nurse practitioner interventionist accompanied the teams on the rounds for 2 mo.	Usual care	QOL: No effect on depression, pain, or being deceased in hospital nor in number of hospital stays during last 90d (primary outcomes).	Only deceased were included. Massive efforts to train staff in 14 NHs.	Moderate 5
Verreault et al., 2018, Canada ⁴⁴	cRCT, quasi-experimental	4 NHs, N=193, terminal dementia, mean age at death 89y, F=73%	Training staff in PC: physicians (3h), nurses (7h), nurse aids (3.5h). Clinical monitoring of pain. Early and systematic communication with families about EOL care issues. Use of a nurse facilitator to implement and monitor the intervention.	Usual care	QOD: Family Perception of Care Score was significantly higher in the intervention group than in the usual care group (157.3 vs 149.1; p=0.04) (primary outcome).. The Comfort Assessment and Symptom Management scores were also significantly higher in the intervention group (primary outcome).	The design and randomization of study wards are unclear. Only deceased patients were analyzed.	Low 4
Husebø et al., 2019, Norway "COSMOS" ^{47,48}	cRCT	33 NHs with 67 units, N=545, 94% dementia, mean age 87y, F=74%	Staff training and assistance in implementing a multicomponent – communication (including advance planning), systematic pain management, medication review, organization of activities – intervention.	Usual care	QOL: After decline in QOL in intervention group at 4 mo, there was a significant improvement between 4 and 9 mo (QUALIDEM, EQ-VAS). The difference in changes between two groups from baseline to 9 mo was not significant in QUALID, QUALIDEM or EQ-VAS (primary outcomes).	Drugs were deprescribed more, the ADL function was better sustained, and staff distress improved in the I group.	High 10
Sævareid et al., 2019, Norway ^{49,50}	cRCT	8 wards, N=154, mean age 87y, F=70%	Training of project teams that trained other facility staff (train-the-trainer) on promoting and communicating about ACP and written templates to make advance care plans.	Usual care	More advance plans and discussions (primary outcome) in I. I residents had better concordance of received treatment with treatment preferences. HSP: no difference.	Most of the outcomes were based on unvalidated questionnaire items.	Moderate 7
Van den Block et al., 2019, EU, "PACE" ^{51,52}	cRCT	78 NHs in 7 countries, decedents N=551/984 pre/post, mean age 86y, F=65%	Implementation of PACE – a six step staff training program by facility and country coordinators. Steps: (1) ACP; (2) resident assessment; (3) monthly multidisciplinary palliative care meetings; (4) improving pain and depression management; (5) care in last days of life; (6) care after death.	Usual care	QOD: CAD-EOLD: no difference between groups reported by staff or proxy (primary outcome), SWC-EOLD: no difference reported by proxy, QOD-LTC significantly better scores in I than in C (staff reported). No clinically significant improvement in staff knowledge (primary outcome).	No outcomes related to residents who remained alive. Very low response rate in some countries.	High 8

Study, year, country	Design	Participants, age, mean age, females (F) %	Intervention	Control	Outcomes (primary outcome as stated by original study)	Comments	Methodological quality
Liu et al., 2020, Forbat et al., 2020, Australia ^{53–55}	cRCT (stepped-wedge design)	12 care homes with 1700 residents, 537 deaths overall, 471 analyzed, 30% dementia, mean age 85y, F=64%	Monthly 1h “palliative care needs rounds”, case conferencing, and case-based training of staff focused on anticipatory planning.	Usual care, all I wards served first as controls	QOD: reported by staff: QODD (range 0–160) scores increased by 8.1 points during intervention in all facilities. HSP: Hospitalizations and hospital days (primary outcome) decreased in I vs C.	Only deceased residents were analyzed. 88% of deceased residents had complete data. No ITT analysis or outcomes related to living residents.	Moderate 7

ACP = advance care planning, AD = advance directive, ADL = activities of daily living, BEHAVE-AD = Behavioral Pathology in Alzheimer’s Disease, C = control arm, CAD = Canadian dollar, CAD-EOLD = Comfort Assessment in Dying End-Of-Life in Dementia scale, (c)RCT = (cluster) randomized controlled trial, d = days, DS-DAT = Discomfort Scale for Dementia of the Alzheimer’s Type, ED = emergency department, EOL = end-of-life, EOLD = End-Of-Life in Dementia scale, EQ-5D = EuroQoL 5D, EQ-VAS = EuroQoL Visual Analogue Scale, FPCS = Family Perception of Care Score, h = hours, HSP = hospitalizations, HUI = Health Utilities Index, I = intervention arm, ITT = intention-to-treat, LTCF = long-term care facility, mo = months, NA = not applicable, NH = nursing home, PACE = Palliative Care for Older People, PC = palliative care, PCPC = palliative care planning coordinator, POMS = Psychiatric Outpatient Mood Scale, QOD = quality of dying, QODD = quality of death and dying, QOD-LTC = QOD in long-term care, QOL = quality of life, QUALID = Quality of Life in Late-Stage Dementia, QUALIDEM = a dementia-specific QOL assessment instrument, RACF = residential aged care facility, RN = registered nurse, SM-EOLD = Symptom Management EOLD scale, SWC = satisfaction with care, SWC-EOLD = Satisfaction With Care EOLD scale, y = years

Table 3. Training characteristics and outcome effects of high- and moderate-quality trials

	Resident characteristics			Outcome effects				Characteristics of training intervention				
Study, year, country	Mean age, years	Dem-tia %	Deceased % / follow-up time	QOL	HSP	QOD	SWC	Learn-ing methods: AI, C, CBL, D/S, e-L, F, L, RP, TT, V, WS	Major contents, intensity, and length	Complexity of intervention	Role of training in intervention	Other intervention components
High-quality trials												
Connolly et al., 2015, New Zealand ^{33,34}	50% ≥75y	na	21%/ 14 mo	na	NS	na	na	C, CBL, D/S, F, L(?)	GNS visited and coached nurses and caregivers in ACP. Meetings: 3×1h/month (3 mo); gerontology training: monitoring and benchmarking residents' QOC and giving feedback ×3 over 9 mo.	Moderate	Major	Facility-level assessment of areas of needs and care plans created from baseline data by researchers, GNS, and facility leaders.
Agar et al., 2017, Australia ^{35,36}	85y	100 %	100%*/ 18 mo	NS	NS	NS	NS	C, CBL, TT	Registered nurse was trained 35h to be a PCPC in each NH. PCPC worked 2d in each NH; organized training in person-centered care and case conferencing for 18 mo for staff and families.	Moderate	Major	The PCPC identified residents nearing EOL and facilitated implementation of palliative care plans.
Hanson et al., 2017, USA ^{38,39}	86y	100 %	20%/ 9 mo	na	+	NS	NS	Video for proxies, L(?) for staff	Videos for family members/proxies to aid ACP decisions. 1h training for staff members on the structured goals of care discussions (ACP) with proxies. Retraining until fidelity.	Simple	Minor	2-part intervention, 1 st part was a video and print copy of decision aid for proxies. Staff training had minor role.
Kane et al., 2017, USA ³⁷	80–82y	na	na/ 26 mo	na	NS	na	na	C, e-L, F, TT	Two facilitators trained from each facility to be a “champion” and to train other staff and implement INTERACT principles. Initial intensive training program for 10wk and follow-up webinars monthly. Webinars for all staff. Study team had monthly phone calls with NH champions.	Complex	Moderate	Champions promoted complex INTERACT tool kit (30 clinical and educational tools) in their NH. Quality improvement project.

Study, year, country	Resident characteristics			Outcome effects				Characteristics of training intervention				
	Mean age, years	Deceased %	Deceased % / follow-up time	QOL	HSP	QOD	SWC	Learning methods: AI, C, CBL, D/S, e-L, F, L, RP, TT, V, WS	Major contents, intensity, and length	Complexity of intervention	Role of training in intervention	Other intervention components
High-quality trials												
Husebø et al., 2019, Norway ^{47,48}	87y	94%	21%/ 9 mo	NS	na	na	na	D/S, TT, L, RP	2d seminar for staff. At least 2 nurses from each unit (“ambassadors”) were trained in all 4 domains of intervention. Main contents were ACP and pain management. Ambassadors organized lunch meetings several times per week for staff over 4 mo.	Moderate	Major	Medication review and organization of weekly activities for residents.
Van den Block et al., 2019, EU	86y	67–72%	100%*/ 17 mo	na	na	NS	NS	C, TT, WS	Trainers of each country were trained in 1wk international WS. They trained and supported NH facilitators and staff in implementing monthly PACE steps over 6 mo. Intensity of actual staff training unclear.	Complex	Major	Country trainers visited /contacted NHs every 7–10d. ACP promoted. Mainly EOL topics, also general pain and depression management.
Moderate-quality trials												
Molloy et al., 2000, Canada	79–85y	~62%	26%/ 18 mo	na	+	na	NS	TT, C, D/S(?), V, WS	2d workshops to 3 facilitators who trained and coached the staff, residents, and proxies to complete ADs. TT approach for staff training, and in-service training, intensity unclear.	Moderate	Moderate	Focus on completing ADs.
Arendts et al., 2018, Australia ^{42,43}	90y	?	? / 12–32 mo	NS	NS	na	na	C, L(?), TT	Nurse practitioners acted as care coordinators to residents and provided staff training.	Moderate	Minor	Care coordination as major intervention, intervention team developed and provided a tool kit to be used by all staff.
Boogaard et al., 2018, the Netherlands ^{40,41}	86y	100%	100%*/ 10 mo	na	na	NS	NS	F	Feedback was given to facilities based on family ratings of SWC-EOLD and CAD-EOLD. Facilities discussed feedback but no information on how intensive/often these meetings were. Facilities either followed improvement suggestions or made their own.	Simple	Minor	Intervention offered minor support. Level of adherence not measured. Only deceased patients were taken into analyses. Primary outcome was the same as feedback tool.

Study, year, country	Resident characteristics			Outcome effects					Characteristics of training intervention			
	Mean age, years	De-mentia %	Deceased % / follow-up time	QOL	HSP	QOD	SWC	Learning methods: AI, C, CBL, D/S, e-L, F, L, RP, TT, V, WS	Major contents, intensity, and length	Complexity of intervention	Role of training in intervention	Other intervention components
Moderate-quality trials												
Temkin-Greener et al., 2018, USA ^{45,46}	86y	58%	100%*/ 12 mo	NS	NS	NS	na	AI, C, L(?), WS	NS palliative care guidelines and tools developed with staff in a Delphi process. Team development (TeamSTEPP), ELNEC, EOL care materials available, geropalliative care nurse accompanied teams on the rounds for 2 mo.	Complex	Major	Multicomponent educational intervention activating and also coaching NH teams + taking into account high staff turnover. Adherence to intervention low.
Sævareid et al., 2019, Norway ^{49,50}	87y	73%	25%/ 9 mo	na	NS	na	na	AI, D/S, RP, TT	ACP guide created in cooperation with project teams selected from each I ward. Project teams received 2d seminar and trained the rest of the staff. Aim was to guide and document ACP conversations.	Moderate	Moderate	Research team provided supervision of project teams, information to the ward staff, written information to patients and next of kin as well as template to document ACP.
Liu et al., 2020, Forbat et al., 2020, Australia ⁵³⁻⁵⁵	85y	30%	32% / 18 mo	na	+	+	na	C, CBL	External palliative care nurse rounded with the ward staff discussing selected patients and offering expert consultations when needed 1h/mo. Short briefing on how to select residents at high risk of dying for the rounds. Checklist to guide practice briefing notes were provided to staff.	Moderate	Moderate	Selected residents were discussed with facility staff and access to palliative care consultants. Case-based conferences arranged for NHs, focus on completing advance care plans.

+ = favors intervention

ACP = advance care planning, AD = advance directive, CAD-EOLD = Comfort Assessment in Dying End-Of-Life in Dementia scale, d = days, ELNEC = end-of-life nursing education, EOL = end-of-life, EOLD = End-Of-Life in Dementia scale, GNS = gerontology nurse specialist, h = hours, HSP = hospitalization or hospital days, I = intervention arm, mo = months, na = not applicable, NH = nursing home, NS = nonsignificant, PACE = Palliative Care for Older People, PCPC = palliative care planning coordinator, QOC = quality of communication, QOD = quality of dying, QOL = quality of life, SWC = satisfaction with care, SWC-EOLD = Satisfaction With Care EOLD scale, wk = week, y = years

Learning methods: AI = learners' active involvement in developing palliative care; C = coaching, CBL = case-based learning, D/S = (small group) discussions/seminars, e-L = e-learning, F = feedback, L = lectures, RP = role play, TT = train-the-trainer, V = video, WS = workshops

*Only included deceased residents