<https://helda.helsinki.fi>

High Discontinuation Rates of Peroral ASA Treatment for CRSwNP : A Real-World Multicenter Study of 171 N-ERD Patients

Laulajainen-Hongisto, Anu

2020

Laulajainen-Hongisto , A , Turpeinen , H , Vento , S , Numminen , J , Sahlman , J , Kauppi , P , Virkkula , P , Hytönen , M & Toppila-Salmi , S 2020 , ' High Discontinuation Rates of Peroral ASA Treatment for CRSwNP : A Real-World Multicenter Study of 171 N-ERD Patients ' , Journal of allergy and clinical immunology: In practice , vol. 8 , no. 10 , pp. 3565-3574 . <https://doi.org/10.1016/j.jaip.2020.06.063>

<http://hdl.handle.net/10138/336049>

<https://doi.org/10.1016/j.jaip.2020.06.063>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Journal Pre-proof

High discontinuation rates of peroral ASA treatment for CRSwNP. A real-world multi-center study of 171 N-ERD patients

A. Laulajainen-Hongisto, MD, PhD, H. Turpeinen, MD, PhD, S.I. Vento, MD, PhD, J. Numminen, MD, PhD, J. Sahlman, MD, PhD, P. Kauppi, MD, PhD, P. Virkkula, MD, PhD, M. Hytönen, MD, PhD, S. Toppila-Salmi, MD, PhD

PII: S2213-2198(20)30707-8

DOI: <https://doi.org/10.1016/j.jaip.2020.06.063>

Reference: JAIP 2987

To appear in: *The Journal of Allergy and Clinical Immunology: In Practice*

Received Date: 7 May 2020

Revised Date: 23 June 2020

Accepted Date: 29 June 2020

Please cite this article as: Laulajainen-Hongisto A, Turpeinen H, Vento S, Numminen J, Sahlman J, Kauppi P, Virkkula P, Hytönen M, Toppila-Salmi S, High discontinuation rates of peroral ASA treatment for CRSwNP. A real-world multi-center study of 171 N-ERD patients, *The Journal of Allergy and Clinical Immunology: In Practice* (2020), doi: <https://doi.org/10.1016/j.jaip.2020.06.063>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology

High discontinuation rates of peroral ASA treatment for CRSwNP. A real-world multi-center study of 171 N-ERD patients

Laulajainen-Hongisto A, MD, PhD^{1,2*}, Turpeinen H, MD, PhD¹, Vento SI, MD, PhD², Numminen J, MD, PhD³, Sahlman J, MD, PhD⁴, Kauppi P, MD, PhD¹, Virkkula P, MD, PhD², Hytönen M, MD, PhD², Toppila-Salmi S, MD, PhD^{1,5}

¹Inflammation Centre, Skin and Allergy Hospital, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

²Department of Otorhinolaryngology – Head and Neck Surgery, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

³Department of Ear and Oral diseases, Tampere University Hospital

⁴Department of Otorhinolaryngology, Kuopio University Hospital, Kuopio, Finland

⁵Medicum, Haartman Institute, University of Helsinki, Helsinki, Finland

* Corresponding author

Anu Laulajainen-Hongisto, MD PhD

Kasarmikatu 11-13

P.O.Box 263

FIN-00029 HUS

Helsinki, Finland

Tel +358 50 4271494

anu.laulajainen-hongisto@hus.fi

23

24 Laulajainen-Hongisto A: anu.laulajainen-hongisto@hus.fi; Turpeinen H: heikki.turpeinen@helsinki.fi;

25 Vento SI: seija.vento@hus.fi; Numminen J: Jura.Numminen@pshp.fi; Sahlman J:

26 Johanna.Sahlman@kuh.fi; Kauppi P: paula.kauppi@hus.fi; Virkkula P: Paula.Virkkula@hus.fi

27 Hytönen M: maija.hytonen@hus.fi; Toppila-Salmi S: sanna.k.salmi@hus.fi

28

29 **Acknowledgements:**

30 This study was supported in part by research grants from the Ahokas Foundation, the Finnish Society
31 of Allergology and Immunology, the Finnish ORL-HNS Foundation, Paulo Foundation, State funding
32 for university-level health research (TYH2018103), Tampere Tuberculosis Foundation, the Jane and
33 Aatos Erkko Foundation, the Väinö and Laina Kivi Foundation, and the Orion Research Foundation.

34

35 **Conflicts of interest:**

36 STS reports consultancies for ERT, Novartis, Sanofi Pharma, Mylan Laboratories and Roche Products
37 and a grant of GSK, outside the submitted work. ALH has received research funding from Orion
38 Research Foundation, outside the submitted work. All other authors declare no conflicts of interest.

39 Abstract word count: 231

40 Text word count: 3721

41

Abstract

Background: Non-steroidal anti-inflammatory drug (NSAID) exacerbated respiratory disease (N-ERD) consists of chronic rhinosinusitis with nasal polyposis (CRSwNP), asthma and NSAID intolerance. Acetylsalicylic acid (ASA) treatment after desensitization (ATAD) is a treatment option for uncontrolled N-ERD.

Objective: To evaluate peroral ATAD's long-term effectiveness on CRSwNP disease control.

Methods: Retrospective patient record data collection (patient characteristics, sinus surgeries prior to ATAD, ATAD, follow-up data [2019]) from 171 N-ERD patients (102 ATAD patients, 69 controls with CRSwNP+N-ERD without ATAD) undergoing tertiary hospital consultation at 2001-2017. Outcome measurements were ATAD discontinuation, revision sinus surgery, corticosteroid and antibiotic courses for airway infections 2016-2019. Associations were analyzed by survival and nonparametric methods.

Results: The ATAD group had more tissue eosinophilia, symptoms and sinus surgeries prior to ATAD than others. The ATAD discontinuation rate was 63%, independent of ATAD dose or duration, usually due to side-effects. Compared to the N-ERD group without ATAD, ATAD (mean duration 2.9 years) did not affect the revision ESS rate ($p=0.21$, by log rank test) or the number of peroral corticosteroid courses/year ($p>0.05$, by Mann-Whitney U-test) during the follow-up (mean 7.6 years) despite the dose or duration of ATAD.

Conclusion: The discontinuation rate of ATAD was high (63%), and ATAD did not affect revision sinus surgery rate, nor need of peroral corticosteroids during follow-up. However, the remaining 37% of the ATAD group did continue the treatment, indicating they may have benefited from ATAD.

64

65 **Highlights box:**

66 1. What is already known about this topic? ATAD is a treatment option for uncontrolled N-ERD.

67 2. What does this article add to our knowledge? Yet, in our population, most patients did not benefit
68 from ATAD and discontinued it.69 3. How does this study impact current management guidelines? ATAD patients must be followed and
70 their treatment response evaluated frequently.

71

72 **Key words:**73 Asthma, Aspirin desensitization, Aspirin treatment after desensitization, Chronic rhinosinusitis with
74 nasal polyposis, Non-steroidal anti-inflammatory drug-exacerbated respiratory disease

75

76 **Abbreviations:**

77 AD=Aspirin desensitization

78 AR=Allergic rhinitis

79 ASA=Acetylsalicylic acid

80 ATAD=Aspirin treatment after desensitization

81 CRS=Chronic rhinosinusitis

82 CRSwNP=Chronic rhinosinusitis with nasal polyposis

- 83 CT=Computed tomography
- 84 ENT=Ear, nose and throat
- 85 ESS = endoscopic sinus surgery
- 86 LM=Lund-Mackay
- 87 N-ERD=NSAID-exacerbated respiratory disease
- 88 NSAID=Non-steroidal anti-inflammatory drug
- 89 OCS=Oral corticosteroid(s)
- 90 QoL=Quality of Life
- 91
- 92
- 93
- 94
- 95
- 96
- 97
- 98
- 99

Introduction

Non-steroidal anti-inflammatory drug (NSAID) exacerbated respiratory disease (N-ERD), an inflammatory disease of the airways, usually includes a triad of chronic rhinosinusitis with nasal polyposis (CRSwNP), asthma, and hypersensitivity to acetylsalicylic acid (ASA) and other NSAIDs (1, 2). N-ERD is characterized by abnormalities of cyclooxygenase pathway, severe eosinophilic hyperplastic inflammation, and tissue remodeling with fibrosis both in paranasal sinuses and lower airways (1, 3, 4). The prevalence of N-ERD is estimated to be 9% in asthmatics. N-ERD is acquired, it is slightly more common in females and has a median age of onset around 30 years (5, 6).

Chronic rhinosinusitis (CRS) is a chronic inflammation of the nose and paranasal sinuses lasting for ≥ 12 weeks (7). Nasal endoscopy and computed tomography (CT) are used in CRS diagnostics (8-11). CRS affects 6-11% of the Western world's population, causing substantial costs with a severe impact on quality of life (QoL) (12-14). CRS with and without nasal polyps are considered CRS phenotypes with different etiologies and pathomechanisms, or different degrees of inflammation (7). CRSwNP is treated with saline irrigations, nasal steroids, oral steroids, and oral antimicrobials. Surgery is considered if conservative treatment is not sufficient (2, 7). CRSwNP often leads to recurrent surgeries, especially in N-ERD patients (15, 16).

A sudden worsening of pre-existing CRS symptoms is called a CRS exacerbation (7). Although no clear definition of a CRS exacerbation exists, an escalation in CRS symptoms and management (acute symptoms of paranasal sinuses, and systemic use of antibiotics or corticosteroids) is usually considered one (17-19). CRS exacerbations are associated with poor asthma control (20). Asthma exacerbation is

an acute or sub-acute worsening of symptoms and lung functions, or in some cases the initial asthma presentation (21).

Most N-ERD patients have moderate to severe asthma (2). Compared to other asthmatics, N-ERD patients more often have severe asthma, and are more likely to need high dose inhaled corticosteroid treatment or steroid bursts (3). Also the risk of uncontrolled asthma, asthma related healthcare visits, hospitalizations, and intubations are increased in N-ERD patients (3, 5).

In a patient with confirmed asthma and CRSwNP, a clear history of multiple reactions with respiratory symptoms within two hours after NSAID ingestion may be sufficient for N-ERD diagnosis (2). However, challenge tests are used to confirm NSAID hypersensitivity in unclear cases, for research purposes, or to determine the provocation dose of ASA before oral desensitization (2). Contraindications for ASA challenge, ASA desensitization (AD), and ASA treatment after desensitization (ATAD) include: history of anaphylactic reactions precipitated by NSAIDs, uncontrolled asthma (Forced expiratory volume in one second [FEV1] <70% of the predicted value), current respiratory tract infection or asthma exacerbation, renal failure, gastrointestinal bleeding, pregnancy, and current treatment with β -blocker (2).

AD followed by ATAD is considered in N-ERD patients with insufficient response to pharmacological treatment, high recurrence of nasal polyps (NP) leading to recurrent surgeries, insufficient control of asthma symptoms with standard medications, need to reduce corticosteroid dose, or in patients who need ASA or NSAID treatment (2, 22). Some studies suggest endoscopic sinus surgery (ESS) prior to ATAD (23), or surgical removal of NPs 3-4 weeks prior to AD/ATAD (1, 2, 24-26). AD can be performed as an extension of ASA challenge in outpatient settings (2, 27). According to the European Academy of Allergy and Clinical Immunology (EAACI) position paper, a patient undergoing oral AD

first receives ASA 20-40 mg, the dose is gradually increased with 90-120 minute intervals on the first day in three steps to 60-100 mg, and on the second day in three steps to 325 mg (2). AD is followed by ATAD, with an effective daily ASA maintenance dose of 300-1300 mg (2, 28, 29). Intranasal ASA has also been used in ASA challenge and AD (30, 31). According to the 2020 European Position Paper on Rhinosinusitis and Nasal Polyposis (EPOS), the most commonly used AD/ATAD protocol is the Scripps clinics oral AD protocol in which, after gradual increases, a peroral dose of ASA 625mg is administered twice daily (30, 31). According to the literature, ASA could be discontinued for 48 hours without losing the desensitization (1, 25).

There is a lack of information regarding the benefits of peroral ATAD on rhinosinusitis and asthma control, and on revision ESS rates. The purpose of this retrospective, controlled, real-world multicenter study was to evaluate the effect of peroral ATAD, and its duration, on the need of revision ESS, antibiotics, and systemic corticosteroids during the follow up. We hypothesized that peroral ATAD is well tolerated and that ATAD decreases ESS rates and use of antibiotics and oral corticosteroid (OCS) courses.

Methods

Subjects

This retrospective follow-up study was carried out at the Skin and Allergy Hospital and Department of Otorhinolaryngology of Helsinki University Hospital, and at the Departments of Otorhinolaryngology of Tampere and Kuopio University Hospitals, between 2001 and 2018. Approval was obtained from the ethical committee of Hospital Districts of Helsinki and Uusimaa and Pirkanmaa (No. 31/13/03/00/2015). This study involved 171 N-ERD subjects (102 peroral ATAD patients, 69 controls

with CRSwNP+N-ERD without ATAD) undergoing surgical and/or ATAD consultation due to CRS between 2001 and 2017. Inclusion criteria were: N-ERD (e.g. a patient-record documented history of wheeze/cough/naso-ocular symptoms after ingestion of NSAID $\pm$ positive ASA challenge test result) and nasal endoscopic signs of NPs. CRSwNP was diagnosed according to the EPOS criteria (7). In Finland, N-ERD has routinely been diagnosed only by clinical examination and anamnestic information regarding NSAID intolerance. Due to the retrospective nature of this study, ASA challenge test was thus not required as a diagnostic criterion. In the control group (N=69), an additional inclusion criterion was ESS performed within one year of the consultation visit as a sign of uncontrolled CRSwNP. Exclusion criteria were unavailable or incomplete data of ATAD. All subjects were regularly monitored by otorhinolaryngologists at university hospital due to uncontrolled CRSwNP, all used maximal conservative therapy including regular intranasal steroids, and nasal lavage. Data regarding the use of OCS and antibiotic courses for CRSwNP, prior to ATAD, were included in our analysis. However, the use of additional therapies was not included in the analysis. Asthma was diagnosed according to GINA (Global Initiative for Asthma) (21), based on typical history, and lung function tests spirometry and peak expiratory flow (PEF) findings; at least 15% improvement with bronchodilator test in spirometry [in forced expiratory flow volume in one second (FEV1) or in forced vital capacity (FVC)] and/or 20% diurnal variation in PEF monitoring or 15% bronchodilator response in PEF monitoring or positive methacholine challenge test (i.e. moderate to severe bronchial hyperresponsiveness). In the ATAD group, N-ERD diagnosis was based on a positive reaction (wheeze and/or naso-ocular symptom) after intake of ASA at the hospital. In the control group, the N-ERD diagnosis was based on any positive history of wheeze/cough or naso-ocular symptoms after intake of NSAID.

The background data (age, gender, smoking, allergic rhinitis, asthma, N-ERD, history of recurrent exacerbations and use of systemic corticosteroid courses, duration of symptoms, symptoms, previous ESS, baseline endoscopic NP score, baseline LM score of sinus CT scans) were collected from patient records. Baseline ESS was defined as ESS performed within 12 months from the sinus CT scans taken during the baseline visit of sinus surgical consultation in control group, or the last ESS performed ≤ 12 months before start of ATAD in the ATAD group. Previous ESS was defined as ESS performed before the baseline visit. Follow-up ESS was defined as ESS performed after start with ATAD in the ATAD group, and in control group as ESS performed after the baseline ESS. Follow-up data were collected in 2018-2020; mean (min-max) 7.6 (0.9-16.3) years after start of ATAD, or after the consultation visit of the control group.

This follow-up data included peroral ASA-desensitization (dose, duration, side-effects, reason of possible discontinuation); revision ESS during follow-up (primary endpoint); information regarding purchased, doctor-prescribed OCS and antibiotic courses (due to CRS exacerbation and/or asthma) during 2016-2019, was obtained from the national electronic prescription system (secondary and tertiary endpoints, respectively). Discontinuation of peroral ASA desensitization as an endpoint, and the effect of peroral ASA maintenance dose and additional continuous OCS on all endpoints were also monitored.

We formed four patient groups based on ASA maintenance dose and adjuvant OCS therapy: No ATAD; ATAD with a peroral maintenance dose of <500 mg/day; ATAD with a peroral maintenance dose of ≥ 500 mg/day; ATAD with a peroral maintenance dose of ≥ 500 mg/day + adjuvant continuous OCS. The ATAD treatment was started by pulmonologists (five) or otorhinolaryngologists (two) specialized in treating N-ERD patients. Since the setting of this study is retrospective and observational, the choice of ATAD dose was made by these highly specialized and experienced

clinicians based on their clinical evaluation, and it was not affected by this study. In analyses of different endpoints, subjects with incomplete follow-up information were excluded. We also formed a variable of peroral ASA-maintenance dose*ATAD-duration (years) taking into account the total peroral ASA dose received before time of discontinuation.

Statistics

Statistical analysis was carried out by the SPSS Base 15.0 Statistical Software Package (SPSS Inc., Chicago, IL, USA). The discontinuation rate of ASA, revision ESS rate, antibiotic courses and OCS courses during the follow-up were signs of interest. Associations between ATAD dose and duration were assessed by Fisher's exact test (dichotomous), Kruskal-Wallis and Mann Whitney U tests (continuous). Differences between control and ASA-dose groups were also assessed by plotting the Kaplan-Meier survival curves and performing Log rank tests. Cox's proportional hazards models adjusted for previous ESS, baseline ESS, time (≤ 60 , >60 days) between baseline ESS and start of ATAD, and regular OCS use during follow-up, were used to evaluate the hazard ratio (HR) of revision ESS rate between ATAD and control group, and ASA dose groups. P-values $< .05$ were considered statistically significant.

Results

Patient characteristics

The subject characteristics are shown in Table 1. Groups did not differ between mean age, gender, duration of symptoms, history of allergic rhinitis (AR), asthma and smoking, baseline endoscopic NP score, or total Lund-Mackay score of sinus CT scans at the time of surgical consultation (Table 1). Compared to the control group, ATAD group had higher tissue eosinophilia, higher number of previous ESS, and higher proportion of patients using OCS (Table 1).

ATAD's discontinuation-rate

Of the ATAD group, 64 (62.7%) discontinued on average (min –max) 2.1 (0-9.3) years after ATAD start. Five patients had immediate severe adverse events (dyspnea $\pm$ skin/GI-symptoms) after ingestion of peroral ASA 35-50 mg, and did not continue ASA treatment at home (Table 1). Twenty-three (22.5%) patients reported long-term side effect(s) (tinnitus, GI pain/reflux, worsening of asthma) as reason for discontinuation, 28 (27.5%) had lack of effect; and 7 (6.9%) discontinued due to other reason(s) (such as pregnancy, operation or trauma) (Table 1). None discontinued due to no further need of ATAD because of good disease control. The peroral ASA dose was decreased to a lower, cardiological dose in six patients (500 mg $\rightarrow$ 250 mg in two patients; 200-300 mg $\rightarrow$ 100 mg in three patients, and 150 mg $\rightarrow$ 75 mg in one patient) after in average one year: in four cases due to not responding to a higher ASA dose, and in one case each due to tinnitus or hemorrhage.

The maximum ASA dose mean (min-max) was 322 (50-1500) mg perorally. The ASA maintenance dose mean (min-max) 293 (50-750) mg perorally was the dose that the patient was able to use at home, and the ATAD groups were formed based on this dose. There was no statistically significant difference between the ATAD groups in ASA discontinuation-rate ($p=0.39$ by log rank test, Figure 1). When stratifying the model by the baseline ESS, the results remained similar ($p>0.05$, data not shown). In the

group of patients who were able to continue ATAD, the mean (min-max) follow-up time (= ATAD duration time) was 4.3 (0.9-1.9) years.

Revision ESS rate during the follow-up

The data of follow-up surgeries was available for 167 of 171 patients. Of all patients, the revision ESS was performed on 62 (36.3%) patients, on average (min-max) 5.5(0.5-15) years after the baseline ESS (n=141) or start of ATAD in subjects without baseline ESS (n=30). Compared to the N-ERD group without ATAD, ATAD did not affect the revision ESS rate ($p=0.21$, by log rank test). Moreover, there was no statistically significant difference between the ATAD dose groups in revision ESS rates during the follow up ($p=0.42$ by log rank test, Figure 2A). When comparing the pooled group $ASA \geq 500mg \pm OCS$ to the $ASA < 500mg$ group, or to the control group, there was no statistically significant difference in the revision ESS rate (Figures 2B-C). The mean (min-max) time between baseline ESS and start of ATAD was 100 (3-359) days. The median time between baseline ESS and start of ATAD was significantly shorter in $ASA \geq 500mg + OCS$ group than in the other ATAD groups (Table 1). For survival models, three groups were formed: ATAD without baseline ESS, ATAD started ≤ 60 days after ESS, and ATAD started > 60 days after ESS. The revision ESS-rate did not differ between these three groups (Figure 2D). When observing the group that started ATAD ≤ 60 days after ESS, there was no difference between $ASA \geq 500mg \pm OCS$ group and $ASA < 500mg$ group in the revision ESS rate (Figure 2E).

The mean (min-max) value of $ASA\text{-maintenance dose} \times ATAD\text{-duration (years)}$ was 941 (3.3-4630) $mg \times years$. This total dose of ASA was divided into three groups: $ASA > 500 mg \times years$, $ASA 250-500$

mg*years, and ASA <250 mg*years. The revision ESS-rate did not differ between these three groups, or between controls (Figure 2F).

All patients, except those with immediate adverse events of ATAD at home, were observed in Cox's proportional hazards models. Compared to controls, revision ESS rate was not significantly associated with ATAD maintenance dose [adjusted HR (aHR) 0.99, 95% confidence interval (CI) 0.998-1.001, P=0.54]. Compared to controls, revision ESS rate was not statistically significantly associated with ASA-maintenance dose*ATAD-duration mg*years (aHR 1.00, 0.999-1.000, p=0.18).

Antibiotics and peroral corticosteroid courses during the follow-up

The data of purchased OCS courses during years 2016-2020 was available for 140 patients, 58 (41%) purchased at least one OCS course/year due to N-ERD. The number of patients purchasing over ≥ 2 antibiotic courses/year due to N-ERD was 16/138 (12%). The proportion of patients using continuous OCS due to N-ERD was 8 (4.7%), and 3 (1.8%) due to other diseases (granulomatous polyangitis, cardiac sarcoidosis and inflammatory bowel disease) (Table 1). Five patients started a biological therapy due to N-ERD after ATAD discontinuation (Table 1).

The number of purchased antibiotic or OCS courses /year during the follow-up, did not differ significantly between the control and ATAD groups, or between the ATAD dose groups ($p > 0.08$, Table 1). Results remained similar when excluding the patients who got immediate adverse events of ASA, and those receiving continuous OCS ($p > 0.1$ by Kruskal-Wallis test, Figure 3). Pair-wise comparisons showed an insignificant trend that the number of purchased OCS courses /year was higher in ASA group < 500 mg, compared to the control group ($p = 0.073$, Figure 3A). The number of purchased OCS courses /year was higher in the group who had received ASA 250-500 mg*year compared to the

control group or the group who had received ASA ≥ 500 mg*year ($p=0.007$, $p=0.018$ respectively, Figure 3B).

Responder analysis

The ATAD patients were analyzed in two groups based on their treatment response and ability to continue ATAD. Discontinuing patients were defined as those who discontinued ASA due to lack of effect or due to side effect. Responders were defined as being able to continue ATAD and not needing revision ESS during follow-up. The total follow-up time was longer in the discontinued group ($p=0.021$, Table 2). The other baseline or follow-up data did not differ between responders and non-responders/discontinuing patients (Table 2).

Discussion

The discontinuation-rate of ATAD was very high (63%) in this retrospective, real-world, multicenter study performed in the Finnish population. While 37% of the patients continued ATAD, we did not see clear dose-response effects of ATAD in respect to revision ESS or purchased antibiotic or OCS courses during the follow-up.

As in the literature (1, 2), also our study patients with N-ERD usually had a triad of CRSwNP, asthma and hypersensitivity to NSAIDs. ATAD is considered in uncontrolled N-ERD patients with difficult to treat symptoms and insufficient response to other pharmacological or surgical treatments (2, 22). Our study findings were in line with this; the ATAD group was more symptomatic, had higher tissue eosinophilia and more prior ESS, compared with the control group consisting of patients with severe CRSwNP and N-ERD not undergoing ATAD. The groups were otherwise very similar, especially in regard to LM score and endoscopic NP score, making study comparisons reliable. Since AD was performed in the ATAD group, their diagnosis of N-ERD can be considered confirmed also by ASA challenge, whereas NSAID intolerance diagnosis in the control group was based on anamnestic information.

Most studies concerning AD and ATAD and their benefits are, like ours, retrospective. The existing prospective, randomized, placebo controlled ATAD studies suggest that ATAD has a beneficial effect on symptoms, and rhinosinusitis and asthma control (32, 33). Ibrahim et al. (2014) reported in a retrospective study of 111 patients, that 73% of those in ASA maintenance therapy, had general improvement in their symptoms (32). They reported improvement in symptoms, QoL, reduced need for rescue therapy for both upper and lower airway symptoms, and improvement in sense of smell or taste

(32). In their double-blind, placebo controlled study comparing results of ATAD in asthmatics with (n=20) or without (n=14) ASA intolerance, Swierczynska-Krepa et al. (2014) report a beneficial effect on the rhinosinusitis and asthma control of those with ASA intolerance (33). Esmailzadeh et al. (2015) reported clinical efficacy of ATAD (ASA 650mg twice daily for 1 month, followed by 325mg twice daily) in improving symptoms, need for medication, QoL, sinus opacification and lung functions in their randomized, double-blind, placebo controlled study of 34 patients with N-ERD (34). Mortazavi et al. (2017) reported improvements in QoL, symptoms, and FEV1 values in their randomized, double-blind clinical study including 38 patients (35). In our study, 37% of the patients continued ATAD, probably benefitting from it. Our study was, however, not able to verify this benefit because symptom scores were not used. It is thus also possible that this is only a placebo effect. This clearly warrants further studies, it would be interesting to find ways to identify the subgroup of patients benefiting from ATAD, possibly using biomarkers.

In our study, ASA doses were lower compared to those usually mentioned in the literature (2, 28-31, 34). However, Fruth et al. (2013) have reported beneficial effects of ASA 100mg daily in patients with N-ERD (36). It seems that our population does not tolerate high doses of ASA. The reason for this is unknown, and could include e.g. genetic factors, clearly warranting further studies. Our results show that compared to controls, ATAD did not affect the need of OCS or antibiotic courses during follow-up, we also did not see clear dose-response effects of ATAD in respect to these.

Previous studies have shown that the following factors associate with the frequent revision ESS: male gender, young adult age, smoking, occupational exposure, radiological inflammatory findings in frontal sinus(es), presence of NPs, asthma, and N-ERD (37). Also tissue eosinophilia is associated with polyp recurrence (38).

CRSwNP often leads to recurrent surgeries, and ASA intolerance is associated with a higher prevalence of uncontrolled CRS after operative treatment (15, 16, 39). Early surgical intervention of CRS may be associated with less sinusitis-related postoperative health care utilization (40). Also the timing of ATAD in relation to ESS has been discussed, and part of the literature suggests ESS prior to ATAD (1, 2, 23-26). Due to the retrospective nature of this study, the timing of ESS in relation to ATAD was not constant. Our results show that compared to controls, ATAD did not affect revision ESS rates, nor did we see dose-response effects of ATAD in respect to this. In our study, there were no statistically significant differences in the revision ESS rates between the controls and different ATAD groups based on ASA dose and time between baseline ESS and ATAD. The group taking $ASA \geq 500mg \pm OCS$ did not have a statistically significantly lower risk for revision ESS in the follow-up. Peroral corticosteroids should usually not be used with ATAD, since it is known that peroral corticosteroid treatment combined with ASA increases the risk of gastrointestinal bleeding, necessitating careful patient selection, additional proton pump inhibitor treatment and monitoring for gastrointestinal side effects (2, 41).

A recent study from the USA reported good long term benefits of ATAD, in this study only 12% had discontinued the treatment (42). We had a much higher discontinuation rate (63%) of ATAD. Treatment responses may vary in different populations, this warrants future studies. Whenever possible, attempts were made to keep the patients on ATAD by treating side effects and lowering ASA doses. The most common reason for ATAD discontinuation in our study was side effects, lack of effect was less common. ATAD discontinuation did not depend on dose or duration.

The optimal ATAD dose, duration or indication has not been validated in randomized, placebo-controlled studies. The diagnostics or medical and surgical treatment of N-ERD were not standardized in our retrospective cohort. Due to real-world effectiveness study and the nature of this cohort the

patient groups were non-randomized. As also in other real-world ATAD studies, despite including all ATAD patients treated at our study hospitals, the number of patients is relatively small. The even smaller number of patients in the subgroups makes statistical analyses between them challenging, this may have affected our results. We acknowledge that, due to the retrospective study setting, the unfortunate lack of disease-specific QoL measurements, symptom scores, and lack of information regarding the sense of smell limit the interpretation of the findings. There might also have been limitations in the variables collected from patient records.

This study provides new data regarding the real-world use of ATAD in three tertiary centers. Our findings point out the need for standardizing N-ERD diagnostics and treatment by international, multicenter, prospective controlled setups.

The discontinuation-rate of ATAD was very high (63%) in this retrospective, real-world, multicenter study. We did not see dose-response effects of ATAD in respect to revision ESS or purchased antibiotic or OCS courses during the follow-up. However, 37% of the ATAD group did choose to continue the ATAD treatment, probably benefitting from it at least to some extent.

Author contributions

AL-H performed data collection and writing of this article. ST-S planned this study, collected & analyzed data and wrote this article. HT, SV, JN, JS, PK, PV, and MH participated in data collection. All authors critically reviewed the final article text.

Journal Pre-proof

References

1. White AAMD, Stevenson DDMD. Aspirin-Exacerbated Respiratory Disease. *New England Journal of Medicine*. 2018;379(11):1060-70.
2. Kowalski ML, Agache I, Bavbek S, Bakirtas A, Blanca M, Bochenek G, et al. Diagnosis and management of NSAID-Exacerbated Respiratory Disease (N-ERD)-a EAACI position paper. *Allergy*. 2018.
3. Mascia K, Haselkorn T, Deniz YM, Miller DP, Bleecker ER, Borish L, et al. Aspirin sensitivity and severity of asthma: evidence for irreversible airway obstruction in patients with severe or difficult-to-treat asthma. *Journal of Allergy & Clinical Immunology*. 2005;116(5):970-5.
4. Roca-Ferrer J, Perez-Gonzalez M, Garcia-Garcia FJ, Pereda J, Pujols L, Alobid I, et al. Low prostaglandin E2 and cyclooxygenase expression in nasal mucosa fibroblasts of aspirin-intolerant asthmatics. *Respirology*. 2013;18(4):711-7.
5. Morales DR, Guthrie B, Lipworth BJ, Jackson C, Donnan PT, Santiago VH. NSAID-exacerbated respiratory disease: a meta-analysis evaluating prevalence, mean provocative dose of aspirin and increased asthma morbidity. *Allergy*. 2015;70(7):828-35.
6. Berges-Gimeno MP, Simon RA, Stevenson DD. The natural history and clinical characteristics of aspirin-exacerbated respiratory disease. *Annals of Allergy, Asthma, & Immunology*. 2002;89(5):474-8.
7. Fokkens WJ, Lund VJ, Mullol J, Bachert C, Alobid I, Baroody F, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2012. *Rhinology Supplement*. 2012;(23)(23):3 p preceding table of contents, 1-298.
8. Stewart MG, Johnson RF. Chronic sinusitis: symptoms versus CT scan findings. *Current Opinion in Otolaryngology & Head & Neck Surgery*. 2004;12(1):27-9.
9. Rosenfeld RM, Andes D, Bhattacharyya N, Cheung D, Eisenberg S, Ganiats TG, et al. Clinical practice guideline: adult sinusitis. *Otolaryngology - Head & Neck Surgery*. 2007;137(3 Suppl):S1-31.
10. Scadding G, Hellings P, Alobid I, Bachert C, Fokkens W, van Wijk RG, et al. Diagnostic tools in Rhinology EAACI position paper. *Clinical and Translational Allergy*. 2011;1(1):2.
11. Stankiewicz JA, Chow JM. Nasal endoscopy and the definition and diagnosis of chronic rhinosinusitis. *Otolaryngology - Head & Neck Surgery*. 2002;126(6):623-7.
12. Ylitalo-Heikkilä M, Virkkula P, Sintonen H, Lundberg M, Roine RP, Hytonen M. Different rhinologic diseases cause a similar multidimensional decrease in generic health-related quality of life. *Clinical Otolaryngology*. 2018;43(6):1487-93.
13. Smith KA, Orlandi RR, Rudmik L. Cost of adult chronic rhinosinusitis: A systematic review. *Laryngoscope*. 2015;125(7):1547-56.
14. Hastan D, Fokkens WJ, Bachert C, Newson RB, Bislimovska J, Bockelbrink A, et al. Chronic rhinosinusitis in Europe--an underestimated disease. A GA2LEN study. *Allergy*. 2011;66(9):1216-23.
15. DeConde AS, Mace JC, Levy JM, Rudmik L, Alt JA, Smith TL. Prevalence of polyp recurrence after endoscopic sinus surgery for chronic rhinosinusitis with nasal polyposis. *The Laryngoscope*. 2017;127(3):550-5.
16. Koskinen A, Salo R, Huhtala H, et al. Factors affecting revision rate of chronic rhinosinusitis. *Laryngoscope Investigative Otolaryngology*. 2016;1(4):96-104.
17. Rank MA, Wollan P, Kita H, Yawn BP. Acute exacerbations of chronic rhinosinusitis occur in a distinct seasonal pattern. *Journal of Allergy & Clinical Immunology*. 2010;126(1):168-9.

18. Phillips KM, Hoehle LP, Bergmark RW, Caradonna DS, Gray ST, Sedaghat AR. Acute Exacerbations Mediate Quality of Life Impairment in Chronic Rhinosinusitis. *The Journal of Allergy & Clinical Immunology in Practice*. 2017;5(2):422-6.
19. Yamasaki A, Hoehle LP, Phillips KM, Feng AL, Campbell AP, Caradonna DS, et al. Association between systemic antibiotic and corticosteroid use for chronic rhinosinusitis and quality of life. *Laryngoscope*. 2018;128(1):37-42.
20. Banoub RG, Phillips KM, Hoehle LP, Caradonna DS, Gray ST, Sedaghat AR. Relationship between chronic rhinosinusitis exacerbation frequency and asthma control. *Laryngoscope*. 2018;128(5):1033-8.
21. GINA. 2018 GINA report: global strategy for asthma management and prevention. 2018 [Available from: <https://ginasthma.org/wp-content/uploads/2018/04/wms-GINA-2018-report-V1.3-002.pdf>.
22. Cortellini G, Caruso C, Romano A. Aspirin challenge and desensitization: how, when and why. *Current Opinion in Allergy & Clinical Immunology*. 2017;17(4):247-54.
23. Adappa ND, Ranasinghe VJ, Trope M, Brooks SG, Glicksman JT, Parasher AK, et al. Outcomes after complete endoscopic sinus surgery and aspirin desensitization in aspirin-exacerbated respiratory disease. *Int Forum Allergy Rhinol*. 2018;8(1):49-53.
24. Stevenson DD, Hankammer MA, Mathison DA, Christiansen SC, Simon RA. Aspirin desensitization treatment of aspirin-sensitive patients with rhinosinusitis-asthma: long-term outcomes. *Journal of Allergy & Clinical Immunology*. 1996;98(4):751-8.
25. Lee RU, Stevenson DD. Aspirin-exacerbated respiratory disease: evaluation and management. *Allergy, asthma & immunology research*. 2011;3(1):3-10.
26. Tajudeen BA, Schwartz JS, Bosso JV. The role of aspirin desensitization in the management of aspirin-exacerbated respiratory disease. *Current Opinion in Otolaryngology & Head & Neck Surgery*. 2017;25(1):30-4.
27. Waldram J, Walters K, Simon R, Woessner K, Waalen J, White A. Safety and outcomes of aspirin desensitization for aspirin-exacerbated respiratory disease: A single-center study. *Journal of Allergy & Clinical Immunology*. 2018;141(1):250-6.
28. Lee JYa, Simon RAB, Stevenson DDc. Selection of aspirin dosages for aspirin desensitization treatment in patients with aspirin-exacerbated respiratory disease. *Journal of Allergy & Clinical Immunology*. 2007;119(1):157-64.
29. Rozsasi A, Polzehl D, Deutschle T, Smith E, Wiesmiller K, Riechelmann H, et al. Long-term treatment with aspirin desensitization: a prospective clinical trial comparing 100 and 300 mg aspirin daily. *Allergy*. 2008;63(9):1228-34.
30. White AA, Stevenson DD. Aspirin desensitization in aspirin-exacerbated respiratory disease. *Immunology & Allergy Clinics of North America*. 2013;33(2):211-22.
31. Fokkens WJ, Lund VJ, Hopkins C, Hellings PW, Kern R, Reitsma S, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2020. *Rhinology*. 2020;58(Suppl S29):1-464.
32. Ibrahim C, Singh K, Tsai G, Huang D, Mazza J, Rotenberg B, et al. A retrospective study of the clinical benefit from acetylsalicylic acid desensitization in patients with nasal polyposis and asthma. *Allergy, Asthma, & Clinical Immunology : Official Journal of the Canadian Society of Allergy & Clinical Immunology*. 2014;10(1):64.
33. SwierczynskaKrepa M, Sanak M, Bochenek G, Strek P, Cmiel A, Gielicz A, et al. Aspirin desensitization in patients with aspirin-induced and aspirin-tolerant asthma: A double-blind study. *Journal of Allergy & Clinical Immunology*. 2014;134(4):883-90.
34. Esmaeilzadeh H, Nabavi M, Aryan Z, Arshi S, Bemanian MH, Fallahpour M, et al. Aspirin desensitization for patients with aspirin-exacerbated respiratory disease: A randomized double-blind placebo-controlled trial. *Clinical Immunology*. 2015;160(2):349-57.

35. Mortazavi N, Esmailzadeh H, Abbasinazari M, Babaie D, Alyasin S, Nabavizadeh H, et al. Clinical and Immunological Efficacy of Aspirin Desensitization in Nasal Polyp Patients with Aspirin-Exacerbated Respiratory Disease. *Iranian Journal of Pharmaceutical Research*.2017;16(4):1639-47.
36. Fruth K, Pogorzelski B, Schmidtmann I, Springer J, Fennan N, Fraessdorf N, et al. Low-dose aspirin desensitization in individuals with aspirin-exacerbated respiratory disease. *Allergy*.2013;68(5):659-65.
37. Brescia G, Marioni G, Franchella S, Ramacciotti G, Giacomelli L, Marino F, et al. A prospective investigation of predictive parameters for post-surgical recurrences in sinonasal polyposis. *European archives of oto-rhino-laryngology : official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS) : affiliated with the German Society for Oto-Rhino-Laryngology - Head and Neck Surgery*. 2016;273(3):655-60.
38. Nakayama T, Yoshikawa M, Asaka D, Okushi T, Matsuwaki Y, Otori N, et al. Mucosal eosinophilia and recurrence of nasal polyps - new classification of chronic rhinosinusitis. *Rhinology*. 2011;49(4):392-6.
39. van der Veen J, Seys SF, Timmermans M, Levie P, Jorissen M, Fokkens WJ, et al. Real-life study showing uncontrolled rhinosinusitis after sinus surgery in a tertiary referral centre. *Allergy*. 2017;72(2):282-90.
40. Benninger MS, Sindwani R, Holy CE, Hopkins C. Early versus delayed endoscopic sinus surgery in patients with chronic rhinosinusitis: impact on health care utilization. *Otolaryngology - Head & Neck Surgery*. 2015;152(3):546-52.
41. Masclee GMC, Valkhoff VE, Coloma PM, de Ridder M, Romio S, Schuemie MJ, et al. Risk of upper gastrointestinal bleeding from different drug combinations. *Gastroenterology*. 2014;147(4):784-92.e9.
42. Walters KM, Waldram JD, Woessner KM, White AA. Long-term Clinical Outcomes of Aspirin Desensitization With Continuous Daily Aspirin Therapy in Aspirin-exacerbated Respiratory Disease. *American Journal of Rhinology & Allergy*.2018;32(4):280-6.

Figure legends:

Figure 1. Predictive effect of ASA (Acetylsalicylic acid) desensitization (ATAD) dose to discontinuation. Data of NSAID-exacerbated respiratory disease patients undergoing surgical and ASA desensitization consultation (2001-2017), able to start ATAD, were used with follow-up data of endoscopic sinus surgery (ESS). Those (5) with immediate adverse events were excluded from analysis. Comparisons between ASA; OCS = oral corticosteroid(s). P-values by log rank test.

Figure 2. Predictive effect of ATAD (ASA desensitization [AD] followed by ASA treatment after AD) to the time until revision endoscopic sinus surgery (ESS) and/or nasal polypectomy was performed according to the Kaplan-Meier method. Data of CRSwNP patients undergoing CRS-surgical consultation in 2001-17 were used with follow-up data of ESS.

Figure 3. Number of purchased prescribed oral corticosteroids (OCS)/year (for CRSwNP and/or asthma). Data analyzed from N-ERD patients undergoing surgical consultation due to CRS 2001-2017 with follow-up data of prescribed medication from electronic database 2016-2020. ASA= Acetylsalicylic acid. Data excluded from analysis for 5 patients with immediate adverse events and 11 receiving OCS. P-values by Mann Whitney U test. Only p-values <0.1 are shown.

536 **Table 1.** Patient background and follow-up characteristics.

	No ATAD	ATAD			p ¹	p ²
		ATAD <500mg/day	ATAD ≥500mg/day	ATAD ≥500mg/day+ OCS		
	n=69	n=67	n=27	n=8		
Female gender, n (%)	40 (58.0)	44 (65.7)	15 (55.6)	5 (62.5)	.74	.63
Age, median (Q1-Q3)	47.0 (31.4-58.8)	46.9 (37.9-55.8)	45.5 (39.5-52.5)	48.3 (41.0-53.5)	1.0	.90
Current smokers, n (%)						
No	52 (80.0)	56 (84.8)	22 (95.7)	7 (87.5)	.35	.27
Current	13 (20.0)	10 (15.2)	1 (4.3)	1 (12.5)		
AR, n (%)						
No	31 (44.9)	26 (39.4)	11 (40.7)	4 (50.0)	.88	.64
Yes	38 (55.1)	40 (60.6)	16 (59.3)	4 (50.0)		
Asthma, n (%)						
No	3 (4.3)	1 (1.5)	1 (3.8)	0 (0.0)	.67	.40
Yes	66 (95.7)	66 (98.5)	25 (96.2)	8 (100.0)		
Duration of symptoms (y), median (Q1-Q3)	10 (4-23)	13 (8-25)	12.5 (8.5-15.75)	27.5 (15-39)	.069	.19
A history of ≥ 1 OCS course /y						
No	35 (54.7)	30 (46.2)	9 (34.6)	2 (25.0)	.21	0.11
Yes	29 (45.3)	35 (53.8)	17 (65.4)	6 (75.0)		
A history of ≥ 4 antibiotic course /y						
No	30 (58.8)	34 (55.7)	19 (70.4)	6 (75.0)	.52	0.86
Yes	21 (41.2)	27 (44.3)	8 (29.6)	2 (25.0)		
NP eosinophilia						
< 30%	15 (37.5)	5 (14.7)	0 (0)	0 (0)	.065	0.19
≥ 30%	25 (62.5)	29 (85.3)	4 (100)	1 (100)		
Endoscopic NP score, median (Q1-Q3)	5 (4-6)	5 (4-6)	4.5 (3.5-6)	4 (3.5-6)	.58	.29
Total LM score of sinus CT scans, median (Q1-Q3)	16 (14-22.5)	20 (15-22)	19.5 (14-21)	15.5 (13.25-18.5)	.36	.52

Number of previous ESS, median (Q1-Q3)	1 (0-2)	2 (2-5)	2 (1-4)	3.5 (2-5)	<.001	<.001
Baseline CRS-surgery in 1 year after the first visit (before ATAD)					<.001	<.001
No surgery	0 (0)	20 (32.8)	4 (16.0)	1 (12.5)		
Polypectomy ±ESS without opening of ethmoidal cells	26 (37.7)	20 (32.8)	6 (24.0)	3 (37.5)		
Polypectomy ±ESS with partial opening of ethmoidal cells	29 (42.0)	16 (26.2)	6 (24.0)	1 (12.5)		
Polypectomy ±ESS with total opening of ethmoidal cells	14 (20.3)	5 (8.2)	9 (36.0)	3 (37.5)		
Time between baseline ESS and start of ATAD, median (Q1-Q3)		137 (51-250)	45 (19-100)	25 (18-48)	<.001	-
Duration of ATAD (y), median (Q1-Q3)	-	2.2 (0.6-4.2)	2.4 (1.0-4.2)	5.3 (3.0-6.4)	.10	-
Reason for discontinuation of ATAD						
Total adverse events, n (%)	-	40 (60.6)	20 (74.1)	4 (50.0)	.37	-
Immediate adverse event ³		5 (23.9)	11 (40.7)	1 (12.5)		
Long-term side effects ⁴		16 (7.5)	0 (0)	0 (0)		
No effect/disease relapse		16 (23.9)	4 (14.8)	3 (37.5)		
Other ⁵		4 (6.0)	3 (11.1)	0 (0)		
Nr of antibiotic courses /year during 2016-19, median (Q1-Q3)	0.5 (0-1.3)	0.5 (0-2)	0.3 (0-1.1)	0.5 (0-1)	.56	.86
Nr of OCS courses /year during 2016-19, median (Q1-Q3)	0 (0-1)	0.5 (0-1)	0.5 (0.5-1.3)	0 (0-0.5)	.084	.11
Nr of patients using continuous OCS, n (%)						
Due to any reason	1 (1.4)	2 (3)	0 (0)	8 (100)	<.001	.029
Due to N-ERD	0 (0)	0 (0)	0 (0)	8 (100)	<.001	.022
Due to other disease ⁶	1 (1.4)	2 (3)	0 (0)	0(0)	.80	1.00
Start with biological therapy (after ATAD discontinuation)	0 (0)	1 (1.5)	1 (1.5)	3 (37.5)	<.001	.082
Revision ESS in 5 years, n (%)						
No	34 (58.6)	34 (59.6)	13 (68.4)	6 (85.7)	.54	.60
Yes	24 (41.4)	23 (40.4)	6 (31.6)	1(14.3)		

AD= aspirin desensitization, ATAD= AD followed by ASA treatment after desensitization, AR= allergic rhinitis, ASA=aspirin, CRS= chronic rhinosinusitis, ESS= endoscopic sinus surgery, N-ERD = patient-reported NSAID-exacerbated respiratory disease, NP=nasal polyps, OCS= oral corticosteroid(s). P values by Fisher's exact test (dichotomous variables) or Kruskal Wallis and Mann Whitney U test (continuous variables). Q1= 25% percentile, Q3= 75% percentile. Bold text indicates a statistically significant difference with a P value less than 0.05. ¹p-value by comparing all 4 groups. ²p-value by comparing no ATAD and ATAD groups, ³Severe acute dyspnea ±skin/gastrointestinal (GI) symptoms. ⁴tinnitus, GI pain/reflux, worsening of asthma. ⁵Operation, trauma, pregnancy/lactation. ⁶granulomatous polyangitis, cardiac sarcoidosis and inflammatory bowel disease.

544 **Table 2.** Comparison of baseline and follow up data of ATAD patients. ATAD patients were divided into two
 545 groups: those who discontinued ASA, and those who continued ASA and were not re-operated during the
 546 follow-up (=responders).

	ATAD discontinued N=57	ATAD continues & no revision ESS N=36	p
Female gender, n (%)	37 (64.9)	23 (63.9)	1.0
Age, median (Q1-Q3)	44.6 (37.4-55.5)	46.4 (39.8-54.9)	.66
Current smoker, n (%)	7 (12.7)	4 (11.1)	1.0
AR, n (%)	33 (58.9)	20 (55.6)	.83
Asthma, n (%)	55 (98.2)	35 (97.2)	1.0
Duration of symptoms (y), median (Q1-Q3)	13 (8-20)	11 (3-30)	.20
≥ 1 OCS course /y	35 (64.8)	17 (47.2)	.13
≥ 4 antibiotic course /y	20 (37.7)	14 (41.2)	.82
NP eosinophilia ≥ 30%	19 (95.0)	12 (75.0)	.15
Baseline endoscopic NP score ≥ 4, n (y)	37 (78.7)	27 (81.8)	.78
Baseline total LM score ≥ 16, n (y)	41 (75.9)	20 (60.6)	.15
≥ 3 previous CRS-surgeries, n (%)	28 (49.1)	17 (47.2)	1.0
Baseline polypectomy± ESS before start of ATAD	36 (63.2)	30 (83.3)	.059
Baseline total ethmoidectomy before start of ATAD	7 (12.3)	9 (25.7)	.16
Time between baseline ESS and start of ATAD ≤ 60 days	16 (50.0)	14 (48.3)	1.0
Maximum dose of ASA mg/day, median (Q1-Q3)	250 (100-500)	250 (100-500)	.46
Maintenance dose of ASA mg/day, median (Q1-Q3)	250 (100-500)	250 (100-500)	.21
Duration of ATAD (y), median (Q1-Q3)	1.0 (0.3-2.6)	4.3 (3.6-5.6)	<.001
Total Follow-up time (y), median (Q1-Q3)	6.3 (4.6-8.0)	4.8 (4.0-6.3)	.021
> 2 antibiotic courses /year during 2016-19, n (%)	6 (12.8)	2 (6.9)	.70
> 1 OCS course /year during 2016-19, n (%)	11 (23.4)	7 (22.6)	1.0
Continuous OCS during the follow-up	3 (5.3)	6 (16.7)	.084
Start with biological therapy (after ATAD discontinuation)	4 (7.0)	1 (2.8)	.65
Revision ESS in 5 years, n (%)	25 (50.0)	0 (0)	<.001

AD=aspirin desensitization, AR=allergic rhinitis, ASA=aspirin, ATAD=AD followed by ASA treatment after desensitization, CRS=chronic rhinosinusitis, ESS=endoscopic sinus surgery, NP=nasal polyps; OCS= oral corticosteroid(s). P values by Fisher's exact test (dichotomous variables) or Kruskal Wallis and Mann Whitney U test (continuous variables). Q1= 25% percentile, Q3= 75% percentile. Bold text indicates a statistically significant difference with a p value less than 0.05. None of the ATAD patients discontinued ATAD due to good response. Only patients with available follow-up data were included. Those who continued ASA but were re-operated (n=6) were excluded.

Time until revision ESS (years)

Proportion without revision ESS