<https://helda.helsinki.fi>

Chlorophyll a fluorescence illuminates a path connecting plant molecular biology to Earth-system science

Porcar-Castell, Albert

2021-08-09

Porcar-Castell , A , Malenovsky , Z , Magney , T , Van Wittenberghe , S , Fernandez-Marin , B , Maignan , F , Zhang , Y , Maseyk , K , Atherton , J , Albert , L P , Robson , T M , Zhao , F , Garcia-Plazaola , J-I , Ensminger , I , Rajewicz , P A , Grebe , S , Tikkanen , M , Kellner , J R , Ihalainen , J A , Rascher , U & Logan , B 2021 , ' Chlorophyll a fluorescence illuminates a path connecting plant molecular biology to Earth-system science ' , Nature plants , vol. 7 , p y n o . 8 , p p . 998 1009 . <https://doi.org/10.1038/s41477-021-00980-4>

<http://hdl.handle.net/10138/339926>

<https://doi.org/10.1038/s41477-021-00980-4>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Chlorophyll-a fluorescence illuminates a path connecting plant molecular biology to Earth-system science

Albert Porcar-Castell^{1*}, Zbyněk Malenovský², Troy Magney³, Shari Van Wittenberghe^{1,4}, Beatriz Fernández-Marín⁵, Fabienne Maignan⁶, Yongguang Zhang⁷, Kadmiel Maseyk⁸, Jon Atherton¹, Loren P. Albert^{9, 10}, Thomas Matthew Robson¹¹, Feng Zhao¹², Jose-Ignacio Garcia-Plazaola¹³, Ingo Ensminger¹⁴, Paulina A. Rajewicz¹, Steffen Grebe¹⁵, Mikko Tikkanen¹⁵, James R. Kellner^{9, 16}, Janne A. Ihalainen¹⁷, Uwe Rascher¹⁸, Barry Logan¹⁹

¹Optics of Photosynthesis Laboratory, Institute for Atmospheric and Earth System Research/Forest Sciences, Viikki Plant Science Center, University of Helsinki, Helsinki, Finland.

²School of Geography, Planning, and Spatial Sciences, College of Sciences Engineering and Technology, University of Tasmania, Private Bag 76, Hobart, TAS 7001, Australia.

³Department of Plant Sciences, University of California, Davis. Davis, CA, 95616 United States of America.

⁴Laboratory of Earth Observation, University of Valencia, C/Catedrático José Beltrán, 2, 46980 Paterna, Spain.

⁵Department of Botany, Ecology and Plant Physiology, University of La Laguna (ULL), Tenerife 38200, Spain.

⁶Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France.

⁷International Institute for Earth System Sciences, Nanjing University, Nanjing, Jiangsu 210023, China.

⁸School of Environment, Earth and Ecosystem Sciences, The Open University, Milton Keynes MK7 6AA, United Kingdom.

- 26 ⁹Institute at Brown for Environment and Society, Brown University, Providence, RI 02912, United
27 States of America.
- 28 ¹⁰Biology Department, West Virginia University, Morgantown, WV 26506-6300, United States of
29 America.
- 30 ¹¹Organismal and Evolutionary Biology, Viikki Plant Science Centre (ViPS), Faculty of Biological
31 and Environmental Science, 00014, University of Helsinki, Finland.
- 32 ¹²School of Instrumentation Science and Opto-Electronics Engineering, Beihang University,
33 Beijing, 100083, China.
- 34 ¹³Department of Plant Biology and Ecology, University of the Basque Country (UPV/EHU),
35 Bilbao, Spain.
- 36 ¹⁴Department of Biology, Graduate Programs in Cell & Systems Biology and Ecology &
37 Evolutionary Biology, University of Toronto, 3359 Mississauga Road, Mississauga, ON L5L 1C6,
38 Canada.
- 39 ¹⁵Molecular Plant Biology, University of Turku, FI-20520 Turku, Finland.
- 40 ¹⁶Department of Ecology and Evolutionary Biology, Brown University, Providence RI 02912,
41 United States of America.
- 42 ¹⁷Nanoscience Center, Department of Biological and Environmental Science, University of
43 Jyväskylä, Jyväskylä 40014, Finland.
- 44 ¹⁸Institute of Bio- and Geosciences, Plant Sciences (IBG-2), Forschungszentrum Jülich GmbH,
45 Jülich, Germany.
- 46 ¹⁹Biology Department, Bowdoin College, Brunswick, Maine, United States of America.
- 47
- 48 *Corresponding author: joan.porcar@helsinki.fi
- 49
- 50

51 **For decades, the dynamic nature of chlorophyll-a fluorescence (ChlaF) has provided insight**
52 **into the biophysics and ecophysiology of the light reactions of photosynthesis from the**
53 **subcellular to leaf scales. Recent advances in remote sensing methods now enable detection of**
54 **ChlaF induced by sunlight across a range of larger scales, using instruments mounted on**
55 **towers above plant canopies to Earth-orbiting satellites. This signal is referred to as *solar-***
56 ***induced fluorescence* (SIF) and its application promises to overcome spatial constraints on**
57 **studies of photosynthesis, opening new research directions and opportunities in ecology,**
58 **ecophysiology, biogeochemistry, agriculture and forestry. However, to unleash the full**
59 **potential of SIF, intensive cross-disciplinary work is required to harmonize these new**
60 **advances with the rich history of biophysical and ecophysiological studies of ChlaF, fostering**
61 **the development of next-generation plant physiological and Earth system models. Here, we**
62 **introduce the scale-dependent link between SIF and photosynthesis, with an emphasis on**
63 **seven remaining scientific challenges, and present a roadmap to facilitate future collaborative**
64 **research towards new SIF applications.**

65 When illuminated, chlorophyll-a molecules weakly emit light in the 650-850 nm range; that is, they
66 fluoresce. Steady state^{1,2} and time-resolved fluorescence spectroscopy^{3,4}, as well as pulse-amplitude
67 modulated (PAM) fluorescence^{5,6} have long been used by biophysicists, molecular biologists and
68 ecophysiolgists to elucidate the structure and function of the photosynthetic apparatus⁷⁻⁹. These
69 techniques are regarded as active because the measured ChlaF originates from a controlled light
70 source, and accordingly have largely^{10,11} been restricted to measurements at the subcellular and leaf
71 levels.

72 Interest in passive remote sensing methods capable of retrieving solar-induced ChlaF across a
73 continuum of spatial scales emerged more than two decades ago¹². These seminal activities led to
74 the first demonstrations of tower-based^{13,14} and satellite¹⁵ SIF measurements over terrestrial
75 ecosystems. The opportunity to remotely detect an energy flux (Box 1) that arises directly from

76 within the photosynthetic process spurred the rapid development of measurement techniques,
77 retrieval protocols, and models for estimating and interpreting SIF across scales. As reviewed in
78 Mohammed et al.¹² and Aasen et al.¹⁶, SIF can now be measured from an expanding number of
79 sensors mounted on towers^{17,18}, drones^{19,20}, aircraft^{21,22} and satellites with ever-improving spatial
80 and temporal resolution^{23,24}. So far, all satellite SIF retrievals have been serendipitous, relying on
81 instruments originally designed to measure atmospheric gases. The first satellite mission designed
82 specifically for the measurement of SIF is the ESA FLuorescence EXplorer (FLEX) mission, which
83 is set to launch in 2024²⁵.

84 SIF methods are rapidly breaking through the scale bottleneck of traditional ChlaF measurements,
85 opening up a range of new opportunities to study photosynthesis across the continuum of spatial
86 scales from the leaf, through plant canopies, and up to the globe. With SIF we now have the
87 potential to illuminate the path connecting plant molecular biology to Earth-system science.
88 However, before the full potential of multiscale SIF observations can be realized, a number of
89 challenges must be overcome. Extracting the information embedded in the SIF signal requires a
90 fundamental understanding and a quantitative description of the processes that connect measured
91 ChlaF with photosynthesis (Fig.1), as well as their variation across space and time (Fig. 2). In this
92 Perspective, we present these challenges and propose a roadmap of activities to facilitate future
93 research. Finally, we discuss key emerging SIF applications that can benefit from cross-disciplinary
94 expertise.

95 **Challenge 1: $APAR_g$.** The common denominator between ChlaF and the photosynthetic uptake of
96 CO_2 is the flux of photosynthetically active radiation absorbed by photosynthetic pigments, or
97 $APAR_g$ (where the g stands for green), which provides the foundation for the mechanistic
98 connection between SIF and photosynthesis. $APAR_g$ is the product of the incoming
99 photosynthetically active radiation (PAR) and the fraction of this PAR absorbed by photosynthetic
100 pigments ($fAPAR_g$) (Fig.1). Importantly, although the absorption of radiation by leaves and plant

101 canopies can be quantified using radiometric sensors either coupled to an integrating sphere²⁶ (e.g.
102 leaf absorptance profile in Fig.1) or mounted above and below a plant canopy²⁷, these
103 measurements also include a significant and dynamic contribution from non-photosynthetic
104 pigments and other canopy elements. While inaccuracies in the estimation of APAR_g do not disrupt
105 the relationship between SIF and photosynthesis, accurate quantification of the energy flux entering
106 the photosynthetic process is essential for a mechanistic interpretation of SIF and remains a
107 challenge.

108 ***Challenge 2: Distribution of excitation energy between PSII and PSI and their ChlaF emissions.***

109 APAR_g is absorbed mostly by chlorophyll-a and chlorophyll-b associated with either photosystem
110 II (PSII) or photosystem I (PSI) reaction centres. Interestingly, while both types of chlorophyll have
111 the capacity to fluoresce, essentially all chlorophyll fluorescence *in vivo* originates from
112 chlorophyll-a due to the efficient transfer of excitation energy from chlorophyll-b to chlorophyll-a
113 within light harvesting antennae²⁸. Likewise, although both photosystems emit ChlaF, ChlaF from
114 PSII typically dominates the signal, especially in the red region of the emission spectrum² (Fig.1),
115 and exhibits greater variation in quantum yield in response to photochemical and non-
116 photochemical processes^{7,29}. The dynamic nature of PSII ChlaF explains the widespread application
117 of PAM fluorescence to probe the energy partitioning between photochemical and non-
118 photochemical processes or to estimate the rate of linear electron transport (LET) in PSII³⁰.
119 However, the estimation of LET requires knowledge of the distribution of absorption between the
120 photosystems (i.e. the use of an energy partitioning factor), which is rarely measured and often
121 assumed to be 0.5⁶. Although biochemical and biophysical methods to assess the stoichiometry and
122 antenna sizes of PSI and PSII do exist³¹⁻³³, these methods only provide a relative assessment of the
123 energy distribution; absolute quantification requires the combination of simultaneous ChlaF and
124 820 nm absorption measurements to probe the energy partitioning in PSII and PSI, respectively,
125 along with photosynthetic gas exchange measurements³⁴. Overall, the evidence gathered to date

126 suggests that neither the distribution of excitation energy between PSII and PSI nor the contribution
127 of ChlaF from PSI to SIF remain constant over time, between species or within canopy light
128 gradients^{35,36}. Questions remain: how large is this variability? What controls it? And what is its
129 significance for the interpretation of SIF? Answers to these questions await the development of
130 versatile field methods and protocols (e.g. based on rapid optical measurements³⁷) to enable
131 the characterization of these factors across a wide range of conditions.

132 **Challenge 3: Energy partitioning in PSII.** Energy absorbed in PSII is partitioned between three
133 main processes: a) photochemical quenching (PQ) of excitation energy, promoting linear electron
134 transport, b) non-photochemical quenching (NPQ), which includes both regulated and sustained
135 forms of thermal dissipation, and c) emission of ChlaF. The quantum yield of a process, e.g. ChlaF
136 emission (Φ_F), can be expressed as the ratio of the rate constant associated to that process relative
137 to the sum of all rate constants. Importantly, the rate constants associated to PQ and NPQ are highly
138 dynamic, which allows plants to regulate the flow of energy through PSII and to protect against
139 light-induced damage^{38, 39}. During the growing season, the rate constants of PQ and NPQ vary over
140 time-scales of seconds to minutes in response to the redox dynamics of the quinone acceptor pool
141 and induction and relaxation of regulated thermal dissipation, respectively. Outside of the growing
142 season, or during periods of profound environmental stress, rate constants can be affected by
143 photoinhibition of PQ and the induction of sustained NPQ. Accordingly, changes in the quantum
144 yield of ChlaF (Φ_F) reflect the combined effect of PQ and NPQ dynamics and a quantitative
145 connection between Φ_F and Φ_P (the quantum yield of photochemistry) cannot be established
146 without knowledge of either PQ or NPQ^{8,40}. PAM fluorescence uses saturating light pulses to solve
147 the energy partitioning and estimate Φ_P ; an approach that is not feasible during SIF measurements,
148 precluding partitioning from SIF alone.

149 Under certain conditions, either NPQ or PQ can dominate the relationships between Φ_F and Φ_P ,
150 resulting in the emergence of a positive or negative relationship respectively. For example, under

151 low light intensities – when regulatory NPQ remains inactive - the relationship between Φ_P and Φ_F
152 is negative under the action of PQ, which exerts opposite effects on (i.e. decouples) Φ_P and Φ_F .
153 Under high light - when PQ tends to saturate and NPQ is highly active - the relationship between
154 Φ_P and Φ_F turns to positive under the action of NPQ, which competes for excitation with both (i.e.
155 couples) Φ_P and Φ_F ^{12,52,53}. The latter case can explain the seasonal correlation between Φ_P and Φ_F
156 observed at the leaf^{41,42} (Fig. 2) and canopy scales¹⁸, in response to the modulation of sustained
157 NPQ that protects the foliage from the harmful combination of excessive light and low
158 temperatures^{43,44}. Despite the positive relationship between Φ_P and Φ_F that emerges in response to
159 certain stress conditions, the quantitative treatment of the energy partitioning in PSII requires the
160 use of mechanistic models and remains one of the core challenges to the interpretation of SIF^{40,45,46}.

161 **Challenge 4: Alternative energy sinks.** Photosynthetic linear electron transport provides reducing
162 power for a range of metabolic processes beyond CO₂ assimilation via the Calvin cycle, including
163 chlororespiration⁴⁷, photorespiration⁴⁸, nitrogen, sulphur and oxygen reduction (the latter known as
164 the Mehler reaction in the water-water cycle⁴⁹), and the synthesis of volatile organic compounds⁵⁰.
165 Importantly, the dynamics of these ‘non-assimilatory’ electron sinks can affect ChlF in a manner
166 not directly correlated with CO₂ assimilation. In particular, because alternative energy sinks can
167 have a protective function by sustaining LET under conditions when CO₂ assimilation is impaired⁵¹,
168 they could influence the capacity of SIF to detect certain plant stress responses. Therefore, it is
169 critical to address the extent that these dynamics decouple SIF from GPP, in particular during plant
170 stress. As with Challenge 2, answering this question will benefit from the development of versatile
171 field methods and protocols to promote the widespread characterization of these factors across a
172 wide range of conditions.

173 **Challenge 5: Leaf and canopy ChlaF scattering, reabsorption and measurement geometry.**

174 Although the lighter and darker green stripes seen on an athletic field may give the impression of
175 different chlorophyll contents, they are an optical reflection effect created when the grass is bent in

176 a particular direction during mowing. SIF measurements over plant canopies are similarly affected
177 by the distribution of leaves, canopy architecture and measurement geometry^{27,52}. The amount and
178 distribution of chlorophyll within a leaf (influenced by photosystem and thylakoid structure,
179 chloroplast distribution, and internal leaf morphology), as well as the amount and geometrical
180 arrangement of leaves and other non-photosynthetic material within a plant canopy (influenced by
181 branch/stem architecture) drive $APAR_g$, connecting SIF and photosynthesis at the leaf and canopy
182 scales, respectively. Once emitted, ChlaF photons travel through the same leaf and canopy
183 structures, where some of the ChlaF photons are reabsorbed (Fig. 1 and Fig. 2 “spectral dynamics”).
184 As a result, spatial and temporal variations in leaf biochemistry, leaf morphology, and canopy
185 architecture, as well as foliage illumination and viewing geometry, influence the probability of
186 ChlaF photons contributing to a SIF measurement (known as the escape probability). These factors
187 decouple the total emitted ChlaF from the measured SIF, and by extension from photosynthesis.
188 Physically-based radiative transfer models, which simulate the movement of photons through leaves
189 and plant canopies (Box 2), can be used to provide a quantitative framework to investigate and
190 account for the impact of these factors on $APAR_g$ and SIF observations^{27,53}. Although spatially
191 explicit RTM approaches are already available (see Supplementary Video 1 and 2), advances in the
192 parametrization of within-leaf and canopy drivers of SIF - e.g. canopy gradients in foliar
193 morphology, pigment contents (Challenge 1) or ChlaF contribution from PSI (Challenge 2) - remain
194 areas of active development.

195 ***Challenge 6: Atmospheric absorption and scattering.*** Atmospheric gases, aerosols and other
196 particles absorb and scatter ChlaF photons traveling from a plant canopy to a remote detector. The
197 extent of atmospheric absorption and scattering of SIF depends on the retrieval wavelength, the
198 distance between target and sensor, and the properties of the atmosphere (Box 1). In particular, SIF
199 retrieval methods based on the in-filling of atmospheric gas absorption bands, such as the O2-A or
200 O2-B bands (Fig. 1), face the challenge that the gas absorption feature used for the SIF retrieval

simultaneously attenuates the ChlaF signal as it travels towards the detector. This effect requires a correction even for short-distance measurements from canopy towers and drones⁵⁴. Although an atmospheric RTM can be used to characterize and correct for these effects, its application requires site-specific measurements of atmospheric profile parameters for model input, which remains an operational challenge⁵⁵.

Challenge 7: Integrating SIF controls across space and time. A final challenge, and perhaps the most relevant, is the contextualization of the interpretation of SIF (including the previous six challenges) within the spatial and temporal domain of the measurements (Fig. 2). Temporally, ChlaF dynamics have been used to investigate the energy transfer within photosystems (femto-picosecond scale)^{7,56}, the redox status of the donor and acceptor sides of the photosystem (microsecond-millisecond scale)^{3,4}, and the variations in PQ and NPQ (seconds-to-seasonal scale)^{38,39,44}. Spatially, the intensity and spectral properties of SIF are also controlled by factors that regulate both APAR_g and ChlaF scattering and reabsorption within a leaf or plant canopy^{57,58} (Fig. 2, “spectral dynamics”). When ChlaF is measured as SIF across coarser spatial and longer temporal scales, the signal carries information that aggregates an expanding assortment of physical and biological factors⁵⁹⁻⁶¹. New controls may appear while the effects of others may be subordinated, strengthening (via ‘couplers’; Fig. 2) or disrupting (via ‘decouplers’; Fig. 2) the relationship between SIF and GPP.

For example, tower-based SIF studies reveal a strong seasonal linear relationship between canopy SIF and ecosystem GPP across a wide range of ecosystems^{17,18,61}, consistent with the coupling action of APAR_g and NPQ described above. Yet, the sensitivity, strength and linearity of the seasonal SIF-GPP relationship is not universal and has been found to depend on additional physical and physiological decoupling factors, such as sun-vegetation-sensor geometries^{62,63}, vegetation canopy structure^{52,64}, or photosynthetic pathway (C3 vs. C4)^{27,65}, with contrasting responses to different environmental stressors^{66,67}. Clearly, integrating and disentangling the relationship

226 between SIF and GPP across species, space, time and in response to environmental stress, remains
227 still a challenge that calls for comprehensive field studies.

228 **Roadmap towards a consistent interpretation of SIF**

229 The time for multiscale SIF measurements is already here (Fig. 3). Yet, converting these data into
230 meaningful information and new applications still requires effort dedicated to scaling and
231 standardizing methods for SIF interpretation, with particular attention to the seven challenges
232 described above. This process requires accounting for the influence of 1) instrumental, 2)
233 atmospheric, 3) structural and 4) physiological factors to unlock the quantitative association
234 between measured SIF and photosynthesis (Fig. 4). Addressing these challenges requires new data,
235 protocols and models to interpret SIF and bridge the gap between molecular processes, i.e.
236 photosynthesis, and satellite imagery.

237 At the leaf level, new instruments and techniques employing optical bandpass filters have been
238 developed to record fluorescence spectral dynamics under both natural or controlled illumination,
239 temperature, and CO₂ concentration^{16,68-71}. Such spectral approaches, combined with foliar pigment
240 analysis, photosynthetic gas exchange, and PAM ChlaF measurements, provide new insights into
241 the connection between SIF and photosynthesis dynamics of leaves^{42,69,72}. Going forward,
242 mechanistically modeling the link between SIF and GPP (**Challenges 1-4**) will require the
243 combination of field campaigns covering full growing seasons, multiple species and stress
244 responses with detailed experimentation under highly controlled conditions, for example using
245 *Arabidopsis* mutants with altered photochemical properties^{9,73}. In particular, the development of
246 versatile field instrumentation and protocols for the estimation of APAR_g (**Challenge 1**), energy
247 distribution between PSII and PSI - including the ChlaF contribution from PSI - (**Challenge 2**), or
248 the quantification of alternative energy sinks (**Challenge 4**), is key to resolving the spatial and
249 temporal influences of these factors on SIF.

250 The synergistic use of complementary data streams can also help to constrain the modelling of
 251 photosynthesis and support SIF interpretation. For example, leaf and canopy reflectance data can
 252 inform us on the chlorophyll content in the leaf or the amount of leaves in the canopy⁷⁴, relating to
 253 $APAR_g$ (**Challenge 1**). In addition, reflectance data have been used to explore the regulatory
 254 dynamics of NPQ⁷⁵, which could contribute to resolving energy partitioning in PSII (**Challenge 3**).
 255 This approach is feasible due to the spectral change that accompanies the operation of the
 256 xanthophyll cycle⁷⁶ - by which violaxanthin is converted to antheraxanthin and zeaxanthin in a
 257 process that modulates NPQ^{38,77} - as well as the seasonal dynamics of leaf carotenoid and
 258 chlorophyll contents⁷⁸. These spectral changes, which have been captured by the photochemical
 259 reflectance index (PRI)^{77,79} or the Chlorophyll/Carotenoid Index (CCI)⁸⁰, are now being revisited
 260 and investigated in depth across the whole VIS-NIR region alongside with SIF dynamics^{71,76}.
 261 Clearly, as in the case of SIF, careful use of canopy and atmospheric RTMs will be needed to
 262 disentangle these subtle physiologically-induced reflectance changes from those of a dynamic
 263 background⁵⁵. In addition to synergies with spectral reflectance, use of thermal imaging⁸¹, radar⁸²,
 264 or multispectral laser scanning methods⁸³ offer interesting possibilities to constrain the carbon
 265 reactions of photosynthesis by providing independent information on plant water status (**Challenge**
 266 **4**). Likewise, leaf and ecosystem-level measurements of carbonyl sulfide (COS) uptake by
 267 vegetation can provide an independent source of information on stomatal conductance in vascular
 268 plants⁸⁴, which could be highly relevant for the development and validation of ecosystem-level SIF-
 269 GPP models.

270 Process-based and radiative transfer models are required to integrate physical and physiological
 271 mechanisms operating at different scales (**Challenge 7**), providing excellent frameworks for
 272 multidisciplinary collaborations to connect molecular-level with Earth-system processes. Clearly, as
 273 our mechanistic understanding of the connection between SIF and GPP increases (**Challenges 1-4**),
 274 so will the accuracy of process-based models. For example, the integration of the Farquhar-

275 Caemmerer-Berry⁸⁵ biochemical model of photosynthesis into dynamic land-surface models (e.g.,
276 ORCHIDEE⁸⁶ or BETHY⁸⁷) provides a gateway for assimilating satellite SIF data and improving
277 the accuracy of GPP estimations^{88,89}. In addition, SIF resides at the core of a new generation of
278 photosynthesis models that emphasize the light reactions^{45,90}. In the case of RTMs with established
279 SIF capabilities (Box 2), further improvements can be achieved by coupling with new techniques
280 measuring detailed 3D structures. Leaf RTMs would benefit from including variations in leaf
281 morphology, thylakoid structure, or the spectral signatures of PSI and PSII. The 3D
282 parameterization of canopy RTMs via lidar-based reconstruction methods^{91,92}, coupled to non-
283 imaging^{17,19} and imaging proximal/airborne SIF measurements (Fig. 3)^{93,94}, offers excellent
284 opportunities to integrate and resolve the diversity of factors that control SIF across space and time
285 (**Challenge 7**). Drone-based measurements could serve to investigate and model the impact of
286 atmospheric properties on SIF retrieval approaches, by hovering at different distances above the
287 target⁵⁴ (**Challenge 6**). Finally, less accurate but simpler alternative methods for separating the
288 physiological and structural influences on the SIF signal have been recently proposed based on the
289 theory of vegetation canopy near-infrared spectral invariants^{95,96}. Whether this or other correction
290 methods are applicable to canopy SIF acquisitions across scales, especially observations at very
291 high spatial resolutions (Fig. 3) should be further investigated.

292 Equally critical for the consistent interpretation of SIF is the establishment of a global network and
293 database of leaf and ecosystem-level SIF measurements covering different biomes, and supporting
294 model development as well as airborne and satellite calibration/validation activities. While regional
295 SIF networks are starting to emerge in North America, Europe, and Asia, their global connectivity
296 should be a priority to promote the adoption of standards for instrument calibrations and long-term
297 monitoring operations (Fig. 4).

298 Our roadmap for resolving the seven SIF challenges will only succeed through multidisciplinary
299 collaboration involving specialists from across molecular biology, plant physiology, optical physics

300 and remote sensing. Together, the characterization and modeling of the interplay between structural,
301 optical and functional dynamics of leaves and plant canopies, can turn our crops and forests into
302 observable field laboratories.

303 **Emerging and potential SIF applications**

304 Satellite SIF data are already providing new insight into photosynthetic dynamics at the global
305 scale^{97,98}. Likewise, with the advent of multiscale SIF measurements (Fig. 3), and as the remaining
306 challenges are overcome (Fig. 4), a new range of SIF applications unfolds across fields of
307 biochemistry, biophysics, ecology, ecophysiology, biogeochemistry, agriculture and forestry (Fig.
308 5). Equally important, the continuum of scales at which SIF can be measured provides a focal point
309 to promote and strengthen the interaction between research communities, from plant molecular
310 biology to Earth-system science. Here, we outline four examples of potential and emerging SIF
311 applications.

312 ***Spatial and 3D photosynthesis.*** Photosynthetic CO₂ assimilation can be measured using infrared
313 gas analyzers, either coupled to chambers or enclosures at the leaf, shoot, and whole-plant level⁹⁹,
314 or with a sonic anemometer at the ecosystem level using the eddy covariance approach¹⁰⁰. These
315 methods, however, lack detailed spatial information. Spatial measurements of photosynthesis, in
316 terms of photochemical rates of the light reactions, require the use of imaging systems that, to date,
317 have remained restricted to the scale of leaves or small-sized plants, e.g. PAM imaging methods¹⁰¹.
318 SIF measurements have potential to fill this scale gap. For example, SIF imaging (Fig. 3) could be
319 benchmarked with eddy-covariance methods to reveal the spatial variability of photosynthesis
320 within the footprint of ecosystem eddy covariance measurements, allowing us to investigate the
321 influence of microenvironment, understory and vertical canopy structure, or the interplay between
322 biological and functional diversity within the ecosystem. Likewise, SIF imaging could be applied to

323 resolve photosynthesis dynamics in 3D, helping to advance our understanding of the interaction
324 between plant structure and function^{102,103}.

325 ***Physiological phenotyping and pre-visual stress detection.*** Spatial and temporal variations in plant
326 morphological traits (e.g., canopy height, leaf area, and plant growth) have been widely used as
327 markers for field phenotypic variability and to investigate long-term plant stress responses.
328 However, these traits are insufficiently responsive to rapid plant physiological changes. This makes
329 them ill-suited for physiological phenotyping (i.e. breeding plant phenotypes displaying specific
330 physiological responses to the environment), or pre-visual stress detection and subsequent
331 optimization of water, pesticide and fertilizer use. The current phenotyping focus has, therefore,
332 shifted towards measurements in the visible and infrared spectral ranges, where reflectance changes
333 can be associated with specific physiological and biochemical traits¹⁰⁴ or used for early-stress
334 detection¹⁰⁵. In this context, emerging SIF imaging systems have already provided promising results
335 for applications in precision agriculture and detection of pest infestations^{93,106}. In the near future,
336 these methods could also support precision forestry applications related to seedling production or
337 tree-scale forest management.

338 ***Functional plant diversity and spatial ecology.*** Functional diversity is a fundamental component of
339 the biodiversity concept¹⁰⁷. As a global network for monitoring biodiversity through remotely
340 sensed plant functional traits is being developed¹⁰⁸, SIF could become one of the new essential
341 variables for mapping functional diversity across ecosystem and landscape scales, given the wide
342 range of biochemical and physiological factors that SIF is sensitive to (Fig. 2) in relation to plant
343 productivity. For example, SIF has been shown to convey spatial information on leaf mass and
344 chlorophyll content¹⁰⁹, and other functional plant traits¹¹⁰ in various forest ecosystems. Additionally
345 and importantly, the combination of high-resolution structural, spectral and SIF data is potentially
346 the only viable option to investigate ecosystem functions that have remained hidden from our
347 observational abilities, such as photosynthetic phenology in evergreen forests¹⁸, cryptogamic

348 biocrusts¹¹¹ and spatially fragmented Antarctic mosses⁷⁴. Together with spatial photosynthesis, SIF
349 could also offer unique opportunities for studies in spatial ecology^{111,112}, where plant environmental
350 responses and biotic interactions could leave their imprint on SIF.

351 ***Carbon and water cycle studies.*** The carbon and water cycles of terrestrial ecosystems are
352 intricately connected via stomatal regulation and total leaf area. Because both canopy
353 evapotranspiration and canopy SIF dynamics are strongly controlled by leaf area, and since ChlaF
354 can also decrease with stomatal closure - via increased NPQ in response to water stress^{113, 114}; tower
355 and satellite SIF have been preliminarily used to investigate canopy conductance and plant
356 transpiration^{115,116}. No doubt, better constraints on transpiration and photosynthetic dynamics in
357 land-surface models will be achieved as the mechanistic basis of SIF is elucidated across scales
358 (Challenges 1-7), and the integration of SIF with other remote sensing datasets increases, such as
359 land-surface temperature¹¹⁵, surface soil moisture⁸⁹, radar-measured vegetation optical depth
360 characterizing canopy structure and water content¹¹⁷, or column-averaged atmospheric CO₂⁹². New
361 knowledge of photosynthesis at the ecosystem and regional scales will bring further insight into the
362 large-scale interactions between environmental drivers and plant productivity, and feedbacks
363 between the biosphere and atmosphere.

364 **Concluding remarks**

365 The SIF signal gathers a wealth of physiological, biochemical, and structural information as it
366 travels from the photosystems to the top of canopy and beyond (Fig. 2). This can leave the
367 impression that SIF is, to use the classic analogy, the ‘Swiss Army Knife’ of photosynthesis
368 measurements. Critically, the variation in SIF caused by physical and biotic factors is entangled in
369 the spatiotemporal domain, and our capacity to disentangle it into useful informative components
370 requires further attention. Historically, photosynthesis research has been a multidisciplinary
371 endeavor, with breakthroughs in the 20th century emerging from collaboration between chemists,

372 biologists and physicists. We are now entering a new era of multiscale observations of
373 photosynthesis which requires the interdisciplinary research environment to flourish further, this
374 time to resolve the mechanistic connection between SIF and GPP and to scale it across space and
375 time. The technology to measure SIF is developing at a faster pace than our capacity to interpret the
376 acquired data. With the challenges, roadmap and unfolding opportunities introduced here we hope
377 to encourage more scientists to join the multidisciplinary quest to reveal the true potential of SIF
378 observation.

379

380 **References**

- 381 1. Genty, B., Wonders, J. & Baker, N. R. Non-photochemical quenching of Fo in leaves is
382 emission wavelength dependent: consequences for quenching analysis and its interpretation.
383 *Photosynth. Res.* **26**, 133-139 (1990).
- 384 2. Franck, F., Juneau, P. & Popovic, R. Resolution of the photosystem I and photosystem II
385 contributions to chlorophyll fluorescence of intact leaves at room temperature. *Biochim.*
386 *Biophys. Acta-Bioenergetics* **1556**, 239-246 (2002).
- 387 3. Neubauer, C. & Schreiber, U. The polyphasic rise of chlorophyll fluorescence upon onset of
388 strong continuous illumination: I. Saturation characteristics and partial control by the
389 photosystem II acceptor side. *Zeitschrift für Naturforschung C* **42**, 1246-1254 (1987).
- 390 4. Strasser, R. J., Tsimilli-Michael, M. & Srivastava, A. Analysis of the chlorophyll a
391 fluorescence transient. In: Papageorgiou G.C., Govindjee (eds) Chlorophyll a Fluorescence.
392 Advances in Photosynthesis and Respiration, vol 19. Springer, Dordrecht (2004).
- 393 5. Schreiber, U., Schliwa, U. & Bilger, W. Continuous recording of photochemical and non-
394 photochemical chlorophyll fluorescence quenching with a new type of modulation
395 fluorometer. *Photosynth. Res.* **10**, 51-62 (1986).

6. Maxwell, K. & Johnson, G. N. Chlorophyll fluorescence—a practical guide. *J. Exp. Bot.* **51**, 659-668 (2000).
7. Govindjee, E. 63 Years since Kautsky-chlorophyll-a fluorescence. *Aust. J. Plant Physiol.* **22**, 131-160 (1995).
8. Porcar-Castell, A. *et al.* Linking chlorophyll a fluorescence to photosynthesis for remote sensing applications: mechanisms and challenges. *J. Exp. Bot.* **65**, 4065-4095 (2014).
9. Tikkanen, M., Rantala, S., Grieco, M. & Aro, E. Comparative analysis of mutant plants impaired in the main regulatory mechanisms of photosynthetic light reactions-From biophysical measurements to molecular mechanisms. *Plant Physiol. Biochem.* **112**, 290-301 (2017).
10. Kolber, Z. *et al.* Measuring photosynthetic parameters at a distance: laser induced fluorescence transient (LIFT) method for remote measurements of photosynthesis in terrestrial vegetation. *Photosynth. Res.* **84**, 121-129 (2005).
11. Keller, B. *et al.* Genotype specific photosynthesis x environment interactions captured by automated fluorescence canopy scans over two fluctuating growing seasons. *Front. Plant Sci.* **10**, 1482 (2019).
12. Mohammed, G. H. *et al.* Remote sensing of solar-induced chlorophyll fluorescence (SIF) in vegetation: 50 years of progress. *Remote Sens. Environ.* **231**, 111177 (2019).
13. Evain, S., Camenen, L. & Moya, I. Three-channel detector for remote sensing of chlorophyll fluorescence and reflectance from vegetation. In: M. Leroy (ed.), 8th International symposium: physical measurements and signatures in remote sensing, pp. 395-400. Aussois, CNES, France (2001).
14. Louis, J. *et al.* Remote sensing of sunlight-induced chlorophyll fluorescence and reflectance of Scots pine in the boreal forest during spring recovery. *Remote Sens. Environ.* **96**, 37-48 (2005).

- 421 15. Guanter, L. *et al.* Estimation of solar-induced vegetation fluorescence from space
422 measurements. *Geophys. Res. Lett.* **34** (2007).
- 423 16. Aasen, H. *et al.* Sun-induced chlorophyll fluorescence II: review of passive measurement
424 setups, protocols, and their application at the leaf to canopy level. *Remote Sensing* **11**, 927
425 (2019).
- 426 17. Yang, X. *et al.* Solar-induced chlorophyll fluorescence that correlates with canopy
427 photosynthesis on diurnal and seasonal scales in a temperate deciduous forest. *Geophys.*
428 *Res. Lett.* **42**, 2977-2987 (2015).
- 429 18. Magney, T. S. *et al.* Mechanistic evidence for tracking the seasonality of photosynthesis
430 with solar-induced fluorescence. *PNAS* **116**, 11640-11645 (2019).
- 431 19. Bendig, J., Malenovsky, Z., Gautam, D. & Lucieer, A. Solar-Induced Chlorophyll
432 Fluorescence Measured From an Unmanned Aircraft System: Sensor Etaloning and Platform
433 Motion Correction. *IEEE Trans. Geosci. Remote Sens.* (2019).
- 434 20. Vargas, J. Q. *et al.* Unmanned aerial systems (UAS)-based methods for solar induced
435 chlorophyll fluorescence (SIF) retrieval with non-imaging spectrometers: state of the art.
436 *Remote Sens.* **12**, 1624 (2020).
- 437 21. Rascher, U. *et al.* Sun-induced fluorescence—a new probe of photosynthesis: First maps from
438 the imaging spectrometer HyPlant. *Global Change Biol.* **21**, 4673-4684 (2015).
- 439 22. Frankenberg, C. *et al.* The Chlorophyll Fluorescence Imaging Spectrometer (CFIS),
440 mapping far red fluorescence from aircraft. *Remote Sens. Environ.* **217**, 523-536 (2018).
- 441 23. Frankenberg, C. *et al.* New global observations of the terrestrial carbon cycle from GOSAT:
442 Patterns of plant fluorescence with gross primary productivity. *Geophys. Res. Lett.* **38**
443 (2011).
- 444 24. Köhler, P. *et al.* Global Retrievals of Solar-Induced Chlorophyll Fluorescence at Red
445 Wavelengths With TROPOMI. *Geophys. Res. Lett.* **47**, e2020GL087541 (2020).

25. Drusch, M. *et al.* The fluorescence explorer mission concept - ESA's earth explorer 8. *IEEE Trans. Geosci. Remote Sens.* **55**, 1273-1284 (2016).
26. Olascoaga, B., Mac Arthur, A., Atherton, J. & Porcar-Castell, A. A comparison of methods to estimate photosynthetic light absorption in leaves with contrasting morphology. *Tree Physiol.* **36**, 368-379 (2016).
27. Zhang, Z. *et al.* Assessing bi-directional effects on the diurnal cycle of measured solar-induced chlorophyll fluorescence in crop canopies. *Agric. For. Meteorol.* **295**, 108147 (2020).
28. Bittner, T., Irrgang, K., Renger, G. & Wasielewski, M. R. Ultrafast excitation energy transfer and exciton-exciton annihilation processes in isolated light harvesting complexes of photosystem II (LHC II) from spinach. *J. Phys. Chem.* **98**, 11821-11826 (1994).
29. Kalaji, H. M. *et al.* Frequently asked questions about chlorophyll fluorescence, the sequel. *Photosynth. Res.* **132**, 13-66 (2017).
30. Genty, B., Briantais, J. & Baker, N. R. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta-General Subjects* **990**, 87-92 (1989).
31. Anderson, J. M., Chow, W. S. & Goodchild, D. J. Thylakoid membrane organisation in sun/shade acclimation. *Funct. Plant Biol.* **15**, 11-26 (1988).
32. Ballottari, M., Dall'Osto, L., Morosinotto, T. & Bassi, R. Contrasting behavior of higher plant photosystem I and II antenna systems during acclimation. *J. Biol. Chem.* **282**, 8947-8958 (2007).
33. Schreiber, U., Klughammer, C. & Kolbowski, J. Assessment of wavelength-dependent parameters of photosynthetic electron transport with a new type of multi-color PAM chlorophyll fluorometer. *Photosynth. Res.* **113**, 127-144 (2012).

34. Laisk, A. *et al.* A computer-operated routine of gas exchange and optical measurements to
diagnose photosynthetic apparatus in leaves. *Plant, Cell Environ.* **25**, 923-943 (2002).
35. Pfündel, E. Estimating the contribution of photosystem I to total leaf chlorophyll
fluorescence. *Photosynthesis Res.* **56**, 185-195 (1998).
36. Peterson, R. B. *et al.* Fluorescence F_o of photosystems II and I in developing C3 and C4
leaves, and implications on regulation of excitation balance. *Photosynth. Res.* **122**, 41-56
37. Pfündel, E. E. Simultaneously measuring pulse-amplitude-modulated (PAM) chlorophyll
fluorescence of leaves at wavelengths shorter and longer than 700 nm. *Photosynth. Res.*, 1-
14 (2021).
38. Demmig-Adams, B. & Adams III, W. W. Photoprotection in an ecological context: the
remarkable complexity of thermal energy dissipation. *New Phytol.* **172**, 11-21 (2006).
39. Porcar-Castell, A. A high-resolution portrait of the annual dynamics of photochemical and
non-photochemical quenching in needles of *Pinus sylvestris*. *Physiol. Plant.* **143**, 139-153
(2011).
40. Van der Tol, C., Berry, J. A., Campbell, P. & Rascher, U. Models of fluorescence and
photosynthesis for interpreting measurements of solar-induced chlorophyll fluorescence. *J.*
Geophys. Res.: Biogeosciences **119**, 2312-2327 (2014).
41. Springer, K. R., Wang, R. & Gamon, J. A. Parallel seasonal patterns of photosynthesis,
fluorescence, and reflectance indices in boreal trees. *Remote Sens.* **9**, 691 (2017).
42. Zhang, C. *et al.* Do all chlorophyll fluorescence emission wavelengths capture the spring
recovery of photosynthesis in boreal evergreen foliage? *Plant, Cell Environ.* **42**, 3264-3279
(2019).
43. Ensminger, I. *et al.* Intermittent low temperatures constrain spring recovery of
photosynthesis in boreal Scots pine forests. *Glob. Change Biol.* **10**, 995-1008 (2004).

44. Verhoeven, A. Sustained energy dissipation in winter evergreens. *New Phytol.* **201**, 57-65 (2014).
45. Gu, L., Han, J., Wood, J. D., Chang, C. Y. & Sun, Y. Sun-induced Chl fluorescence and its importance for biophysical modeling of photosynthesis based on light reactions. *New Phytol.* **223**, 1179-1191 (2019).
46. Raczka, B. *et al.* Sustained nonphotochemical quenching shapes the seasonal pattern of solar-induced fluorescence at a high-elevation evergreen forest. *J. Geophys. Res.: Biogeosciences* **124**, 2005-2020 (2019).
47. Nixon, P. J. Chlororespiration. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **355**, 1541-1547 (2000).
48. Ogren, W. L. Photorespiration: pathways, regulation, and modification. *Annu. Rev. Plant Physiol.* **35**, 415-442 (1984).
49. Asada, K. The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons. *Annu. Rev. Plant Biol.* **50**, 601-639 (1999).
50. Morfopoulos, C. *et al.* A model of plant isoprene emission based on available reducing power captures responses to atmospheric CO₂. *New Phytol.* **203**, 125-139 (2014).
51. Maseyk, K., Lin, T., Cochavi, A., Schwartz, A. & Yakir, D. Quantification of leaf-scale light energy allocation and photoprotection processes in a Mediterranean pine forest under extensive seasonal drought. *Tree Physiol.* **39**, 1767-1782 (2019).
52. Migliavacca, M. *et al.* Plant functional traits and canopy structure control the relationship between photosynthetic CO₂ uptake and far-red sun-induced fluorescence in a Mediterranean grassland under different nutrient availability. *New Phytol.* **214**, 1078-1091 (2017).
53. Kallel, A. FluLCVRT: Reflectance and fluorescence of leaf and canopy modeling based on Monte Carlo vector radiative transfer simulation. *J. Quant. Spectrosc. Radiat. Transf.* **253**, 107183 (2020).

54. Sabater, N. *et al.* Compensation of oxygen transmittance effects for proximal sensing retrieval of canopy-leaving sun-induced chlorophyll fluorescence. *Remote Sens.* **10**, 1551 (2018).
55. Sabater, N., Kolmonen, P., Van Wittenberghe, S., Arola, A. & Moreno, J. Challenges in the atmospheric characterization for the retrieval of spectrally resolved fluorescence and PRI region dynamics from space. *Remote Sens. Environ.* **254**, 112226 (2021).
56. Iermak, I., Vink, J., Bader, A. N., Wientjes, E. & van Amerongen, H. Visualizing heterogeneity of photosynthetic properties of plant leaves with two-photon fluorescence lifetime imaging microscopy. *Biochim. Biophys. Acta-Bioenergetics* **1857**, 1473-1478 (2016).
57. Romero, J. M., Cordon, G. B. & Lagorio, M. G. Modeling re-absorption of fluorescence from the leaf to the canopy level. *Remote Sens. Environ.* **204**, 138-146 (2018).
58. Magney, T. S. *et al.* Disentangling changes in the spectral shape of chlorophyll fluorescence: Implications for remote sensing of photosynthesis. *J. Geophys. Res.: Biogeosciences* **124**, 1491-1507 (2019).
59. Murchie, E. H. *et al.* Measuring the dynamic photosynthome. *Ann. Bot.* **122**, 207-220 (2018).
60. Magney, T. S., Barnes, M. L. & Yang, X. On the covariation of chlorophyll fluorescence and photosynthesis across scales. *Geophys. Res. Lett.* **47**, e2020GL091098 (2020).
61. Yang, P., van der Tol, C., Campbell, P. K. & Middleton, E. M. Unraveling the physical and physiological basis for the solar-induced chlorophyll fluorescence and photosynthesis relationship using continuous leaf and canopy measurements of a corn crop. *Biogeosciences* **18**, 441-465 (2021).
62. Liu, X. *et al.* Downscaling of solar-induced chlorophyll fluorescence from canopy level to photosystem level using a random forest model. *Remote Sens. Environ.* **231**, 110772 (2019).

63. Joiner, J. *et al.* Systematic Orbital Geometry-Dependent Variations in Satellite Solar-Induced Fluorescence (SIF) Retrievals. *Remote Sens.* **12**, 2346 (2020).
64. Dechant, B. *et al.* Canopy structure explains the relationship between photosynthesis and sun-induced chlorophyll fluorescence in crops. *Remote Sens. Environ.* **241**, 111733 (2020).
65. He, L. *et al.* From the Ground to Space: Using Solar-Induced Chlorophyll Fluorescence to Estimate Crop Productivity. *Geophys. Res. Lett.* **47**, e2020GL087474 (2020).
66. Ač, A. *et al.* Meta-analysis assessing potential of steady-state chlorophyll fluorescence for remote sensing detection of plant water, temperature and nitrogen stress. *Remote Sens. Environ.* **168**, 420-436 (2015).
67. Wohlfahrt, G. *et al.* Sun-induced fluorescence and gross primary productivity during a heat wave. *Sci. Rep.* **8**, 1-9 (2018).
68. Van Wittenberghe, S., Alonso, L., Verrelst, J., Moreno, J. & Samson, R. Bidirectional sun-induced chlorophyll fluorescence emission is influenced by leaf structure and light scattering properties: A bottom-up approach. *Remote Sens. Environ.* **158**, 169-179 (2015).
69. Magney, T. S. *et al.* Connecting active to passive fluorescence with photosynthesis: A method for evaluating remote sensing measurements of Chl fluorescence. *New Phytol.* **215**, 1594-1608 (2017).
70. Rajewicz, P. A., Atherton, J., Alonso, L. & Porcar-Castell, A. Leaf-level spectral fluorescence measurements: comparing methodologies for broadleaves and needles. *Remote Sens.* **11**, 532 (2019).
71. Van Wittenberghe, S., Alonso, L., Malenovský, Z. & Moreno, J. In vivo photoprotection mechanisms observed from leaf spectral absorbance changes showing VIS-NIR slow-induced conformational pigment bed changes. *Photosynth. Res.* **142**, 283-305 (2019).

72. Meeker, E. W., Magney, T. S., Bambach, N., Momayyezi, M. & McElrone, A. J. Modification of a gas exchange system to measure active and passive chlorophyll fluorescence simultaneously under field conditions. *AoB Plants* **13**, plaa066 (2021).
73. Acebron, K. *et al.* Diurnal dynamics of nonphotochemical quenching in *Arabidopsis npq* mutants assessed by solar-induced fluorescence and reflectance measurements in the field. *New Phytol.* (2020).
74. Malenovský, Z., Lucieer, A., King, D. H., Turnbull, J. D. & Robinson, S. A. Unmanned aircraft system advances health mapping of fragile polar vegetation. *Methods Ecol. Evol.* **8**, 1842-1857 (2017).
75. Atherton, J., Nichol, C. J. & Porcar-Castell, A. Using spectral chlorophyll fluorescence and the photochemical reflectance index to predict physiological dynamics. *Remote Sens. Environ.* **176**, 17-30 (2016).
76. Van Wittenberghe, S. *et al.* Combined dynamics of the 500–600 nm leaf absorption and chlorophyll fluorescence changes in vivo: Evidence for the multifunctional energy quenching role of xanthophylls. *Biochim. Biophys. Acta-Bioenergetics* **1862**, 148351 (2021).
77. Gamon, J. A. *et al.* Remote sensing of the xanthophyll cycle and chlorophyll fluorescence in sunflower leaves and canopies. *Oecologia* **85**, 1-7 (1990).
78. Filella, I. *et al.* PRI assessment of long-term changes in carotenoids/chlorophyll ratio and short-term changes in de-epoxidation state of the xanthophyll cycle. *Int. J. Remote Sens.* **30**, 4443-4455 (2009).
79. Peñuelas, J., Filella, I. & Gamon, J. A. Assessment of photosynthetic radiation-use efficiency with spectral reflectance. *New Phytol.* **131**, 291-296 (1995).
80. Gamon, J. A. *et al.* A remotely sensed pigment index reveals photosynthetic phenology in evergreen conifers. *PNAS* **113**, 13087-13092 (2016).

81. Costa, J. M., Grant, O. M. & Chaves, M. M. Thermography to explore plant-environment interactions. *J. Exp. Bot.* **64**, 3937-3949 (2013).
82. Konings, A. G., Rao, K. & Steele-Dunne, S. C. Macro to micro: microwave remote sensing of plant water content for physiology and ecology. *New Phytol.* **223**, 1166-1172 (2019).
83. Junttila, S. *et al.* Terrestrial laser scanning intensity captures diurnal variation in leaf water potential. *Remote Sens. Environ.* **255**, 112274 (2021).
84. Whelan, M. E. *et al.* Two Scientific Communities Striving for a Common Cause: innovations in carbon cycle science. *Bull. Am. Meteorol. Soc.* (2020).
85. Farquhar, G. D., von Caemmerer, S. v. & Berry, J. A. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* **149**, 78-90 (1980).
86. Bacour, C. *et al.* Improving estimates of gross primary productivity by assimilating solar-induced fluorescence satellite retrievals in a terrestrial biosphere model using a process-based SIF model. *J. Geophys. Res.: Biogeosciences* **124**, 3281-3306 (2019).
87. Norton, A. J. *et al.* Estimating global gross primary productivity using chlorophyll fluorescence and a data assimilation system with the BETHY-SCOPE model. *Biogeosciences* **16**, 3069-3093 (2019).
88. Thum, T. *et al.* Modelling sun-induced fluorescence and photosynthesis with a land surface model at local and regional scales in northern Europe. *Biogeosciences* **14**, 1969-1987 (2017).
89. Qiu, B., Chen, J. M., Ju, W., Zhang, Q. & Zhang, Y. Simulating emission and scattering of solar-induced chlorophyll fluorescence at far-red band in global vegetation with different canopy structures. *Remote Sens. Environ.* **233**, 111373 (2019).
90. Johnson, J. E. & Berry, J. A. The role of Cytochrome b₆ f in the control of steady-state photosynthesis: a conceptual and quantitative model. *Photosynth. Res.*, 1-36 (2021).

91. Janoutová, R. *et al.* Influence of 3D spruce tree representation on accuracy of airborne and satellite forest reflectance simulated in DART. *Forests* **10**, 292 (2019).
92. Liu, W. *et al.* Simulating solar-induced chlorophyll fluorescence in a boreal forest stand reconstructed from terrestrial laser scanning measurements. *Remote Sens. Environ.* **232**, 111274 (2019).
93. Pinto, F. *et al.* Sun-induced chlorophyll fluorescence from high-resolution imaging spectroscopy data to quantify spatio-temporal patterns of photosynthetic function in crop canopies. *Plant, Cell Environ.* **39**, 1500-1512 (2016).
94. Siegmann, B. *et al.* The high-performance airborne imaging spectrometer HyPlant—From raw images to top-of-canopy reflectance and fluorescence products: Introduction of an automatized processing chain. *Remote Sens.* **11**, 2760 (2019).
95. Yang, P., van der Tol, C., Campbell, P. K. & Middleton, E. M. Fluorescence Correction Vegetation Index (FCVI): A physically based reflectance index to separate physiological and non-physiological information in far-red sun-induced chlorophyll fluorescence. *Remote Sens. Environ.* **240**, 111676 (2020).
96. Zeng, Y. *et al.* A radiative transfer model for solar induced fluorescence using spectral invariants theory. *Remote Sens. Environ.* **240**, 111678 (2020).
97. Green, J. K. *et al.* Large influence of soil moisture on long-term terrestrial carbon uptake. *Nature* **565**, 476 (2019).
98. Wang, S. *et al.* Urban– rural gradients reveal joint control of elevated CO₂ and temperature on extended photosynthetic seasons. *Nat. Ecol. Evo.* **3**, 1076-1085 (2019).
99. Long, S. P., Farage, P. K. & Garcia, R. L. Measurement of leaf and canopy photosynthetic CO₂ exchange in the field. *J. Exp. Bot.* **47**, 1629-1642 (1996).

100. Baldocchi, D. D. Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future. *Glob. Change Biol.* **9**, 479-492 (2003).
101. Kaiser, Y. I., Menegat, A. & Gerhards, R. Chlorophyll fluorescence imaging: a new method for rapid detection of herbicide resistance in *Alopecurus myosuroides*. *Weed Res.* **53**, 399-406 (2013).
102. Sievänen, R., Godin, C., DeJong, T. M. & Nikinmaa, E. Functional–structural plant models: a growing paradigm for plant studies. *Ann. Bot.* **114**, 599-603 (2014).
103. Damm, A., Paul-Limoges, E., Kükenbrink, D., Bachofen, C. & Morsdorf, F. Remote sensing of forest gas exchange: Considerations derived from a tomographic perspective. *Glob. Change Biol.* **26**, 2717-2727 (2020).
104. Ensminger, I. Fast track diagnostics: Hyperspectral reflectance differentiates disease from drought stress in trees. *Tree Physiol.* **40**, 1143-1146 (2020).
105. Mutka, A. M. & Bart, R. S. Image-based phenotyping of plant disease symptoms. *Frontiers in plant science* **5**, 734 (2015).
106. Zarco-Tejada, P. J. *et al.* Previsual symptoms of *Xylella fastidiosa* infection revealed in spectral plant-trait alterations. *Nat. Plants* **4**, 432-439 (2018).
107. Díaz, S. & Cabido, M. Vive la différence: plant functional diversity matters to ecosystem processes. *Trends Ecol. Evol.* **16**, 646-655 (2001).
108. Skidmore, A. K. *et al.* Environmental science: Agree on biodiversity metrics to track from space. *Nat. News* **523**, 403 (2015).
109. Tagliabue, G. *et al.* Sun–induced fluorescence heterogeneity as a measure of functional diversity. *Remote Sens. Environ.* **247**, 111934 (2020).

110. Pacheco-Labrador, J. *et al.* Multiple-constraint inversion of SCOPE. Evaluating the potential of GPP and SIF for the retrieval of plant functional traits. *Remote Sens. Environ.* **234**, 111362 (2019).
111. Smith, W. K. *et al.* Remote sensing of dryland ecosystem structure and function: Progress, challenges, and opportunities. *Remote Sens. Environ.* **233**, 111401 (2019).
112. Kellner, J. R., Albert, L. P., Burley, J. T. & Cushman, K. C. The case for remote sensing of individual plants. *Am. J. Bot.* **106**, 1139-1142 (2019).
113. Flexas, J. *et al.* Steady-state chlorophyll fluorescence (Fs) measurements as a tool to follow variations of net CO₂ assimilation and stomatal conductance during water-stress in C₃ plants. *Physiol. Plant.* **114**, 231-240 (2002).
114. Marrs, J. K. *et al.* Solar-induced fluorescence does not track photosynthetic carbon assimilation following induced stomatal closure. *Geophys. Res. Lett.* **47**, e2020GL087956 (2020).
115. Maes, W. H. *et al.* Sun-induced fluorescence closely linked to ecosystem transpiration as evidenced by satellite data and radiative transfer models. *Remote Sens. Environ.* **249**, 112030 (2020).
116. Shan, N. *et al.* A model for estimating transpiration from remotely sensed solar-induced chlorophyll fluorescence. *Remote Sens. Environ.* **252**, 112134 (2021).
117. Liu, W. *et al.* Simulating solar-induced chlorophyll fluorescence in a boreal forest stand reconstructed from terrestrial laser scanning measurements. *Remote Sens. Environ.* **232**, 111274 (2019).
118. Albert, L. P. *et al.* Stray light characterization in a high-resolution imaging spectrometer designed for solar-induced fluorescence. Proc. SPIE 10986, Algorithms, Technologies, and Applications for Multispectral and Hyperspectral Imagery XXV, 109860G (2019).

119. Meroni, M. *et al.* Remote sensing of solar-induced chlorophyll fluorescence: Review of methods and applications. *Remote Sens. Environ.* **113**, 2037-2051 (2009).
120. Cendrero-Mateo, M. P. *et al.* Sun-induced chlorophyll fluorescence III: Benchmarking retrieval methods and sensor characteristics for proximal sensing. *Remote Sens.* **11**, 962 (2019).
121. Vilfan, N. *et al.* Extending Fluspect to simulate xanthophyll driven leaf reflectance dynamics. *Remote Sens. Environ.* **211**, 345-356 (2018).
122. Yang, P., Prikaziuk, E., Verhoef, W. & van der Tol, C. SCOPE 2.0: A model to simulate vegetated land surface fluxes and satellite signals. *Geosci. Model Dev. Discuss.*, 1-26 (2020).
123. Gastellu-Etchegorry, J. *et al.* DART: recent advances in remote sensing data modeling with atmosphere, polarization, and chlorophyll fluorescence. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **10**, 2640-2649 (2017).

Corresponding author: Albert Porcar-Castell (joan.porcar@helsinki.fi)

Acknowledgements: This perspective idea originated during the Fluorescence Across Space and Time (FAST) Workshop, which took place in Hyytiälä Forestry Research Station (SMEARII, Finland) during February 2019. We thank the following participants for active discussions during the workshop: Juliane Bendig, Kukka-Maaria Erkkilä, Noda Hibiki, Laura V. Junker-Frohn, Valentyna Kuznetsova, Hannakaisa Lindqvist, Paul Näthe, Jaakko Oivukkamaki, Neus Sabater, Twinkle Solanki, Tea Thum, Shan Xu and Chao Zhang. We also thank Barry Osmond and Josep Peñuelas for valuable comments to the manuscript, to Nuria Altimir for improving graphic design of Figs 1 and 5, and to Bastian Siegmann for the preparation of the HyPlant image in Fig. 3. The

709 Academy of Finland (Project # 288039 and 319211) is acknowledged for the financial support. ZM
710 was supported by the Australian Research Council (FT160100477), TM was supported by the
711 National Aeronautics and Space Administration (80NSSC19M0129), and SVW was supported by
712 the Generalitat Valenciana and the European Social Fund (APOSTD/2018/162). Headwall SIF
713 images from LPA and JRK were supported by grants from the Institute at Brown for Environment
714 and Society at Brown University.

715 **Author contributions:** APC conceived the original idea and wrote the manuscript with ZM, TM,
716 BL, SVW, BFM, FM, YZ, KM with comments and contributions from all co-authors. In addition,
717 these authors had special contribution to the following parts: Fig.1 (APC, ZM and SVW), Fig.2
718 (APC, BFM, TM and SVW), Fig. 3 (LPA, UR and JRK), Fig. 4. (APC, ZM, UR, BFM), Fig. 5
719 (JIGP, JA, ZM, IE), Box 1 (TM, APC), Box 2 (ZM, APC), Supplementary information (ZM, FZ).

720

721

722

723

724

725

726

727

728

729

730 **Figure Legends**

731 **Figure 1. From incoming radiation to observed SIF and photosynthesis: mechanistic**

732 **challenges.** Solar radiation reaching the top of the atmosphere (TOA) is partly absorbed and
733 scattered by atmospheric gases and particles, decreasing its intensity as it reaches the bottom of the
734 atmosphere (BOA), generating specific absorption features. Part of the radiation is absorbed by
735 photosynthetic pigments in vegetation and leaves ($fAPAR_g$) (**Challenge 1**), associated with either
736 photosystem I (PSI) or photosystem II (PSII), which contribute with differential dynamics and
737 spectral properties to overall SIF emission (**Challenge 2**). Within each photosystem, energy is
738 further partitioned into three dynamic processes (**Challenge 3**): i) photochemistry (leading mainly
739 to linear (LET) or cyclic (CET) electron transport, the latter involving PSI only), ii) thermal energy
740 dissipation, and iii) ChlaF. Photosynthetic energy (expressed for simplicity in terms of NADPH
741 equivalents) is further partitioned between alternative energy sinks and gross photosynthesis (A_G),
742 and again between gross primary productivity (GPP) and photorespiration (P_R), with dynamics that
743 are not necessarily seen by SIF (**Challenge 4**). Notably, because it is only possible to measure the
744 net flux of CO_2 from a leaf or ecosystem, i.e. net photosynthesis or net primary productivity (NPP),
745 the rate of daytime respiration (R_D) must be known or estimated. In turn, because emitted ChlaF
746 overlaps with the absorption spectra of leaves and plant canopies, some SIF photons - especially
747 those in the red wavelengths - are re-absorbed within the canopy (**Challenge 5**). Emitted ChlaF is
748 further scattered and absorbed by aerosols and gases in the atmosphere (**Challenge 6**).

749

750 **Figure 2. The connection between SIF and GPP across space and time.** The relationship
751 between SIF and GPP is affected by multiple factors as we move across spatial and temporal scales.
752 Some factors exert a similar effect on SIF and GPP, keeping them positively correlated - we call
753 these couplers. Other factors differentially affect SIF and GPP - we call these decouplers. Factors

driving the dynamics of NPQ and APAR will tend to keep SIF and GPP coupled both across space
 and time, whereas factors adding variation to the energy partitioning between ChlF and GPP, or
 influencing the reabsorption of ChlF, will tend to decouple SIF from GPP (see examples in the
 figure). Note how the shape of the ChlF spectrum (“**Spectral dynamics**”) changes across scales in
 response to reabsorption within the chloroplast, leaf and canopy, measurable as SIF only within
 discrete wavelengths at the canopy and ecosystem levels (Box 1). Equally important to our
 understanding of the spatial context of the factors that couple/decouple SIF to GPP is understanding
 their temporal range of action (lower panels). For example, the rapid (**second/minute**) decrease in
 ChlaF upon saturating illumination of dark acclimated leaves reflects the dynamics of NPQ⁷⁶.
 Similar dynamics can be seen under natural conditions at the **diurnal/seasonal** scale in Scots pine
 needles, as the quantum yield of fluorescence (ΦF) responds to PQ and NPQ (redrawn from Porcar-
 Castell³⁹). Here, SIF was estimated for illustrative purposes as $SIF (r.u.) = PAR \times 0.8 \times 0.5 \times \Phi F$,
 where 0.8 and 0.5 are estimates for $fAPAR_g$ and the fraction of radiation absorbed by PSII.
 Likewise, **interannual** dynamics at the regional-to-local scales²⁴ can reflect changes in canopy
 structure, physiological stress responses or other functional traits. Ultimately, the challenge of
 integrating and disentangling the impact of these couplers/decouplers across space, time, species
 and plant stress responses remains (**Challenge 7**).

771

Figure 3. State-of-the-art SIF imaging methods allow for the observation of SIF across a continuum
 of scales: from the leaf-to-individual (top row) to the individual-to-landscape (bottom row). Panel A
 shows an RGB image of a senescing maple tree next to an oak tree with green leaves. Panel B
 shows the SIF image of the same trees retrieved in the O2A band at 760 nm (SIF760) using a
 commercial, off-the-shelf imaging spectrometer¹¹⁸ mounted on a tripod some meters away and after
 applying a filter to exclude non-vegetation pixels (pixels with an normalized difference vegetation
 index (NDVI) < 0.65). As expected, the green and photosynthetically active oak emitted SIF at

779 higher magnitude (Panel C) than the senescing maple. Similarly, panels D-E present an airborne
780 RGB and SIF760 map obtained with data from the HyPlant sensor collected at an altitude of 680 m
781 above ground⁹⁴. The scene shows several plots within an experimental apple tree plantation at the
782 agricultural research site Campus Klein-Altendorf (University of Bonn, Germany), where apple tree
783 varieties of different ages were growing in a typical row structure. Single tree crowns were
784 segmented by overlaying the SIF images with a 3D surface map and all pixels that were related to a
785 background signal (defined as ground level + 30 cm) were excluded. The image visualizes the
786 signal of individual trees, where each pixel corresponds to an area of 1x1 meters and thus the small
787 clusters represent the signal of an individual tree.

788

789 **Figure 4.** A roadmap towards the standardized interpretation of SIF. The critical steps, data sources
790 and methods that will be required to overcome the seven challenges are introduced to allow for a
791 consistent interpretation of spectral observations in terms of leaf, canopy and ecosystem traits.

792

793 **Figure 5.** Potential and emerging SIF applications illustrated in the form of a “SIF-city” metro plan,
794 where different colors denote five fields of plant science. Identified research applications (metro
795 stops) are causally connected in individual communication lines, but the final trajectories and
796 number of stops will depend on how the field of SIF research evolves over the next years. The red-
797 colored stops denote the application topics elaborated in Section 3.

798

799

800

801

802

Box 1 | Principle of solar-induced fluorescence (SIF) retrieval

SIF measurements take place outdoors, under ambient sunlight. Accordingly, when pointing a spectroradiometer towards a leaf or plant canopy to make a SIF measurement, we face the challenge that vegetation is highly reflective in the near infrared (NIR) wavelengths, and the signal is dominated by reflected light. The retrieval of SIF from the background reflected radiation is made possible thanks to the spectral properties of incoming light.

The solar spectrum, as measured above a plant canopy, is not continuous; rather, radiation is strongly attenuated within so-called Fraunhofer absorption lines and telluric absorption bands originating from absorption by gases in the Sun’s photosphere or the Earth’s atmosphere, respectively (see Fig. 1 and an idealized spectral feature in **A**). These features are exploited by the Fraunhofer line depth (FLD) methods¹¹⁹ where at least four spectral measurements, usually more¹²⁰, are required: the irradiance of the incoming sunlight and the apparent reflected radiance (called apparent, as it includes also SIF), inside and outside of the spectral absorption feature (E_{in}/E_{out} and L_{in}/L_{out} , respectively). Since SIF contributes photons similarly both inside and outside the spectral feature (**B**), the relative contribution of SIF to reflected radiation is significantly greater inside the spectral feature, causing an increase in the apparent reflectance (**C**). This increase is proportional to the amount of SIF and can be used to construct a system of equations to retrieve SIF.

Although not mutually exclusive, SIF measurements are often conducted using either the Fraunhofer or Telluric absorption bands, which involve some trade-offs:

- **Fraunhofer lines** (multiple lines across the SIF spectrum). The advantage of these retrievals lies in their lower sensitivity to atmospheric properties, which is practical for remote measurements as well as applications with variable target-to-sensor distances (e.g., multiangular tower measurements). The main disadvantage is that they require spectrometers with extremely high spectral resolutions and generally require longer periods of signal integration.
- **Telluric bands** (mainly oxygen absorption bands B and A, centered around 687-692 nm; and 759-770 nm, respectively). Since these bands are broader, measurements do not require as high spectral resolution and can be also conducted with shorter integration time, which can be especially suitable for some applications (e.g. drone-based observations). Their main disadvantage is that attention must be paid to corrections for atmospheric absorption (Challenge 6).

804
805
806
807
808
809
810
811
812

Box 2 | Radiative transfer models (RTMs)

- **Forward mode.** When the required inputs are provided, RTMs are capable of simulating leaf and canopy SIF together with reflected and emitted optical and thermal radiance. Once successfully validated by independent measurements, RTMs can be used in the forward mode to investigate the sensitivity of outputs, (i.e., surface reflectance and SIF) to different structural, biochemical, and physiological inputs, extending our mechanistic understanding of reflected and emitted photons' propagation across scales.
- **Inverse mode.** RTMs can be also inverted (i.e., run backwards) to estimate from laboratory, field and remote sensing spectral data those leaf and canopy traits that match measured reflectance and SIF data.

- **1D models.** 1D leaf RTMs assume that leaf constituents are horizontally homogeneously distributed in vertically stacked plate structures, and hence require only basic morphological and biochemical inputs (e.g., pigment contents driving PAR absorption and within-leaf reabsorption, the intrinsic PSII and PSI fluorescence spectra, and the dynamics in the quantum yield of fluorescence as the mechanistic link to photosynthesis). This simplicity, however, ignores potentially important factors, such as within-leaf heterogeneity or chloroplast movements. As with the 1D leaf construct, 1D canopy RTMs assume that vegetation can be represented by horizontally homogeneous layers filled with leaves of a predefined size, density and geometry (angular distribution), which allows for minimal model inputs and a relatively straightforward application. The 1D architecture has its uses for spatially homogeneous canopies (e.g., crops).
- **3D models.** Structurally complex leaves and spatially heterogeneous plant communities (e.g., forests and savannas) require 3D representations. 3D leaf RTMs can model optical interactions within a genuine 3D digital representation of leaf interior reconstructed, for example, with imaging tomography or confocal microscopy. As demonstrated in the Supplementary Videos 1 and 2, 3D RTM solutions also exist for spatially diversified plant canopies, allowing for accurate physical simulations of APAR_g and SIF fluxes in complex canopies.

813

814

815

EXCITATION

Solar spectral irradiance
[W m⁻² nm⁻¹]

PHOTOSYNTHETIC ENERGY PARTITIONING

Biochemistry & Biophysics

Molecular regulation
of photosynthesis

Investigating
drivers
of photosynthesis

3D photosynthesis:
Plant structure &
function

Detecting plant
functional traits

Phenotyping
& stress
detection

Breeding &
testing of
new varieties

Agriculture & Forestry

Resource-use
optimized
management

Pre-visual
pest &
disease
detection

Precision
agriculture
& forestry

Biogeochemistry

Land cover/use
change detection

Multiscale carbon
& water cycles

Biosphere-atmosphere
feedbacks

Impacts of climate
change & extreme
events

Plant Ecophysiology

Experimental
modeling

Functional
dynamics in
extreme &
remote biomes

Phenology &
circadian controls

Functional
& spatial
diversity

Ecology