<https://helda.helsinki.fi>

Exposing wind stress as a driver of fine-scale variation in plant communities

Momberg, Mia

2021-05

Momberg , M , Hedding , D W , Luoto , M & le Roux , P C 2021 , ' Exposing wind stress as a driver of fine-scale variation in plant communities ' , Journal of Ecology , vol. 109 , no. 5 , pp. 2121-2136 . <https://doi.org/10.1111/1365-2745.13625>

<http://hdl.handle.net/10138/340195>

<https://doi.org/10.1111/1365-2745.13625>

---

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Journal of Ecology

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

MS MIA MOMBERG (Orcid ID : 0000-0002-8901-9271)

DR DAVID W HEDDING (Orcid ID : 0000-0002-9748-4499)

Article type : Research Article

Editor: Robert Björk

Exposing wind stress as a driver of fine-scale variation in plant communities

Mia Momberg¹, David W. Hedding², Miska Luoto³ and Peter C. le Roux¹

¹University of Pretoria, South Africa ²University of South Africa, South Africa ³University of Helsinki, Finland

M. Momberg (miamomberg@gmail.com)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/1365-2745.13625](https://doi.org/10.1111/1365-2745.13625)

This article is protected by copyright. All rights reserved

25

26

27

28

29

30

31

32

33

34 **Abstract**

35 1. The effects of temperature and precipitation, and the impacts of changes in these climatic
36 conditions, on plant communities have been investigated extensively. The roles of other climatic
37 factors are, however, comparatively poorly understood, despite potentially also strongly
38 structuring community patterns. Wind, for example, is seldom considered when forecasting
39 species responses to climate change, despite having direct physiological and mechanical impacts
40 on plants. It is, therefore, important to understand the magnitude of potential impacts of changing
41 wind conditions on plant communities, particularly given that wind patterns are shifting globally.

42 2. Here, we examine the relationship between wind stress (i.e. a combination of wind exposure
43 and wind speed) and species richness, vegetation cover and community composition using fine-
44 scale, field-collected data from 1440 quadrats in a windy sub-Antarctic environment.

45 3. Wind stress was consistently a strong predictor of all three community characteristics, even
46 after accounting for other potentially ecophysiological important variables, including pH,
47 potential direct incident solar radiation, winter and summer soil temperature, soil moisture, soil
48 depth, and rock cover. Plant species richness peaked at intermediate wind stress, and vegetation
49 cover was highest in plots with the greatest wind stress. Community composition was also related
50 to wind stress, and, after the influence of soil moisture and pH, had a similar strength of effect as
51 winter soil temperature.

4. *Synthesis*: Wind conditions are, therefore, clearly related to plant community characteristics in this ecosystem that experiences chronic winds. Based on these findings, wind conditions require greater attention when examining environment-community relationships, and changing wind patterns should be explicitly considered in climate change impact predictions.

Keywords: community composition, fine-scale, plant community, species richness, vegetation cover, wind

Introduction

Recent changes in climate have impacted all aspects of ecological communities, including species richness, total abundance, biomass, cover and species composition (e.g. Menéndez et al., 2006; Kardol et al., 2010; Liu et al., 2018). Given the implications of these community-level changes for conservation, ecosystem services and human health, predicting how further changes in climate will affect communities is a key challenge in ecology (Kolstad and Johansson, 2011; Pecl et al., 2017). While some trends, like species range shifts, have been observed fairly consistently (e.g. Morueta-Holme et al., 2015; Freeman et al., 2018), exceptions have also been noted (e.g. Lenoir et al., 2010; Zhang et al., 2019), suggesting that some key factors or processes still need to be accounted for. Indeed, many studies forecasting ecological responses to climate change consider only temperature and precipitation as climatic variables, and do not account for other components of climate. For example, the ten climatic variables most commonly used in plant species distribution modelling studies all just quantify aspects of temperature and precipitation (Gardner et al., 2019).

Other climatic factors may, however, also be influential in determining individual species distributions and, as a result, community characteristics (Barton, 2017; Cherry and Barton, 2017; Maclean, 2020). Therefore, to improve the quality of the predictions of the ecological impacts of climate change, it is important to consider a more comprehensive set of climatic variables that directly affect plant performance (Gardner et al., 2019). Indeed, other variables, such as snow cover duration and solar radiation, can have profound effects both on individual species and on communities (see Bennie et al., 2008; Austin and Van Niel, 2011; Niittynen and Luoto, 2018). For example, decreasing snow cover duration in the Arctic has a larger effect on species' probability of becoming locally extinct than rising temperatures alone (Niittynen et al., 2018). Thus, a more thorough understanding of the impacts of climatic factors, extending beyond conventional

measures of temperature and precipitation, is required in order to more accurately predict how communities may be affected by changing climates (Mod et al., 2016).

Wind is a climatic variable that is currently underexplored as a driver of community patterns (see Table 1), despite representing an important stress and disturbance for many different taxa (Watanabe and Hasumi, 2005; Bintanja et al., 2014; Combes and Matano, 2018). It is relatively well understood how extreme winds (e.g. hurricanes and tornadoes) affect individuals, species, and communities (Mitchell, 2012; Møller, 2013; Behie et al., 2014; Xuan and Chang, 2014). However, these extreme conditions are limited in time and space (Nyberg et al., 2007; Lugo, 2008), and the impacts of chronic (i.e. continuous) wind conditions may be important but have received little focused attention. Exposure to wind has clear physiological impacts on plants, whereby, for example, plants typically close their stomata during windy conditions to reduce the rate of transpiration, consequently leading to lower rates of photosynthesis (Grace, 1977; de Langre, 2008). In addition, winds may desiccate the soil, creating a moisture stress for plants (Bertiller et al., 1996; Fitzgerald and Kirkpatrick, 2017), and redistribute litter which has an effect on soil temperature and nutrient content (Fahnestock et al., 2000). Wind may also have a range of mechanical impacts on plants, with, for example, strong winds potentially tearing leaves, causing abrasion and desiccation (Hadley and Smith, 1983; Hadley and Smith, 1986; de Langre, 2008; Gardiner et al., 2016), uprooting individuals (Yang et al., 2014), and causing flowers and fruits to be shed (e.g. Lahav and Zamet, 1999). As a result, lower wind speeds are typically associated with faster plant growth rates and higher productivity (e.g. Bang et al., 2010), although physiological responses to wind conditions may be species-specific (Onoda and Anten, 2011). Further, wind can shape the outcome of plant-plant interactions (severity-interaction-relationships; le Roux and McGeoch, 2010). Therefore, wind has the potential to directly affect vegetation patterns, from altering individual species' dominance (Okitsu and Ito, 1984) to impacting the distribution of different plant communities (Williams and Ashton, 1987; Lynch and Kirkpatrick, 1995; see Table 1 for an illustrative summary of what is currently known about the impacts of wind on vegetation). Consequently, differences in chronic wind conditions between sites and/or years has considerable potential to drive spatio-temporal variations in plant communities.

Wind stresses and disturbances may vary greatly over short distances (Katsaprakakis and Christakis, 2012; Chiras, 2017), and could therefore potentially contribute to fine-scale variation in ecological communities. For example, wind velocity can vary around individual plants

(Combrinck et al., 2020) and models that do not account for fine-scale variation in wind patterns do not accurately predict seed dispersal (Lönnell et al., 2015). Variability in wind speed has also, for instance, been related to fine-scale differences in plant height, species richness, growth form diversity, and community type (Whitehead, 1954; Whitehead, 1959; Wilson, 1959). Nevertheless, wind is seldom considered as a driver of fine-scale variation in community patterns (see review by Gardner et al., 2019), and despite technological advances that have improved measurement and modelling of wind conditions, little work has recently examined the influence of wind on plant communities (although, see, e.g. Fitzgerald and Kirkpatrick, 2017; Sun et al., 2019; Sparacino et al., 2020; and Table 1). In addition, the reciprocal effect of vegetation on wind patterns, has also attracted limited attention (although see, e.g., Combrinck et al., 2020).

Based on our current understanding of the ecological impacts of wind, variation in prevailing wind conditions at fine spatial scales is likely to affect multiple measures of plant community structure. As wind exposure increases, vegetation cover generally declines due to wind pruning and slower growth rates (Bang et al., 2010; Gardiner et al., 2016), as evident, for example, on New Zealand's off-shore islands (McGlone, 2002). Species richness may also decline in increasingly windy microclimates, especially in species-poor environments. However, in more species-rich communities, species may instead show a pattern of replacement along a wind stress gradient due to inter-specific differences in wind tolerance (i.e. more wind-tolerant species replace taller and/or more competitive species that are less tolerant of wind stress; e.g. Burke et al., 1989; Fernández-Palacios and Nicolás, 1995). Under such a scenario species richness would be predicted to remain relatively constant while species composition shifts in response to greater wind stress (i.e. a type of Gleasonian distribution; Gleason, 1939).

The paucity of studies investigating the effects of wind on vegetation was already noted six decades ago (Wilson, 1959; see also Whitehead, 1954; Whitehead, 1959), but this issue has yet to be explicitly and thoroughly addressed (see e.g. Sutherland et al., 2017). Furthermore, understanding how wind, and changes in wind patterns, affects biodiversity is increasingly relevant as, over the last decade, mean wind speeds have accelerated globally, with the largest changes happening in the Southern Ocean (Young and Ribal, 2019; Zeng et al., 2019), and current predictions suggesting that this trend will continue in the long-term (Jeong and Sushama, 2019; Zeng et al., 2019). Therefore, in this study, the impact of wind on fine-scale vegetation characteristics was assessed by examining the relationship between wind stress (i.e. a combination

of wind exposure and wind speed) and 1) vascular plant species richness, 2) species cover, and 3) species composition in a wind exposed environment, after accounting for other potentially ecophysiological important variables (Mod et al., 2016).

147

148 **Materials and methods**

149 *Study site*

150 This study was conducted on the isolated sub-Antarctic Marion Island (46°54' S, 37°45' E). The
151 sub-Antarctic provides an ideal study system to examine the ecological consequences of chronic
152 winds as the region experiences consistently strong daily wind conditions (Pendlebury and Barnes-
153 Keoghan, 2007; le Roux, 2008). Terrestrial habitats of Marion Island exhibit broad gradients of
154 wind stress due to their complex topography, with sites ranging from sheltered to exposed often
155 separated by only short distances. Marion Island lies approximately halfway between Africa and
156 Antarctica, covers an area of 293 km², and rises to 1240 m a.s.l. (Boelhouwers et al., 2008). The
157 island is situated in the “roaring forties”, the region between 40 and 50°S, that experiences strong
158 and relatively consistent westerly winds (Pendlebury and Barnes-Keoghan, 2007; le Roux, 2008)
159 which can affect pedogenesis, vegetation growth, soil frost formation and aeolian transport of
160 particles (Hedding et al., 2015). Gale force winds blow on >100 days per annum on Marion Island
161 (with relatively consistent directionality; le Roux, 2008). The island has a hyper-oceanic climate
162 (characterized by cool temperatures with little diurnal and seasonal variation) and high humidity,
163 with near complete cloud cover and precipitation on most days (le Roux, 2008). The mean daily
164 maximum and minimum temperatures are 8.7 °C and 3.2 °C respectively, and the mean annual
165 precipitation is c. 1800 mm (recorded at the island’s meteorological station which is ~1 km away
166 from the study site; average from 2008 - 2018).

167 *Data collection*

168 Data were collected from early 2016 to early 2017 on the eastern side of Marion Island, with
169 sampling focused in a topographically-, geologically- and biotically-heterogenous area
170 (comprising a smooth pre-glacial lava flow north of the Van den Boogaard river and a rugged
171 post-glacial lava flow south of the river; Fig. 1). A survey approach was used that has previously
172 been applied in other high-altitude and high-latitude systems (e.g. le Roux et al., 2013b;

173 Kemppinen et al., 2019; Niittynen et al., 2020a), where vegetation composition and abiotic
174 conditions are measured across steep environmental gradients within a single landscape unit
175 (avoiding differences in macroclimate and regional species pools). Nine grids of 8 x 20 m were
176 sampled, each comprising 160 contiguous quadrats of 1 m² (resulting in 1440 quadrats sampled in
177 total; Figure 1, Table 2). These grids encompassed fellfield, mire vegetation, and fern-dominated
178 slopes, and contained 18 vascular plant species. Grids were located to sample the full range of
179 conditions within the site, covering the range of vegetation types, vegetation cover and topography
180 present. The distance between grids was 70 - 915 m, with the grids covering an altitudinal range of
181 c. 40 to 100 m a.s.l. Individual grids, in turn, were orientated to encompass as much local
182 environmental variability as possible.

183 Within each of the 1 m² quadrats several abiotic and biotic variables were measured. The
184 cover and identity of all vascular plant species in each quadrat were recorded and then used to
185 determine the species richness and composition for each quadrat. Percentage plant cover, rock
186 cover (i.e. the cover of rocks or boulders large enough to inhibit plant growth) and bare soil cover
187 were visually estimated. Soil depth was measured using a thin metal rod (diameter of 8 mm) at
188 three points randomly located in each quadrat, and an average soil depth then calculated for the
189 quadrat (where soil depth exceeded 60 cm, a value of 70 cm was assigned). The slope and aspect
190 of each quadrat were recorded and these values were then used to calculate potential annual direct
191 incident radiation (PDIR; McCune and Keon, 2002; McCune, 2007). Soil samples were taken
192 from 16 - 32 quadrats per grid and used to determine soil pH in the laboratory using the CaCl₂
193 method (Hendershot et al., 2008). Soil pH was then interpolated to the unsampled quadrats using
194 bilinear interpolation (Bovik, 2009).

195 Instantaneous soil temperature and soil moisture measurements were taken in each quadrat
196 in June, July and October 2016 and January and April 2017. These readings were only taken on
197 days when there had been at least 24 hours since the last rainfall event. Volumetric soil moisture
198 content was measured using a handheld time-domain soil moisture meter (using 7.5 cm probes;
199 FieldScout TDR 300; Spectrum Technologies), and soil temperature was measured using a
200 corkscrew thermometer (EXTECH Instruments, TM40) at a depth of 5 cm. To test whether the
201 order in which measurements were taken had an effect on soil moisture or temperature readings
202 (i.e. testing for an effect of time of the day), the grid which was measured first was revisited at the
203 end of the measurement period and partly remeasured (20 quadrats). The average difference in

204 instantaneous soil moisture readings between the initial and final measurements in resampled
205 quadrats was 0.8%, and, therefore, the raw data for soil moisture readings were used in analyses.
206 Soil temperature, however, showed a significant difference (t-test: $p < 0.001$) between the
207 temperature measured at the start of the measurement time period versus the end of the
208 measurement period for the quadrats which were remeasured. Therefore, to account for the effect
209 of the measurement time on soil temperature, readings were corrected using a linear adjustment.
210 This correction was based on soil temperature data logged in 3 – 9 cells per grid using temperature
211 loggers (Thermochron iButton DS1921G; Maxim Integrated), with data from the specific day on
212 which instantaneous measurements were made being used (following le Roux et al., 2013a).

213 Soil temperatures in the winter months were positively correlated (June and July), while
214 there was minimal correlation between winter and summer temperatures (Figure A1). Therefore,
215 one measurement of summer soil temperature (January; mid-summer) and one measurement of
216 winter soil temperature (June; mid-winter) were used in all subsequent analyses. Soil moisture was
217 significantly positively correlated across all months (Figure A2), and, therefore, moisture data
218 from only one month (October) were included in further analyses.

219 Wind speed and direction were measured at 17 locations across Marion Island using two
220 sonic anemometers (Gill Windsonic: Gill Instruments, UK), mounted at 0.5 m and 1 m above
221 ground at each of the locations. The anemometers measured wind speed and direction in the
222 horizontal plane (two-dimensional) at a 0.01 m.s^{-1} resolution, with measurement accuracy of 2%
223 (at 12 m.s^{-1}). Wind speed and direction were sampled at 0.5 Hz, with mean direction and speed
224 logged every 10 minutes using a CR300 datalogger (Campbell Scientific, USA). Both the
225 anemometers and the logger were powered using a 12 V DC battery continuously charged using a
226 25 W solar panel. Wind direction data for the study site showed one clear and dominant wind
227 direction, and, therefore, only the dominant wind direction was extracted from the wind station
228 located closest to the study site ($< 700 \text{ m}$ from the furthest grid) over a 12-month period (April
229 2018 - March 2019). Wind at the study site predominantly comes from a north-westerly direction
230 (68 % of readings were from $260\text{--}330^\circ$ from N), with the strongest wind speeds being even more
231 limited to the north-westerly sector (88 % of all wind speeds greater than 11 m.s^{-1} were recorded
232 between $260\text{--}330^\circ$; Figure 1; in agreement with long-term data from the island's weather station;
233 le Roux, 2008).

234 Photographic images of each grid were taken using an unmanned aerial vehicle (UAV; DJI
235 Phantom 4 Pro fitted with a 20-megapixel camera) at a ground sample distance of 0.7 cm during
236 April 2019. The aerial images from the UAV were processed in Agisoft Photoscan using the
237 principle of structure-from-motion to produce a point dense cloud which in turn was used to
238 generate a Digital Surface Model (DSM) at 2 cm resolution, which captures the elevation of the
239 surface. From the DSM's, a within-grid wind exposure metric was calculated for each 1 m²
240 quadrat using the Wind Effect module in SAGA-GIS (Conrad et al., 2015), specifying the wind
241 direction as North-West (with the majority of the wind coming from 292 - 338°; based on the data
242 from the wind station at the site). The minimum, maximum, and mean of the wind effect values
243 were extracted for each quadrat. Wind effect is a dimensionless index, where values below 1
244 indicate areas that are sheltered from wind and values above 1 indicate areas that are exposed to
245 wind in terms of the specified wind direction (Böhner and Antonić, 2009). To account for
246 differences in coarser-scale wind stress between grids (since some grids were more wind-exposed
247 than others; Fig. 1), the mean simulated wind speed for each grid was extracted from a
248 computational fluid dynamics (CFD) model of Marion Island (maximum 50 m resolution) using
249 ANSYS Fluent 2019R3 (Ansys, USA). The CFD model uses a full-scale digital elevation model of
250 Marion Island (DRDLR, 2019) and simulates air flow over the topology by iteratively solving a
251 set of partial differential equations (the Reynolds-Averaged Navier Stokes Equations; see Versteeg
252 and Malalasekera, 2007 for a detailed consideration of CFD; and Cindori et al., 2018 for a recent
253 implementation). A westerly wind was assumed as the free-stream condition with a reference
254 speed of 5 m.s⁻¹ at 1.5 m above ground, based on measured data from the wind stations. The model
255 includes considerations for the atmospheric boundary layer and the effect of the Coriolis force
256 (Breedt et al., 2018). This estimate of grid-level wind speed was multiplied by quadrat-level wind
257 index indices (for minimum, maximum, mean, and range wind stress) to estimate wind stress
258 values that are comparable across and within grids (i.e. accounting for grids differing considerably
259 in landscape-level wind stress). The wind stress metric provides relative values, enabling
260 comparisons between sites within this particular study system, where higher values indicate higher
261 wind stress. Here wind stress represents the combined influence of wind exposure to the dominant
262 wind direction and the simulated mean wind speed at each grid when the wind is blowing from the
263 dominant wind direction. These values, therefore, provide an estimate of the mean physical and
264 mechanical strain that plants experience (i.e. representing chronic wind stress). The resulting wind

265 stress metrics were highly correlated ($r > |0.8|$; Figure A3), and, therefore, only maximum wind
266 stress was used in subsequent analyses.

267 *Statistical analyses*

268 None of the predictor variables in the final dataset were strongly or significantly correlated
269 with each other ($r < |0.8|$ and Variance Inflation Factor < 3.1 ; Figure A4). The relationship
270 between species richness (and cover) and predictor variables was analysed using three different
271 statistical methods to reduce uncertainties due to modelling approach: generalized linear models
272 (GLZ; Müller, 2012), generalized additive models (GAM; Wood and Augustin, 2002), and
273 generalized boosted regression models (GBM; Friedman et al., 2000; Friedman, 2001). These
274 three methods were chosen due to their differing flexibility and complexity. All of the statistical
275 methods were run assuming a Poisson (for species richness) or quasibinomial (for vegetation
276 cover; which was significantly over-dispersed) distribution.

277 For all three statistical methods, a first model was run with PDIR, soil depth, rock cover,
278 winter temperature, summer temperature, moisture, and pH as predictor variables (hereafter
279 referred to as the “simple model”). A second model was then run with all of the above variables,
280 as well as the maximum wind stress (hereafter referred to as the “full model”). The GLZs included
281 quadratic terms for all predictor variables to allow for non-linear relationships. For GAMs, the
282 initial degree of smoothness for each predictor was set to four. The tree complexity was set to 6
283 for GBMs and the tree threshold to 1000. To determine whether the addition of wind stress
284 improved the performance of the models, each pair of simple and full models from GAMs and
285 GLZs were compared using a likelihood ratio test (for species richness) and an F-test (for
286 vegetation cover). Similar model comparisons are not possible for GBMs, but response curves and
287 variable importance could be compared between GBMs and the other two methods. Variable
288 importance for GAMs and GLZs was calculated by comparing the Pearson correlation between
289 predictions made on the original data and predictions made on the data where the predictor
290 variable of interest has been randomly shuffled (following Niittynen and Luoto, 2018). The
291 calculations of variable importance were calculated 10 times and the mean importance value
292 reported.

293 Species composition was modelled using non-metric multidimensional scaling (nMDS)
294 and permutational multivariate analysis of variance (PERMANOVA) based on species occurrence.

Two dimensions, 200 random starts and the standard transformation (Wisconsin double standardization) were used in nMDS analyses. Nestedness analyses were also conducted on the species composition data to determine whether species were nested along the wind stress gradient (Ulrich, 2009). The species-site matrix was first ordered by maximum wind stress, and the nestedness metric based on Overlap and Decreasing Fill (NODF) was calculated. NODF is less prone to type I statistical errors and is insensitive to matrix size and shape, and, therefore, provides a more conservative approach than other nestedness metrics (Almeida-Neto et al., 2008). Finally, the nestedness analysis was complemented by calculating the overall beta diversity between quadrats using the Sørensen dissimilarity index, with the contribution of species turnover (measured as Simpson dissimilarity) and nestedness (measured as the nestedness-resultant fraction of the Sørensen dissimilarity index) being partitioned to explain the observed differences in community composition (Baselga and Orme, 2012).

All analyses were run in R statistical software, version 3.5.0 (R Core Team, 2018), using additional functions from the vegan (Oksanen et al., 2018), mgcv (Wood, 2006), spind (Carl et al., 2018), and betapart (Baselga et al., 2018) libraries.

310

311 **Results**

Wind stress ranged from 4.5 to 8.5, with grid one experiencing the overall lowest wind stress, and grid four the highest wind stress (Figure 2). Species richness varied between zero and eight species per 1 m² quadrat, and between 9 and 15 species per grid (Figure 3), while vascular plant cover varied between a minimum of zero and a maximum of 100 % (Figure A5).

The full model, including wind stress, performed significantly better than the simple model for species richness in the GAM, and marginally significantly so in the GLZ. For vegetation cover the full model performed significantly better in both statistical approaches (GLZ and GAM; likelihood ratio test and F-test, $p < 0.05$; Table A1). The full model for species richness improved the deviance explained by 0.8 % for the GLZ and by 3.5 % for the GAM relative to the simple models that did not include a measure of wind stress. For vegetation cover, the full model improved the deviance explained by 1.4 % for the GLZ and by 1.4 % for the GAM (Table A1).

Both GLZ and GAM models for species richness contained maximum wind stress within the top five predictor variables (i.e. on the basis of variable importance; Table 3; wind stress was the second most important predictor in the GBM model, Table A2). Response curves from the GAM model for species richness showed that species richness has a complex relationship with maximum wind stress, with the highest species richness observed at intermediate maximum wind stress (Figure 4, with raw data shown in Figure A6; similar response curves were observed from the GLZ model, Fig. A8; and for the GBM model, Fig. A10). The addition of wind stress to the model changed the shape of the response curve between species richness and PDIR (from negative to nearly horizontal), winter soil temperature (the slope of the positive response becomes less steep), and summer soil temperature (remains hump-shaped, but high temperatures result in a smaller drop in species richness). Species richness also showed non-linear relationships with both rock cover and pH (which were also consistently in the top five predictors), where species richness peaked at c. 60% rock cover and 4.6 pH (Figure 4).

Vegetation cover was significantly predicted by maximum wind stress, and wind stress was the second most important predictor when modelling vegetation cover for both the GLZ and the GAM (Table 3; and third most important in the GBM model, Table A2). The response curves for vegetation cover showed that maximum wind stress had a valley-shaped relationship with vegetation cover (Figure 5 and Figure A7; see Fig. A9 and A11 for GLZ and GBM response curves which showed similar patterns). Rock cover had a strong negative relationship with vegetation cover, and was the most important predictor in both statistical approaches (Table 3).

All eight of the predictor variables in the full model contributed significantly to explaining variation in species composition, and together explained 33.9 % of the variation (the simple model explained 32.4 %). Wind stress was the fourth most important predictor in explaining species composition, after soil moisture, pH, and winter soil temperature. Species composition was weakly nested by wind stress, with NODF = 31.0, whereas NODF = 57.9 when the species matrix is arranged to maximize nestedness. This result was mirrored by the partitioning of beta diversity into species turnover and nestedness, where an overall Sørensen dissimilarity index value of 0.996 (indicating high dissimilarity in the species composition between quadrats) was predominantly driven by species turnover (0.994), with nestedness having a much smaller effect (0.003).

352

353 Discussion

354 Wind stress was consistently prominent in explaining fine-scale spatial variation in species
355 richness, vegetation cover, and species composition, as evidenced both by models including wind
356 stress performing better than models without wind stress and by wind stress repeatedly having
357 among the highest relative importance values. These results agree with findings across broader
358 scales, where, for example, wind protection is important in determining the distribution of high-
359 altitude forests and tree sapling growth (Sparacino et al., 2020), and wind patterns improved the
360 accuracy of future temperature forecasts and resulting predictions of climate refugia (Ashcroft et
361 al., 2009).

362 Both species richness and species cover were significantly related to wind stress. Species
363 richness responded non-linearly to wind stress, with the highest species richness at intermediate
364 maximum wind stress. This suggests that in ecosystems with high wind stress, such as sub-
365 Antarctic islands, areas that do not have sheltered microsites have a constrained species richness,
366 and that the largest number of species co-occur where the maximum stress conditions in an area
367 are neither entirely sheltered nor exposed. Vegetation cover follows the same trend as species
368 richness, where maximum wind stress is one of the key predictors. Wind stress and vegetation
369 cover have a non-linear relationship (i.e. both low and high maximum wind stress values are
370 correlated with higher vegetation cover), but the highest vegetation cover is found in microsites
371 with the highest maximum wind stress. It is possible that in this system where bryophytes can be
372 co-dominant (e.g. in both the wettest habitats; Smith and Steenkamp, 2001; Smith et al., 2001),
373 that at high wind speeds evapotranspiration causes too great a moisture stress for bryophytes
374 (which are poikilohydric and are highly dependent on external water availability; Vitt et al., 2014),
375 allowing vascular plants to achieve greater cover.

376 Species composition was also significantly related to wind stress, and differed strongly
377 between quadrats, driven chiefly by species replacement. The low nestedness exhibited by
378 vascular plant species along the wind stress gradient, which matches the observation that species
379 richness did not decline uniformly with increasing wind stress, suggests that species may be
380 replacing each other as wind conditions become progressively more stressful. For example, on
381 Marion Island, the alien grass *Agrostis stolonifera*, is limited to sites which are sheltered due to its
382 lack of supportive tissue, while the indigenous congeneric *A. magellanica* grows in more wind
383 exposed locations (Pammenter et al., 1986). Further, presumably due to the wind-sheltered nature

384 of streambanks, *Agrostis stolonifera* disproportionately invades these communities, but is absent
385 from windier habitats (Gremmen et al., 1998). Irrespective of the mechanism driving the influence
386 of wind on community composition, these results agree with previous studies in other temperature-
387 limited systems that have also identified wind (acting, e.g., via erosion) as a driver of variation in
388 species composition (Vonlanthen et al., 2006a; le Roux and Luoto, 2014). Therefore, in terms of
389 species richness, cover and composition, wind is an important driver of plant community structure
390 at fine-scales in an environment that is exposed to chronic winds.

391 The other abiotic variables that were strongly related to vascular plant richness, cover and
392 composition exhibited patterns consistent with other studies conducted at high altitude and/or
393 latitude sites. For example, pH was a consistently strongly correlated with species richness, in
394 agreement with findings from other temperature-limited systems (e.g. Vonlanthen et al., 2006b;
395 Filibeck et al., 2019). Soil moisture played the largest role in driving community composition, and
396 this variable has been highlighted as a key driver of species composition in other low-energy
397 systems (le Roux et al., 2013a). Indeed, wind and soil moisture may potentially have an interactive
398 relationship, with strong winds desiccating upper soil layers (Bertiller et al., 1996). Because wind
399 stress was related to plant community properties even after accounting for these other abiotic
400 variables, these results clearly highlight the importance of wind in driving variation in floral
401 communities.

402

403 ***Wind as a globally underexplored environmental driver***

404 While wind may be important in high latitude and high altitude regions (e.g. Whitehead, 1954;
405 Wilson, 1959 and this study), and experiments have shown it to have widespread effects on plant
406 growth in other regions too (Wilson, 1959), it is still partly unclear if the patterns observed here
407 can be generalized to other systems. At a minimum, it is likely that wind will be important in (1)
408 areas with strong winds (chronic and/or extreme winds) as it is a mechanical stress, (2) areas with
409 fine and loose material available for aeolian transport (e.g. due to erosion, deposition and wind
410 scouring in dune environments; Nylén et al., 2015; Nylén and Luoto, 2015), (3) dry areas where
411 wind enhances evaporation and the potential for plants to experience greater moisture stress
412 (Zhang et al., 2007), (4) areas with thin soils which are more susceptible to desiccation (Bertiller
413 et al., 1996), and (5) Arctic and alpine areas where it affects snow drift (Sturm et al., 2001; Dadic

et al., 2010). Wind may also have an indirect effect by interacting with other climatic factors (see e.g. Kullman and Loyer, 2005; Ashcroft et al., 2009). As a result, the large recent changes in global wind patterns may have direct consequences for species distributions and interactions between species (Young et al., 2011; Young and Ribal, 2019), as well as indirect impacts (e.g. by affecting seed dispersal; Kling and Ackerly, 2020).

Predictions of the ecological impacts of changing wind conditions may be complicated by species- and site-specific responses. For example, wind may differentially impact on groups of species within communities. Since understory species are less exposed, for example in forest communities, it is expected that canopy species will show the greatest responses to changes in wind (as observed for changes in temperature; Mau et al., 2018). Further, species growing on ridge crests or slopes will likely be more affected than those growing in protected valleys, since changes to wind speed and/or direction in wind exposed locations will be less buffered by other vegetation or topographic features. An added challenge in making predictions on the impacts of changing wind conditions is that altered wind patterns may contribute to the development of “novel climates”, representing climatic conditions not currently present elsewhere (*sensu* Williams and Jackson, 2007; Dahinden et al., 2017). Plants and animals may lack adaptations to changes in wind conditions; similar to some species’ inability to adapt fast enough to recent changes in temperature and precipitation (e.g. Parmesan and Hanley, 2015; Gómez-Ruiz and Lacher Jr, 2019; Radchuk et al., 2019). Therefore, the impacts of changes in wind patterns on plant communities will likely be dependent on interactions with microclimatic and/or topographic conditions, and will vary between habitat types and species groups.

To further understand the ecological impact of wind, future studies should investigate not only community-level metrics (e.g. as in this study), but also the responses of the individual species that comprise the community. Species may, for example, respond to wind stress through trading-off resources between different processes or traits, as seen in response to changes in other climatic factors (Gandin et al., 2011; Liu et al., 2013). Indeed, recent evidence has shown that, for example, some plant species respond to higher wind speeds by increasing their foliar silicon content (which may provide protection against mechanical stress; Song et al., 2020). Functional traits have been used to examine how species react to changes in other environmental variables (e.g. Bjorkman et al., 2018; Niittynen et al., 2020b; Thomas et al., 2020), and traits related to resource allocation (e.g. specific leaf area) and leaf toughness (e.g. leaf dry matter content and

445 tensile strength) may provide useful metrics for quantifying functional variation within species in
446 response to spatio-temporal variation in wind patterns.

447 Here we show that spatial variation in wind stress has a defining impact on vegetation
448 communities in an ecosystem that experiences chronic winds. Therefore, temporal variation in
449 wind patterns, which are currently shifting due to global climate change (Young et al., 2011;
450 Young and Ribal, 2019; Zeng et al., 2019), may also likely affect plant community characteristics
451 through time, in the same way that wind currently affects their spatial patterning. The growing
452 availability of wind data at a coarse spatial scale needs to be (1) examined in the context of
453 landscape-level biological patterns (i.e. related to topography), and (2) refined to also describe
454 variation in wind characteristics at the scales affecting individual plants and local communities
455 (Maclean, 2020). Wind has been an understudied climatic driver of biotic communities for more
456 than 60 years (Wilson, 1959), and, especially in light of the global challenge of climate change, it
457 is now time to advance our understanding of where, and to what extent, wind acts as a driver of
458 vegetation patterns at multiple spatial scales.

459

460 **Acknowledgements**

461 This research was supported by the National Research Foundation's South African National
462 Antarctic Programme (grant number 93077, 110726 and 110723) and was conducted under
463 permits from the Prince Edward Islands Management Committee (PEIMC1/2013). MM received
464 funding from the South African National Research Foundation. We thank E. Mostert, N. Mhlongo,
465 J. Schoombie and J. van Berkel for field assistance. We thank P. Niittynen for advice on statistical
466 analyses. The authors declare no conflict of interest.

467

468 **Data availability statement**

469 Data will available from the Dryad Digital Repository from 1 January 2022:
470 <https://doi.org/10.5061/dryad.547d7wm7k> (Momberg et al., 2021).

471

472 **Author contributions**

473 PCLR, ML and MM conceived the ideas and designed methodology; MM, PCLR and DWH
474 collected and processed the data; MM and PCLR analysed the data, with inputs from DWH and
475 ML; MM led the writing of the manuscript. All authors contributed critically to the drafts and gave
476 final approval for publication.

477

478 **References**

- 479 Almeida-Neto, M., Guimarães, P., Guimarães Jr, P. R., Loyola, R. D. & Ulrich, W. (2008). A
480 consistent metric for nestedness analysis in ecological systems: reconciling concept and
481 measurement. *Oikos* **117**: 1227-1239.
- 482 Ashcroft, M. B., Chisholm, L. A. & French, K. O. (2009). Climate change at the landscape scale:
483 Predicting fine-grained spatial heterogeneity in warming and potential refugia for
484 vegetation. *Global Change Biology* **15**: 656-667.
- 485 Austin, M. P. & Van Niel, K. P. (2011). Improving species distribution models for climate change
486 studies: Variable selection and scale. *Journal of Biogeography* **38**: 1-8.
- 487 Bang, C., Sabo, J. L. & Faeth, S. H. (2010). Reduced wind speed improves plant growth in a
488 desert city. *PloS One* **5**: e11061.
- 489 Barton, B. T. (2017). Beyond global warming: Putting the “climate” back into “climate change
490 ecology”. *Food Webs* **13**: 51-52.
- 491 Baselga, A. & Orme, C. D. L. (2012). betapart: An R package for the study of beta diversity.
492 *Methods in Ecology and Evolution* **3**: 808-812.
- 493 Baselga, A., Orme, D., Villeger, S., De Bortoli, J. & Leprieur, F. (2018). betapart: Partitioning
494 beta diversity into turnover and nestedness components. *R package version 1.5.1*.
- 495 Behie, A. M., Kutz, S. & Pavelka, M. S. (2014). Cascading effects of climate change: Do
496 hurricane-damaged forests increase risk of exposure to parasites? *Biotropica* **46**: 25-31.
- 497 Bennie, J., Huntley, B., Wiltshire, A., Hill, M. O. & Baxter, R. (2008). Slope, aspect and climate:
498 Spatially explicit and implicit models of topographic microclimate in chalk grassland.
499 *Ecological Modelling* **216**: 47-59.
- 500 Bertiller, M. B., Zaixso, P., Irisarri, M. & Brevedan, E. R. (1996). The establishment of *Festuca*
501 *pallenscens* in arid grasslands in Patagonia (Argentina): The effect of soil water stress.
502 *Journal of Arid Environments* **32**: 161-171.

503 Bintanja, R., Severijns, C., Haarsma, R. & Hazeleger, W. (2014). The future of Antarctica's
 504 surface winds simulated by a high-resolution global climate model: 2. Drivers of
 505 21st century changes. *Journal of Geophysical Research: Atmospheres* **119**: 7160-7178.

506 Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Normand, S., Rüger, N., Beck, P. S. A.,
 507 Blach-Overgaard, A., Blok, D., Cornelissen, J. H. C., Forbes, B. C., Georges, D., Goetz, S.
 508 J., Guay, K. C., Henry, G. H. R., HilleRisLambers, J., Hollister, R. D., Karger, D. N.,
 509 Kattge, J., Manning, P., Prevéy, J. S., Rixen, C., Schaepman-Strub, G., Thomas, H. J. D.,
 510 Vellend, M., Wilmking, M., Wipf, S., Carbognani, M., Hermanutz, L., Lévesque, E.,
 511 Molau, U., Petraglia, A., Soudzilovskaia, N. A., Spasojevic, M. J., Tomaselli, M., Vowles,
 512 T., Alatalo, J. M., Alexander, H. D., Anadon-Rosell, A., Angers-Blondin, S., Beest, M. t.,
 513 Berner, L., Björk, R. G., Buchwal, A., Buras, A., Christie, K., Cooper, E. J., Dullinger, S.,
 514 Elberling, B., Eskelinen, A., Frei, E. R., Grau, O., Grogan, P., Hallinger, M., Harper, K. A.,
 515 Heijmans, M. M. P. D., Hudson, J., Hülber, K., Iturrate-Garcia, M., Iversen, C. M.,
 516 Jaroszynska, F., Johnstone, J. F., Jørgensen, R. H., Kaarlejärvi, E., Klady, R., Kuleza, S.,
 517 Kulonen, A., Lamarque, L. J., Lantz, T., Little, C. J., Speed, J. D. M., Michelsen, A.,
 518 Milbau, A., Nabe-Nielsen, J., Nielsen, S. S., Ninot, J. M., Oberbauer, S. F., Olofsson, J.,
 519 Onipchenko, V. G., Rumpf, S. B., Semenchuk, P., Shetti, R., Collier, L. S., Street, L. E.,
 520 Suding, K. N., Tape, K. D., Trant, A., Treier, U. A., Tremblay, J.-P., Tremblay, M., Venn,
 521 S., Weijers, S., Zamin, T., Boulanger-Lapointe, N., Gould, W. A., Hik, D. S., Hofgaard,
 522 A., Jónsdóttir, I. S., Jorgenson, J., Klein, J., Magnusson, B., et al. (2018). Plant functional
 523 trait change across a warming tundra biome. *Nature* **562**: 57-62.

524 Boelhouwers, J. C., Meiklejohn, K. I., Holness, S. D. & Hedding, D. W. 2008. Geology,
 525 geomorphology and climate change. In: Chown, S. L. & Froneman, P. W. (eds.) *The*
 526 *Prince Edward Islands: Land-sea interactions in a changing ecosystem*. Stellenbosch:
 527 SUN PRESS.

528 Böhner, J. & AntoniĆ, O. 2009. Land-surface parameters specific to topo-climatology. In: Hengl,
 529 T. & Reuter, H. I. (eds.) *Geomorphology: Concepts, software, applications*. Elsevier.

530 Bovik, A. C. 2009. Basic gray level image processing. In: Bovik, A. C. (ed.) *The Essential Guide*
 531 *to Image Processing*. Boston: Academic Press.

532 Breedt, H. J., Craig, K. J. & Jothiprakasam, V. D. (2018). Monin-Obukhov similarity theory and
 533 its application to wind flow modelling over complex terrain. *Journal of Wind Engineering*
 534 *and Industrial Aerodynamics* **182**: 308-321.

- 535 Burke, I. C., Reiners, W. A. & Olson, R. K. (1989). Topographic control of vegetation in a
536 mountain big sagebrush steppe. *Vegetatio* **84**: 77-86.
- 537 Carl, G., Levin, S. C. & Kühn, I. (2018). Spind: An R package to account for spatial
538 autocorrelation in the analysis of lattice data. *Biodiversity Data Journal* **6**: e20760.
- 539 Cherry, M. & Barton, B. (2017). Effects of wind on predator-prey interactions. *Food Webs* **13**: 92-
540 97.
- 541 Chiras, D. 2017. *Power from the wind: A practical guide to small-scale energy production*.
542 Vancouver: New Society Publishers.
- 543 Cindori, M., Juretić, F., Kozmar, H. & Džijan, I. (2018). Steady RANS model of the homogeneous
544 atmospheric boundary layer. *Journal of Wind Engineering and Industrial Aerodynamics*
545 **173**: 289-301.
- 546 Combes, V. & Matano, R. P. (2018). The Patagonian shelf circulation: Drivers and variability.
547 *Progress in Oceanography* **167**: 24-43.
- 548 Combrinck, M. L., Harms, T. M., McGeoch, M. A., Schoombie, J. & le Roux, P. C. (2020). Wind
549 and seed: a conceptual model of shape-formation in the cushion plant *Azorella selago*.
550 *Plant and Soil*.
- 551 Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann,
552 V. & Böhner, J. (2015). System for Automated Geoscientific Analyses (SAGA) v. 2.1.4.
553 *Geoscientific Model Development* **8**: 1991-2007.
- 554 Dadic, R., Mott, R., Lehning, M. & Burlando, P. (2010). Wind influence on snow depth
555 distribution and accumulation over glaciers. *Journal of Geophysical Research: Earth*
556 *Surface* **115**.
- 557 Dahinden, F., Fischer, E. M. & Knutti, R. (2017). Future local climate unlike currently observed
558 anywhere. *Environmental Research Letters* **12**: 084004.
- 559 de Langre, E. (2008). Effects of wind on plants. *Annual Review of Fluid Mechanics* **40**: 141-168.
- 560 Department of Rural Development and Land Reform (DRDLR), 2019: 1m Digital Elevation
561 Model – Marion Island, Directorate: National Geo-spatial Information, Department of
562 Rural Development and Land Reform, Republic of South Africa.
- 563 Fahnestock, J. T., Povirk, K. L. & Welker, J. M. (2000). Ecological significance of litter
564 redistribution by wind and snow in arctic landscapes. *Ecography* **23**: 623-631.
- 565 Fernández-Palacios, J. M. & Nicolás, J. P. (1995). Altitudinal pattern of vegetation variation on
566 Tenerife. *Journal of Vegetation Science* **6**: 183-190.

- 567 Filibeck, G., Sperandii, M. G., Bazzichetto, M., Mancini, L. D., Rossini, F. & Cancellieri, L.
568 (2019). Exploring the drivers of vascular plant richness at very fine spatial scale in sub-
569 Mediterranean limestone grasslands (Central Apennines, Italy). *Biodiversity and*
570 *Conservation* **28**: 2701-2725.
- 571 Fitzgerald, N. B. & Kirkpatrick, J. B. (2017). Wind distortion in alpine and Subantarctic plants is
572 constant among life forms but does not necessarily reflect prevailing wind direction.
573 *Arctic, Antarctic, and Alpine Research* **49**: 521-535.
- 574 Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V. & Fitzpatrick, J. W. (2018). Climate change
575 causes upslope shifts and mountaintop extirpations in a tropical bird community.
576 *Proceedings of the National Academy of Sciences* **115**: 11982.
- 577 Friedman, J., Hastie, T. & Tibshirani, R. (2000). Additive logistic regression: A statistical view of
578 boosting *The Annals of Statistics* **28**: 337-407.
- 579 Friedman, J. H. (2001). Greedy function approximation: A gradient boosting machine. *The Annals*
580 *of Statistics* **29**: 1189-1232.
- 581 Gandin, A., Gutjahr, S., Dizengremel, P. & Lapointe, L. (2011). Source-sink imbalance increases
582 with growth temperature in the spring geophyte *Erythronium americanum*. *Journal of*
583 *Experimental Botany* **62**: 3467-3479.
- 584 Gardiner, B., Berry, P. & Moulia, B. (2016). Review: Wind impacts on plant growth, mechanics
585 and damage. *Plant Science* **245**: 94-118.
- 586 Gardner, A. S., Maclean, I. M. D. & Gaston, K. J. (2019). Climatic predictors of species
587 distributions neglect biophysiological meaningful variables. *Diversity and Distributions*
588 **25**: 1318-1333.
- 589 Gleason, H. A. (1939). The individualistic concept of the plant association. *The American Midland*
590 *Naturalist* **21**: 92-110.
- 591 Gómez-Ruiz, E. P. & Lacher Jr, T. E. (2019). Climate change, range shifts, and the disruption of a
592 pollinator-plant complex. *Scientific Reports* **9**: 14048.
- 593 Grace, J. 1977. *Plant Response to Wind*. London: Academic Press.
- 594 Gremmen, N. J. M., Chown, S. L. & Marshall, D. J. (1998). Impact of the introduced grass
595 *Agrostis stolonifera* on vegetation and soil fauna communities at Marion Island, sub-
596 Antarctic. *Biological Conservation* **85**: 223-231.

- 597 Hadley, J. L. & Smith, W. K. (1983). Influence of wind exposure on needle desiccation and
598 mortality for timberline conifers in Wyoming, U.S.A. *Arctic and Alpine Research* **15**: 127-
599 135.
- 600 Hadley, J. L. & Smith, W. K. (1986). Wind effects on needles of timberline conifers: Seasonal
601 influence on mortality. *Ecology* **67**: 12-19.
- 602 Hedding, D. W., Nel, W. & Anderson, R. L. (2015). Aeolian processes and landforms in the sub-
603 Antarctic: preliminary observations from Marion Island. *Polar Research* **34**: 26365.
- 604 Hendershot, W. H., Lalonde, H. & Duquette, H. 2008. Soil reaction and exchangeable acidity. *In*:
605 Carter, M. R. & Gregorich, E. G. (eds.) *Soil sampling and methods of analysis*. 2 ed. Boca
606 Raton: CRC Press.
- 607 Holtmeier, F.-K. & Broll, G. (2010). Wind as an ecological agent at treelines in North America,
608 the Alps, and the European subarctic. *Physical Geography* **31**: 203-233.
- 609 Jeong, D. I. & Sushama, L. (2019). Projected changes to mean and extreme surface wind speeds
610 for North America based on regional climate model simulations. *Atmosphere* **10**: 497.
- 611 Kardol, P., Campany, C. E., Souza, L., Norby, R. J., Weltzin, J. F. & Classen, A. T. (2010).
612 Climate change effects on plant biomass alter dominance patterns and community
613 evenness in an experimental old-field ecosystem. *Global Change Biology* **16**: 2676-2687.
- 614 Katsaprakakis, D. A. & Christakis, D. G. 2012. Wind parks design, including representative case
615 studies. *In*: Sayigh, A. (ed.) *Comprehensive Renewable Energy*. Oxford: Elsevier.
- 616 Kemppinen, J., Niittynen, P., Aalto, J., Roux, P. C. I. & Luoto, M. (2019). Water as a resource,
617 stress and disturbance shaping tundra vegetation. *Oikos* **128**: 811-822.
- 618 Kirkpatrick, J. B., Bridle, K. L. & Lynch, A. J. J. (2002). Changes in alpine vegetation related to
619 geomorphological processes and climatic change on Hill One, Southern Range, Tasmania,
620 1989 - 2000. *Australian Journal of Botany* **50**: 753-759.
- 621 Kling, M. M. & Ackerly, D. D. (2020). Global wind patterns and the vulnerability of wind-
622 dispersed species to climate change. *Nature Climate Change* **10**: 868-875.
- 623 Kolstad, E. W. & Johansson, K. A. (2011). Uncertainties associated with quantifying climate
624 change impacts on human health: A case study for diarrhea. *Environmental Health*
625 *Perspectives* **119**: 299-305.
- 626 Kullman, L. & Loyer, N. (2005). Wind-conditioned 20th century decline of Birch treeline
627 vegetation in the Swedish Scandes. *Arctic* **58**: 286-294.

- 628 Lahav, E. & Zamet, D. (1999). Flowers, fruitlets and fruit drop in avocado trees *Revista Chapingo*
629 *Serie Horticultura* **5**: 95-100.
- 630 le Roux, P. C. 2008. Climate and climate change. *In*: Chown, S. L. & Froneman, P. W. (eds.) *The*
631 *Prince Edward Islands: Land-sea interactions in a changing ecosystem*. Stellenbosch:
632 SUN PRESS.
- 633 le Roux, P. C., Aalto, J. & Luoto, M. (2013a). Soil moisture's underestimated role in climate
634 change impact modelling in low-energy systems. *Global Change Biology* **19**: 2965-2975.
- 635 le Roux, P. C., Lenoir, J., Pellissier, L., Wisz, M. S. & Luoto, M. (2013b). Horizontal, but not
636 vertical, biotic interactions affect fine-scale plant distribution patterns in a low-energy
637 system. *Ecology* **94**: 671-682.
- 638 le Roux, P. C. & Luoto, M. (2014). Earth surface processes drive the richness, composition and
639 occurrence of plant species in an arctic–alpine environment. *Journal of Vegetation Science*
640 **25**: 45-54.
- 641 le Roux, P. C. & McGeoch, M. A. (2010). Interaction intensity and importance along two stress
642 gradients: Adding shape to the stress-gradient hypothesis. *Oecologia* **162**: 733-745.
- 643 Lenoir, J., Gégout, J.-C., Guisan, A., Vittoz, P., Wohlgemuth, T., Zimmermann, N. E., Dullinger,
644 S., Pauli, H., Willner, W. & Svenning, J.-C. (2010). Going against the flow: Potential
645 mechanisms for unexpected downslope range shifts in a warming climate. *Ecography* **33**:
646 295-303.
- 647 Liu, H., Mi, Z., Lin, L., Wang, Y., Zhang, Z., Zhang, F., Wang, H., Liu, L., Zhu, B., Cao, G.,
648 Zhao, X., Sanders, N. J., Classen, A. T., Reich, P. B. & He, J.-S. (2018). Shifting plant
649 species composition in response to climate change stabilizes grassland primary production.
650 *Proceedings of the National Academy of Sciences* **115**: 4051.
- 651 Liu, Y.-H., Offler, C. & Ruan, Y.-L. (2013). Regulation of fruit and seed response to heat and
652 drought by sugars as nutrients and signals. *Frontiers in Plant Science* **4**: 282.
- 653 Lönnell, N., Norros, V., Sundberg, S., Rannik, Ü., Johansson, V., Ovaskainen, O. & Hylander, K.
654 (2015). Testing a mechanistic dispersal model against a dispersal experiment with a wind-
655 dispersed moss. *Oikos* **124**: 1232-1240.
- 656 Lugo, A. E. (2008). Visible and invisible effects of hurricanes on forest ecosystems: An
657 international review. *Austral Ecology* **33**: 368-398.
- 658 Lynch, A. J. J. & Kirkpatrick, J. B. (1995). Pattern and process in alpine vegetation and landforms
659 at Hill One, Southern Range, Tasmania. *Australian Journal of Botany* **43**: 537-554.

- 660 Maclean, I. M. D. (2020). Predicting future climate at high spatial and temporal resolution. *Global*
 661 *Change Biology* **26**: 1003-1011.
- 662 Mau, A. C., Reed, S. C., Wood, T. E. & Cavaleri, M. A. (2018). Temperate and tropical forest
 663 canopies are already functioning beyond their thermal thresholds for photosynthesis.
 664 *Forests* **9**: 47.
- 665 McCune, B. (2007). Improved estimates of incident radiation and heat load using non-parametric
 666 regression against topographic variables. *Journal of Vegetation Science* **18**: 751-754.
- 667 McCune, B. & Keon, D. (2002). Equations for potential annual direct incident radiation and heat
 668 load. *Journal of Vegetation Science* **13**: 603-606.
- 669 McGlone, M. S. (2002). The Late Quaternary peat, vegetation and climate history of the Southern
 670 Oceanic Islands of New Zealand. *Quaternary Science Reviews* **21**: 683-707.
- 671 McIntire, E. J. B., Piper, F. I. & Fajardo, A. (2016). Wind exposure and light exposure, more than
 672 elevation-related temperature, limit tree line seedling abundance on three continents.
 673 *Journal of Ecology* **104**: 1379-1390.
- 674 Menéndez, R., Megías, A. G., Hill, J. K., Braschler, B., Willis, S. G., Collingham, Y., Fox, R.,
 675 Roy, D. B. & Thomas, C. D. (2006). Species richness changes lag behind climate change.
 676 *Proceedings of the Royal Society B: Biological Sciences* **273**: 1465-1470.
- 677 Mitchell, S. J. (2012). Wind as a natural disturbance agent in forests: a synthesis. *Forestry: An*
 678 *International Journal of Forest Research* **86**: 147-157.
- 679 Mod, H. K., Scherrer, D., Luoto, M., Guisan, A. & Scheiner, S. (2016). What we use is not what
 680 we know: environmental predictors in plant distribution models. *Journal of Vegetation*
 681 *Science* **27**: 1308-1322.
- 682 Møller, A. P. (2013). Long-term trends in wind speed, insect abundance and ecology of an
 683 insectivorous bird. *Ecosphere* **4**: art6.
- 684 Momberg, M., Hedding, D. W., Luoto, M., le Roux, P. C. (2021). Exposing wind stress as a driver
 685 of fine-scale variation in plant communities. Dryad Dataset,
 686 <https://doi.org/10.5061/dryad.547d7wm7k>.
- 687 Morueta-Holme, N., Engemann, K., Sandoval-Acuña, P., Jonas, J. D., Segnitz, R. M. & Svenning,
 688 J.-C. (2015). Strong upslope shifts in Chimborazo's vegetation over two centuries since
 689 Humboldt. *Proceedings of the National Academy of Sciences* **112**: 12741.
- 690 Müller, M. 2012. Generalized linear models. *Handbook of Computational Statistics*. Springer.

- 691 Niittynen, P., Heikkinen, R., Aalto, J., Guisan, A., Kemppinen, J. & Luoto, M. (2020a). Fine-scale
692 tundra vegetation patterns are strongly related to winter thermal conditions. *Nature Climate*
693 *Change*.
- 694 Niittynen, P., Heikkinen, R. & Luoto, M. (2020b). Decreasing snow cover alters functional
695 composition and diversity of Arctic tundra. *Proceedings of the National Academy of*
696 *Sciences* **117**: 202001254.
- 697 Niittynen, P., Heikkinen, R. K. & Luoto, M. (2018). Snow cover is a neglected driver of Arctic
698 biodiversity loss. *Nature Climate Change* **8**: 997-1001.
- 699 Niittynen, P. & Luoto, M. (2018). The importance of snow in species distribution models of arctic
700 vegetation. *Ecography* **41**: 1024-1037.
- 701 Nyberg, J., Malmgren, B. A., Winter, A., Jury, M. R., Kilbourne, K. H. & Quinn, T. M. (2007).
702 Low Atlantic hurricane activity in the 1970s and 1980s compared to the past 270 years.
703 *Nature* **447**: 698-701.
- 704 Nylén, T., Hellemaa, P. & Luoto, M. (2015). Determinants of sediment properties and organic
705 matter in beach and dune environments based on boosted regression trees. *Earth Surface*
706 *Processes and Landforms* **40**: 1137-1145.
- 707 Nylén, T. & Luoto, M. (2015). Influence of patch size and connectivity on beach and dune species
708 in land-uplift coasts. *Plant Ecology & Diversity* **9**: 35-44.
- 709 Okitsu, S. & Ito, K. (1984). Vegetation dynamics of the Siberian dwarf pine (*Pinus pumila* Regel)
710 in the Taisetsu mountain range, Hokkaido, Japan. *Vegetatio* **58**: 105-113.
- 711 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
712 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E. & Wagner, H.
713 (2018). vegan: Community Ecology Package. *R package version 2.5-2* [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
714 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan).
- 715 Onoda, Y. & Anten, N. P. R. (2011). Challenges to understand plant responses to wind. *Plant*
716 *Signaling & Behavior* **6**: 1057-1059.
- 717 Pammenter, N. W., Drennan, P. M. & Smith, V. R. (1986). Physiological and anatomical aspects
718 of photosynthesis of two *Agrostis* species at a sub-Antarctic island. *New Phytologist* **102**:
719 143-160.
- 720 Parmesan, C. & Hanley, M. E. (2015). Plants and climate change: Complexities and surprises.
721 *Annals of Botany* **116**: 849-864.

722 Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I.-C., Clark, T. D.,
723 Colwell, R. K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S., Frusher, S., Garcia, R.
724 A., Griffis, R. B., Hobday, A. J., Janion-Scheepers, C., Jarzyna, M. A., Jennings, S.,
725 Lenoir, J., Linnetved, H. I., Martin, V. Y., McCormack, P. C., McDonald, J., Mitchell, N.
726 J., Mustonen, T., Pandolfi, J. M., Pettorelli, N., Popova, E., Robinson, S. A., Scheffers, B.
727 R., Shaw, J. D., Sorte, C. J. B., Strugnell, J. M., Sunday, J. M., Tuanmu, M.-N., Vergés,
728 A., Villanueva, C., Wernberg, T., Wapstra, E. & Williams, S. E. (2017). Biodiversity
729 redistribution under climate change: Impacts on ecosystems and human well-being.
730 *Science* **355**: eaai9214.

731 Pendlebury, S. & Barnes-Keoghan, I. (2007). Climate and climate change in the sub-Antarctic.
732 *Papers and Proceedings of the Royal Society of Tasmania* **141**: 67-81.

733 R Core Team 2018. R: A language and environment for statistical computing. Vienna, Austria: R
734 Foundation for Statistical Computing.

735 Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P.,
736 Adriaensen, F., Ahola, M. P., Arcese, P., Miguel Avilés, J., Balbontin, J., Berg, K. S.,
737 Borrás, A., Burthe, S., Clobert, J., Dehnhard, N., de Lope, F., Dhondt, A. A., Dingemanse,
738 N. J., Doi, H., Eeva, T., Fickel, J., Filella, I., Fossøy, F., Goodenough, A. E., Hall, S. J. G.,
739 Hansson, B., Harris, M., Hasselquist, D., Hickler, T., Joshi, J., Kharouba, H., Martínez, J.
740 G., Mihoub, J.-B., Mills, J. A., Molina-Morales, M., Moksnes, A., Ozgul, A., Parejo, D.,
741 Pilard, P., Poisbleau, M., Rousset, F., Rödel, M.-O., Scott, D., Senar, J. C., Stefanescu, C.,
742 Stokke, B. G., Kusano, T., Tarka, M., Tarwater, C. E., Thonicke, K., Thorley, J., Wilting,
743 A., Tryjanowski, P., Merilä, J., Sheldon, B. C., Pape Møller, A., Matthysen, E., Janzen, F.,
744 Dobson, F. S., Visser, M. E., Beissinger, S. R., Courtiol, A. & Kramer-Schadt, S. (2019).
745 Adaptive responses of animals to climate change are most likely insufficient. *Nature*
746 *Communications* **10**: 3109.

747 Smith, V. R. & Steenkamp, M. (2001). Classification of the terrestrial habitats on Marion Island
748 based on vegetation and soil chemistry. *Journal of Vegetation Science* **12**: 181-198.

749 Smith, V. R., Steenkamp, M. & Gremmen, N. J. M. (2001). Terrestrial habitats on sub-Antarctic
750 Marion Island: Their vegetation, edaphic attributes, distribution and response to climate
751 change. *South African Journal of Botany* **67**: 641-654.

- 752 Song, Y.-B., Hu, Y.-K., Pan, X., Liu, G.-F., Xiong, W., Dong, M. & Cornelissen, J. H. C. (2020).
 753 Association of leaf silicon content with chronic wind exposure across and within
 754 herbaceous plant species. *Global Ecology and Biogeography* **29**: 711-721.
- 755 Sparacino, J., Renison, D., Devegili, A. M. & Suarez, R. (2020). Wind protection rather than soil
 756 water availability contributes to the restriction of high-mountain forest to ravines. *New*
 757 *Forests* **51**: 101-117.
- 758 Sturm, M., Liston, G. E., Benson, C. S. & Holmgren, J. (2001). Characteristics and growth of a
 759 snowdrift in Arctic Alaska, U.S.A. *Arctic, Antarctic, and Alpine Research* **33**: 319-329.
- 760 Sun, S.-S., Liu, X.-P., He, Y.-H., Wei, S.-L., Zhang, L.-M., Lv, P., Bao, C., Wang, M.-M. &
 761 Cheng, L. (2019). Responses of annual herb plant community characteristics to increased
 762 precipitation and reduced wind velocity in semiarid sandy grassland. *Ecology and*
 763 *Evolution* **0**: 1-11.
- 764 Sutherland, W. J., Barnard, P., Broad, S., Clout, M., Connor, B., Côté, I. M., Dicks, L. V., Doran,
 765 H., Entwistle, A. C., Fleishman, E., Fox, M., Gaston, K. J., Gibbons, D. W., Jiang, Z.,
 766 Keim, B., Lickorish, F. A., Markillie, P., Monk, K. A., Pearce-Higgins, J. W., Peck, L. S.,
 767 Pretty, J., Spalding, M. D., Tonneijck, F. H., Wintle, B. C. & Ockendon, N. (2017). A 2017
 768 horizon scan of emerging issues for global conservation and biological diversity. *Trends in*
 769 *Ecology & Evolution* **32**: 31-40.
- 770 Thom, D. & Seidl, R. (2016). Natural disturbance impacts on ecosystem services and biodiversity
 771 in temperate and boreal forests. *Biological Reviews of the Cambridge Philosophical*
 772 *Society* **91**: 760-781.
- 773 Thomas, H. J. D., Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Kattge, J., Diaz, S.,
 774 Vellend, M., Blok, D., Cornelissen, J. H. C., Forbes, B. C., Henry, G. H. R., Hollister, R.
 775 D., Normand, S., Prevéy, J. S., Rixen, C., Schaepman-Strub, G., Wilmking, M., Wipf, S.,
 776 Cornwell, W. K., Beck, P. S. A., Georges, D., Goetz, S. J., Guay, K. C., Rüger, N.,
 777 Soudzilovskaia, N. A., Spasojevic, M. J., Alatalo, J. M., Alexander, H. D., Anadon-Rosell,
 778 A., Angers-Blondin, S., te Beest, M., Berner, L. T., Björk, R. G., Buchwal, A., Buras, A.,
 779 Carbognani, M., Christie, K. S., Collier, L. S., Cooper, E. J., Elberling, B., Eskelinen, A.,
 780 Frei, E. R., Grau, O., Grogan, P., Hallinger, M., Heijmans, M. M. P. D., Hermanutz, L.,
 781 Hudson, J. M. G., Johnstone, J. F., Hülber, K., Iturrate-Garcia, M., Iversen, C. M.,
 782 Jaroszynska, F., Kaarlejarvi, E., Kulonen, A., Lamarque, L. J., Lantz, T. C., Lévesque, E.,
 783 Little, C. J., Michelsen, A., Milbau, A., Nabe-Nielsen, J., Nielsen, S. S., Ninot, J. M.,

784 Oberbauer, S. F., Olofsson, J., Onipchenko, V. G., Petraglia, A., Rumpf, S. B., Shetti, R.,
 785 Speed, J. D. M., Suding, K. N., Tape, K. D., Tomaselli, M., Trant, A. J., Treier, U. A.,
 786 Tremblay, M., Venn, S. E., Vowles, T., Weijers, S., Wookey, P. A., Zamin, T. J., Bahn,
 787 M., Blonder, B., van Bodegom, P. M., Bond-Lamberty, B., Campetella, G., Cerabolini, B.
 788 E. L., Chapin, F. S., Craine, J. M., Dainese, M., Green, W. A., Jansen, S., Kleyer, M.,
 789 Manning, P., Niinemets, Ü., Onoda, Y., Ozinga, W. A., Peñuelas, J., Poschlod, P., et al.
 790 (2020). Global plant trait relationships extend to the climatic extremes of the tundra biome.
 791 *Nature Communications* **11**: 1351.

792 Ulrich, W. (2009). Nestedness analysis as a tool to identify ecological gradients. *Ecological*
 793 *Questions* **11**: 27-34.

794 Versteeg, H. K. & Malalasekera, W. 2007. *An introduction to computational fluid dynamics: The*
 795 *finite volume method*. Pearson Education Limited.

796 Vitt, D., Crandall-Stotler, B. & Wood, A. 2014. Bryophytes: Survival in a dry world through
 797 tolerance and avoidance. In: Rajakaruna, N., Boyd, R. & Harris, T. (eds.) *Plant Ecology*
 798 *and Evolution in Harsh Environments*. Hauppauge: Nova Publishers.

799 Vonlanthen, C., Bühler, A., Veit, H., Kammer, P. & Eugster, W. (2006a). Alpine plant
 800 communities: A statistical assessment of their relation to microclimatological, pedological,
 801 geomorphological, and other factors. *Physical Geography* **27**: 137-154.

802 Vonlanthen, C. M., Kammer, P., Eugster, W., Bühler, A. & Veit, H. (2006b). Alpine vascular
 803 plant species richness: The importance of daily maximum temperature and pH. *Plant*
 804 *Ecology* **184**: 13-25.

805 Watanabe, E. & Hasumi, H. (2005). Arctic sea ice response to wind stress variations. *Journal of*
 806 *Geophysical Research: Oceans* **110**: C11007.

807 Whitehead, F. H. (1954). A study of the relation between growth form and exposure on Monte
 808 Maiella, Italy. *Journal of Ecology* **42**: 180-186.

809 Whitehead, F. H. (1959). Vegetational changes in response to alterations of surface roughness on
 810 M. Maiella, Italy. *Journal of Ecology* **47**: 603-606.

811 Williams, J. W. & Jackson, S. T. (2007). Novel climates, no-analog communities, and ecological
 812 surprises. *Frontiers in Ecology and the Environment* **5**: 475-482.

813 Williams, R. J. & Ashton, D. H. (1987). The composition, structure and distribution of heathland
 814 and grassland communities in the subalpine tract of the Bogong High Plains, Victoria.
 815 *Australian Journal of Ecology* **12**: 57-71.

- 816 Wilson, J. W. (1959). Notes on wind and its effects in arctic-alpine vegetation. *Journal of Ecology*
817 **47**: 415-427.
- 818 Wood, S. N. 2006. *Generalized Additive Models: An Introduction with R*. Boca Raton: Chapman
819 and Hall.
- 820 Wood, S. N. & Augustin, N. H. (2002). GAMs with integrated model selection using penalized
821 regression splines and applications to environmental modelling. *Ecological Modelling* **157**:
822 157-177.
- 823 Xuan, Z. & Chang, N.-B. (2014). Modeling the climate-induced changes of lake ecosystem
824 structure under the cascade impacts of hurricanes and droughts. *Ecological Modelling* **288**:
825 79-93.
- 826 Yang, M., Défossez, P., Danjon, F. & Fourcaud, T. (2014). Tree stability under wind: Simulating
827 uprooting with root breakage using a finite element method. *Annals of Botany* **114**: 695-
828 709.
- 829 Young, I. R. & Ribal, A. (2019). Multiplatform evaluation of global trends in wind speed and
830 wave height. *Science* **364**: 548.
- 831 Young, I. R., Zieger, S. & Babanin, A. V. (2011). Global trends in wind speed and wave height.
832 *Science* **332**: 451.
- 833 Zeng, Z., Ziegler, A. D., Searchinger, T., Yang, L., Chen, A., Ju, K., Piao, S., Li, L. Z. X., Ciais,
834 P., Chen, D., Liu, J., Azorin-Molina, C., Chappell, A., Medvigy, D. & Wood, E. F. (2019).
835 A reversal in global terrestrial stilling and its implications for wind energy production.
836 *Nature Climate Change* **9**: 979-985.
- 837 Zhang, X., Zhang, B., Feeley, K. J., Wang, G. G., Zhang, J. & Zhai, L. (2019). Ecological
838 contingency in species shifts: downslope shifts of woody species under warming climate
839 and land-use change. *Environmental Research Letters* **14**: 114033.
- 840 Zhang, Y., Liu, C., Tang, Y. & Yang, Y. (2007). Trends in pan evaporation and reference and
841 actual evapotranspiration across the Tibetan Plateau. *Journal of Geophysical Research:*
842 *Atmospheres* **112**.
- 843 Zheng, M., Song, J., Ru, J., Zhou, Z., Zhong, M., Jiang, L., Hui, D. & Wan, S. (2020). Effects of
844 grazing, wind erosion, and dust deposition on plant community composition and structure
845 in a temperate steppe. *Ecosystems*.

846

869 **Table 1** A summary of key papers highlighting the types of research questions that have been examined,
 870 and illustrating broadly the currently knowledge in this field.

	Study location	Ecosystem	Spatial scale	Result	Reference
Richness	Mongolia, China	Semi-arid temperate steppe	4 x 4 m plots	Wind erosion did not affect species richness, while dust deposition (due to wind erosion) reduced species richness.	Zheng et al. (2020)
Vegetation cover	Hill One, Southern range, Tasmania	Alpine	1.5 x 1.0 m plots	Vegetation cover decreased over 11 years, mostly due to erosion, caused by an increase in wind speed.	Kirkpatrick et al. (2002)
	Sonoran Desert, Arizona, USA	Desert, desert remnants and urban sites	20 individually potted plants at each site	Reduced wind speed increased cover of individuals in desert and desert remnants. In urban sites reduced wind speed had no effect on cover.	Bang et al. (2010)
	Mongolia, China	Semi-arid temperate steppe	4 x 4 m plots	Wind erosion decreased total plant cover. Dust deposition by wind resulted in higher vegetation cover.	Zheng et al. (2020)
	Swiss Alps	Alpine	1 x 1 m plots	Wind speed was an important driver of species composition; $R^2 > 0.35$.	Vonlanthen et al. (2006a)
Distribution of vegetation types	North America, the Alps, European subarctic	Alpine	Review	Wind determines the treeline position in three different ecosystems and can override the role of heat deficiency.	Holtmeier and Broll (2010)
	Swedish Scandes	Alpine	10 x 10 m plots	Wind determines the birch treeline and increasing wind circulation constrains upslope treeline expansion in response to warming temperatures.	Kullman and Loyer (2005)
	5 mountain regions from 3 continents	Alpine	50 m ² plots	Wind exposure was the main determinant of the number of seedlings present at the treeline, with increased wind exposure correlated with fewer tree seedlings.	McIntire et al. (2016)

	Central Argentina	Forest patches	396 individual saplings	Reduced wind speeds likely restrict the occurrence of forest to ravines.	Sparacino et al. (2020)
Review	Multiple locations	Boreal & temperate forest		A review of the impact of disturbance agents on natural vegetation showed 146 studies of wind as a disturbance agent versus 677 for fire. More than 87% of these studies are from Europe or North America. There was no difference between the effects of wind, fire and bark beetles on biodiversity.	Thom and Seidl (2016)

Table 2 Vegetation and abiotic characteristics of each grid. See Fig. 1 for the location of the sampling grids. VWC: volumetric water content

Grid	Vascular plant cover (mean; %)	Total vascular plant richness	Summer temperature (°C)			Winter temperature (°C)			Soil moisture range (% VWC)	Wind stress	
			Minimum	Mean	Maximum	Minimum	Mean	Maximum		Mean maximum	Absolute maximum
1	72	9	2.92	6.28	8.12	1.62	4.02	5.92	76.2	5.50	6.06
2	65	15	2.91	6.35	7.71	2.66	4.70	6.46	41.2	6.52	7.02
3	65	12	5.56	7.70	12.86	2.61	4.83	6.21	69.2	7.62	8.22
4	55	12	2.72	5.77	11.22	2.19	4.10	5.49	68.6	8.22	8.53
5	48	10	4.42	5.50	6.62	3.58	5.34	6.28	48.7	6.22	6.46
6	67	9	4.24	6.12	10.54	2.48	4.81	6.48	38.7	6.10	6.60
7	64	9	4.60	6.66	8.10	4.13	5.73	7.63	48.1	6.18	6.46
8	25	11	3.83	5.02	6.33	0.00	1.06	2.45	27.6	7.18	7.64
9	19	10	4.72	5.97	7.22	0.36	1.70	3.26	53.4	7.48	7.60

Table 3 Model fit and variable importance for all variables when predicting spatial variation in species richness and vegetation cover in the full model. The five most important predictors for each approach are highlighted in bold and significant predictors are indicated with an asterisk. GAM = generalized additive model; GLZ = generalized linear model. See Table A2 for results from the generalized boosted regression models.

	Richness		Cover	
	GAM	GLZ	GAM	GLZ
Deviance explained (%)	24.9	19.5	67.7	66.9
Relative importance (%)				
Rock cover	16.7*	12.6*	94.15*	94.63*
Wind stress: maximum	33.8*	6.0	3.00*	2.88*
Soil pH	19.4*	16.0*	0.04	0.06
Soil depth	12.2*	13.8	0.17	0.21
Temperature: summer	1.5	1.7	0.90*	0.12
Temperature: winter	13.8*	44.5*	0.49	0.64
PDIR	0.0	1.3	0.70*	0.52*
Soil moisture	2.6	4.0	0.65*	0.94*

Figure legends

Figure 1 Study site location on Marion Island and study design. a: Marion Island, with contour intervals of 150 m and study location indicated by the red block; b: the location of all 9 study grids (each 8 x 20 m; the distance from grid 1 to grid 9 is ~900 m; Google Earth imagery), with the Van den Boogaard river indicated in blue; c: biotic and abiotic data were sampled from 160 quadrats within each grid, resulting in a total of 1440 quadrats (each 1 x 1 m in size; indicated here with the white frame); d: drone imagery was obtained for each grid, from which a digital surface model was created and used to calculate the wind exposure of each quadrat; e: wind speed for each grid was obtained from a computational fluid dynamics model of island-scale wind speed; f: wind exposure (calculated from drone imagery products) and wind speed were combined to create a wind stress metric for each quadrat.

Figure 2 Variation in maximum wind stress in all nine study grids (see Fig. 1 for location of the grids) indicated for each 1 m² quadrat. Cooler colours indicate low wind stress and warmer colours indicate higher wind stress.

Figure 3 Species richness across all nine study grids (see Fig. 1 for location of the grids), with vascular plant species richness indicated for each 1 m² quadrat. Cooler colours indicate low species richness and warmer colours indicate higher richness.

Figure 4 Individual predictor variables' response curves for species richness in GAM models.

Tick marks on the x-axis indicate observed values. Solid black lines represent the response curves from the simple model (excluding wind stress predictor variables), while dashed red lines represent the response curves from the full model.

Figure 5 Individual predictor variables' response curves for vegetation cover in GAM models.

Tick marks on the x-axis indicate observed values. Solid black lines represent the response curves in the simple model (excluding wind stress predictor variables), while dashed red lines represent the response curves in the full model.

Figure 6 a) Non-metric multidimensional scaling (nMDS) ordination plot showing variables impact on species composition (Stress = 0.18), symbols indicate sampled quadrats; b) Variable importance, as assessed by marginal R^2 values, from the PERMANOVA. Temp = soil temperature.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6