<https://helda.helsinki.fi>

MED27 Variants Cause Developmental Delay, Dystonia, and Cerebellar Hypoplasia

Meng, Linyan

2021-04

Meng , L , Isohanni , P , Shao , Y , Graham , B H , Hickey , S E , Brooks , S , Suomalainen , A , Joset , P , Steindl , K , Rauch , A , Hackenberg , A , High , F A , Armstrong-Javors , A , Mencacci , N E , Gonzalez-Latapi , P , Kamel , W A , Al-Hashel , J Y , Bustos , B , Hernandez , A , Krainc , D , Lubbe , S J , Van Esch , H , De Luca , C , Ballon , K , Ravelli , C , Burglen , L , Qebibo , L , Calame , D G , Mitani , T , Marafi , D , Pehlivan , D , Saadi , N W , Sahin , Y , Maroofian , R , Efthymiou , S , Houlden , H , Maqbool , S , Rahman , F , Gu , S , Posey , J E , Lupski , J R , Hunter , J , Wangler , M F , Carroll , C J & Yang , Y 2021 , ' MED27 Variants Cause Developmental Delay, Dystonia, and Cerebellar Hypoplasia ' , Annals of Neurology , vol. 89 , no. 4 , pp. 828-833 . <https://doi.org/10.1002/ana.26019>

<http://hdl.handle.net/10138/340557>

<https://doi.org/10.1002/ana.26019>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

***MED27* variants cause Developmental Delay, Dystonia, and Cerebellar Hypoplasia**

Running head: *MED27* associated with novel neurodevelopmental syndrome

Linyan Meng, PhD^{1,2*}, Pirjo Isohanni, MD, PhD^{3,4}, Yunru Shao, MS^{1,5,6}, Brett H. Graham, MD, PhD⁷, Scott E. Hickey, MD⁸, Stephanie Brooks, MS⁸, Anu Suomalainen, PhD³, Pascal Joset, PhD⁹, Katharina Steindl, MD⁹, Anita Rauch, PhD⁹, Annette Hackenberg, MD¹⁰, Frances A. High, MD, PhD^{11, 12}, Amy Armstrong-Javors, MD^{12, 13}, Niccolò E. Mencacci, MD, PhD^{14,15}, Paulina González-Latapi, MD¹⁴, Walaa A. Kamel, MD^{16,17}, Jasem Y. Al-Hashel, MD^{16,18}, Bernabé I. Bustos, PhD^{14,15}, Alejandro V. Hernandez, BS¹⁴, Dimitri Krainc, MD, PhD^{14,15}, Steven J. Lubbe, PhD^{14,15}, Hilde Van Esch, MD¹⁹, Chiara De Luca, MD¹⁹, Katleen Ballon, MD²⁰, Claudia Ravelli, MD²¹, Lydie Burglen, MD, PhD^{22,23}, Leila Qebibo, MD²², Daniel G. Calame, MD, PhD^{1,6,24}, Tadahiro Mitani, MD, PhD¹, Dana Marafi, MD^{1,25}, Davut Pehlivan, MD^{1,6,24}, Nebal Waill Saadi, MD^{26,27}, Yavuz Sahin, MD²⁸, Reza Maroofian, PhD²⁹, Stephanie Efthymiou, MD²⁹, Henry Houlden, MD²⁹, Shazia Maqbool, MD³⁰, Fatima Rahman, MD³⁰, Shen Gu, PhD³¹, Jennifer E Posey, MD, PhD¹, James R. Lupski, MD, PhD, DSc (hon)^{1,5,6,32}, Jill V. Hunter, MD^{33,34}, Michael F. Wangler, MD, PhD^{1,4,35}, Christopher J Carroll, PhD^{3,36}, Yaping Yang, PhD^{1,2* **}

1. Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA.
2. Baylor Genetics, Houston, TX, USA.
3. Research Programs Unit, Stem Cells and Metabolism, Faculty of Medicine, University of Helsinki, Helsinki, Finland
4. Department of Child Neurology, Children's Hospital, University of Helsinki and Helsinki University Hospital, Helsinki, Finland
5. Department of Pediatrics, Baylor College of Medicine, Houston, TX, USA.

6. Texas Children's Hospital, Houston, TX, USA.
7. Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, Indiana, USA
8. Department of Pediatrics, The Ohio State University College of Medicine, Division of Genetic & Genomic Medicine, Nationwide Children's Hospital, Columbus, OH, USA
9. Institute of Medical Genetics, University of Zurich, Schlieren-Zurich, Switzerland
10. Department of Pediatric Neurology, University Children's Hospital Zurich, University of Zurich, Switzerland
11. Division of Medical Genetics, Massachusetts General Hospital, Boston, MA
12. Harvard Medical School, Boston, MA
13. Department of Pediatric Neurology, Massachusetts General Hospital, Boston, MA
14. Ken and Ruth Davee Department of Neurology, Feinberg School of Medicine, Northwestern University, Chicago, IL
15. Simpson Querrey Center for Neurogenetics, Feinberg School of Medicine, Northwestern University, Chicago, IL
16. Department of Neurology, Ibn Sina Hospital, Kuwait City, Kuwait
17. Department of Neurology, Faculty of Medicine, Beni-Suef University, Egypt
18. Department of Medicine, Faculty of Medicine, Kuwait University, Kuwait
19. Centre for Human Genetics, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium
20. Centre for Developmental Disabilities, University Hospitals Leuven, Herestraat 49, 3000 Leuven, Belgium
21. Pediatric neurology Department, Neurogenetics Reference Center, I-motion Institut, AP-HP.Sorbonne Université, Armand Trousseau Hospital, Paris, France

22. Cerebellar Malformations and Congenital diseases Reference Center and Neurogenetics Lab, Department of Genetics, AP-HP.Sorbonne Université, Armand Trousseau Hospital, Paris, France
23. Developmental Brain Disorders Laboratory, Imagine Institute, INSERM UMR 1163, Paris, France
24. Division of Neurology and Developmental Neuroscience, Department of Pediatrics, BCM, Houston, TX 77030, USA
25. Department of Pediatrics, Faculty of Medicine, Kuwait University, P.O. Box 24923, 13110 Safat, Kuwait
26. College of Medicine, Baghdad University, Iraq
27. Children Welfare Teaching Hospital, Baghdad, Iraq
28. Department of Medical Genetics, Genoks Genetic Center, Ankara, Turkey
29. Department of Neuromuscular Disorders, UCL Institute of Neurology, London WC1N 3BG, UK
30. Development and Behavioural Pediatrics Department, Institute of Child Health and The Children Hospital, Lahore, Pakistan
31. School of Biomedical Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong S.A.R
32. Human Genome Sequencing Center, Baylor College of Medicine, Houston, Texas, 77030, USA
33. EB Singleton Department of Pediatric Radiology, Texas Children's Hospital, TX, USA
34. Department of Radiology, Baylor College of Medicine, Houston, TX, USA
35. Jan and Dan Neurological Research Institute, Texas Children's Hospital, Houston, TX, USA.
36. Molecular and Clinical Sciences Research Institute, St. George's, University of London, Cranmer Terrace, London, SW17 0RE, UK.

* co-corresponding authors

** Current address: AiLife Diagnostics, 1920 Country Place Suite 100, Pearland, TX, USA.

Linyan Meng,

Department of Molecular and Human Genetics

Baylor College of Medicine

Houston, TX 77030.

Tel: 281.849.8871

Email: lmeng@bcm.edu

Yaping Yang

Department of Molecular and Human Genetics

Baylor College of Medicine

Houston, TX 77030

Email: yapingy@ailifeus.com

Number of characters in the title: 77

Number of characters in the running head: 50

Number of words in the Abstract: 96

Number of words in the Introduction: 162

Number of words in the Discussion: 378

Number of words in the body of manuscript: 1409

Number of figures: 2

Number of colored figures: 0

Number of tables: 1

Key words: *MED27*, neurodevelopmental syndrome

ABSTRACT

The Mediator multiprotein complex functions as a regulator of RNA polymerase II-catalyzed gene transcription. In this study, exome sequencing (ES) detected biallelic putative disease-causing variants in *MED27*, encoding Mediator Complex Subunit 27, in sixteen patients from eleven families with a novel neurodevelopmental syndrome. Patient phenotypes are highly homogeneous including global developmental delay, intellectual disability, axial hypotonia with distal spasticity, dystonic movements, and cerebellar hypoplasia. Seizures and cataracts were noted in severely affected individuals. Identification of multiple patients with biallelic *MED27* variants supports the critical role of MED27 in normal human neural development, particularly for the cerebellum.

INTRODUCTION

The Mediator complex acts as a bridge between transcription factors at enhancers and the basal transcriptional machinery at specific promoters, thereby stabilizing the preinitiation complex and stimulating promoter release^{1,2}. It is also involved in additional aspects of transcriptional regulation, including mRNA and noncoding RNA processing, chromatin remodeling and epigenetic regulation³. *MED12* was the first human disease gene discovered in the Mediator complex, associated with Opitz-Kaveggia syndrome (MIM: 305450) and Lujan-Fryns syndrome (MIM: 309520)^{4,5}. Since then, a total of six genes in the complex have been reported as disease genes mostly in association with neurodevelopmental disorders. However, the majority of the Mediator genes remain without disease associations.

Here, we report a novel neurodevelopmental syndrome with a unique phenotype of global developmental delay, axial hypotonia with appendicular spasticity, dystonic movements, cerebellar hypoplasia, epilepsy, and cataracts. Exome sequencing detected biallelic disease-causing variants in *MED27* (MIM: 605044), a eukaryotic specific subunit thought to represent an ortholog of budding yeast Med3^{6,7}.

PATIENTS AND METHODS

Patients were identified through GeneMatcher and all subjects were examined by the referring physicians⁸. Clinical information including medical notes, facial photos, and MRI images were collected, critically reviewed and compared. Informed consents were obtained from the legal guardians of all subjects. The study was performed in accordance with the guidelines specified by the Institutional Review Boards and Ethics Committees at each institution.

Exome sequencing (ES) on each patient was performed in commercial or academic laboratories per each laboratory's protocol. Targeted Sanger sequencing was performed in probands and

available relatives to validate variant segregation. Variant filtering and prioritization were performed by assessing variant characteristics including general population frequency, variant severity, *in silico* prediction, inheritance modeling, pathway analysis, and family segregation analysis. *MED27* variants were annotated on reference sequence NM_004269.3.

RESULTS

Prior to ES analysis, most reported patients had extensive clinical, metabolic, and genetic investigations, yet had not received a molecular diagnosis. ES analysis followed by targeted Sanger sequencing revealed biallelic variants, either compound heterozygous or homozygous alleles, in *MED27* in sixteen affected individuals (Supplementary Table 1). A total of eleven unique *MED27* variants were identified, including frameshift (3/11), canonical splice-site (1/11), and missense variants (7/11) (Fig 1, Supplementary Table 2). Three recurrent variants [c.776C>T (p.Pro259Leu), c.839C>T (p.Pro280Leu), and c.871G>A (p.Gly291Ser)] affecting CpG sites were identified in multiple families with different ethnic backgrounds. All *MED27* variants are either absent or rare in the Genome Aggregation Database (gnomAD v2). The variants occurred at residues that are extremely conserved from human to drosophila with GERP++ RS score greater than 5.0⁹. Multiple *in silico* programs including PolyPhen2, SIFT¹, Mutation Taster and CADD (Combined Annotation Dependent Depletion) support the deleterious effect of these variants (Supplementary Table 2).^{10,11,12,13} Notably, six out of seven missense variants are located in close proximity near the C-terminal end of the protein (Fig 1).

The clinical phenotypes of each subject are summarized in Table 1 and Supplementary Table 3. The reported cohort consists of 13 pediatric patients (ages 0 to 13 years) and three adult patients (ages 26, 36, and 42 years). Consanguinity was noted by historical report in five of the eleven families. Except for one subject born preterm (11-2), all were born full term with no perinatal complications. Patient 10-2, who was previously published in a large arthrogryposis

cohort study, had a dual molecular diagnosis with homozygous pathogenic variants in both *MED27* and *COG6*, the latter of which is associated with a congenital disorder of glycosylation type III (MIM: 614576).¹⁴ The severe clinical symptoms and early death of this individual were most likely attributable to the *COG6* variant and may have masked the *MED27*-related phenotype. Therefore, this individual was excluded from clinical analysis.

Global developmental delay, ranging from mild to profound, and intellectual disability were observed in all patients. Eight patients were severely affected and were non-verbal and unable to sit or walk independently. Among them, two patients (2, 3-2) also had motor regression. Five patients were moderately delayed developmentally but achieved ambulation and some expressive language. Two siblings (6-1, 6-2) were reported to have normal development until 8-9 years old when both demonstrated progressive difficulties with ambulation, speech articulation, writing, and school performance. Motor and cognitive symptoms progressed until the late teenage years and recent neurocognitive testing demonstrated moderate intellectual disability.

Axial hypotonia was noted in 93.3% (14/15) of patients. Appendicular spasticity and dystonic movements were seen in 86.7% (13/15). These symptoms were especially prominent in two siblings (6-1, 6-2) who experienced generalized dystonia and moderate dysarthria due to involvement of the oromandibular muscles.

Brain MRI demonstrated cerebellar hypoplasia involving the vermis more than the cerebellar hemispheres in 86.7% of patients (Fig 2). Multiple severely affected patients (3-1, 3-2, 4, 5) had strikingly severe vermian hypoplasia. In some patients, additional brain abnormalities were observed, including hypomyelination (5), cerebral atrophy (3-1, 3-2), thin corpus callosum (3-1, 3-2, 5), and enlarged ventricles (5, 7). Progressive atrophy involving the cerebrum, cerebellum

and/or basal ganglia was seen in four patients (2, 3-1, 4 and 5). Microcephaly was present in 28.6% of patients (4/14, 2, 3-1, 3-2, 5).

Epilepsy was present in 60.0% (9/15) of patients with an age of onset ranging from 20 days to 5 years. Reported seizure types were varied and included focal motor seizures (3/7; 43%), generalized tonic-clonic seizures (2/7; 29%), hemifacial clonic seizures (2/7; 29%), generalized myoclonic seizures (1/7; 14%), epileptic (infantile) spasms (1/7; 14%), atonic seizures (1/7; 14%) and atypical absence seizures (1/7; 14%). Epilepsy was drug-resistant in 3/9 (33%) and drug-responsive in 5/9 (56%). One patient with seizures was not treated with anti-epileptic drugs (AEDs). Two of the subjects with drug-resistant epilepsy experienced multiple seizures daily. AEDs that were trialed included valproate (5/9), levetiracetam (3/9), clobazam (2/9), gabapentin (1/9), carbamazepine (1/9), phenobarbital (1/9), topiramate (1/9), and vigabatrin (1/9). The combination of valproate and levetiracetam or clobazam was reported to be effective in three patients (8-1, 8-2, 11-1).

Cataracts were present in 66.7% (10/15) of patients. Four reported mature cataracts and two had posterior cataracts. Feeding difficulties were present in seven patients, with one individual (2) requiring G-tube placement. Dysmorphic features were reported in some patients (Supplementary Table 1 and 3), though no recognizable facial gestalt or pattern was appreciated.

DISCUSSION

We report sixteen patients with a novel autosomal recessive disease due to pathogenic variants in *MED27* consisting of global developmental delay, axial hypotonia, spastic tetraplegia, dystonia, cerebellar hypoplasia, seizures, and cataracts (Table 1). Missense variants were more commonly identified than frameshift and splicing variants. Three missense variants associated

with milder phenotypes [c.188T>G (p.Val63Gly), c.776C>T (p.Pro259Leu), and c.878C>T (p.Pro293Leu)]. The c.188T>G variant is distinct as it is the only missense variant located outside the C-terminal region where all other missense variants clustered (Fig 1).

The Mediator complex is composed of 25 (yeast) or 30 (human) subunits that form four modules: head, middle, tail, and CDK8 kinase ^{2, 15}. MED27 is a metazoan-specific Mediator subunit sitting at the junction of the head and tail modules of the complex¹⁵. *Med27/Crsp34* loss-of-function in zebrafish disrupts dopaminergic amacrine cells and serotonergic neurons at 2.5 dpf (days post-fertilization) ¹⁶. Mutant embryos had a reduction of head, eye, and jaw size, and died around 6 dpf. In flies, *Med27* knockout caused lethality at the pupal stage ^{17, 18}. Similarly, chickens carrying homozygous *MED27* insertional truncating mutations were born at less than expected Mendelian ratios, suggesting partial embryonic lethality in homozygotes¹⁹. Although the specific biological function of MED27 remains unknown, it clearly plays an essential role in early embryonic and neuronal development.

Similar to the effect of knockout mutations, homozygous C-terminal Flag-tagged *Med27* mutations were also lethal in fruit flies, suggesting a critical role of the MED27 C-terminal domain.¹⁸ In the patients reported here, 6/7 missense variants clustered near the C-terminus of the MED27 protein. One study indicated MED27 C-terminal domain has a C2-H2 zinc finger motif ⁶. MED27 interacts extensively with multiple subunits in the head module, including MED17 ²⁰. Cryo-electron microscopy of the *S. pombe* head module reveals that Med27 connects Med18/20 with Med17 ¹⁵. Interestingly, compared with other Mediator complex-associated diseases, *MED27* and *MED17*-related diseases are most similar. Both are autosomal recessive and characterized by psychomotor developmental delay, spasticity, seizures, progressive microcephaly and cerebellar atrophy (Supplementary Table 4). One intriguing hypothesis would be that variants in MED27 disrupt its interaction with other Mediator

complex subunits, such as MED17, leading to similar disease phenotypes. Additional studies on the functional consequence of *MED27* variants are needed to further address the molecular mechanisms underlying the disease.

Acknowledgements: N.E.M. is supported by a Parkinson's foundation grant. P.I. is supported by Foundation for Pediatric Research. D.K. is supported by the Simpson Querrey Center for Neurogenetics. Biospecimens used in the analyses presented in this article were obtained from the Northwestern University Movement Disorders Center (MDC) Biorepository. As such, the investigators within MDC Biorepository contributed to the design and implementation of the MDC Biorepository and/or provided data and collected biospecimens but did not participate in the analysis or writing of this report. MDC Biorepository investigators include (Tanya Simuni, MD; Dimitri Krainc, MD PhD; Opal Puneet MD, PhD; Cindy Zadikoff, MD; Onur Melen, MD; Danny Bega, MD; Roneil G. Malkani, MD; Steven Lubbe, PhD; Niccolo E. Mencacci, MD, PhD; Christina Zelano, PhD; Joanna Blackburn, MD; Firas Wehbe, MD, PhD; Lisa Kinsley, MS, CGC; Tina Ward, MS). A gift from the Malkin family generously supported the work of the MDC biorepository. We thank Finnish Institute for Molecular Medicine for whole-exome sequencing and Anu Harju and Max Pohjanpelto (University of Helsinki) for technical assistance. J.R.L. is supported by a grant from the National Human Genome Research Institute (NHGRI) and National Heart, Lung, and Blood Institute (NHBLI) to the Baylor-Hopkins Center for Mendelian Genomics (BHCMG, UM1 HG006542); a National Institute of Neurological Disorders and Stroke (NINDS) grant (R35NS105078); and an Muscular Dystrophy Association (MDA) grant (512848). T.M. is supported by the Uehara Memorial Foundation. D.M. is supported by a Medical Genetics Research Fellowship Program through the United States National Institute of Health (T32 GM007526-42). D.P. is supported by a Clinical Research Training Scholarship in Neuromuscular Disease partnered by the American Brain Foundation (ABF) and Muscle Study

Group (MSG), and International Rett Syndrome Foundation (IRSF grant #3701-1). J.E.P. was supported by NHGRI K08 HG008986.

Author contributions

L.M. and Y.Y. contributed to study conception and design. L.M. drafted the manuscript text and prepared the figures. All authors contributed to patient clinical data and exome sequencing data acquisition and analysis, and manuscript review and revision.

Potential Conflicts of Interest:

L.M. is employee of Baylor Genetics, in which exome sequencing of patient 1 was performed. The Department of Molecular and Human Genetics at Baylor College of Medicine receives revenue from clinical genetic testing conducted at Baylor Genetics (BG) Laboratories. Other authors have no potential conflicts to report.

References

1. Jeronimo C, Robert F. The Mediator Complex: At the Nexus of RNA Polymerase II Transcription. *Trends Cell Biol.* 2017 Oct;27(10):765-83.
2. Soutourina J. Transcription regulation by the Mediator complex. *Nat Rev Mol Cell Biol.* 2017 Dec 6.
3. Yin JW, Wang G. The Mediator complex: a master coordinator of transcription and cell lineage development. *Development.* 2014 Mar;141(5):977-87.
4. Risheg H, Graham JM, Jr., Clark RD, et al. A recurrent mutation in MED12 leading to R961W causes Opitz-Kaveggia syndrome. *Nat Genet.* 2007 Apr;39(4):451-3.
5. Schwartz CE, Tarpey PS, Lubs HA, et al. The original Lujan syndrome family has a novel missense mutation (p.N1007S) in the MED12 gene. *J Med Genet.* 2007 Jul;44(7):472-7.
6. Bourbon HM. Comparative genomics supports a deep evolutionary origin for the large, four-module transcriptional mediator complex. *Nucleic Acids Res.* 2008 Jul;36(12):3993-4008.
7. Boubé M, Joulia L, Cribbs DL, Bourbon HM. Evidence for a mediator of RNA polymerase II transcriptional regulation conserved from yeast to man. *Cell.* 2002 Jul 26;110(2):143-51.
8. Sobreira N, Schiettecatte F, Valle D, Hamosh A. GeneMatcher: a matching tool for connecting investigators with an interest in the same gene. *Hum Mutat.* 2015 Oct;36(10):928-30.
9. Davydov EV, Goode DL, Sirota M, Cooper GM, Sidow A, Batzoglou S. Identifying a high fraction of the human genome to be under selective constraint using GERP++. *PLoS Comput Biol.* 2010 Dec 2;6(12):e1001025.
10. Adzhubei IA, Schmidt S, Peshkin L, et al. A method and server for predicting damaging missense mutations. *Nat Methods.* 2010 Apr;7(4):248-9.
11. Vaser R, Adusumalli S, Leng SN, Sikic M, Ng PC. SIFT missense predictions for genomes. *Nat Protoc.* 2016 Jan;11(1):1-9.
12. Schwarz JM, Rodelsperger C, Schuelke M, Seelow D. MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods.* 2010 Aug;7(8):575-6.
13. Rentzsch P, Witten D, Cooper GM, Shendure J, Kircher M. CADD: predicting the deleteriousness of variants throughout the human genome. *Nucleic Acids Res.* 2019 Jan 8;47(D1):D886-D94.
14. Pehlivan D, Bayram Y, Gunes N, et al. The Genomics of Arthrogryposis, a Complex Trait: Candidate Genes and Further Evidence for Oligogenic Inheritance. *Am J Hum Genet.* 2019 Jul 3;105(1):132-50.
15. Tsai KL, Yu X, Gopalan S, et al. Mediator structure and rearrangements required for holoenzyme formation. *Nature.* 2017 Apr 13;544(7649):196-201.
16. Durr K, Holzschuh J, Filippi A, et al. Differential roles of transcriptional mediator complex subunits Crsp34/Med27, Crsp150/Med14 and Trap100/Med24 during zebrafish retinal development. *Genetics.* 2006 Oct;174(2):693-705.
17. Gokcezade J, Sienski G, Duchek P. Efficient CRISPR/Cas9 plasmids for rapid and versatile genome editing in *Drosophila*. *G3 (Bethesda).* 2014 Sep 17;4(11):2279-82.
18. Li-Kroeger D, Kanca O, Lee PT, et al. An expanded toolkit for gene tagging based on MiMIC and scarless CRISPR tagging in *Drosophila*. *Elife.* 2018 Aug 9;7.
19. Tsujino K, Okuzaki Y, Hibino N, et al. Identification of transgene integration site and anatomical properties of fluorescence intensity in a EGFP transgenic chicken line. *Dev Growth Differ.* 2019 Sep;61(7-8):393-401.
20. Tsai KL, Tomomori-Sato C, Sato S, Conaway RC, Conaway JW, Asturias FJ. Subunit Architecture and Functional Modular Rearrangements of the Transcriptional Mediator Complex. *Cell.* 2014 Jul 17;158(2):463.

FIGURES

Figure 1 Nucleotide and amino acid changes in patients with *MED27* biallelic variants.

(A) Schematic representation of *MED27* gene structure (not drawn to scale) and nucleotide position of eleven variants identified in eleven families. (B) Alignment of human *MED27* protein sequence with other model organisms including mouse, rat, pig, bovine, *Xenopus*, zebrafish and *Drosophila*.

Figure 2 Brain imaging of patients with *MED27* biallelic variants.

Brain imaging from ten patients. Sagittal T1 +/- coronal T2 brain MRI imaging of nine patients (A-K, M). Sagittal brain computed tomography (CT) imaging is provided for patient 9 (L). (A) Patient 1 (2 years) showing mild cerebellar vermician hypoplasia (ARROW). (B) Patient 2 (2 years 11 months) showing normal-appearing corpus callosum and cerebellar vermician hypoplasia (ARROW). (C) Patient 3-1 (1 year) showing thin corpus callosum (ARROWHEAD) and severe cerebellar hypoplasia (ARROW). (D) Patient 3-1 (2 years 5 months), showing thin corpus callosum (ARROWHEAD) and progressive cerebellar atrophy (ARROW). (E) Patient 3-2 (11 years 9 months) showing thin corpus callosum (ARROWHEAD) and cerebellar vermician hypoplasia (ARROW) (F) Patient 4 (1 year), showing cerebellar vermician hypoplasia (ARROW).

(G) Patient 4 (2 years 5 months) showing progressive cerebellar atrophy (ARROW) and cortical gyral simplification. (H) Patient 5 (1 year 2 months), showing thin corpus callosum (ARROWHEAD), severe cerebellar hypoplasia (ARROW) with flattening of the pons and hypomyelination. (I) Patient 6-1 (34 years) showing mild cerebellar vermian hypoplasia (ARROW). (J) Patient 7 (1 year) showing cerebellar vermis hypoplasia (ARROW). (K) Patient 8-1 (6 years 3 months) showing cerebellar vermian hypoplasia (ARROW). (L) Patient 9 showing cerebellar hypoplasia (ARROW) (M) Patient 11-1 showing cerebral atrophy and normal cerebellum.

Table 1 Clinical features of patients with biallelic *MED27* variants

Clinical findings	Severe (n=8)	Moderate (n=5)	Family 6 (n=2)	Overall*
Gender				5M/10F
Intellectual disability	100% (7/7)	100% (5/5)	100% (2/2)	100% (14/14)
Central hypotonia	100% (8/8)	80.0% (4/5)	100% (2/2)	93.3% (14/15)
Distal limb spasticity/dystonic movement	100% (8/8)	60.0% (3/5)	100% (2/2)	86.7% (13/15)
Delayed motor/speech development	100% (8/8)	100% (5/5)	0% (0/2)	86.7% (13/15)
Cerebellar hypoplasia	100% (8/8)	75.0% (3/4)	50.0% (1/2)	85.7% (12/14)
Cataracts	87.5% (7/8)	20.0% (1/5)	100% (2/2)	66.7% (10/15)
Epilepsy	87.5% (7/8)	40.0% (2/5)	0% (0/2)	60.0% (9/15)
Feeding issues	42.8% (3/7)	60.0% (3/5)	100% (2/2)	57.1% (8/14)
Microcephaly	50.0% (4/8)	0.0% (0/4)	0% (0/2)	28.6% (4/14)
Developmental regression	25.0% (2/8)	20.0% (1/5)	100% (2/2)	33.3% (5/15)

*Patient 10-2 had dual diagnosis with variants in both *MED27* and *COG6*. Therefore the case is excluded in this summary of *MED27*-

associated phenotypes.

Key: M, male; F, female

Supplementary Table 1 Family pedigrees, facial images of individuals with biallelic *MED27* variants and Sanger sequencing data

Family	Pedigree	Facial photos	<i>MED27</i> variants (Sanger sequencing)
1		<p>Individual 1 at 7y6m</p>	<p>Individual 1</p> <p>Mother</p> <p> c.776C>T c.188T>G </p>
2		<p>Individual 2 at 3y10m</p>	<p>Individual 2</p> <p>Mother</p> <p>Father</p> <p> c.682-2A>G c.871G>A </p>

3	<p> c.839C>T/+ c.839C>T/+ 3-1 +/+ 3-2 c.839C>T/ c.839C>T c.839C>T/ c.839C>T </p>		<p> Individual 3-1 Individual 3-2 Mother Father Unaffected sibling c.839C>T </p>
4	<p> c.725T>C/+ c.725T>C/+ nd 4 c.725T>C/ c.725T>C </p>		<p> Individual 4 Mother Father c.725T>C </p>
5	<p> c.565_566del/+ c.839C>T/+ 5 c.565_566del/ c.839C>T </p>	<p>Individual 5 at 1y2m</p>

6	<p>nd nd</p> <p>6-1 6-2</p> <p>c.871G>A/ c.871G>A c.871G>A/ c.871G>A</p>		<p>Individual 6-1</p> <p>c.871G>A</p>
7	<p>c.695C>T/+ c.298_302del/+</p> <p>7</p> <p>c.298_302del/ c.695C>T</p>	<p>Individual 7 at 2y3m</p>	<p>Individual 7</p> <p>Mother</p> <p>Father</p> <p>c. 298_302delIAAAC c. 695C>T</p>
8	<p>nd c.871G>A/+</p> <p>8-1 8-2</p> <p>c.392_393del/ c.871G>A c.392_393del/ c.871G>A</p>	<p>Individual 8-1 at 8y</p> <p>Individual 8-2 at 6y</p>

9	<p>c.839C>T/+ c.839C>T/+</p> <p>c.839C>T/ c.839C>T</p>	<p>Individual 9</p> <p>Mother</p> <p>Father</p> <p>c.839C>T</p>	

10	<p> $c.878C>T/+$ $c.878C>T/+$ 10-1 10-2 $c.878C>T/$ $c.878C>T/$ $c.878C>T$ $c.878C>T^*$ </p>	<p>Individual 10-1</p>	<p> Individual 10-1 Individual 10-2 Mother Father $c.878C>T$ </p>
11	<p> $c.776C>T/+$ $c.776C>T/+$ 11-1 11-2 $c.776C>T/$ $c.776C>T/$ $c.776C>T$ $c.776C>T$ +/+ </p>	<p>Individual 11-1 at 9y5m</p> <p>Individual 11-2 at 6y9m</p>	<p> Individual 11-1 Mother Father Individual 11-2 Unaffected Sibling $c.776C>T$ </p>

Patients and/or legal guardians have consented for publication of facial images.

Supplementary Table 2 *MED27* Variant characteristics identified in this study

Family	Exon/I ntron	cDNA	Protein	Variant type	GERP+ + RS	PolyP hen2	SIFT	Mutation Taster	CADD Phred score	gnomAD database
1	Exon 1	c.188T>G	p.Val63Gly	Missense	5.07	D	D	D	28.2	2/245554
7	Exon 2	c.298_302del	p.Lys100Serfs*4	Frameshift	NA	NA	NA	NA	NA	0
8	Exon 3	c.392_393del	p.Gln131Leufs*7	Frameshift	NA	NA	NA	NA	NA	0
5	Exon 4	c.565_566del	p.M189Afs*21	Frameshift	NA	NA	NA	NA	NA	0
2	Intron 5	c.682-2A>G	N/A	Splice site	5.62	NA	NA	NA	34	2/239080
7	Exon 6	c.695C>T	p.Ser232Phe	Missense	5.62	P	D	D	31	0
4	Exon 7	c.725T>C	p.Val242Ala	Missense	5.52	B	D	D	26.3	0
1,11	Exon 7	c.776C>T	p.Pro259Leu	Missense	5.52	D	D	D	34	0
3,5,9	Exon 8	c.839C>T	p.Pro280Leu	Missense	5.3	P	D	D	34	14/241660
2,6,8	Exon 8	c.871G>A	p.Gly291Ser	Missense	5.3	D	D	D	27.5	3/235830
10	Exon 8	c.878C>T	p.Pro293Leu	Missense	5.46	D	D	D	29.9	0

Reference sequence: NM_004269.3; D: probably damaging; P: possibly damaging; B: benign

Supplementary Table 3 Summary of clinical features in patients with biallelic *MED27* variants.

Individual	1	2	3-1	3-2	4	5	6-1	6-2
Disease severity	Moderate	Severe	Severe	Severe	Severe	Severe	Family 6	Family 6
Gender	Male	Male	Female	Female	Female	Male	Female	Female
Ethnicity	Hispanic	Western European	Finnish	Finnish	Turkish/Syrian	European, French Canadian	Kuwaiti	Kuwaiti
Consanguinity	No	No	No	No	No	No	Yes	Yes
Birth history	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal
Age at measurement	10y 7m	5y 10m	11y, (passed at 13y)	8m (passed at 5y)	2y 11m	20m	36y	42y
Height	23%ile (-0.8SD)	52%ile (-0.1SD)	<1%ile (-3SD)	20%ile (-0.84SD)	49%ile (-0.03SD)	35%ile (-0.39SD)	NA	<1%ile (-2.8 SD)
Weight	14%ile (-1.1SD)	12%ile (-1.3SD)	43%ile (-0.18SD)	18%ile (-0.92SD)	27%ile (-0.61SD)	33%ile (-0.44SD)	NA	23%ile (-0.74SD)
FOC	2%ile (-1.99SD)	<1%ile (-3.3SD) (at 33m)	<1%ile (-5SD)	<1%ile (-3.5SD)	5%ile (-1.61SD)	<1%ile (-4.6SD)	Normocephalic	Normocephalic
Microcephaly	-	+	+	+	-	+	-	-
Feeding issues	Difficult swallowing	Swallowing issue, G-tube placement	-	-	severe swallowing issue	Dysphagia	Occasional choking	Occasional choking
Development assessment	Chronological age 10.8y, intelligence <4y, expressive language 23m, receptive language 3y, fine motor 3.5y, visual perceptual ability 4y	NA	NA	NA	NA, felt to be 6m at chronological age of 6y	NA	NA	NA
Motor delay	+, delayed, able to walk	+, cannot sit/walk	+, cannot sit/walk	+, achieved walking but regressed and lost ability to move	+, sit with support, cannot walk	+, poor head control, cannot sit/walk	-	-
Language delay	+, delayed, 10 words at 40 m	+, non-verbal	+, non-verbal	+, non-verbal	+, non-verbal	+, non-verbal (early)	-, progressive dysarthria since age 8	-, progressive dysarthria since age 5
Intellectual disability	Moderate	Profound	Severe	Severe	Profound	NA (early)	Moderate	Moderate

Developmental regression	-	+, at 7y, motor function is worsening	+, onset around 6y	-	-	-	+	+
Central hypotonia	-	+	+	+	+	+	Mild	Mild
Distal spasticity	-	+	+	+	+	+	+	+
Dystonic movement	-, Unsteady gait	+, tongue protrusion	+	+	+, hands	+, tongue protrusion	+, tongue, mandibular, dysarthria, posture	+, tongue, mandibular, dysarthria, posture
Epilepsy	-, seizure like activity in infancy, normal EEG	+, onset 2y	+, onset 3y	+, onset 2y	+, onset 2-3y,	+, onset 8m, abnormal EEG, consistent with modified hypsarrhythmia	-	-
History of Drug-resistant epilepsy	NA	-	+, (at 6 y then became seizure free on AEDs at 11y)	-	+	+	NA	NA
Seizure type(s) and frequency	NA	Generalized tonic-clonic seizure	N/A	N/A	Myoclonic during sleep, atypical absences during daytime; multiple daily.	Focal motor (tonic), absence, epileptic (infantile) spasms; multiple daily.	NA	NA
AEDs	NA	GBP and low dose LEV	VPA, TPM and PB	VPA	N/A	Steroids, ACTH, Vigabatrin, LEV and CBZ	NA	NA
Brain MRI cerebellar atrophy	mild	+	+	+	+, progressive	+	mild	-
Brain MRI others	-	progression of diffuse cerebral atrophy and progression of abnormal signal and atrophy in basal ganglia	cerebral atrophy, thin corpus callosum	cerebral atrophy, thin corpus callosum	cortical gyral simplification	hypomyelination with a thin corpus callosum, progressive <i>ex vacuo</i> ventriculomegaly	-	-
Cataracts	-	+, left mature cataract, right incomplete cataract	+, congenital, operated at 7y, type NA	+, congenital, operated at 5m, type NA	+, mature	+, developed at 23m, type not specified	+, mature	+, mature
Dysmorphic features	Large cupped ears with	Prominent eyebrows	-	-	narrow forehead, frontal upsweep	Large ears, slightly upturned	-	-

	thickened helices, long eyelashes, frontal bossing, upswept frontal hair				of hair, synophrys, highly arched broad eyebrows, large eyes, blue sclerae, prominent nasal bridge, anteverted nares, exaggerated cupid bow, widely spaced teeth, long eyelashes, long hyperextensible and broad fingers, hypertrichosis	nose, long eyelashes, full lips		
Other issues	Mild obstructive sleep apnea; Significant drooling, behavioral concerns; Dental concern	Born with loud breathing and diagnosed with laryngomalacia as infant	Peripheral neuropathy both in lower and upper limbs	N/A	Drooling	Severe central sleep apnea, moderate obstructive sleep apnea, laryngomalacia, milk protein allergy, constipation, bilateral hearing loss	Urinary urgency with frequent episodes of incontinence	Urinary urgency with frequent episodes of incontinence
MED27 (NM_004269.3)	c.776C>T (p.Pro259Leu) presumed paternal; c.188T>G (p.Val63Gly) maternal	c.682-2A>G paternal; c.871G>A (p.Gly291Ser) maternal	Homozygous c.839C>T (p.Pro280Leu)	Homozygous c.839C>T (p.Pro280Leu)	Homozygous c.725T>C (p.Val242Ala)	c.565_566del(p. Met189Alafs*21) paternal; c.839C>T (p.Pro280Leu) maternal	Homozygous c.871G>A (p.Gly291Ser)	Homozygous c.871G>A (p.Gly291Ser)
Additional diagnosis								

Individual	7	8-1	8-2	9	10-1	10-2*	11-1	11-2	Total
Disease severity	Severe	Severe	Severe	Moderate	Moderate	NA	Moderate	Moderate	8 severe; 5 moderate
Gender	Female	Female	Female	Female	Male	Female	Male	Female	5M/11F
Ethnicity	North-western European	Algerian	Algerian	Iraqi	Turkish	Turkish	Pakistani	Pakistani	
Consanguinity	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Birth history	Normal	Normal	Normal	Normal	Normal	Polyhydramnios, SGA, respiratory distress	Normal	Preterm	
Age at measurement	2y 3m	7y 4m	5y 11m	10m	20 y	16 d, (passed at 3m)	9y 5m	6yr 9mo	
Height	2%ile (-2.04SD)	18%ile (-0.3SD)	20%ile (-0.82SD)	NA	25%ile (-0.67SD)	50%ile (0SD)	44%ile (-0.16SD)	21%ile (-0.82SD)	
Weight	41%ile (-0.24SD)	14%ile (-0.65SD)	29%ile (-0.57SD)	>99%ile ($\geq +3.0SD$)	3%ile (-1.88SD)	25%ile (-0.67SD)	75%ile (+0.67SD)	25%ile (-0.68SD)	
FOC	18%ile (-0.92SD)	12%ile (-1.16SD)	13%ile (-1.12SD)	NA	<3%ile, <-1.88SD	<3%ile, <-1.88SD	53%ile (+0.08SD)	27%ile (-0.61SD)	4/14
Microcephaly	-	-	-	-	-	-	-	-	4/14
Feeding issues	NA	-	-	+	-	+	-	Mild	8/14
Development assessment	Bailey scale: chronological age 16m, cognition 7m	NA	NA	NA	NA, felt to be 5-6y level at chronological age of 20y	NA	Protage guide: Chronological age 9.5y, Cognition 66.4m, Socialization 57.8m, Self-help 61.5m, Motor 61.6m, Language 48m	Protage guide: Chronological age 7.1y, Cognition 67m, Socialization 59m, Self-help 68m, Motor 66m, Language 59m	
Motor delay	+, cannot sit/walk	+	+, cruising	+, walk with support	+, able to walk	NA	+, delayed, able to walk	+, delayed, able to walk	13/15
Language delay	+, non-verbal	+, delayed	+, non-verbal	+, delayed, one word at 22m	+, delayed, able to read	NA	+, delayed	+, delayed	13/15
Intellectual disability	Severe	Profound	Profound	Moderate	Moderate	N/A	Moderate	Mild	14/14

Developmental regression	-	-	-	-	+	NA	-	-	5/15
Central hypotonia	+	+	+	+	+	+	Mild	Mild	14/15
Distal spasticity	+	+	+	+	-	-	+	mild	13/15
Dystonic movement	+, tongue protrusion, stereotypical movements	+	+	+	-	NA	+, tongue fasciculations	+, tongue fasciculations	13/15
Epilepsy	-, disturbed EEG	+, onset 5y, abnormal EEG	+, onset 5y, abnormal EEG	+, onset at 18m; normal EEG	-	-	+, onset at 20d	-	9/15
History of Drug-resistant epilepsy	NA	-	-	Unable to assess	NA	NA	-	NA	3/15
Seizure type(s) and frequency	NA	Hemifacial clonic seizures; none while on AEDs	Hemifacial clonic; none while on AEDs	Focal motor with impaired awareness	NA	NA	Atonic and right- sided focal; once every week till 6Y when he experienced a prolonged GTC; Last seizure was 7Y on AEDs	NA	
Current AEDs	NA	VPA and CLB effective; LEV previously used and only partially effective	VPA and CLB effective; LEV previously used and only partially effective	none	NA	NA	VPA and LEV for last 3yrs (effective)	NA	
Brain MRI cerebellar atrophy	mild	+	+	+	mild	mild	-	NA	12/14
Brain MRI others	-	-	-	-	-	-	cerebral atrophy with ischemic demyelination in parietal and occipital white matter	-	
Cataracts	-	+, congenital	+, congenital	-	+, posterior cataracts	+, posterior cataracts	-	-	10/15

Dysmorphic features	frontal bossing, sparse medial eyebrow, wide nasal bridge, hypoplasia of the primary teeth, Deep palmar creases	no (long narrow face)	-	-	micrognathia	-	-	Hypertelorism, flat nasal bridge, low set ears, broad nose, convergent squint	
Other issues	Drooling	Drooling	Drooling	NA	Knee and wrist contractures, dysmetria	Hand and feet contractures	Micropenis with normal testosterone levels, mild drooling	Drooling	
MED27 (NM_004269.3)	c.695C>T (p.Ser232Phe) paternal; c.298_302del (p.Lys100Serfs*4) maternal	c.871G>A (p.Gly291Ser) maternal; c.392_393del (p.Gln131Leufs Ter7) presumed paternal	c.871G>A (p.Gly291Ser) maternal; c.392_393del (p.Gln131Leufs Ter7) presumed paternal	Homozygous c.839C>T (p.Pro280Leu)	Homozygous c.878C>T: p.Pro293Leu; both parents are heterozygous	Homozygous c.878C>T: p.Pro293Leu; both parents are heterozygous	Homozygous c.776C>T, p.Pro259Leu	Homozygous c.776C>T, p.Pro259Leu	
Additional diagnosis						Homozygous COG6 c.726del (p.Cys242Trpfs*7)			

*Patient has dual diagnosis with variants in both *MED27* and *COG6*, not counted towards phenotype summary in Table 1

AED: antiepileptic drug; CBZ: carbamazepine; CLB: clobazam; EMG: electroneuromyography; GBP: gabapentin; LEV: levetiracetam; NA: not available; NCS: nerve conduction studies; PB: phenobarbital; SSEP: somato-sensory evoked potentials; TPM: topiramate; VPA: valproate

Supplementary Table 4 Summary of Mediator subunits known to be associated with human diseases

Gene	Module	Inheritance	Disease	Phenotype description
<i>MED12</i>	Kinase	XL	Lujan-Fryns syndrome [MIM: 309520]; Ohdo syndrome, X-linked [MIM: 300895]; Opitz-Kaveggia syndrome [MIM: 305450]	dysmorphic features, intellectual disability, relative macrocephaly, hypotonia, and constipation, partial agenesis of the corpus callosum
<i>MED13</i>	Kinase	AD	Intellectual developmental disorder 61 [MIM: 618009]	intellectual disability and/or developmental delays, including speech delays or disorders, autism
<i>MED13L</i>	Kinase	AD	Mental retardation and distinctive facial features with or without cardiac defects (MIM: 616789); Transposition of the great arteries, dextro-looped 1 (MIM: 608808)	delayed psychomotor development, poor speech acquisition, distinctive dysmorphic facial features, variable penetrance of cardiac malformations
<i>MED17</i>	Head	AR	Microcephaly, postnatal progressive, with seizures and brain atrophy (MIM: 613668)	microcephaly, postnatal progressive, with seizures and brain atrophy
<i>MED23</i>	Tail	AR	Mental retardation, autosomal recessive 18 (MIM: 614249)	mild to moderate ID, no dysmorphic
<i>MED25</i>	Unassigned	AR	Basel-Vanagait-Smirin-Yosef syndrome (MIM616449)	severely delayed psychomotor development resulting in intellectual disability, as well as variable eye, brain, cardiac, and palatal abnormalities.
<i>MED27</i>	Head/ Tail	AR	Current report	global developmental delay, central hypotonia, spasticity, dystonic movements, cerebellar hypoplasia, epilepsy, and cataract

AD, autosomal dominant; AR, autosomal recessive; XL, X-linked