

Non-Abelian gauge fields as dark matter

Christian Gross,^a Oleg Lebedev^a and Yann Mambrini^b

^a*Department of Physics and Helsinki Institute of Physics,
Gustaf Hällströmin katu 2, FI-00014 Helsinki, Finland*

^b*Laboratoire de Physique Théorique, CNRS – UMR 8627,
Université de Paris-Sud 11, F-91405 Orsay Cedex, France*

E-mail: christian.gross@helsinki.fi, oleg.lebedev@helsinki.fi,
yann.mambrini@th.u-psud.fr

ABSTRACT: $SU(N)$ Lie algebras possess discrete symmetries which can lead naturally to stable vector dark matter (DM). In this work, we consider the possibility that the dark $SU(N)$ sector couples to the visible sector through the Higgs portal. We find that minimal CP -conserving hidden “Higgs sectors” entail stable massive gauge fields which fall into the WIMP category of dark matter candidates. For $SU(N)$, $N > 2$, DM consists of three components, two of which are degenerate in mass. In all of the cases, there are substantial regions of parameter space where the direct and indirect detection as well as relic abundance constraints are satisfied.

KEYWORDS: Cosmology of Theories beyond the SM, Higgs Physics, Beyond Standard Model

ARXIV EPRINT: [1505.07480](https://arxiv.org/abs/1505.07480)

Contents

1	Introduction	1
2	Massive U(1) and SU(2) gauge fields as vector DM	2
2.1	Hidden U(1)	2
2.2	Hidden SU(2)	3
3	Extension to SU(3)	4
3.1	Breaking SU(3) by tensor fields	4
3.2	Breaking SU(3) by triplets and $Z_2 \times Z'_2$	5
3.3	Explicit example	6
3.4	Detailed study for $v_{3,4} = 0$	8
3.4.1	Masses	9
3.4.2	Couplings	11
4	Generalisation to arbitrary SU(N)	12
5	Dark matter phenomenology	13
5.1	U(1) dark matter	14
5.2	SU(2) dark matter	17
5.3	SU(3) dark matter	17
6	Summary and conclusions	18
A	Hidden SU(3) vector-scalar couplings for $v_3 = 0$	20

1 Introduction

The puzzle of dark matter remains an outstanding problem of particle physics. One of the more attractive approaches to this problem exploits the fact that weakly interacting massive particles (WIMPs) in thermal equilibrium produce the relic dark matter abundance in the right ballpark. Particles of this type appear in many extensions of the Standard Model. In this work, we explore the possibility that the Standard Model is connected to the dark sector through the Higgs portal [1–3],

$$V_{\text{portal}} = \lambda_{h\phi} H^\dagger H \phi^\dagger \phi, \tag{1.1}$$

where H is the Higgs field and ϕ is the “hidden Higgs”, that is, the field responsible for breaking the gauge symmetry of the hidden sector. We assume the dark sector to be endowed with U(1) or SU(N) gauge symmetry. In that case, the massive gauge fields can

play the role of WIMP-type dark matter quite naturally. Indeed, they are weakly coupled to the Standard Model and, owing to inherent discrete symmetries, can be stable.

The U(1) case was considered in ref. [4] where the stabilising Z_2 symmetry was found to be related to charge conjugation. The SU(2) example was worked out in ref. [5] and the DM stability was attributed to a custodial SO(3). In this work, we extend these ideas to larger SU(N) Lie groups and uncover the nature of the underlying discrete symmetries. We focus on weakly coupled theories, although confining hidden sectors represent a viable alternative [6, 7]. Finally, we perform a comprehensive study of DM phenomenology in all of these cases, which includes direct and indirect DM detection as well as an analysis of the DM relic abundance. Our main conclusion is that massive hidden gauge fields serve as viable and attractive DM candidates.

2 Massive U(1) and SU(2) gauge fields as vector DM

In this section, we review the cases of massive U(1) and SU(2) gauge fields as vector DM and identify the underlying symmetries leading to stability of vector dark matter. In what follows we assume that the hidden sector consists only of gauge fields and of scalar multiplets which are necessary to make these gauge fields massive.

2.1 Hidden U(1)

An Abelian gauge sector provides the simplest example of the vector DM model endowed with a natural Z_2 symmetry [4]. In this case, the Z_2 corresponds to the charge conjugation symmetry.

Consider a U(1) gauge theory with a single charged scalar ϕ ,

$$\mathcal{L}_{\text{hidden}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu\phi)^\dagger D^\mu\phi - V(\phi), \tag{2.1}$$

where $F_{\mu\nu}$ is the field strength tensor of the gauge field A_μ and $V(\phi)$ is the scalar potential. For easier comparison to the non-Abelian case, we take the charge of ϕ to be $+1/2$. Suppose at the minimum of the scalar potential ϕ develops a VEV, $\langle\phi\rangle = 1/\sqrt{2}\tilde{v}$. The imaginary part of ϕ gets eaten by the gauge field which now acquires the mass $m_A = \tilde{g}\tilde{v}/2$, where \tilde{g} is the gauge coupling. The real part of ϕ remains as a degree of freedom. Denoting it by ρ and normalising it canonically, $\phi = 1/\sqrt{2}(\rho + \tilde{v})$, we get the following gauge-scalar interactions:

$$\Delta\mathcal{L}_{s-g} = \frac{\tilde{g}^2}{4}\tilde{v}\rho A_\mu A^\mu + \frac{\tilde{g}^2}{8}\rho^2 A_\mu A^\mu. \tag{2.2}$$

The system possesses the Z_2 symmetry

$$A_\mu \rightarrow -A_\mu, \tag{2.3}$$

which is the usual charge conjugation symmetry. In terms of the original scalar field, this symmetry acts as $\phi \rightarrow \phi^*$ and $A_\mu \rightarrow -A_\mu$, which is preserved by both the Lagrangian and the vacuum. The Z_2 makes the massive gauge field stable. Note that this symmetry applies

to a sequestered U(1) which has no tree level mixing with the hypercharge, in which case no mixing is generated radiatively either.

Interactions with the visible sector proceed through the Higgs portal coupling

$$\mathcal{L}_{\text{portal}} = -\lambda_{h\phi} |H|^2 |\phi|^2, \tag{2.4}$$

which also leads to the Higgs mixing with ρ . In the unitary gauge, the Higgs field is given by $H^T = (0, v + h)/\sqrt{2}$. The fields ρ and h are then to be expressed in terms of the mass eigenstates $h_{1,2}$ as follows:

$$\begin{aligned} \rho &= -h_1 \sin \theta + h_2 \cos \theta, \\ h &= h_1 \cos \theta + h_2 \sin \theta, \end{aligned} \tag{2.5}$$

where the mixing angle θ is constrained by various experiments, most notably, by LEP and LHC. The upper bound on $\sin \theta$ depends on the mass of the heavier state h_2 and is around 0.3 for m_{h_2} of the order of a TeV, see e.g. figure 3 of ref. [8] for details. The lighter state h_1 is identified with the 125 GeV Higgs, while the mass of the second state can vary in a wide range.

As the Higgs portal necessarily preserves the Z_2 symmetry, A_μ is a viable DM candidate. All the relevant scattering processes proceed through an exchange of h_1 and h_2 , which include DM annihilation into the SM particles as well as DM scattering on nucleons.

2.2 Hidden SU(2)

The U(1) considerations can easily be extended to SU(2), albeit with a modification of the stabilising symmetry. Consider an SU(2) gauge theory with one doublet ϕ ,

$$\mathcal{L}_{\text{hidden}} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + (D_\mu \phi)^\dagger D^\mu \phi - V(\phi), \tag{2.6}$$

where $a = 1, 2, 3$. The potential is assumed to have a minimum at a nonzero VEV of ϕ . In the unitary gauge, ϕ takes the form

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \rho + \tilde{v} \end{pmatrix}, \tag{2.7}$$

with ρ being a real field and \tilde{v} being the VEV. Denoting the gauge coupling by \tilde{g} , this leads to the gauge boson mass $m_A = \tilde{g}\tilde{v}/2$. The scalar-gauge and gauge-gauge field interactions are given by

$$\begin{aligned} \Delta\mathcal{L}_{\text{s-g}} &= \frac{\tilde{g}^2}{4} \tilde{v} \rho A_\mu^a A^{a\mu} + \frac{\tilde{g}^2}{8} \rho^2 A_\mu^a A^{a\mu}, \\ \Delta\mathcal{L}_{\text{g-g}} &= -\tilde{g} \epsilon^{abc} (\partial_\mu A_\nu^a) A^{\mu b} A^{\nu c} - \frac{\tilde{g}^2}{4} \left((A_\mu^a A^{\mu a})^2 - A_\mu^a A_\nu^a A^{\mu b} A^{\nu b} \right). \end{aligned} \tag{2.8}$$

Although the triple gauge vertex breaks the parity of the previous section, it follows that the system possesses a $Z_2 \times Z'_2$ symmetry,

$$\begin{aligned} Z_2 : A_\mu^1 &\rightarrow -A_\mu^1, \quad A_\mu^2 \rightarrow -A_\mu^2, \\ Z'_2 : A_\mu^1 &\rightarrow -A_\mu^1, \quad A_\mu^3 \rightarrow -A_\mu^3. \end{aligned} \tag{2.9}$$

As a result, all three A_μ^a fields are stable and can play the role of dark matter. While the above symmetry is sufficient to ensure stability of DM, it actually generalises in this simple case to a custodial $SO(3)$ [5]. As we will see, for larger $SU(N)$ groups, it is the discrete symmetry that plays a crucial role. The first Z_2 is associated with a gauge transformation, while the Z'_2 generalises the charge conjugation symmetry, i.e. it corresponds to complex conjugation of the group elements.

As before, the dark sector couples to the visible one through the Higgs portal, $\lambda_{h\phi}|H|^2|\phi|^2$. The discussion of the Higgs mixing with ρ of the previous section applies here as well. Clearly, the $Z_2 \times Z'_2$ symmetry is preserved by the Higgs portal and the hidden gauge fields couple to the visible sector only in pairs.

3 Extension to $SU(3)$

While the $SU(2)$ case is straightforward, larger $SU(N)$ groups exhibit more complicated breaking patterns. In a phenomenologically viable set-up, the symmetry must be broken completely to avoid the existence of massless fields (barring confinement). One possibility would be to break $SU(3)$ by two scalar multiplets in the fundamental representation, i.e. triplets. One may also explore other options involving $SU(3)$ tensors. As we detail below, our conclusion is that a single irreducible representation with two indices cannot break $SU(3)$ completely, which leaves the two triplet option as the minimal one.

3.1 Breaking $SU(3)$ by tensor fields

The lowest order $SU(3)$ tensor whose generic VEV can break $SU(3)$ completely is the symmetric tensor ϕ_{ij} , that is **6** of $SU(3)$. Gauge transformations act on ϕ_{ij} as

$$\phi \rightarrow U\phi U^T. \tag{3.1}$$

By virtue of Takagi's matrix decomposition, this allows one to bring ϕ_{ij} to the diagonal form,

$$\phi = \begin{pmatrix} \phi_1 & 0 & 0 \\ 0 & \phi_2 & 0 \\ 0 & 0 & \phi_3 \end{pmatrix}, \tag{3.2}$$

where ϕ_1, ϕ_2, ϕ_3 are real up to an overall complex phase. If the VEVs of ϕ_1, ϕ_2 and ϕ_3 are all different, $SU(3)$ is broken completely. However, when some of them coincide, the residual gauge group is at least $SO(2)$.

In order to determine what VEVs are possible, let us write down the most general gauge invariant potential for ϕ_{ij} . It is easy to convince oneself that the potential has the form

$$V = m^2 \text{Tr}\phi^\dagger\phi + \lambda_1 \text{Tr}(\phi^\dagger\phi)^2 + \lambda_2 \left(\text{Tr}\phi^\dagger\phi\right)^2 + (\mu \text{Det}\phi + \text{h.c.}), \tag{3.3}$$

where m^2 can be negative. The minima of this potential determine the ϕ_i VEVs. Acting upon V with the operator $\phi_i\partial/\partial\phi_i$ (no summation over i), one finds that all nonzero $\langle\phi_i\rangle$ satisfy the same equation. This implies that $\langle\phi_i\rangle$ are either degenerate or zero. For the case

$\mu = 0$, such behaviour has been noticed in [9, 10]. We find (analytically and numerically) that it persists in the case of nonzero μ as well. As a result, the residual gauge symmetry is at least $SO(2)$ and the corresponding gauge bosons remain massless.¹ Thus, the model with a single symmetric tensor is unrealistic.

One may also consider the possibility of breaking $SU(3)$ by an antisymmetric tensor. By virtue of Youla’s decomposition, it can be gauge-transformed to the block-diagonal form

$$\phi = \begin{pmatrix} 0 & \phi_1 & 0 \\ -\phi_1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \tag{3.4}$$

Therefore, the residual gauge symmetry is at least $Sp(1)=SU(2)$, which can also be understood via the equivalence between the antisymmetric tensor and the (anti)fundamental representation. Again we find that the model is unrealistic.

Similarly, complete $SU(3)$ breaking cannot be achieved by a VEV of an adjoint scalar. In this case, the group rank is preserved and thus there are massless gauge fields.

The next simplest option is to combine the symmetric and antisymmetric tensors. This system would have 9 complex degrees of freedom, while two $SU(3)$ triplets only have 6. Thus $SU(3)$ breaking with two triplets is minimal and sufficient for our purposes.

3.2 Breaking $SU(3)$ by triplets and $Z_2 \times Z'_2$

Misaligned VEVs of two triplets break $SU(3)$ completely. This breaking pattern can be understood in stages: the first triplet VEV reduces the symmetry to $SU(2)$, while the second breaks the remaining $SU(2)$. VEVs misaligned in $SU(3)$ space represent a generic situation, that is, they result from the minimisation of a general scalar potential consistent with $SU(3)$ symmetry. Therefore, such a breaking pattern is phenomenologically viable.

Before going into details, let us identify the Lie group discrete symmetries which eventually lead to DM stability. One way to find them is to analyse the $SU(3)$ structure constants (using the usual Gell-Mann basis),

$$\begin{aligned} f^{123} &= 1, \\ f^{147} &= -f^{156} = f^{246} = f^{257} = f^{345} = -f^{367} = \frac{1}{2}, \\ f^{458} &= f^{678} = \frac{\sqrt{3}}{2}. \end{aligned} \tag{3.5}$$

Identifying the transformation properties of the generators with those of the gauge fields, we define the “parity” transformation as

$$A_\mu^a \rightarrow \eta(a)A_\mu^a. \tag{3.6}$$

It is easy to see that the structure constants are invariant if the parities are

$$\begin{aligned} Z_2 : \eta(a) &= -1 \text{ for } a = 1, 2, 4, 5, \\ \eta(a) &= +1 \text{ for } a = 3, 6, 7, 8, \end{aligned} \tag{3.7}$$

¹Ref. [11] has considered $\langle \phi \rangle$ proportional to the unit matrix, which entails unbroken $SO(3)$. The corresponding gauge bosons may be confined in glueballs.

and also

$$Z'_2 : \quad \begin{aligned} \eta(a) &= -1 \quad \text{for } a = 1, 3, 4, 6, 8, \\ \eta(a) &= +1 \quad \text{for } a = 2, 5, 7. \end{aligned} \quad (3.8)$$

One may notice that the first Z_2 acts on the off-diagonal generators with nonzero entries in the first row, while the second reflects the real $SU(3)$ generators. In section 4, we will show that these symmetries generalise to arbitrary $SU(N)$ and that the first Z_2 is a gauge transformation, whereas the second corresponds to an outer automorphism of the group, i.e. complex conjugation of the group elements.

These symmetries are inherited by the Yang-Mills Lagrangian. If CP is conserved, they are also preserved in the matter sector leading to stable dark matter. Below we study the relevant interactions in detail.

3.3 Explicit example

Consider an SM extension by two complex fields ϕ_1 and ϕ_2 transforming as triplets of hidden gauge $SU(3)$. The Lagrangian of the model is

$$\mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{portal}} + \mathcal{L}_{\text{hidden}}, \quad (3.9)$$

where

$$-\mathcal{L}_{\text{SM}} \supset V_{\text{SM}} = \frac{\lambda_H}{2} |H|^4 + m_H^2 |H|^2, \quad (3.10a)$$

$$-\mathcal{L}_{\text{portal}} = V_{\text{portal}} = \lambda_{H11} |H|^2 |\phi_1|^2 + \lambda_{H22} |H|^2 |\phi_2|^2 - (\lambda_{H12} |H|^2 \phi_1^\dagger \phi_2 + \text{h.c.}), \quad (3.10b)$$

$$\mathcal{L}_{\text{hidden}} = -\frac{1}{2} \text{tr}\{G_{\mu\nu} G^{\mu\nu}\} + |D_\mu \phi_1|^2 + |D_\mu \phi_2|^2 - V_{\text{hidden}}. \quad (3.10c)$$

Here, $G_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + i\tilde{g}[A_\mu, A_\nu]$ is the field strength tensor of the $SU(3)$ gauge fields A_μ^a , $D_\mu \phi_i = \partial_\mu \phi_i + i\tilde{g} A_\mu \phi_i$ is the covariant derivative of ϕ_i , H is the Higgs doublet, and the most general renormalisable hidden sector scalar potential is given by

$$\begin{aligned} V_{\text{hidden}}(\phi_1, \phi_2) &= m_{11}^2 |\phi_1|^2 + m_{22}^2 |\phi_2|^2 - (m_{12}^2 \phi_1^\dagger \phi_2 + \text{h.c.}) \\ &\quad + \frac{\lambda_1}{2} |\phi_1|^4 + \frac{\lambda_2}{2} |\phi_2|^4 + \lambda_3 |\phi_1|^2 |\phi_2|^2 + \lambda_4 |\phi_1^\dagger \phi_2|^2 \\ &\quad + \left[\frac{\lambda_5}{2} (\phi_1^\dagger \phi_2)^2 + \lambda_6 |\phi_1|^2 (\phi_1^\dagger \phi_2) + \lambda_7 |\phi_2|^2 (\phi_1^\dagger \phi_2) + \text{h.c.} \right]. \end{aligned} \quad (3.11)$$

Using $SU(3)$ gauge freedom, 5 real degrees of freedom of ϕ_1 and 3 real degrees of freedom of ϕ_2 can be removed. Therefore, in the unitary gauge ϕ_1, ϕ_2 read

$$\phi_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 0 \\ v_1 + \varphi_1 \end{pmatrix}, \quad \phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 + \varphi_2 \\ (v_3 + \varphi_3) + i(v_4 + \varphi_4) \end{pmatrix}, \quad (3.12)$$

where the v_i are real VEVs and φ_{1-4} are real scalar fields. Analogously, we express the Higgs field in the unitary gauge as $H^T = (0, v + h)/\sqrt{2}$.

fields	$Z_2 \times Z'_2$
$h, \varphi^1, \varphi^2, \varphi^3, A_\mu^7$	(+, +)
A_μ^2, A_μ^5	(-, +)
A_μ^1, A_μ^4	(-, -)
$\varphi^4, A_\mu^3, A_\mu^6, A_\mu^8$	(+, -)

Table 1. $Z_2 \times Z'_2$ parities of the scalars and dark gauge bosons.

In what follows, we make two assumptions which are crucial for stability of vector dark matter:

- the scalar potential is CP invariant
- the VEVs of ϕ_1, ϕ_2 are real.

The first condition implies that the scalar couplings are real, while the second assumes that no spontaneous CP violation occurs ($v_4 = 0$). As a result, CP -even and CP -odd fields do not mix.

In this case, the $Z_2 \times Z'_2$ symmetry extends to the Higgs sector as well. Under the first Z_2 all φ_i are even, while the second Z'_2 reflects φ_4 and leaves the other fields intact. As we detail in section 4, this assignment follows from the explicit form of the first Z_2 as a gauge transformation and the fact that the second Z'_2 acts as complex conjugation. In any case, these are explicit symmetries of the Lagrangian and the vacuum. The full list of the parities is presented in table 1. Clearly, the lightest states with non-trivial parities cannot decay to the Standard Model particles.

We now discuss the Lagrangian in more detail, starting with the covariant derivatives of ϕ_1 and ϕ_2 ,

$$|D_\mu \phi_i|^2 = |\partial_\mu \phi_i|^2 + i\tilde{g}A_\mu^a \left((\partial^\mu \phi_i)^\dagger T^a \phi_i - \text{h.c.} \right) + \tilde{g}^2 A_\mu^a A^{\mu b} \phi_i^\dagger T^a T^b \phi_i. \quad (3.13)$$

Inserting the parametrization (3.12) with $v_4 = 0$, we get the kinetic terms for the scalars

$$|\partial_\mu \phi_1|^2 + |\partial_\mu \phi_2|^2 = \frac{1}{2} \sum_{i=1}^4 (\partial_\mu \varphi_i)^2, \quad (3.14)$$

the mass terms for the gauge fields, the mixing terms as well as the gauge-scalar interactions. Let us first discuss the terms quadratic in the fields. The third term on the r.h.s. of eq. (3.13) contains the mass terms for the gauge bosons,

$$\mathcal{L} \supset \frac{1}{2} \left(\vec{A}_{(1,4)}^T \mathcal{M}_{(1,4)} \vec{A}_{(1,4)} + \vec{A}_{(2,5)}^T \mathcal{M}_{(2,5)} \vec{A}_{(2,5)} + \vec{A}_{(3,6,8)}^T \mathcal{M}_{(3,6,8)} \vec{A}_{(3,6,8)} + \mathcal{M}_{(7)} A_\mu^7 A^{\mu 7} \right), \quad (3.15)$$

where we have used the shorthand notation $\vec{A}_{(1,4)}^T \equiv (A_\mu^1, A_\mu^4)$ and similarly for the other gauge bosons. The mass-squared matrices are

$$\mathcal{M}_{(1,4)} = \mathcal{M}_{(2,5)} = \frac{\tilde{g}^2}{4} \begin{pmatrix} v_2^2 & v_2 v_3 \\ v_2 v_3 & v_1^2 + v_3^2 \end{pmatrix},$$

$$\begin{aligned} \mathcal{M}_{(3,6,8)} &= \frac{\tilde{g}^2}{4} \begin{pmatrix} v_2^2 & -v_2 v_3 & -v_2^2/\sqrt{3} \\ -v_2 v_3 & v_1^2 + v_2^2 + v_3^2 & -v_2 v_3/\sqrt{3} \\ -v_2^2/\sqrt{3} & -v_2 v_3/\sqrt{3} & (4v_1^2 + v_2^2 + 4v_3^2)/3 \end{pmatrix}, \\ \mathcal{M}_{(7)} &= \frac{\tilde{g}^2}{4} (v_1^2 + v_2^2 + v_3^2). \end{aligned} \quad (3.16)$$

The second term on the r.h.s. of eq. (3.13) contains the $A_\mu \partial^\mu \varphi$ mixing terms

$$\mathcal{L} \supset \frac{\tilde{g} v_2}{2} A^{\mu 6} \partial_\mu \varphi_4 - \frac{\tilde{g} v_3}{\sqrt{3}} A^{\mu 8} \partial_\mu \varphi_4 + \frac{\tilde{g} v_3}{2} A^{\mu 7} \partial_\mu \varphi_2 - \frac{\tilde{g} v_2}{2} A^{\mu 7} \partial_\mu \varphi_3. \quad (3.17)$$

In general, terms of the type $\kappa_{ai} A_\mu^a \partial^\mu \varphi^i$ can be removed by the field redefinition

$$\tilde{A}_\mu^a = A_\mu^a + \partial_\mu Y^a, \quad \text{where } Y^a \equiv (\mathcal{M}^{-1})_{ab} \kappa_{bi} \varphi^i. \quad (3.18)$$

We then have

$$\begin{aligned} & -\frac{1}{2} \text{tr}\{G_{\mu\nu} G^{\mu\nu}\} + \frac{1}{2} (\partial_\mu \varphi^i)^2 + \frac{1}{2} \mathcal{M}_{ab} A_\mu^a A^{\mu b} + \kappa_{ai} A_\mu^a \partial^\mu \varphi^i = \\ & -\frac{1}{2} \text{tr}\{\tilde{G}_{\mu\nu} \tilde{G}^{\mu\nu}\} + \frac{1}{2} \gamma_{ij} (\partial_\mu \varphi^i) (\partial^\mu \varphi^j) + \frac{1}{2} \mathcal{M}_{ab} \tilde{A}_\mu^a \tilde{A}^{\mu b}, \end{aligned} \quad (3.19)$$

where

$$\tilde{G}_{\mu\nu} = \partial_\mu \tilde{A}_\nu - \partial_\nu \tilde{A}_\mu + i\tilde{g}[\tilde{A}_\mu - \partial_\mu Y, \tilde{A}_\nu - \partial_\nu Y], \quad (3.20a)$$

$$\gamma_{ij} = \delta_{ij} - \kappa_{ia}^T \mathcal{M}_{ab}^{-1} \kappa_{bj}, \quad (3.20b)$$

and we have defined $Y = Y^a T^a$. The kinetic terms for the vector fields are still canonically normalised, unlike those for the scalar fields. The latter can be normalised canonically by a further field redefinition

$$\tilde{\varphi}^i = \omega_{ik} \varphi^k, \quad \text{where } (\omega^T \omega)_{ij} = \gamma_{ij}. \quad (3.21)$$

Note that the term $-\frac{1}{2} \text{tr}\{\tilde{G}_{\mu\nu} \tilde{G}^{\mu\nu}\}$ includes additional couplings of the gauge fields to the scalars. The resulting vertices are obtained by replacing A_μ^a in the triple and quartic gauge boson terms by $-\partial_\mu Y^a$ such that interactions of the type $(\tilde{A})^3 \partial Y, (\tilde{A})^2 (\partial Y)^2$, etc. arise. The analysis of the general case is very cumbersome, so further we will focus on the simple case of $v_3 = 0$ which retains all the relevant physics. In that case, the couplings involving ∂Y play no role in the DM phenomenology.

3.4 Detailed study for $v_{3,4} = 0$

As long as v_1 and v_2 are nonzero, $SU(3)$ is broken completely. Hence it suffices to consider the case $v_3 = 0$, which simplifies the analysis. Then the only mixing term among the gauge bosons is $A_\mu^3 A^{\mu 8}$ and the only gauge-scalar mixing terms are $A_\mu^6 \partial^\mu \varphi_4, A_\mu^7 \partial^\mu \varphi_3$.

In what follows, we restrict ourselves to the case $v_3 = 0$.

3.4.1 Masses

Gauge boson masses. The gauge boson mass eigenstates are

$$\begin{pmatrix} A'_\mu{}^3 \\ A'_\mu{}^8 \end{pmatrix} = \begin{pmatrix} \cos \alpha A_\mu^3 + \sin \alpha A_\mu^8 \\ \cos \alpha A_\mu^8 - \sin \alpha A_\mu^3 \end{pmatrix}, \quad \text{where} \quad \tan 2\alpha = \frac{\sqrt{3}v_2^2}{2v_1^2 - v_2^2}, \quad (3.22)$$

and the masses are²

$$m_{A'^3}^2 = \frac{\tilde{g}^2 v_2^2}{4} \left(1 - \frac{\tan \alpha}{\sqrt{3}}\right), \quad m_{A'^8}^2 = \frac{\tilde{g}^2 v_1^2}{3} \frac{1}{1 - \frac{\tan \alpha}{\sqrt{3}}}, \quad (3.23)$$

while the other gauge boson masses can be read off directly from eq. (3.16) for $v_3 = 0$:

$$m_{A^1}^2 = m_{A^2}^2 = \frac{\tilde{g}^2}{4} v_2^2, \quad m_{A^4}^2 = m_{A^5}^2 = \frac{\tilde{g}^2}{4} v_1^2, \quad m_{A^6}^2 = m_{A^7}^2 = \frac{\tilde{g}^2}{4} (v_1^2 + v_2^2). \quad (3.24)$$

For $v_2 < v_1$, the light fields are $A_\mu^{1,2}$ and $A_\mu'^3$, with the latter being the lightest. It is instructive to consider the case $v_2^2 \ll v_1^2$, so that $\tan \alpha$ is small and positive. Then $A_\mu'^3$ is slightly lighter than A_μ^1 and A_μ^2 by a factor of $(1 - \frac{\tan \alpha}{\sqrt{3}})^{1/2}$, while the other five dark gauge bosons are all much heavier, by a factor of order v_1/v_2 . The mass degeneracy between A_μ^1 and A_μ^2 persists at loop level by symmetry arguments (see section 4).

One can easily verify that the heavier states A_μ^{4-7} and $A_\mu'^8$ all decay into the light states and the SM particles. The decay proceeds via emission (off-shell or on-shell) of the CP even scalars which couple to the SM Higgs and thus to all other SM fields.

The 3 lightest states all have different $Z_2 \times Z_2'$ parities such that they cannot decay into each other by emitting SM particles. The only scalar with negative parity is φ_4 , however it is generally heavy (see below) and does not contribute to the above decay. Hence, $A_\mu^{1,2}$ and $A_\mu'^3$ are stable.³

Gauge boson – scalar mixing. According to eq. (3.18), the $A_\mu \partial^\mu \varphi$ mixing terms are removed by the redefinition (which does not affect the gauge boson masses)

$$\begin{aligned} \tilde{A}_\mu^6 &= A_\mu^6 + \partial_\mu Y^6, & \text{where} & \quad Y^6 = \frac{2}{\tilde{g}} \frac{v_2}{v_1^2 + v_2^2} \varphi^4, \\ \tilde{A}_\mu^7 &= A_\mu^7 + \partial_\mu Y^7, & \text{where} & \quad Y^7 = -\frac{2}{\tilde{g}} \frac{v_2}{v_1^2 + v_2^2} \varphi^3, \end{aligned} \quad (3.25)$$

while, according to eq. (3.21), the canonically normalised scalars are

$$\tilde{\varphi}^3 = \frac{v_1}{\sqrt{v_1^2 + v_2^2}} \varphi^3, \quad \tilde{\varphi}^4 = \frac{v_1}{\sqrt{v_1^2 + v_2^2}} \varphi^4. \quad (3.26)$$

²Note that $\frac{\tan \alpha}{\sqrt{3}} = \frac{v_2^2}{4v_1^2} + \mathcal{O}\left(\frac{v_2^6}{v_1^6}\right)$, $\cos \alpha = 1 - \frac{3v_2^4}{32v_1^4} + \mathcal{O}\left(\frac{v_2^8}{v_1^8}\right)$ and $\sin \alpha = \frac{\sqrt{3}v_2^2}{4v_1^2} + \mathcal{O}\left(\frac{v_2^6}{v_1^6}\right)$.

³The parities allow for a decay of one DM component into two others, however this is forbidden kinematically.

Scalar masses. The general scalar potential has many parameters. To make our discussion more transparent, let us assume the symmetry $\phi_2 \rightarrow -\phi_2$ which does not affect the essence of our considerations and requires $m_{12}^2 = \lambda_{H12} = \lambda_6 = \lambda_7 = 0$. In this case, it turns out that the potential has no local minima with all of v, v_1, v_2, v_3 being nonzero such that setting $v_3 = 0$ is actually required. It should be noted that in practice we are considering the limit in which the above quantities are very small but nonzero such that the decay channels for heavy particles, e.g. $\tilde{\varphi}_3$, into the SM fields are open.

The VEVs v, v_1, v_2 can be expressed in a compact form using the matrix

$$\mathbf{\Lambda} \equiv \begin{pmatrix} \lambda_H & \lambda_{H11} & \lambda_{H22} \\ \lambda_{H11} & \lambda_1 & \lambda_3 \\ \lambda_{H22} & \lambda_3 & \lambda_2 \end{pmatrix} \quad (3.27)$$

as well as the matrices $\mathbf{\Lambda}_{ij}$ defined as $(-1)^{i+j}$ times the matrix obtained by deleting the i -th row and j -th column of $\mathbf{\Lambda}$, i.e. $\det \mathbf{\Lambda}_{ij}$ is the (i, j) -cofactor of $\mathbf{\Lambda}$. One finds

$$\begin{aligned} v^2 &= -2(m_H^2 \det \mathbf{\Lambda}_{11} + m_{11}^2 \det \mathbf{\Lambda}_{21} + m_{22}^2 \det \mathbf{\Lambda}_{31}) / \det \mathbf{\Lambda}, \\ v_1^2 &= -2(m_H^2 \det \mathbf{\Lambda}_{12} + m_{11}^2 \det \mathbf{\Lambda}_{22} + m_{22}^2 \det \mathbf{\Lambda}_{32}) / \det \mathbf{\Lambda}, \\ v_2^2 &= -2(m_H^2 \det \mathbf{\Lambda}_{13} + m_{11}^2 \det \mathbf{\Lambda}_{23} + m_{22}^2 \det \mathbf{\Lambda}_{33}) / \det \mathbf{\Lambda}. \end{aligned} \quad (3.28)$$

The mass terms for the scalars are

$$-\mathcal{L} \supset \frac{1}{2} \Phi^T \mathbf{m}^2 \Phi + \frac{1}{4} (\lambda_4 + \lambda_5) (v_1^2 + v_2^2) \tilde{\varphi}_3^2 + \frac{1}{4} (\lambda_4 - \lambda_5) (v_1^2 + v_2^2) \tilde{\varphi}_4^2, \quad (3.29)$$

where $\Phi^T \equiv (h, \varphi_1, \varphi_2)$ and

$$\mathbf{m}^2 = \begin{pmatrix} \lambda_H v^2 & \lambda_{H11} v v_1 & \lambda_{H22} v v_2 \\ \lambda_{H11} v v_1 & \lambda_1 v_1^2 & \lambda_3 v_1 v_2 \\ \lambda_{H22} v v_2 & \lambda_3 v_1 v_2 & \lambda_2 v_2^2 \end{pmatrix}. \quad (3.30)$$

Note that $\tilde{\varphi}_3$ and $\tilde{\varphi}_4$ are generally heavier than A^1, A^2, A^3 since their masses involve v_1 (unless $\lambda_{4,5}$ are very small).

The matrix \mathbf{m}^2 is positive definite if and only if $\det \mathbf{\Lambda} > 0$, $\det \mathbf{\Lambda}_{33} \equiv \lambda_H \lambda_1 - \lambda_{H11}^2 > 0$ and $\lambda_H > 0$. It can be diagonalised by an orthogonal transformation $O^T \mathbf{m}^2 O = \text{diag}(m_{H_1}^2, m_{H_2}^2, m_{H_3}^2)$, where

$$O = \begin{pmatrix} c_{12} c_{13} & s_{12} & c_{12} s_{13} \\ -c_{13} c_{23} s_{12} - s_{13} s_{23} & c_{12} c_{23} & -c_{23} s_{12} s_{13} + c_{13} s_{23} \\ -c_{23} s_{13} + c_{13} s_{12} s_{23} & -c_{12} s_{23} & c_{13} c_{23} + s_{12} s_{13} s_{23} \end{pmatrix} \quad (3.31)$$

and we have used the abbreviation $s_{ij} \equiv \sin \theta_{ij}$, $c_{ij} \equiv \cos \theta_{ij}$. Instead of providing the most general formulae, let us focus on a simplified case. Suppose that the (12) and (23) entries of \mathbf{m}^2 are much smaller than the other matrix elements, in other words, that λ_{H11} and λ_3 are very small. In this case, the Higgs mixing with φ_2 is the dominant one. The reason behind this choice is that the DM constituents A^1, A^2 and A^3 all have a significant coupling

to φ_2 , see table 3. Given that the mixing between the Higgs and φ_2 is substantial, this facilitates DM annihilation into the SM fields. Clearly, similar considerations also apply to the general case with all the angles θ_{ij} being significant.

Assuming small $\mathbf{m}_{12}^2, \mathbf{m}_{23}^2$, one finds

$$\theta_{12} \approx p_{32} \mathbf{m}_{12}^2 + q \mathbf{m}_{23}^2, \tag{3.32a}$$

$$\theta_{23} \approx p_{21} \mathbf{m}_{23}^2 + q \mathbf{m}_{12}^2, \tag{3.32b}$$

$$\tan 2\theta_{13} \approx \frac{2\lambda_{H22}vv_2}{\lambda_2v_2^2 - \lambda_Hv^2}, \tag{3.32c}$$

where $p_{ij} = (\mathbf{m}_{ii}^2 - \mathbf{m}_{jj}^2)/s$, $q = \mathbf{m}_{13}^2/s$ and $s = (\mathbf{m}_{13}^2)^2 + (\mathbf{m}_{11}^2 - \mathbf{m}_{22}^2)(\mathbf{m}_{22}^2 - \mathbf{m}_{33}^2)$. The mass eigenstates are

$$\begin{pmatrix} h_1 \\ \mathcal{H} \\ h_2 \end{pmatrix} \equiv O^T \Phi \approx \begin{pmatrix} c_{13}h - s_{13}\varphi_2 \\ \varphi_1 \\ c_{13}\varphi_2 + s_{13}h \end{pmatrix} - \theta_{12} \begin{pmatrix} c_{13}\varphi_1 \\ -h \\ s_{13}\varphi_1 \end{pmatrix} - \theta_{23} \begin{pmatrix} s_{13}\varphi_1 \\ \varphi_2 \\ -c_{13}\varphi_1 \end{pmatrix}. \tag{3.33}$$

and the mass-squared eigenvalues are

$$m_{h_1, h_2}^2 \approx \frac{1}{2}(\lambda_2v_2^2 + \lambda_Hv^2) \mp \frac{\lambda_2v_2^2 - \lambda_Hv^2}{2 \cos 2\theta_{13}} \tag{3.34a}$$

$$m_{\mathcal{H}}^2 \approx \lambda_1v_1^2. \tag{3.34b}$$

The eigenstates h_1, h_2 are typically the lighter ones, while \mathcal{H} is heavy (unless λ_1 is very small). In our analysis of DM phenomenology, we retain only the former states. In summary, the relevant light fields are the DM components A^1, A^2 and A'^3 as well as the mediators h_1, h_2 which link the dark sector to the SM fermions and gauge bosons.

3.4.2 Couplings

The full list of the couplings is not necessary for our DM studies. The important couplings are those with two gauge bosons and one or two scalars at the vertex. In terms of the variables \tilde{A}_μ^a and $\tilde{\varphi}^i$, most of these are listed in table 3 and table 2. For our applications, the couplings of h_1 and h_2 are obtained from these tables using the relation (3.33), in which one may neglect θ_{12} and θ_{23} .

We focus on the case $v_1 \gg v_2$ so that DM consists of A^1, A^2 and A'^3 . Other fields with non-trivial parities decay into these states and the SM particles. For instance, the processes $\tilde{\varphi}_4 \rightarrow A^1A^2 + \text{SM}$ and $\tilde{\varphi}_4 \rightarrow A'^3 + \text{SM}$ are allowed. (When v_1 and v_2 are close, the composition of DM depends on the mass splittings.) DM annihilation and scattering proceeds through an exchange of h_1 and h_2 . Therefore, only the vertices involving these fields play a significant role.

4 Generalisation to arbitrary SU(N)

SU(N) is broken completely by generic VEVs of $N - 1$ fields ϕ_i in the fundamental representation. The ϕ_i 's can be gauge-transformed to the form

$$\phi_1 = \begin{pmatrix} 0 \\ 0 \\ \dots \\ 0 \\ \rho_1 \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} 0 \\ 0 \\ \dots \\ \rho_2^{(1)} \\ \rho_2^{(2)} e^{i\xi_2} \end{pmatrix}, \quad \dots, \quad \phi_{N-1} = \begin{pmatrix} 0 \\ \rho_{N-1}^{(1)} \\ \dots \\ \rho_{N-1}^{(N-2)} e^{i\xi_{N-1}^{(N-3)}} \\ \rho_{N-1}^{(N-1)} e^{i\xi_{N-1}^{(N-2)}} \end{pmatrix}. \quad (4.1)$$

Here the radial fields $\rho_i^{(j)}$ and the phases $\xi_i^{(j)}$ are real. We label the scalars such that the lightest gauge fields are associated with the SU(2) subgroup which gets broken at the last stage by a VEV of $\rho_{N-1}^{(1)}$. We assume that the VEVs as well as the couplings in the scalar potential are all real so that CP is preserved in the hidden sector.

The generalisation of the $Z_2 \times Z'_2$ parity to SU(N) is as follows. The transformation properties of the gauge fields are identified with those of the corresponding SU(N) generators. The basis of the $N(N - 1)$ off-diagonal generators T^{ab}, \tilde{T}^{ab} can be chosen as

$$\begin{aligned} (T^{ab})_{ij} &= \delta_{ia}\delta_{jb} + \delta_{ib}\delta_{ja}, \\ (\tilde{T}^{ab})_{ij} &= -i\delta_{ia}\delta_{jb} + i\delta_{ib}\delta_{ja}, \end{aligned} \quad (4.2)$$

where $a = 1, \dots, N - 1$ and $b = 2, \dots, N$. With the Cartan generators denoted by H^α , the Z_2 associated with complex conjugation of the group elements acts as

$$T^{ab} \rightarrow -T^{ab}, \quad \tilde{T}^{ab} \rightarrow \tilde{T}^{ab}, \quad H^\alpha \rightarrow -H^\alpha. \quad (4.3)$$

This is a well known outer automorphism of SU(N) which entails the corresponding symmetry of the Yang-Mills Lagrangian.

Another Z_2 can be defined by reflecting the off-diagonal generators containing nonzero elements in the first row:

$$\begin{aligned} T^{1a} &\rightarrow -T^{1a}, \quad \tilde{T}^{1a} \rightarrow -\tilde{T}^{1a}, \\ T^{bc} &\rightarrow T^{bc}, \quad \tilde{T}^{bc} \rightarrow \tilde{T}^{bc} \quad (b, c \geq 2), \\ H^\alpha &\rightarrow H^\alpha. \end{aligned} \quad (4.4)$$

It is easy to show that this Z_2 is an inner automorphism. Indeed, it corresponds to the group transformation with

$$U = e^{\frac{i\pi}{N}} \text{diag}(-1, 1, \dots, 1). \quad (4.5)$$

The gauge fields A_μ^{1-3} associated with the upper left SU(2) block

$$T^{12}, \quad \tilde{T}^{12}, \quad H^1 = \text{diag}(1, -1, 0, \dots, 0) \quad (4.6)$$

transform under these parities the same way as did the SU(2) gauge fields under $Z_2 \times Z'_2$ of the previous section. The above gauge transformation of course leaves the Yang-Mills Lagrangian invariant.

These symmetries are preserved by gauge interactions with scalars in our set-up. The Z_2 associated with complex conjugation acts on scalars by reflecting the complex phases, which therefore correspond to odd fields under the transformation (4.3). This symmetry is guaranteed by CP invariance of the Lagrangian and preserved by the vacuum (assuming no spontaneous CP violation). The second Z_2 is a gauge transformation. On vectors of the form $(0, a_1, \dots, a_{N-1})$, it acts as multiplication by an overall constant phase which cancels in all the Lagrangian terms. It is therefore a valid symmetry in the broken phase as well.

As long as the ϕ_i have a zero first component, the interaction vertices contain an even number of T^{1a} and \tilde{T}^{1a} . The gauge fields associated with $a > 2$ are heavier than those corresponding to $a = 2$. By virtue of the vertices involving $T^{12}T^{1k}$ ($k > 2$) and the matter fields, they decay to the lighter fields such that only the final $SU(2)$ block remains stable. (Similar considerations also apply to other heavy gauge fields.) Then DM is composed mostly of the aforementioned A_μ^{1-3} whose stability is enforced by $Z_2 \times Z'_2$.⁴

As in the $SU(3)$ case, DM consists of three components, two of which are degenerate in mass,

$$m_{A^{1'}} = m_{A^{2'}} \neq m_{A^{3'}} , \tag{4.7}$$

where $A_\mu^{1'-3'}$ are the mass eigenstates consisting mostly of A_μ^{1-3} with some admixture of other gauge fields (see the $SU(3)$ example). The degeneracy persists at loop level. This can be seen as follows. The $SU(N)$ Lie algebra possesses a discrete symmetry which interchanges the real and imaginary generators with nonzero entries in the first row,

$$T^{1a} \rightarrow \tilde{T}^{1a} , \quad \tilde{T}^{1a} \rightarrow -T^{1a} , \tag{4.8}$$

while all the other generators remain intact. This is achieved by the group transformation

$$U' = e^{\frac{-i\pi}{2N}} \text{diag}(i, 1, \dots, 1) , \tag{4.9}$$

which can be recognised as the square root of U in (4.5). This gauge transformation acts on ϕ_i with a zero first entry as an overall constant phase multiplication. Since such a phase cancels in all of the Lagrangian terms, (4.8) remains a valid symmetry even in the broken phase.

Consider now the mass matrix for the gauge fields associated with T^{1a} and \tilde{T}^{1a} . By virtue of (4.4), only fields corresponding to T^{1a} and \tilde{T}^{1a} can mix, while (4.3) forbids a mixing between the tilded and untilded fields. The resulting mass matrix for T^{1a} is then identical to that of \tilde{T}^{1a} according to (4.8). Analogous considerations apply to the kinetic terms. Hence the lightest eigenstates have the same mass.

The resulting DM phenomenology is analogous to that for the $SU(3)$ case.

5 Dark matter phenomenology

In what follows, we consider direct and indirect detection as well as relic abundance constraints on vector DM. In the $U(1)$ and $SU(2)$ cases, all the scattering processes are

⁴As before, we take the “phase” fields to be heavy since they get their masses from large VEVs $\langle \phi_i \rangle$, unlike A_μ^{1-3} . In the presence of more than $N - 1$ fundamentals, this logic no longer applies and the light phase fields can constitute DM.

mediated by h_1 and h_2 . For SU(3) and larger groups, further states can contribute. However, we make the simplifying assumption that the Higgs mixing with those states is small and/or such states are heavy. In that case, it suffices to consider an exchange of h_1 and h_2 only. We note that earlier phenomenological analyses of vector dark matter have appeared in various contexts [12]–[21]. The current collider searches for dark matter, e.g. in the form of a monojet plus missing energy, do not set further useful constraints on the model, see e.g. ref. [22].

5.1 U(1) dark matter

Let us start with the U(1) case. The relevant terms in the Lagrangian are

$$\begin{aligned} \mathcal{L} \supset & \frac{1}{2} m_A^2 A_\mu A^\mu + \frac{\tilde{g} m_A}{2} (-h_1 \sin \theta + h_2 \cos \theta) A_\mu A^\mu \\ & + \frac{\tilde{g}^2}{8} (h_1^2 \sin^2 \theta - 2h_1 h_2 \sin \theta \cos \theta + h_2^2 \cos^2 \theta) A_\mu A^\mu. \end{aligned} \quad (5.1)$$

The couplings of h_1 and h_2 to the SM fields are those of the SM Higgs up to the suppression factors of $\cos \theta$ and $\sin \theta$, respectively. A phenomenological analysis of this model in the decoupling limit $m_{h_2} \gg m_{h_1}$, $\sin \theta \rightarrow 0$ can be found in [12] and [4, 13]. Related studies have also appeared in [14, 15].

The DM scattering on nucleons proceeds through the t -channel exchange of $h_{1,2}$ and leads to the following spin-independent cross section (see e.g. [4]),

$$\sigma_{A-N}^{\text{SI}} = \frac{g^2 \tilde{g}^2}{16\pi} \frac{m_N^4 f_N^2}{m_W^2} \frac{(m_{h_2}^2 - m_{h_1}^2)^2 \sin^2 \theta \cos^2 \theta}{m_{h_1}^4 m_{h_2}^4}, \quad (5.2)$$

where m_N is the nucleon mass and $f_N \simeq 0.3$ parametrizes the Higgs-nucleon coupling. One should keep in mind that there is significant uncertainty in f_N and here we use the value somewhat smaller than the one assumed in [4]. As expected, h_1 and h_2 contribute with opposite signs. Since $m_{h_2}^2 \gg m_{h_1}^2$ in realistic cases, the cancellation is not very significant.

The calculation of the DM annihilation cross section is more involved due to a few contributing diagrams (figure 1). To compute the DM relic abundance, we use the software package micrOMEGAs 4.1.8 [23]. It is important to note that the leading s -channel annihilation amplitude due to the $h_{1,2}$ -exchange is proportional to the factor

$$\mathcal{A}^{\text{annih}} \propto \sin \theta \cos \theta \left(\frac{1}{s - m_{h_1}^2} - \frac{1}{s - m_{h_2}^2} \right), \quad (5.3)$$

where the s -parameter can be approximated by $s \simeq 4m_A^2$. Therefore, the amplitude is highly suppressed at $m_A \gg m_{h_2}$. Although other annihilation channels remain available, this makes DM annihilation inefficient for heavy masses and the corresponding parameter space is challenged by the direct detection constraint.

Clearly, the s -channel annihilation becomes very efficient around the resonances, $m_A \simeq m_{h_1}/2$ and $m_A \simeq m_{h_2}/2$. In this case, a very small gauge coupling is sufficient to obtain the right relic abundance. The first resonance is quite narrow due to the small width of the SM




Figure 1. Leading diagrams for vector DM annihilation.

Higgs, whereas the second resonance is broad since many decay channels are available to h_2 . For $m_{h_2} > 2m_{h_1}$, the decay $h_2 \rightarrow h_1 h_1$ becomes important. Its significance depends on $\lambda_{h\phi}$ of eq. (1.1) with $\text{BR}_{h_2 \rightarrow h_1 h_1}$ increasing for larger $\lambda_{h\phi}$ (see the explicit formulae in [8]). The resonance is widened further by the thermal averaging over the DM momentum.

The most important features of DM annihilation are associated with the s -channel. Other channels play a less significant role. Similar considerations apply to the indirect detection constraint due to the gamma ray emission in the process of DM annihilation (FERMI).

The last constraint we impose is of theoretical nature. We require that the theory be perturbative at the TeV scale. One way to enforce it is to demand perturbative unitarity at tree level, for instance, in the process $h_i h_i \rightarrow h_j h_j$. The resulting constraint was estimated in [24] to be

$$\lambda_i < \mathcal{O}(4\pi/3), \tag{5.4}$$

where λ_i are the scalar quartic couplings. We define the quartic couplings involving ϕ by $\Delta V_{\text{quart}} = \lambda_{h\phi} |H|^2 |\phi|^2 + \frac{1}{2} \lambda_\phi |\phi|^4$. They can be expressed in terms of the masses, the gauge coupling and the mixing angle as (see e.g. [8])

$$\begin{aligned} \lambda_{h\phi} &= \tilde{g} \sin 2\theta \frac{m_{h_2}^2 - m_{h_1}^2}{4vm_A}, \\ \lambda_\phi &= \frac{4\lambda_{h\phi}^2 v^2}{\sin^2 2\theta (m_{h_2}^2 - m_{h_1}^2)} \left(\frac{m_{h_2}^2}{m_{h_2}^2 - m_{h_1}^2} - \sin^2 \theta \right), \end{aligned} \tag{5.5}$$

where we have used $\tilde{v} = 2m_A/\tilde{g}$. This implies that both $\lambda_{h\phi}$ and λ_ϕ become large for heavy h_2 or for light dark matter. As a result, eq. (5.4) imposes an important constraint on our model. We further require the standard perturbativity constraint

$$\frac{\tilde{g}^2}{4\pi} < 1, \tag{5.6}$$

which we find less significant for our purposes.

Our results for U(1) DM are presented in figure 2, upper row. We include the constraints from PLANCK [25] (relic abundance), LUX [26] (direct detection), FERMI [27] (indirect detection) and perturbativity of λ_i . The area between the red lines is consistent with the thermal relic DM abundance measured by PLANCK. The LUX data provide the strongest constraint on the allowed parameter space. The FERMI bound is typically relevant for light DM, while the perturbativity bound becomes important for heavy h_2 . In each panel, the mixing angle is chosen such that, for a given m_{h_2} , it is consistent with the




Figure 2. Constraints on the gauge coupling \tilde{g} vs DM mass m_A for U(1) and SU(2). The area between the red lines is favoured by the DM relic abundance, while the regions above the dashed blue, dotted black and green lines are ruled out by direct detection, indirect detection and perturbativity of λ_i , respectively. *Upper row:* U(1) dark matter with $\sin \theta = 0.3, m_{h_2} = 280$ GeV (left), $\sin \theta = 0.3, m_{h_2} = 1$ TeV (center), $\sin \theta = 0.2, m_{h_2} = 3$ TeV (right). *Lower row:* SU(2) dark matter with $\sin \theta = 0.3, m_{h_2} = 280$ GeV (left), $\sin \theta = 0.3, m_{h_2} = 1$ TeV (center), $\sin \theta = 0.2, m_{h_2} = 3$ TeV (right).

LHC and EW precision data [8]. Heavier h_2 imply smaller $\sin \theta$, so we take $\sin \theta = 0.3$ for the left and center panels, and $\sin \theta = 0.2$ for the right panel.

The two dips are associated with the resonant annihilation through h_1 and h_2 . The second resonance gets broader with increasing h_2 due to the increase in $\lambda_{h\phi}$ and availability of the decay $h_2 \rightarrow h_1 h_1$. The area around this resonance constitutes the largest parameter space consistent with all of the constraints. For $m_{h_2} = 280$ GeV, the allowed DM mass range is about 100 GeV; for $m_{h_2} = 1$ TeV, it widens to 1 TeV, and for $m_{h_2} = 3$ TeV, it reaches more than 3 TeV. The resonance is broadened by the thermal averaging over the DM momentum, so even though it appears very broad for $m_{h_2} = 3$ TeV, it is still consistent with perturbativity.

The dip associated with the resonant annihilation through h_1 is quite narrow and does not open up further significant areas of parameter space. Other features of the PLANCK curve are local peaks corresponding to the kinematic opening of additional annihilation




Figure 3. Semiannihilation of vector DM.

channels. For instance, the peak at $m_A \sim 80$ GeV is associated with the W^+W^- final state. There are further visible peaks at $(m_{h_2} + m_{h_1})/2$ and m_{h_2} .

Away from the resonances, there appears to be a further allowed region at $m_A > m_{h_2}$. Since the h_2 production is not suppressed by $\sin\theta$, in this case the t -channel annihilation $AA \rightarrow h_2h_2$ and the quartic interactions dominate. The Planck-allowed strip is dangerously close to the LUX bound, so the conclusion depends strongly on the nucleon-Higgs coupling $f_N \simeq 0.3$, which suffers from substantial uncertainties.

5.2 SU(2) dark matter

Aside from the gauge self-interactions, the Lagrangian (5.1) applies to the SU(2) case as well, up to the summation over the 3 species, $A_\mu A^\mu \rightarrow A_\mu^a A^{a\mu}$. The main change compared to the U(1) case is that the annihilation cross section decreases since only the species with the same group index can annihilate through the Higgs-like states. In order to keep the same relic abundance, one needs to increase the gauge coupling. Since the s -channel annihilation through $h_{1,2}$ often dominates, this amounts approximately to

$$\tilde{g} \rightarrow \sqrt{3}\tilde{g}, \quad (5.7)$$

whereas if the t -channel and the quartic interactions dominate, the rescaling factor is closer to $3^{1/4}$. The direct detection constraint remains the same as in the U(1) case since particles with different group indices scatter the same way on nucleons. This decreases somewhat the allowed parameter space compared to the Abelian case (figure 2).

Non-Abelian DM features a semi-annihilation channel $AA \rightarrow Ah_{1,2}$ (figure 3). In some regimes, for example at large \tilde{g} and small $\sin\theta$, it can even dominate [11] (see also [28, 29]). Using the analytical results of [11], we find that DM semi-annihilation is insignificant in the relevant parameter regions. For example, at $m_A > m_{h_2}$ the $\sin\theta$ -unsuppressed and potentially important channel $AA \rightarrow Ah_2$ opens up, yet it is dominated by $AA \rightarrow h_2h_2$. Also, around the resonances the gauge coupling is rather small which diminishes the relative importance of semi-annihilation.

No firm conclusion can be reached as to whether the region $m_A > m_{h_2}$ is allowed. As stated above, the uncertainties in f_N play a critical role due to the proximity of the Planck band and the LUX bound.

5.3 SU(3) dark matter

In the SU(3) case, DM is composed again of 3 species with two of them being degenerate (A_μ^1, A_μ^2) in mass and the third one being lighter ($A_\mu^{3'}$). This is a result of the mixing

between the gauge bosons corresponding to the Cartan generators of SU(3). Therefore, while the couplings of A_μ^1, A_μ^2 to the Higgs like scalars remain the same as in the SU(2) case, the coupling of $A_\mu^{3'}$ changes. In terms of $m_A \equiv m_{A^{1,2}}$, the relevant Lagrangian reads

$$\begin{aligned} \mathcal{L} \supset & \frac{1}{2} m_A^2 \left(\sum_{a=1,2} A_\mu^a A^{a\mu} + \left(1 - \frac{\tan \alpha}{\sqrt{3}}\right) A_\mu^{3'} A^{3'\mu} \right) \\ & + \frac{\tilde{g} m_A}{2} (-h_1 \sin \theta + h_2 \cos \theta) \left(\sum_{a=1,2} A_\mu^a A^{a\mu} + \left(\cos \alpha - \frac{\sin \alpha}{\sqrt{3}}\right)^2 A_\mu^{3'} A^{3'\mu} \right) \\ & + \frac{\tilde{g}^2}{8} (h_1^2 \sin^2 \theta - 2h_1 h_2 \sin \theta \cos \theta + h_2^2 \cos^2 \theta) \left(\sum_{a=1,2} A_\mu^a A^{a\mu} + \left(\cos \alpha - \frac{\sin \alpha}{\sqrt{3}}\right)^2 A_\mu^{3'} A^{3'\mu} \right). \end{aligned} \tag{5.8}$$

The mass of the lighter DM component is reduced by the factor $(1 - \frac{1}{\sqrt{3}} \tan \alpha)^{1/2}$ compared to that of A_μ^1 and A_μ^2 , while the gauge-scalar couplings decrease by a factor $(\cos \alpha - \frac{1}{\sqrt{3}} \sin \alpha)^2$. Therefore, the lighter state has a smaller annihilation cross section. Note that $\sin \alpha \simeq \frac{\sqrt{3} v_2^2}{4 v_1^2}$ and even a factor of two difference in the triplet VEVs leads to a rather small $\alpha \sim 10^{-1}$. In that case, there is no tangible difference between the SU(2) and the SU(3) analyses.

To understand where differences can appear, it is instructive to consider the limit $v_1 \simeq v_2$. Although in this case there are further relatively light states that can mediate DM annihilation (e.g. $A_\mu^{8'}$), let us consider the simplified example in which only the same states are allowed to contribute in the SU(2) and SU(3) set-ups. The main features (figure 4) are that the gauge coupling must be larger in the SU(3) case in order to allow for efficient annihilation of $A_\mu^{3'}$ and that the resonant dips are slightly shifted due to a different freeze-out temperature (see e.g. [30]). The DM density today is dominated by the lighter component. Since it couples to nucleons weaker than $A_\mu^{1,2}$ do, the direct detection constraint relaxes. Understanding further features of the model would require precise knowledge of the spectrum and the couplings, which we relegate to future work [31].

Finally, one should keep in mind that there exists the coupling $A_\mu^{3'} \tilde{\varphi}_3 \tilde{\varphi}_4$, where $A_\mu^{3'}$ and $\tilde{\varphi}_4$ have the same parities. If $\tilde{\varphi}_4$ were light, that is, $\lambda_4 - \lambda_5 \ll 1$ and/or $v_1 \sim v_2$ (see eq. (3.29)), the decay $A_\mu^{3'} \rightarrow \tilde{\varphi}_4 + \text{SM}$ would occur. In that case, DM would consist of both the vector and scalar components. We defer a detailed study of this scenario to future work [31].

6 Summary and conclusions

In this paper, we have considered the possibility that the hidden sector enjoys SU(N) gauge symmetry and couples to the Standard Model through the Higgs portal. We find that when endowed with a “minimal” matter content, such hidden sectors lead naturally to stable vector dark matter. The underlying Lie group symmetries which stabilise DM are associated with complex conjugation of the group elements and discrete gauge transformations.

We require complete breaking of hidden SU(N) by scalar multiplets to avoid massless states (barring confinement in some cases). That can be done in a minimal fashion by introducing $N - 1$ scalar multiplets in the fundamental representation, which develop generic




Figure 4. Illustration of the main features of SU(3) vs SU(2) DM in the *extreme* case. Here $v_1/v_2 = 1.1$ and $\sin \theta = 0.3, m_{h_2} = 280$ GeV. In the SU(3) case, M_{DM} stands for the mass of the (dominant) lighter component $A_\mu^{3'}$. See figure 2 for further details.

VEVs. If the scalar sector preserves CP , the above-mentioned discrete symmetries of the Lie group generalise to full-fledged symmetries of the model and lead to stable gauge fields. When sufficiently light, they constitute all of dark matter. In this case, DM consists of 3 components associated with an SU(2) subgroup which hosts the lightest gauge fields $A_\mu^{1'}$, $A_\mu^{2'}$ and $A_\mu^{3'}$. Two of them ($A_\mu^{1'}$, $A_\mu^{2'}$) are always degenerate in mass, while for $N = 2$ all 3 components have the same mass.

We have performed phenomenological analyses of U(1), SU(2) and SU(3) gauge field dark matter. We find that there are vast regions of parameter space where all of the relevant constraints are satisfied. In many of these regions, DM annihilation is facilitated by the broad resonances associated with the Higgs-like scalars. We also find that the SU(3) case appears very similar to that of SU(2), unless the scalar VEVs breaking SU(3) are close in magnitude.

Acknowledgments

The authors are indebted to Sasha Pukhov and Genevieve Belanger for their help with the new version of micrOMEGAs 4.1.8. C.G. and O.L. acknowledge support from the Academy of Finland, project “The Higgs boson and the Cosmos”.

This work was also supported by the Spanish MICINN’s Consolider-Ingenio 2010 Programme under grant Multi-Dark **CSD2009-00064**, the contract **FPA2010-17747** and the France-US PICS no. 06482. Y.M. acknowledges partial support from the European Union FP7 ITN INVISIBLES (Marie Curie Actions, PITN- GA-2011- 289442) and the ERC advanced grants Higgs@LHC and MassTeV. This research was also supported in part by the Research Executive Agency (REA) of the European Union under the Grant Agreement PITN-GA2012-316704 (“HiggsTools”).

The authors would like to thank the Instituto de Fisica Teorica (IFT UAM-CSIC) in Madrid for its support via the Centro de Excelencia Severo Ochoa Program under Grant SEV-2012-0249, during the Program ‘‘Identification of Dark Matter with a Cross-Disciplinary Approach’’ where some of the ideas presented in this paper were developed.

A Hidden SU(3) vector-scalar couplings for $v_3 = 0$

The tables below provide a list of most important gauge-scalar couplings. These are relevant to DM phenomenology as well as to understanding the decay channels of the heavier gauge fields.

a	b	i	j	coeff. of $\tilde{A}_\mu^a \tilde{A}^{\mu b} \tilde{\varphi}^i \tilde{\varphi}^j$
4	4	1	1	$\tilde{g}^2/8$
5	5	1	1	
6	6	1	1	
7	7	1	1	
8	8	1	1	$\tilde{g}^2/6$
1	1	2	2	$\tilde{g}^2/8$
2	2	2	2	
3	3	2	2	
6	6	2	2	
7	7	2	2	
8	8	2	2	$\tilde{g}^2/24$
3	8	2	2	$-\frac{1}{4\sqrt{3}}\tilde{g}^2$

a	b	i	j	coeff. of $\tilde{A}_\mu^a \tilde{A}^{\mu b} \tilde{\varphi}^i \tilde{\varphi}^j$
4	4	3	3	$\frac{1}{8}\tilde{g}^2 \frac{v_1^2+v_2^2}{v_1^2}$
5	5	3	3	
6	6	3	3	
7	7	3	3	$\frac{1}{6}\tilde{g}^2 \frac{v_1^2+v_2^2}{v_1^2}$
8	8	3	3	
4	4	4	4	$\frac{1}{8}\tilde{g}^2 \frac{v_1^2+v_2^2}{v_1^2}$
5	5	4	4	
6	6	4	4	
7	7	4	4	
8	8	4	4	$\frac{1}{6}\tilde{g}^2 \frac{v_1^2+v_2^2}{v_1^2}$
1	4	2	3	$\frac{1}{4}\tilde{g}^2 \frac{\sqrt{v_1^2+v_2^2}}{v_1}$
2	5	2	3	
3	6	2	3	$-\frac{1}{4}\tilde{g}^2 \frac{\sqrt{v_1^2+v_2^2}}{v_1}$
6	8	2	3	$-\frac{1}{4\sqrt{3}}\tilde{g}^2 \frac{\sqrt{v_1^2+v_2^2}}{v_1}$
1	5	2	4	$\frac{1}{4}\tilde{g}^2 \frac{\sqrt{v_1^2+v_2^2}}{v_1}$
2	4	2	4	$-\frac{1}{4}\tilde{g}^2 \frac{\sqrt{v_1^2+v_2^2}}{v_1}$
3	7	2	4	
7	8	2	4	$-\frac{1}{4\sqrt{3}}\tilde{g}^2 \frac{\sqrt{v_1^2+v_2^2}}{v_1}$

Table 2. Non-derivative couplings $\tilde{A}_\mu^a \tilde{A}^{\mu b} \tilde{\varphi}^i \tilde{\varphi}^j$.

a	b	i	coeff. of $\tilde{A}_\mu^a \tilde{A}^{\mu b} \tilde{\varphi}^i$
4	4	1	$\frac{1}{4} \tilde{g}^2 v_1$
5	5	1	
6	6	1	
7	7	1	
8	8	1	$\frac{1}{3} \tilde{g}^2 v_1$
1	1	2	$\frac{1}{4} \tilde{g}^2 v_2$
2	2	2	
3	3	2	
6	6	2	
7	7	2	
8	8	2	$\frac{1}{12} \tilde{g}^2 v_2$
3	8	2	$-\frac{1}{2\sqrt{3}} \tilde{g}^2 v_2$

a	b	i	coeff. of $\tilde{A}_\mu^a \tilde{A}^{\mu b} \tilde{\varphi}^i$
1	4	3	$\frac{1}{4} \tilde{g}^2 \frac{v_2 \sqrt{v_1^2 + v_2^2}}{v_1}$
2	5	3	
3	6	3	$-\frac{1}{4} \tilde{g}^2 \frac{v_2 \sqrt{v_1^2 + v_2^2}}{v_1}$
6	8	3	$-\frac{1}{4\sqrt{3}} \tilde{g}^2 \frac{v_2 \sqrt{v_1^2 + v_2^2}}{v_1}$
1	5	4	$\frac{1}{4} \tilde{g}^2 \frac{v_2 \sqrt{v_1^2 + v_2^2}}{v_1}$
2	4	4	$-\frac{1}{4} \tilde{g}^2 \frac{v_2 \sqrt{v_1^2 + v_2^2}}{v_1}$
3	7	4	
7	8	4	$-\frac{1}{4\sqrt{3}} \tilde{g}^2 \frac{v_2 \sqrt{v_1^2 + v_2^2}}{v_1}$

Table 3. Non-derivative couplings $\tilde{A}_\mu^a \tilde{A}^{\mu b} \tilde{\varphi}^i$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] V. Silveira and A. Zee, *Scalar Phantoms*, *Phys. Lett. B* **161** (1985) 136 [[INSPIRE](#)].
- [2] R. Schabinger and J.D. Wells, *A minimal spontaneously broken hidden sector and its impact on Higgs boson physics at the large hadron collider*, *Phys. Rev. D* **72** (2005) 093007 [[hep-ph/0509209](#)] [[INSPIRE](#)].
- [3] B. Patt and F. Wilczek, *Higgs-field portal into hidden sectors*, [hep-ph/0605188](#) [[INSPIRE](#)].
- [4] O. Lebedev, H.M. Lee and Y. Mambrini, *Vector Higgs-portal dark matter and the invisible Higgs*, *Phys. Lett. B* **707** (2012) 570 [[arXiv:1111.4482](#)] [[INSPIRE](#)].
- [5] T. Hambye, *Hidden vector dark matter*, *JHEP* **01** (2009) 028 [[arXiv:0811.0172](#)] [[INSPIRE](#)].
- [6] T. Hambye and M.H.G. Tytgat, *Confined hidden vector dark matter*, *Phys. Lett. B* **683** (2010) 39 [[arXiv:0907.1007](#)] [[INSPIRE](#)].
- [7] K.K. Boddy, J.L. Feng, M. Kaplinghat and T.M.P. Tait, *Self-Interacting Dark Matter from a Non-Abelian Hidden Sector*, *Phys. Rev. D* **89** (2014) 115017 [[arXiv:1402.3629](#)] [[INSPIRE](#)].
- [8] A. Falkowski, C. Gross and O. Lebedev, *A second Higgs from the Higgs portal*, *JHEP* **05** (2015) 057 [[arXiv:1502.01361](#)] [[INSPIRE](#)].
- [9] L.-F. Li, *Group Theory of the Spontaneously Broken Gauge Symmetries*, *Phys. Rev. D* **9** (1974) 1723 [[INSPIRE](#)].
- [10] P. Jetzer, J.M. Gerard and D. Wyler, *Possible Symmetry Breaking Patterns With Totally Symmetric and Antisymmetric Representations*, *Nucl. Phys. B* **241** (1984) 204 [[INSPIRE](#)].
- [11] F. D’Eramo, M. McCullough and J. Thaler, *Multiple Gamma Lines from Semi-Annihilation*, *JCAP* **04** (2013) 030 [[arXiv:1210.7817](#)] [[INSPIRE](#)].

- [12] S. Kanemura, S. Matsumoto, T. Nabeshima and N. Okada, *Can WIMP Dark Matter overcome the Nightmare Scenario?*, *Phys. Rev. D* **82** (2010) 055026 [[arXiv:1005.5651](#)] [[INSPIRE](#)].
- [13] A. Djouadi, O. Lebedev, Y. Mambrini and J. Quevillon, *Implications of LHC searches for Higgs-portal dark matter*, *Phys. Lett. B* **709** (2012) 65 [[arXiv:1112.3299](#)] [[INSPIRE](#)].
- [14] Y. Farzan and A.R. Akbarieh, *VDM: A model for Vector Dark Matter*, *JCAP* **10** (2012) 026 [[arXiv:1207.4272](#)] [[INSPIRE](#)].
- [15] S. Baek, P. Ko, W.-I. Park and E. Senaha, *Higgs Portal Vector Dark Matter: Revisited*, *JHEP* **05** (2013) 036 [[arXiv:1212.2131](#)] [[INSPIRE](#)].
- [16] T. Abe, M. Kakizaki, S. Matsumoto and O. Seto, *Vector WIMP Miracle*, *Phys. Lett. B* **713** (2012) 211 [[arXiv:1202.5902](#)] [[INSPIRE](#)].
- [17] F. Domingo, O. Lebedev, Y. Mambrini, J. Quevillon and A. Ringwald, *More on the Hypercharge Portal into the Dark Sector*, *JHEP* **09** (2013) 020 [[arXiv:1305.6815](#)] [[INSPIRE](#)].
- [18] C.-W. Chiang, T. Nomura and J. Tandean, *Nonabelian Dark Matter with Resonant Annihilation*, *JHEP* **01** (2014) 183 [[arXiv:1306.0882](#)] [[INSPIRE](#)].
- [19] H. Davoudiasl and I.M. Lewis, *Dark Matter from Hidden Forces*, *Phys. Rev. D* **89** (2014) 055026 [[arXiv:1309.6640](#)] [[INSPIRE](#)].
- [20] C.-R. Chen, Y.-K. Chu and H.-C. Tsai, *An Elusive Vector Dark Matter*, *Phys. Lett. B* **741** (2015) 205 [[arXiv:1410.0918](#)] [[INSPIRE](#)].
- [21] M. Duch, B. Grzadkowski and J. Wudka, *Classification of effective operators for interactions between the Standard Model and dark matter*, *JHEP* **05** (2015) 116 [[arXiv:1412.0520](#)] [[INSPIRE](#)].
- [22] J.S. Kim, O. Lebedev and D. Schmeier, *Higgsophilic gauge bosons and monojets at the LHC*, [arXiv:1507.08673](#) [[INSPIRE](#)].
- [23] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, *MicrOMEGAs4.1: two dark matter candidates*, *Comput. Phys. Commun.* **192** (2015) 322 [[arXiv:1407.6129](#)] [[INSPIRE](#)].
- [24] C.-Y. Chen, S. Dawson and I.M. Lewis, *Exploring resonant di-Higgs boson production in the Higgs singlet model*, *Phys. Rev. D* **91** (2015) 035015 [[arXiv:1410.5488](#)] [[INSPIRE](#)].
- [25] PLANCK collaboration, P.A.R. Ade et al., *Planck 2015 results. XIII. Cosmological parameters*, [arXiv:1502.01589](#) [[INSPIRE](#)].
- [26] LUX collaboration, D.S. Akerib et al., *First results from the LUX dark matter experiment at the Sanford Underground Research Facility*, *Phys. Rev. Lett.* **112** (2014) 091303 [[arXiv:1310.8214](#)] [[INSPIRE](#)].
- [27] FERMI-LAT collaboration, M. Ackermann et al., *Searching for Dark Matter Annihilation from Milky Way Dwarf Spheroidal Galaxies with Six Years of Fermi-LAT Data*, [arXiv:1503.02641](#) [[INSPIRE](#)].
- [28] C. Arina, T. Hambye, A. Ibarra and C. Weniger, *Intense Gamma-Ray Lines from Hidden Vector Dark Matter Decay*, *JCAP* **03** (2010) 024 [[arXiv:0912.4496](#)] [[INSPIRE](#)].
- [29] V.V. Khoze, C. McCabe and G. Ro, *Higgs vacuum stability from the dark matter portal*, *JHEP* **08** (2014) 026 [[arXiv:1403.4953](#)] [[INSPIRE](#)].
- [30] K. Griest and D. Seckel, *Three exceptions in the calculation of relic abundances*, *Phys. Rev. D* **43** (1991) 3191 [[INSPIRE](#)].
- [31] C. Gross, O. Lebedev and Y. Mambrini, to appear.