



Neutrino–antineutrino mass splitting in the Standard Model and baryogenesis

Kazuo Fujikawa^a, Anca Tureanu^{b,*}

^a Mathematical Physics Laboratory, RIKEN Nishina Center, Wako 351-0198, Japan

^b Department of Physics, University of Helsinki, P.O. Box 64, FIN-00014 Helsinki, Finland

ARTICLE INFO

Article history:

Received 29 September 2014

Received in revised form 27 January 2015

Accepted 29 January 2015

Available online 9 February 2015

Editor: G.F. Giudice

ABSTRACT

On the basis of a previously proposed mechanism of neutrino–antineutrino mass splitting in the Standard Model, which is Lorentz and $SU(2) \times U(1)$ invariant but non-local to evade the CPT theorem, we discuss the possible implications of neutrino–antineutrino mass splitting on neutrino physics and baryogenesis. It is shown that non-locality within a distance scale of the Planck length, that may not be fatal to unitarity in a generic effective theory, can generate the neutrino–antineutrino mass splitting of the order of the observed neutrino mass differences, which is tested in oscillation experiments, and a non-negligible baryon asymmetry depending on the estimate of sphaleron dynamics. The one-loop order induced electron–positron mass splitting in the Standard Model is shown to be finite and estimated at $\sim 10^{-20}$ eV, well below the experimental bound $< 10^{-2}$ eV. The induced CPT violation in the K -meson in the Standard Model is expected to be even smaller and well below the experimental bound $|m_K - m_{\bar{K}}| < 0.44 \times 10^{-18}$ GeV.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The baryogenesis on the basis of CPT violating proton–antiproton mass splitting [1] and neutrino–antineutrino mass splitting [2] has been discussed some time ago. The possible neutrino–antineutrino mass splitting has also been discussed in connection with the phenomenology of neutrino oscillation [3,4]. On the other hand, an experimental study [5] states that “we have detected no evidence for Lorentz invariance violation in the antineutrino data set.” It has been recently shown that the neutrino–antineutrino mass splitting can be realized in the Standard Model by preserving both the Lorentz invariance and the $SU(2) \times U(1)$ gauge symmetry [6]. This model is based on a Lorentz invariant CPT violation mechanism which also breaks C and CP but preserves T [7]. The basic ingredient to evade the CPT theorem [8] in this scheme is that the theory is assumed to be non-local [9,10], although the original scheme of CPT violation with T violation but preserving C and CP does not generate particle–antiparticle mass splitting [9]. It was emphasized in [10] that the Lorentz invariant CPT breaking scheme, which we adopt in the present study, is a very natural logical possibility. In this paper, we perform a detailed evaluation of this neutrino–antineutrino mass splitting in

the Standard Model and discuss its possible implications on neutrino physics and baryogenesis. It is shown that the non-locality within a distance scale of the Planck length, which may not be fatal to unitarity in a generic effective theory, can generate the neutrino–antineutrino mass splitting of the order of the observed neutrino mass differences and non-negligible baryon asymmetry, depending on the estimate of sphaleron dynamics. The one-loop order electron–positron mass splitting induced by the neutrino–antineutrino mass splitting in the Standard Model is confirmed to be finite and kept well below the experimental bound.

2. The model

We start with a minimal extension of the Standard Model [11] by incorporating the right-handed neutrino

$$\psi_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}, \quad e_R, \nu_R. \quad (1)$$

For simplicity, we consider only a single flavor of leptons. The part of the Standard Model Lagrangian relevant to our discussion is given by

$$\begin{aligned} \mathcal{L} = & i\bar{\psi}_L \gamma^\mu \left(\partial_\mu - igT^a W_\mu^a - i\frac{1}{2}g'Y_L B_\mu \right) \psi_L \\ & + i\bar{e}_R \gamma^\mu (\partial_\mu + ig' B_\mu) e_R + i\bar{\nu}_R \gamma^\mu \partial_\mu \nu_R \end{aligned}$$

* Corresponding author.

E-mail address: anca.tureanu@helsinki.fi (A. Tureanu).

$$+ \left[-\frac{\sqrt{2}m_e}{v} \bar{e}_R \phi^\dagger \psi_L - \frac{\sqrt{2}m_D}{v} \bar{\nu}_R \phi_c^\dagger \psi_L - \frac{m_R}{2} \nu_R^T C \nu_R + \text{h.c.} \right], \quad (2)$$

with $Y_L = -1$, and the Higgs doublet and its $SU(2)$ conjugate

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \quad \phi_c \equiv i\tau_2 \phi^* = \begin{pmatrix} \bar{\phi}^0 \\ -\phi^- \end{pmatrix}. \quad (3)$$

The operator C stands for the charge conjugation matrix for spinors. The term with m_R in the above Lagrangian is the Majorana mass term for the right-handed neutrino. We tentatively assume that the right-handed Majorana mass vanishes, namely, we adopt the Dirac limit $m_R = 0$ with enhanced lepton number conservation. Physically, this assumption amounts to the ansatz that *all the masses arise from the Higgs boson*, which has been discovered recently. Related schemes of neutrino masses have been discussed by various authors in the past [12–14] (for reviews, see [15]).

Our next observation is that the combination

$$\phi_c^\dagger(x) \psi_L(x) \quad (4)$$

is invariant under the full $SU(2)_L \times U(1)$ gauge symmetry. One may thus add a hermitian non-local Higgs coupling with a real parameter μ to the Lagrangian (2),

$$\mathcal{L}_{CPT}(x) = -i \frac{2\sqrt{2}\mu}{v} \int d^4y \Delta_I(x-y) \theta(x^0 - y^0) \times \{ \bar{\nu}_R(x) (\phi_c^\dagger(y) \psi_L(y)) - (\bar{\psi}_L(y) \phi_c(y)) \nu_R(x) \}, \quad (5)$$

without spoiling the basic Lorentz invariance and the $SU(2)_L \times U(1)$ gauge symmetry. Here we defined

$$\Delta_I(x-y) \equiv \delta((x-y)^2 - l^2) - \delta((x-y)^2 - (l')^2) \quad (6)$$

with l and $l' \neq l$ standing for fixed length scales. This factor differs from $\delta((x-y)^2 - l^2)$ used in [6], and this modified $\Delta_I(x-y)$ avoids a quadratic infrared divergence, as is explained later. In our concrete evaluation below, we set $l' = 0$ for simplicity, although a very small $l' \ll l$ may help to make the separation of the future and past light-cones clear without modifying the essence of our analysis.

In the unitary gauge, $\phi^\pm(x) = 0$ and $\phi^0(x) \rightarrow (v + \varphi(x))/\sqrt{2}$, the neutrino mass term (with $m_R = 0$) becomes in terms of the action

$$\begin{aligned} S_{\nu\text{mass}} &= \int d^4x \left\{ -m_D \bar{\nu}(x) \nu(x) \left(1 + \frac{\varphi(x)}{v} \right) \right. \\ &\quad - i\mu \int d^4y \Delta_I(x-y) \theta(x^0 - y^0) \left[\bar{\nu}(x) \left(1 + \frac{\varphi(y)}{v} \right) \right. \\ &\quad \left. \left. \times (1 - \gamma_5) \nu(y) - \bar{\nu}(y) \left(1 + \frac{\varphi(y)}{v} \right) (1 + \gamma_5) \nu(x) \right] \right\} \\ &= \int d^4x \left\{ -m_D \bar{\nu}(x) \nu(x) \left(1 + \frac{\varphi(x)}{v} \right) \right. \\ &\quad - i\mu \int d^4y \Delta_I(x-y) [\theta(x^0 - y^0) - \theta(y^0 - x^0)] \\ &\quad \times \bar{\nu}(x) \nu(y) + i\mu \int d^4y \Delta_I(x-y) \bar{\nu}(x) \gamma_5 \nu(y) \\ &\quad - i\frac{\mu}{v} \int d^4y \Delta_I(x-y) \theta(x^0 - y^0) \\ &\quad \left. \times [\bar{\nu}(x)(1 - \gamma_5) \nu(y) - \bar{\nu}(y)(1 + \gamma_5) \nu(x)] \varphi(y) \right\}, \quad (7) \end{aligned}$$

where we have changed the naming of integration variables $x \leftrightarrow y$ in some of the terms and used $\theta(x^0 - y^0) + \theta(y^0 - x^0) = 1$. The term

$$-i\mu \int d^4x \int d^4y \Delta_I(x-y) [\theta(x^0 - y^0) - \theta(y^0 - x^0)] \bar{\nu}(x) \nu(y) \quad (8)$$

in the action preserves T but has $C = CP = CPT = -1$ and thus gives rise to particle–antiparticle mass splitting [7].

The equation of motion for the free neutrino is given by

$$\begin{aligned} i\gamma^\mu \partial_\mu \nu(x) &= m_D \nu(x) + i\mu \int d^4y \Delta_I(x-y) \\ &\quad \times [\theta(x^0 - y^0) - \theta(y^0 - x^0)] \nu(y) \\ &\quad - i\mu \int d^4y \Delta_I(x-y) \gamma_5 \nu(y). \quad (9) \end{aligned}$$

By inserting an Ansatz for the solution, $\nu(x) = e^{-ipx} U(p)$, into the equation of motion, we obtain

$$\not{p} U(p) = \{ m + i[f_+(p) - f_-(p)] - ig(p^2) \gamma_5 \} U(p), \quad (10)$$

where $f_\pm(p)$ is a Lorentz invariant quantity defined by

$$f_\pm(p) = \mu \int d^4z e^{\pm ipz} \theta(z^0) [\delta((z)^2 - l^2) - \delta(z^2)]. \quad (11)$$

$f_+(p)$ and $f_-(p)$ are inequivalent for time-like p due to the factor $\theta(z^0)$. The parity violating mass term in (10), which is C and CPT preserving and thus does not contribute to the mass splitting, contains the factor

$$g(p^2) = \mu \int d^4z e^{ipz} [\delta((z)^2 - l^2) - \delta(z^2)]. \quad (12)$$

The factor $f_\pm(p)$ is mathematically related to the two-point Wightman function for a free scalar field. To be explicit,

$$\langle 0 | \phi(x) \phi(y) | 0 \rangle = \int d^4p e^{i(x-y)p} \theta(p^0) \delta(p^2 - m^2), \quad (13)$$

and the Wightman function is finite for $x - y \neq 0$ except for the possible cut in the time-like separation, but divergent for the short distance $x - y \rightarrow 0$. This short distance behavior, whose leading term is mass independent, is related to the infrared $p \rightarrow 0$ behavior of $f_\pm(p)$ and $g(p^2)$. Our modification $\Delta_I(x-y) = \delta((z)^2 - l^2) - \delta(z^2)$ thus eliminates the quadratic infrared divergence, which is independent of l , in $f_\pm(p)$ and $g(p^2)$. Also, our mass splitting term is analogous to the discontinuity along the cut in the time-like separation $x - y$ of Wightman function.

For time-like $p^2 > 0$, one may go to the frame where $\vec{p} = 0$. Then the eigenvalue equation is given by

$$p_0 = \gamma^0 [m_D - f(p_0) - ig(p_0^2) \gamma_5], \quad (14)$$

with

$$\begin{aligned} f(p_0) &\equiv -i[f_+(p_0) - f_-(p_0)] \\ &= 4\mu\pi \int_0^\infty dz \left\{ \frac{z^2 \sin[p_0 \sqrt{z^2 + l^2}]}{\sqrt{z^2 + l^2}} - \frac{z^2 \sin[p_0 \sqrt{z^2}]}{\sqrt{z^2}} \right\} \quad (15) \end{aligned}$$

and

$$g(p_0^2) = 4\mu\pi \int_0^\infty dz \left\{ \frac{z^2 \cos[p_0 \sqrt{z^2 + l^2}]}{\sqrt{z^2 + l^2}} - \frac{z^2 \cos[p_0 \sqrt{z^2}]}{\sqrt{z^2}} \right\}. \quad (16)$$

For space-like $p^2 < 0$, one can confirm that the *CPT* violating term vanishes, $f(p) = 0$, by choosing $p_\mu = (0, \vec{p})$.

Since we are assuming that the *CPT* breaking terms are small, we may solve the mass eigenvalue equations iteratively by assuming that the terms with the parameter μ , whose mass dimension is $[\mu] = [M]^3$, are much smaller than $m = m_D$. We then obtain the mass eigenvalues of the neutrino and antineutrino at [6]

$$m_\pm \simeq m_D - i\gamma_5 g(m_D^2) \pm f(m_D). \quad (17)$$

The parity violating mass $-i\gamma_5 g(m_D^2)$ is now transformed away by a suitable global chiral transformation without modifying the last term in (17) to the order linear in the parameter μ . In this way, the neutrino–antineutrino mass splitting is incorporated in the Standard Model by the Lorentz invariant non-local *CPT* breaking mechanism, without spoiling the $SU(2)_L \times U(1)$ gauge symmetry. The Higgs particle φ itself has a tiny *C*, *CP* and *CPT* violating coupling in (7).

3. Evaluation of mass splitting

We now explicitly evaluate the mass splitting in our model. As already mentioned, for space-like momentum $p^2 < 0$ in (10), the *CPT* violating term vanishes, $f(p) = 0$. We thus consider only the time-like momentum $p^2 > 0$ in the following and use a generic notation $f(p)$.

With a uniform large cut-off L of the spatial variable z and after the change of the integration variable as $y = \sqrt{z^2 + l^2}$ in the first term of (15), we have

$$\begin{aligned} f(p) &= 4\pi\mu \left\{ \int_1^{\sqrt{L^2+l^2}} dy \sqrt{y^2 - l^2} \sin[p_0 y] - \int_0^L dy y \sin[p_0 y] \right\} \\ &= -4\pi\mu [\theta(p_0) - \theta(-p_0)] \theta(p^2) \\ &\quad \times \left\{ \int_{|p_0|}^{\sqrt{L^2+l^2}|p_0|} dv \frac{1}{[\sqrt{v^2 - (l|p_0|)^2} + v]} \sin v \right. \\ &\quad \left. + \int_0^l dy y \sin[|p_0|y] - \int_L^{\sqrt{L^2+l^2}} dy y \sin[|p_0|y] \right\}, \quad (18) \end{aligned}$$

which clearly shows that the limiting value $f(\pm 0) = 0$ for any fixed L . We here took into account the fact that $p^2 \geq 0$. It is confirmed that the last term in (18) gives for large L

$$2\pi\mu l^2 [\theta(p_0) - \theta(-p_0)] \theta(p^2) \sin(|p_0|L), \quad (19)$$

which implies that the convergence of the integral (15) is rather subtle. But it is explained later that this term, which is bounded but oscillates for $L \rightarrow \text{large}$, is set to zero following the customary treatment of the two-point Wightman function in (13). The expression (18) is thus re-written for $|p_0|L \rightarrow \infty$

$$\begin{aligned} f(p) &= -4\pi\mu l^2 [\theta(p_0) - \theta(-p_0)] \theta(p^2) \\ &\quad \times \left\{ \int_1^\infty du \frac{1}{2u(\sqrt{u^2 - 1} + u)^2} \sin(|p_0|lu) \right. \\ &\quad \left. - \frac{1}{2} \int_0^1 du \frac{\sin(|p_0|lu)}{u} + \int_0^1 du u \sin(|p_0|lu) \right\} \end{aligned}$$

$$+ \frac{1}{2} \int_0^\infty du \frac{\sin(u)}{u} \Big\}, \quad (20)$$

by adding and subtracting the term $\frac{1}{2} \int_0^\infty du \frac{\sin(u)}{u}$, and using

$$\int_0^\infty du \frac{\sin(u)}{u} = \int_0^\infty du \frac{\sin(|p_0|lu)}{u}$$

for $|p_0|l \neq 0$. This expression shows that our *CPT* violating term is characterized by the quantity

$$\mu l^2, \quad (21)$$

which has the dimension of mass.

For $|p_0|l \ll 1$ but $|p_0| \neq 0$ and using $\int_0^\infty du \frac{\sin(u)}{u} = \pi/2$, we have

$$f(p) \simeq -\pi^2 \mu l^2 [\theta(p_0) - \theta(-p_0)] \theta(p^2), \quad (22)$$

which is Lorentz invariant. Note that the terms linear in $|p_0|l$ cancel out exactly although they do not appear to do so at first glance, and no infrared divergence appears in the *CPT* violating term. (Incidentally, if one chooses the non-local factor with an additional parameter l' in (6), the factor $\pi^2 \mu l^2$ is replaced by $\pi^2 \mu (l^2 - (l')^2)$ in (22).) Thus the mass gap in (17) is given by

$$\Delta m \simeq 2\pi^2 \mu l^2. \quad (23)$$

We now briefly explain the prescription $\sin(p_0 L) = 0$ for $L \rightarrow \infty$ in (19), which is natural in the sense of a distribution

$$\int dp_0 \sin(p_0 L) F(p_0) = 0,$$

for any test function $F(p_0)$, by the Riemann–Lebesgue lemma. As we already mentioned, our non-local factor in (11) is identical to a difference of two two-point Wightman functions in (13) provided that the coordinates and momenta are interchanged. Our factor $\sin(p_0 L)$ appears in the procedure that corresponds to the evaluation (or definition) of the two-point Wightman function (and related propagators) for fixed finite $x - y$, for which one does not usually mention such a subtlety; of course, the Wightman function contains the short distance singularities for $x - y \rightarrow 0$ corresponding to the infrared singularities in our non-local factors, which have no connection with (19) that vanishes for $p_0 = 0$. This disposal of (19) corresponds to the treatment of Wightman function as a distribution, as we illustrated above, or one may simply set the bounded infinitely-oscillating term to zero. Besides, the propagator as such is defined by its action on well-defined initial states in momentum space and thus its treatment as a distribution is justified. In fact, in our actual lowest order applications below we analyze the propagator between well-defined initial and final states in momentum space in the Standard Model.

Our definition of the non-local factor in (6) thus eliminates the infrared singularity in the *CPT* violating term. Our *CPT* violating term $f(p_0)$ is odd in p_0 and $f(\pm 0) = \mp \Delta m/2$ but $f(0) = 0$. The formula for the mass gap is defined for $|p_0|L \rightarrow \infty$ and thus precise $p_0 = 0$ is avoided.

As for the parity violating mass term in (10), we have (in the frame with $\vec{p} = 0$ for $p^2 > 0$)

$$\begin{aligned} g(p^2) &= \mu \int d^4 z e^{ipz} [\delta((z)^2 - l^2) - \delta((z)^2)] \\ &= 4\pi\mu \left\{ \int_1^\infty dy \sqrt{y^2 - l^2} \cos[p_0 y] - \int_0^\infty dy y \cos[p_0 y] \right\}, \quad (24) \end{aligned}$$

which is evaluated as

$$g(p^2) = -4\pi\mu l^2 \left\{ \int_1^\infty du \frac{1}{\sqrt{u^2-1}+u} \cos[p_0 l u] + \frac{\sin[p_0 l]}{p_0 l} + \frac{\cos[p_0 l] - 1}{(p_0 l)^2} \right\}. \quad (25)$$

This result (25) also potentially contains a subtlety as in (19), which is safely set to zero as already explained, and a mild logarithmic (or weaker) divergence at $p_0 = 0$ in the first term. This formula is again well-defined if precise $p_0 = 0$ is excluded.

4. Neutrino–antineutrino mass splitting and baryogenesis

The Lorentz invariant non-local factor

$$[\theta(x^0 - y^0) - \theta(y^0 - x^0)]\delta((x - y)^2 - l^2) \quad (26)$$

used in [6,7] induces infinite non-locality along the light-cone, and a dimensional counting shows that it diverges quadratically in the infrared in momentum space. This infinite non-locality may lead to a severe breaking of unitarity. In contrast, our modified Lorentz invariant non-local factor

$$[\theta(x^0 - y^0) - \theta(y^0 - x^0)] \times [\delta((x - y)^2 - l^2) - \delta((x - y)^2 - (l')^2)], \quad (27)$$

(with $l' = 0$ in practical applications) mostly cancels out the infinite time-like volume effect and eliminates the quadratic infrared divergence completely. In effect, our modified non-local factor induces non-locality which is limited within the fluctuation around the tip of the light-cone characterized by the length scale of the parameter l . This is very welcome from the point of view of unitarity, since one can choose the length scale l at the order of the Planck length and thus at least avoid the issue of unitarity violation in the framework of a generic effective theory, which may be valid at a length scale much larger than the Planck scale.

It is important to analyze if our mass splitting can induce a physically meaningful size of neutrino–antineutrino mass splitting by choosing l of the order of Planck length and $\mu = M^3$ suitably, where M is another mass scale of possible new physics. The natural neutrino mass splitting given by (23) is then

$$2\pi^2\mu l^2 = 2\pi^2 M(M/M_P)^2, \quad (28)$$

which is a kind of see-saw between M and the Planck mass M_P . This neutrino mass splitting may also be regarded as a gravitational effect because of the appearance of the Newton constant $G_N = 1/M_P^2$. If one chooses $M \sim 10^9$ GeV, the mass splitting becomes of the order of the observed neutrino mass (difference) ~ 0.1 eV [16]. This size of the mass scale M is not uncommon in the leptogenesis based on the see-saw scenario [17], for example.

If one assumes $M \sim 1000$ GeV, namely, the scale of the Standard Model, the mass splitting is $\sim 10^{-20}$ eV, which is too small for phenomenological interest. Although our model is for the neutrinos, the presently known experimental limit on the electron–positron mass splitting is [16]

$$\Delta m_e \leq 10^{-8} m_e \sim 10^{-2} \text{ eV}. \quad (29)$$

To generate the corresponding value $\Delta m \sim 10^{-2}$ eV for the neutrinos in our scheme, we need a value slightly smaller than $M \sim 10^9$ GeV.

The neutrino mass splitting $\Delta m = 10^{-1} \sim 10^{-2}$ eV, which is intended to be of the order of $m_D/5$, is generated by $M \simeq$

$10^8 \sim 10^9$ GeV and appears to be allowed by presently available experimental data [3,18]. Our Lorentz invariant model, which is written for the electron but applicable to other leptons also, is concerned with neutrinos so far, but the higher order effect in renormalizable theory is generally expected to give rise to an electron–positron mass splitting of the order $\Delta m_e \sim \alpha \Delta m$, with α the fine structure constant. This induced mass splitting, which is substantially smaller than $\Delta m = 10^{-1} \sim 10^{-2}$ eV, is expected for all the massive charged leptons in the Standard Model. We, however, show in the next section that the induced effect is actually much smaller, $\sim 10^{-20}$ eV. Thus the value $M \simeq 10^8 \sim 10^9$ GeV is interesting for a phenomenological purpose, which does not apparently contradict any experimental data and yet may be measurable in neutrino oscillation experiments in the near future. We tentatively adopt Δm at $10^{-1} \sim 10^{-2}$ eV, which is similar to the magnitude of m_D , in phenomenological discussions.

As for the baryogenesis, it is believed that leptons acquire masses from electroweak symmetry breaking during the electroweak phase transition in the early universe. A neutrino–antineutrino mass difference would then result in a leptonic matter–antimatter asymmetry proportional to the mass difference. This asymmetry is transmitted to the baryon sector through the chiral anomaly [19] and sphaleron processes [20] which preserve $B - L$ but violate $B + L$. This “kinematical” picture implies the asymmetry in the neutrino and antineutrino of the order [1] $(n_\nu - n_{\bar{\nu}})/n_\nu \simeq m_D \Delta m / T^2$, which is, however, too small at the electroweak energy scale to generate the baryon asymmetry via sphaleron processes in our case with $\Delta m = 10^{-1} \sim 10^{-2}$ eV. Besides, this initial asymmetry requires the lepton number non-conservation [1], while the lepton number is conserved in our model without sphaleron effects.

Thus the lepton and quark sectors need to be treated simultaneously in the presence of sphalerons [20] which break $B + L$ by preserving $B - L$. The authors of Ref. [2] discuss a rather elaborate sphaleron dynamics by referring to [21–24], and they conclude that the final baryon number at the energy scale of the weak mass M_W is estimated at

$$\frac{n_B}{n_\gamma} \sim \frac{\Delta m}{M_W}, \quad (30)$$

where n_γ stands for the photon number density. This estimate in the present case with $\Delta m = 10^{-1} \sim 10^{-2}$ eV, namely $n_B/n_\gamma \sim 10^{-12} \sim 10^{-13}$, is smaller than the observed value $n_B/n_\gamma \simeq 10^{-10}$, but it still gives a promising number by considering the crude estimate in our model. The estimate of the generated baryon number is mainly constrained by experimental bounds on neutrino mass differences.

We emphasize that *this equilibrium electroweak baryogenesis does not need CP violation other than for the purpose of producing neutrino–antineutrino mass splitting*. The mechanism to generate (30) by a specific sphaleron dynamics [2], which is very interesting but requires further elaboration, differs from the more conventional baryogenesis [25,26] and also from the leptogenesis [17].

5. Higher order induced effects

We now analyze if the higher order effect due to the neutrino mass splitting is well-controlled in the Standard Model. The propagator of the neutrino in the path integral formulation on the basis of Schwinger’s action principle, which is based on the equations of motion, is given by [7] (see also [27]),

$$\langle T^* \nu(x) \bar{\nu}(y) \rangle = \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)}$$




Fig. 1. (a) Represents the W -boson contribution and (b) represents the charged unphysical Nambu–Goldstone ϕ -boson contribution to the induced mass splitting between electron and positron.

$$\times \frac{i}{\not{p} - m_D + i\epsilon + i\gamma_5 g(p^2) - i[f_+(p) - f_-(p)]}, \quad (31)$$

where $f_{\pm}(p)$ and $g(p^2)$ are defined in (11) and (12), respectively, in connection with the free equation of motion (10).

We wish to examine the large momentum behavior of $f(p) = -i[f_+(p) - f_-(p)]$ defined in (20) by assuming that l is not very small and thus large $|p_0|l$ is physically relevant. By noting that $\frac{1}{2} \int_0^1 du \frac{\sin(|p_0|lu)}{u} \rightarrow \frac{1}{2} \int_0^\infty du \frac{\sin(u)}{u}$ for $|p_0|l \rightarrow \infty$, the expression (20) becomes

$$f(p) = -4\pi \mu l^2 [\theta(p_0) - \theta(-p_0)] \theta(p^2) \times \left\{ \int_1^\infty du \frac{1}{2u(\sqrt{u^2 - 1} + u)^2} \sin(|p_0|lu) + \frac{\sin(|p_0|l)}{(|p_0|l)^2} - \frac{\cos(|p_0|l)}{(|p_0|l)} \right\}. \quad (32)$$

The integral in this expression is shown to approach zero for $|p_0|l \rightarrow \infty$ by the Riemann–Lebesgue lemma, and thus $f(p)$ approaches zero for $|p_0|l \rightarrow \infty$.

The high energy behavior of the parity violating term $g(p^2)$ in (25) is shown to be the same as the above CPT violating term.

The propagator (31) for Minkowski momentum is thus well behaved and the effects of non-locality are mild and limited and, in this sense, T^* product may even be replaced by the canonical T product in (31) [27]. (The logarithmic singularity in $g(p^2)$ at $p_\mu = 0$ may not be serious in the propagator (31) due to the presence of the phase space volume factor d^4p in the numerator.) In the analysis of the renormalization procedure, however, it is customary to consider the Euclidean amplitude obtained from the Minkowski amplitude by Wick rotation. This is because the power counting rule (superficial degree of divergence) of each Feynman amplitude is well defined with Euclidean momenta. Our propagator, which contains trigonometric functions, has undesirable behavior under the Wick rotation such as $\sin p_0 z \rightarrow i \sinh p_4 z$, and an exponentially divergent behavior is generally induced and the effects of non-locality become significant. One might still argue that higher order effects in field theory defined on Minkowski space are in principle analyzed in Minkowski space and, if that is the case, our propagator maintains the ordinary renormalizable behavior. In passing, we mention that an alternative scheme of CPT violation, which breaks Lorentz symmetry but preserves locality, suffers from numerous difficulties related to unitarity and other issues once one attempts to construct physically interesting models with particle–antiparticle mass splitting (i.e., C and CP violations as well) [28].

One may assume that l is of the order of Planck scale, as we did in our phenomenological analysis. Then the higher order effect is expected to be small as long as μl^2 is small in either Minkowski or Euclidean formulation, since it is natural to assume that the loop momenta are cut-off below the Planck scale in generic effective theory and thus $|p_0|l$ may always be assumed to be small even inside the loop diagrams. In this case, our Lorentz invariant CPT violating term is effectively replaced by

$$f(p) = -\pi^2 \mu l^2 [\theta(p_0) - \theta(-p_0)] \theta(p^2), \quad (33)$$

which is similar to a constant mass term except for the CPT violating factor $[\theta(p_0) - \theta(-p_0)] \theta(p^2)$. When this term is inserted into Feynman diagrams in the Standard Model, those Feynman diagrams are expected to show ordinary high energy behavior for a mass insertion, if the naive power counting works.

5.1. Induced electron–positron mass splitting

We now wish to show that the electron–positron mass splitting induced by the above factor $f(p)$ in (33), when inserted into one-loop self-energy diagrams of the electron in the Standard Model, is well controlled. This correction amounts to the replacement of the ordinary momentum space neutrino propagator by

$$\tilde{S}_F(p)_\nu = \frac{i}{\not{p} - m_D + i\epsilon} f(p) \frac{i}{\not{p} - m_D + i\epsilon} \quad (34)$$

in the one-loop self-energy diagrams of the electron in the Standard Model. We show that the electron self-energy correction induced by one-loop W -boson is well convergent, which implies that the momentum flowing through the neutrino propagator is limited by the W -boson mass scale M_W and thus our starting assumption $|p|l \ll 1$ is justified.

We work in the 't Hooft–Feynman gauge ($\xi = 1$ in R_ξ -gauge [29] with Feynman rules given there). Then we have contributions from the two Feynman diagrams shown in Fig. 1, with ϕ standing for the charged unphysical Nambu–Goldstone boson. The W -boson term in Fig. 1a is then given by (ignoring precise numerical factors)

$$g^2 \int \frac{d^4 p}{(2\pi)^4} \left[\gamma^\alpha \frac{(1 - \gamma_5)}{2} \frac{\not{p} + m_D}{p^2 - m_D^2 + i\epsilon} f(p) \frac{\not{p} + m_D}{p^2 - m_D^2 + i\epsilon} \times \gamma^\alpha \frac{(1 - \gamma_5)}{2} \right] \frac{1}{(k - p)^2 - M_W^2 + i\epsilon}, \quad (35)$$

where the numerator is proportional to $2m_D \not{k} f(p) [(1 - \gamma_5)/2]$, and the integral is linearly convergent. This term is proportional to Dirac γ^μ and contributes to the electron kinetic term of the order

$$\alpha [m_D \not{k} / M_W^2] (\mu l^2) [\theta(k^0) - \theta(-k^0)] \theta(k^2) [(1 - \gamma_5)/2],$$

where α is the fine structure constant. To infer this result, it is easier to analyze (35) with $2m_D(kp)f(p)$ in the numerator by replacing γ^μ with k^μ . It is then confirmed that the amplitude is invariant under the change of the signature of \vec{k} but changes the signature under the change of the signature of k^0 of the external electron at its rest frame $\vec{k} = 0$. It vanishes for space-like k^μ , as is seen by choosing the frame with $k^0 \rightarrow \pm 0$, and it is Lorentz invariant. We thus infer

$$\sim \alpha [m_D k^2 / M_W^2] (\mu l^2) [\theta(k^0) - \theta(-k^0)] \theta(k^2)$$

for (35) with $2m_D(kp)f(p)$ in the numerator. One may finally replace k^2 by \not{k} .

The charged ϕ -boson term in Fig. 1b is given by (ignoring precise numerical factors)

$$\begin{aligned} & \frac{g^2}{M_W^2} \int \frac{d^4 p}{(2\pi)^4} \left[m_e \frac{(1-\gamma_5)}{2} - \frac{(1+\gamma_5)}{2} m_D \right] \\ & \times \frac{\not{p} + m_D}{p^2 - m_D^2 + i\epsilon} f(p) \frac{\not{p} + m_D}{p^2 - m_D^2 + i\epsilon} \\ & \times \left[m_D \frac{(1-\gamma_5)}{2} - \frac{(1+\gamma_5)}{2} m_e \right] \frac{1}{(k-p)^2 - M_W^2 + i\epsilon}, \quad (36) \end{aligned}$$

where the deviation from the Standard Model result is given by the Lagrangian (2). The numerator contains two class of terms: the first class has the structure

$$\left[(m_e^2(1-\gamma_5)/2 + m_D^2(1+\gamma_5)/2) / M_W^2 \right] 2m_D \not{p} f(p),$$

which has a linearly convergent behavior as in the case of W -boson contribution and gives rise to a contribution of the order

$$\alpha(1-\gamma_5)/2 \left[m_e^2/M_W^2 \right] \left[m_D \not{k} / M_W^2 \right] (\mu^2) [\theta(k^0) - \theta(-k^0)] \theta(k^2).$$

The second class of terms contains in the numerator a factor $(m_e m_D / M_W^2)(p^2 + m_D^2) f(p)$ which potentially gives rise to a logarithmic divergence. If this divergence should persist, we need a new CPT violating counter term for the electron, which would spoil the renormalizability of the Standard Model. But this term is in fact convergent due to the structure of the CPT violating factor $[\theta(p_0) - \theta(-p_0)]\theta(p^2)$, as

$$\begin{aligned} & \frac{g^2}{M_W^2} \int \frac{d^3 p}{(2\pi)^4} \int_0^\infty dp_0 \frac{(m_e m_D)(p^2 + m_D^2)}{(p^2 - m_D^2 + i\epsilon)^2} \theta(p^2) (\mu^2) \\ & \times \left[\frac{1}{(k-p)^2 - M_W^2 + i\epsilon} - \frac{1}{(k+p)^2 - M_W^2 + i\epsilon} \right]. \quad (37) \end{aligned}$$

This is linearly convergent, and it is independent of the Dirac γ^μ , thus contributing to the mass correction. Remark that this term changes its signature under the change of the signature of k^0 of the external electron at its rest frame $k=0$ and vanishes for the space-like k^μ as is seen by choosing the frame with $k^0 \rightarrow \pm 0$. This term also vanishes at $k^\mu = 0$, and it is estimated at the order of

$$\alpha(m_e m_D / M_W^2) (k^2 / M_W^2) (\mu^2) [\theta(k^0) - \theta(-k^0)] \theta(k^2),$$

which is

$$\alpha(m_e m_D / M_W^2) (m_e^2 / M_W^2) (\mu^2) [\theta(k^0) - \theta(-k^0)] \theta(k^2)$$

near on-shell.

The induced effect on the electron is determined by the eigenvalue equation (by ignoring ordinary corrections in the Standard Model, for simplicity)

$$\not{k}(1 + \alpha A(k)) - (m_e + \alpha B(k)) = 0 \quad (38)$$

or equivalently $\not{k} = m_e + \alpha B(k) - m_e \alpha A(k)$, where $A(k)$ and $B(k)$ stand for the induced effects evaluated above. The induced CPT violating effect on the electron-positron splitting is thus finite and the leading contribution is given by the W -boson at the order,

$$\alpha \left[m_D m_e / M_W^2 \right] (\mu^2) \left[(1-\gamma_5)/2 \right] [\theta(k^0) - \theta(-k^0)] \theta(k^2), \quad (39)$$

which, if we choose our parameter at $\pi^2 \mu^2 = 10^{-1} \sim 10^{-2}$ eV as in the previous section, is $\sim 10^{-20}$ eV and thus well below the present experimental bound. The induced CPT violation is expected to be smaller in the quark sector (as a two-loop effect) than in the charged leptons in the $SU(2) \times U(1)$ invariant theory, and thus much smaller than the well-known limit on the K -meson, $|m_K - m_{\bar{K}}| < 0.44 \times 10^{-18}$ GeV [16].

6. Conclusion

We have demonstrated that a well-defined neutrino-antineutrino mass splitting can be realized in the Standard Model. It has potentially interesting implications on equilibrium electroweak baryogenesis, depending on the subtle details of sphaleron dynamics. The Lorentz invariant CPT breaking mass term in momentum space is effectively represented by $f(p) = -(\Delta m/2)[\theta(p_0) - \theta(-p_0)]\theta(p^2)$. The induced CPT violating effect on the electron-positron mass splitting in the Standard Model is shown to be finite and kept well below the experimental bound. Our analysis suggests that non-local Lorentz invariant CPT violation is natural in the Standard Model and it would be interesting to search for possible neutrino-antineutrino mass splitting in oscillation experiments.

There are some fundamental issues to be mentioned in the present scheme of CPT violation. We have shown that the Standard Model treated as an effective theory non-local in the Planck scale can generate a sizable neutrino-antineutrino mass splitting which can be tested by experiment. We have then shown that the CPT violating term, when treated in lowest order perturbation, gives rise to sensible and interesting predictions for the induced mass splitting of charged fermions. The induced electron-positron mass splitting, for example, is too small to be detected, and in this sense our scheme of neutrino mass splitting is consistent. Our analysis suggests that the CPT violating term treated as an interaction term defined by (33) gives rise to sensible higher order effects, and perturbative unitarity with the original particle spectrum (without the CPT breaking term) is defined. However, our theory which is non-local in time lacks a canonical formulation. When one sums the diagrams with the CPT violating term (33) inserted and defines a theory with particle-antiparticle mass splitting, unitarity and the structure of the vacua in the conventional sense become less obvious. The non-locality in the Planck scale defers the basic issues to some extent but does not solve them. To resolve those issues completely, one may presumably need to go beyond the existing framework of field theory. A physical origin of our non-local CPT violation at the Planck scale, which is most likely related to quantum gravity, also remains to be clarified.

Acknowledgements

We thank M. Chaichian for stimulating discussions. We also thank R. Bufalo for drawing the figures. The support of the Academy of Finland under the Projects Nos. 136539 and 272919, of the Vilho, Yrjö and Kalle Väisälä Foundation, and of JSPS KAKENHI (Grant No. 25400415) is gratefully acknowledged.

References

- [1] A.D. Dolgov, Ya.B. Zeldovich, *Rev. Mod. Phys.* 53 (1981) 1.
- [2] G. Barenboim, L. Borissov, J.D. Lykken, A.Y. Smirnov, *J. High Energy Phys.* 0210 (2002) 001, arXiv:hep-ph/0108199.
- [3] V.D. Barger, S. Pakvasa, T.J. Weiler, K. Whisnant, *Phys. Rev. Lett.* 85 (2000) 5055, arXiv:hep-ph/0005197.
- [4] H. Murayama, T. Yanagida, *Phys. Lett. B* 520 (2001) 263, arXiv:hep-ph/0010178; G. Barenboim, L. Borissov, J. Lykken, *Phys. Lett. B* 534 (2002) 106, arXiv:hep-ph/0201080; S.M. Bilenky, M. Freund, M. Lindner, T. Ohlsson, W. Winter, *Phys. Rev. D* 65 (2002) 073024, arXiv:hep-ph/0112226; G. Barenboim, J.D. Lykken, *Phys. Rev. D* 80 (2009) 113008, arXiv:0908.2993 [hep-ph].
- [5] P. Adamson, et al., MINOS Collaboration, *Phys. Rev. D* 85 (2012) 031101, arXiv:1201.2631 [hep-ex].
- [6] M. Chaichian, K. Fujikawa, A. Tureanu, *Phys. Lett. B* 718 (2012) 178, arXiv:1208.5508 [hep-ph].
- [7] M. Chaichian, K. Fujikawa, A. Tureanu, *Phys. Lett. B* 712 (2012) 115, arXiv:1203.0267 [hep-th]; M. Chaichian, K. Fujikawa, A. Tureanu, *Phys. Lett. B* 718 (2013) 1500, arXiv:1210.0208 [hep-th];

- M. Chaichian, K. Fujikawa, A. Tureanu, *Eur. Phys. J. C* 73 (2013) 2349, arXiv:1205.0152 [hep-th].
- [8] W. Pauli, in: W. Pauli (Ed.), *Niels Bohr and the Development of Physics*, Pergamon Press, New York, 1955;
G. Lüders, *Mat. Fys. Medd. Dan. Vid. Selsk.* 28 (5) (1954) 1.
- [9] M. Chaichian, A.D. Dolgov, V.A. Novikov, A. Tureanu, *Phys. Lett. B* 699 (2011) 177, arXiv:1103.0168 [hep-th].
- [10] M. Duetsch, J.M. Gracia-Bondia, *Phys. Lett. B* 711 (2012) 428, arXiv:1204.2654 [hep-th].
- [11] S. Weinberg, *Phys. Rev. Lett.* 19 (1967) 1264;
A. Salam, in: N. Svartholm (Ed.), *Elementary Particle Theory*, Stockholm, 1968, p. 367.
- [12] L. Wolfenstein, *Nucl. Phys. B* 186 (1981) 147;
S.T. Petcov, *Phys. Lett. B* 110 (1982) 245;
S.M. Bilenky, B. Pontecorvo, *Sov. J. Nucl. Phys.* 38 (1983) 248;
S.M. Bilenky, S.T. Petcov, *Rev. Mod. Phys.* 59 (1987) 671;
C. Giunti, C.W. Kim, U.W. Lee, *Phys. Rev. D* 46 (1992) 3034, arXiv:hep-ph/9205214;
J.P. Bowes, R.R. Volkas, *J. Phys. G* 24 (1998) 1249, arXiv:hep-ph/9804310;
M. Kobayashi, C.S. Lim, *Phys. Rev. D* 64 (2001) 013003, arXiv:hep-ph/0012266;
K. Fujikawa, *Prog. Theor. Phys.* 113 (2005) 1065, arXiv:hep-ph/0407331.
- [13] P. Langacker, *Phys. Rev. D* 58 (1998) 093017, arXiv:hep-ph/9805281;
G. Cleaver, M. Cvetič, J.R. Espinosa, L. Everett, P. Langacker, *Phys. Rev. D* 57 (1998) 2701, arXiv:hep-ph/9705391;
D. Chang, O.C. Kong, *Phys. Lett. B* 477 (2000) 416, arXiv:hep-ph/9912268.
- [14] K.R. Dienes, E. Dudas, T. Gherghetta, *Nucl. Phys. B* 557 (1999) 25, arXiv:hep-ph/9811428;
G.R. Dvali, A.Y. Smirnov, *Nucl. Phys. B* 563 (1999) 63, arXiv:hep-ph/9904211;
N. Arkani-Hamed, S. Dimopoulos, G.R. Dvali, J. March-Russel, *Phys. Rev. D* 65 (2002) 024032, arXiv:hep-ph/9811448;
Z. Berezhiani, L. Bento, *Phys. Rev. Lett.* 96 (2006) 081801, arXiv:hep-ph/0507031.
- [15] S.M. Bilenky, C. Giunti, W. Grimus, *Prog. Part. Nucl. Phys.* 43 (1999) 1, arXiv:hep-th/9812360;
R.N. Mohapatra, A.Y. Smirnov, *Annu. Rev. Nucl. Part. Sci.* 56 (2006) 569, arXiv:hep-ph/0603118.
- [16] J. Beringer, et al., *Particle Data Group*, *Phys. Rev. D* 86 (2012) 010001.
- [17] M. Fukugita, T. Yanagida, *Phys. Lett. B* 174 (1986) 45;
G.F. Giudice, A. Notari, M. Raidal, A. Riotto, A. Strumia, *Nucl. Phys. B* 685 (2004) 89, arXiv:hep-ph/0310123;
W. Buchmüller, R.D. Peccei, T. Yanagida, *Annu. Rev. Nucl. Part. Sci.* 55 (2005) 311, arXiv:hep-ph/0502169.
- [18] P. Adamson, et al., MINOS Collaboration, *Phys. Rev. Lett.* 108 (2012) 191801.
- [19] G. 't Hooft, *Phys. Rev. Lett.* 37 (1976) 8.
- [20] N. Manton, *Phys. Rev. D* 28 (1983) 2019;
F. Klinkhamer, N. Manton, *Phys. Rev. D* 30 (1984) 2212;
V. Kuzmin, V. Rubakov, M. Shaposhnikov, *Phys. Lett. B* 155 (1985) 36.
- [21] V.A. Rubakov, M.E. Shaposhnikov, *Phys. Usp.* 39 (1996) 461, arXiv:hep-ph/9603208.
- [22] J. Boguta, J. Kunz, *Phys. Lett. B* 154 (1985) 407.
- [23] A. Ringwald, *Phys. Lett. B* 213 (1988) 61;
M. Axenides, A. Johansen, H.B. Nielsen, *Nucl. Phys. B* 414 (1994) 53, arXiv:hep-ph/9308299.
- [24] M. Dine, O. Lechtenfeld, B. Sakita, W. Fischler, J. Polchinski, *Nucl. Phys. B* 342 (1990) 381.
- [25] A.D. Sakharov, *JETP Lett.* 5 (1967) 24.
- [26] M. Yoshimura, *Phys. Rev. Lett.* 41 (1978) 281;
A.Yu. Ignatiev, N.V. Krasnikov, V.A. Kuzmin, A.N. Tavkhelidze, *Phys. Lett. B* 76 (1978) 436.
- [27] K. Fujikawa, *Phys. Rev. D* 70 (2004) 085006, arXiv:hep-th/0406128;
As for the Bjorken–Johnson–Low method, see also Appendix in K. Fujikawa, P. van Nieuwenhuizen, *Ann. Phys.* 308 (2003) 78, arXiv:hep-th/0305144.
- [28] O.M. Del Cima, D.H.T. Franco, A.H. Gomes, J.M. Fonseca, O. Piguet, *Phys. Rev. D* 85 (2012) 065023, arXiv:1202.0009 [hep-th].
- [29] K. Fujikawa, B.W. Lee, A.I. Sanda, *Phys. Rev. D* 6 (1972) 2923.