

Review

A hitchhiker's guide to European lake ecological assessment and intercalibration



Sandra Poikane^{a,*}, Sebastian Birk^b, Jürgen Böhmer^c, Laurence Carvalho^d,
 Caridad de Hoyos^e, Hubert Gassner^f, Seppo Hellsten^g, Martyn Kelly^h,
 Anne Lyche Solheimⁱ, Mikko Olin^j, Karin Pall^k, Geoff Phillips^l, Rob Portielje^m,
 David Ritterbuschⁿ, Leonard Sandin^o, Ann-Kristin Schartau^p, Angelo G. Solimini^r,
 Marcel van den Berg^m, Georg Wolfram^s, Wouter van de Bund^a

^a European Commission, Joint Research Centre, Institute for Environment and Sustainability, Ispra, Italy

^b Aquatic Ecology, Faculty of Biology, University of Duisburg-Essen, Universitätsstraße 5, 45141 Essen, Germany

^c Bioforum GmbH, Sudetenstr. 34, DE-73230 Kirchheim/Teck, Germany

^d Centre for Ecology & Hydrology, Bush Estate, Penicuik EH26 0QB, UK

^e Centre for Hydrographic Studies, CEDEX, C/ Paseo Bajo de la Virgen del Puerto 3, Madrid 28005, Spain

^f Federal Agency of Water Management, Institute of Freshwater Ecology, Fisheries Management and Lake Research, Scharfling 18, 5310 Mondsee, Austria

^g SYKE, University of Oulu, P.O. Box 413, 90014 Oulu, Finland

^h Bowburn Consultancy, 11 Montaigne Drive, Bowburn, Durham DH6 5QB, UK

ⁱ Norwegian Institute for Water Research, Gaustadalléen 21, Oslo N-0349, Norway

^j Department Environmental Sciences, University of Helsinki, P.O. Box 65, 00014 Helsinki, Finland

^k Systema GmbH, Environment Agency, Bensasteig 8, 1140 Vienna, Austria

^l Environment Agency, Kings Meadow Road, Reading RG1 8DQ, UK

^m Centre for Water Management, Zuiderwagenplein 2, Lelystad, NL-8200 AA, The Netherlands

ⁿ Institute of Inland Fisheries, Im Königswald 2, 14469 Potsdam, Germany

^o Department of Bioscience, Aarhus University, Vejlsovej 25, P.O. Box 314, DK-8600, Silkeborg, Denmark

^p Norwegian Institute for Nature Research, Gaustadalléen 21, Oslo 0349, Norway

^r Department of Public Health and Infectious Diseases, Sapienza University of Rome, 00185 Rome, Italy

^s DWS Hydro-Ökologie GmbH, Zentagasse 47, Vienna A-105, Austria

ARTICLE INFO

Article history:

Received 17 August 2014

Received in revised form

10 November 2014

Accepted 3 January 2015

Keywords:

Benthic invertebrates

Ecological assessment

Europe

Fish fauna

Lakes

Macrophytes

Phytoplankton

Water Framework Directive

ABSTRACT

The Water Framework Directive is the first international legislation to require European countries to establish comparable ecological assessment schemes for their freshwaters. A key element in harmonising quality classification within and between Europe's river basins is an "Intercalibration" exercise, stipulated by the WFD, to ensure that the good status boundaries in all of the biological assessment methods correspond to similar levels of anthropogenic pressure. In this article, we provide a comprehensive overview of this international comparison, focusing on the assessment schemes developed for freshwater lakes. Out of 82 lake ecological assessment methods reported for the comparison, 62 were successfully intercalibrated and included in the EC Decision on intercalibration, with a high proportion of phytoplankton (18), macrophyte (17) and benthic fauna (13) assessment methods. All the lake assessment methods are reviewed in this article, including the results of intercalibration. Furthermore, the current gaps and way forward to reach consistent management objectives for European lakes are discussed.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

| | |
|---------------------------------------|-----|
| 1. Introduction | 534 |
| 2. Intercalibration methodology | 534 |

* Corresponding author. Tel.: +39 0332 789720.

E-mail address: sandra.poikane@jrc.ec.europa.eu (S. Poikane).

| | | |
|------|--|-----|
| 3. | Lake assessment methods..... | 538 |
| 3.1. | Lake assessment methods: phytoplankton..... | 538 |
| 3.2. | Lake assessment methods: macrophytes..... | 538 |
| 3.3. | Lake assessment methods: phytobenthos..... | 538 |
| 3.4. | Lake assessment methods: benthic invertebrates..... | 538 |
| 3.5. | Lake assessment methods: fish fauna..... | 539 |
| 4. | Lake intercalibration..... | 539 |
| 4.1. | Intercalibration groups..... | 539 |
| 4.2. | Common metrics and benchmarking..... | 539 |
| 4.3. | Boundary harmonisation..... | 539 |
| 5. | Intercalibration gaps..... | 540 |
| 5.1. | Intercalibration gaps: fish fauna, phytobenthos and benthic invertebrates..... | 540 |
| 5.2. | Eastern Continental and Mediterranean regions..... | 540 |
| 5.3. | Anthropogenic pressures addressed by lake assessment methods..... | 541 |
| 5.4. | Heavily modified and artificial water bodies..... | 541 |
| 5.5. | Detection of cyanobacteria blooms..... | 542 |
| 5.6. | Uncertainty..... | 542 |
| 6. | Conclusions..... | 542 |
| | Appendix A. Supplementary data..... | 542 |
| | References..... | 542 |

1. Introduction

Many benefits provided by aquatic ecosystems can only be maintained if the ecosystems are protected from deterioration (Millennium Ecosystem Assessment, 2005; de Groot et al., 2010). This aim requires (1) suitable methods to assess anthropogenic impact on aquatic ecosystems and to evaluate ecological integrity, (2) common management objectives across state boundaries and administrative barriers, and (3) concerted action aimed at halting and reversing degradation on the national and international level (Palmer et al., 2005; Hering et al., 2013).

Many countries have adopted legislation to determine the ecological integrity of surface waters including streams, rivers, lakes, estuaries and coastal waters. The purpose of the US Clean Water Act (CWA) is to “restore and maintain the chemical, physical, and biological integrity of the Nation’s waters.” Also in Australia, a broader, more holistic approach to aquatic ecosystem management is adapted “to maintain and enhance the ecological integrity of freshwater and marine ecosystems” (ANZECC, 2000). Similarly, the South African National Water Act aims at “protecting aquatic and associated ecosystems and their biological diversity”. Still, in many cases, these legislation acts have not fulfilled their ambitions (Doremus and Dan Tarlock, 2013; Adler, 2013), mainly due to a lack of clear guidelines for the assessment of biological integrity (Davies and Jackson, 2006), the insufficient development and quality of bioassessment methods (Adler, 2003; Yoder and Barbour, 2009), a lack of consistent management objectives (Davies and Jackson, 2006; Adler, 2013), and poor comparability of biological data (Cao and Hawkins, 2011; Diamond et al., 2012).

In Europe, the Water Framework Directive (EC, 2000; WFD) establishes a framework for the protection and improvement of inland and coastal waters, which aims to achieve ‘good’ surface water status by 2015 or, at the latest, by 2027. In contrast to other legislations, the WFD provides operational definitions for assessing ecological status, setting management objectives, and harmonising EU Member States’ ecological assessment systems. In short, the WFD is based on the following main principles:

- Biological assessment uses numerical measurements of communities of plants and animals (phytoplankton, aquatic flora, benthic invertebrates and fish fauna) as stipulated in the Directive (e.g., biomass, taxonomic composition, diversity, etc.).
- In biological assessment, the observed condition is compared with the reference status with the result given in five classes:

‘high’ status (no differences to reference conditions), ‘good’ status (slight differences), ‘moderate’ status (moderate differences), ‘poor’ and ‘bad’ statuses (major differences).

- ‘Good’ ecological status represents the target value that all surface water bodies must achieve in the near future. These values (expressed as ‘good’ status class boundaries) are compared and harmonised through the intercalibration exercise, ensuring consistent management objectives across Europe.

Since the adoption of the European Water Framework Directive (WFD) in 2000, huge progress has been made in the ecological assessment of European waters. Many European countries now have a set of assessment tools for indicating the state of Europe’s water resources and for monitoring improvements in relation to investments in river basin management, or deterioration in response to future environmental changes (Birk et al., 2012a; Bruce et al., 2013b). These assessment methods are composed of several metrics (see Tables 1–5), and combination rules are applied to calculate the ecological assessment result for the whole system.

In order to harmonise ecological assessment systems and to ensure a consistent level of ambition in the protection and restoration of surface water bodies across the EU, an intercalibration exercise was launched, involving hundreds of experts from all Member States (Nöges et al., 2009). This exercise led to the development of innovative new approaches to accomplishing this highly complex task (Birk et al., 2013). In total, 230 methods from 28 countries were intercalibrated and published in the EC Decision (EC, 2013). This flagship document sets the harmonised boundaries for the Member States’ national methods for classifying the ecological quality of their rivers, lakes, coastal waters and estuaries.

In this article, we provide an overview of this international comparison, focusing on the assessment schemes developed for freshwater lakes. More specifically, we (1) briefly review the assessment methods developed for lakes focusing on the metrics included and the pressures addressed; (2) describe the intercalibration exercise performed on lake assessment methods; (3) assess the gaps in the lake assessment methods regarding biological communities, pressures addressed and geographical regions.

2. Intercalibration methodology

A step-by-step methodology for the comparison and harmonisation of ecological assessment methods was developed (EC, 2011; Birk et al., 2013). The assessment methods were first checked for their compliance with the WFD requirements - only methods that

Table 1

Overview of the Member States (MS) lake phytoplankton assessment methods (only intercalibrated methods). For detailed description of metrics see supplementary material S2, Table 1.

| MS | Metrics included in national systems | | | Pressures addressed and pressure indicators |
|------------------|--|-------------------------------------|-------------------------|---|
| | Abundance | Taxonomic composition | Sensitivity/tolerance | |
| Austria | CHL-A, TBV | | Brettum index | EUTR: TP |
| Belgium | CHL-A | %CYA | | EUTR: TP, TN |
| Cyprus | CHL-A, TBV | BV-CYA | IGA index | EUTR: TP, land-use, PD |
| Denmark | CHL-A | %CYA, %CHRY | Sensitivity Index | EUTR: TP, TN |
| Estonia | CHL-A _{surf} , CHL-A _{tot} | Phytoplankton community description | PCQ index | EUTR: TP, TN |
| Finland | CHL-A, TBV | %CYA (impact taxa) | TPI index | EUTR: TP |
| Germany | CHL-A, CHL-A _{max} , TBV | Algal class metrics | PTSI index | EUTR: TP, TN |
| Ireland | CHL-A | | IPI index | EUTR: TP, TN |
| Italy-lakes | CHL-A, TBV | | PTI _{OT} index | EUTR: TP |
| Italy-reservoirs | CHL-A, TBV | BV-CYA | MedPTI index | EUTR: TP, land-use, PD |
| Netherlands | CHL-A | | Bloom index | EUTR: TP, TN |
| Norway | CHL-A, TBV | BV-CYA _{max} | PTI _{NO} index | EUTR: TP |
| Poland | CHL-A, TBV | BV-CYA | | EUTR: TP, TN |
| Portugal | CHL-A, TBV | BV-CYA | IGA index | EUTR: TP, land-use, PD |
| Slovenia | CHL-A, TBV | | Brettum index | EUTR: TP |
| Spain | CHL-A, TBV | %CYA | IGA index | EUTR: TP, land-use, PD |
| Sweden | CHL-A, TBV | %CYA | TPI index | EUTR: TP |
| UK | CHL-A, TBV | BV-CYA | PTI _{UK} index | EUTR: TP, TN |

CHL-a, chlorophyll-a concentration; TBV, total biovolume; BV-CYA, biovolume of Cyanobacteria; %CYA, percentage of Cyanobacteria of total biovolume; CHRY, Chrysophyta; EUTR, eutrophication; TP, total phosphorus; TN, total nitrogen; PD, population density.

Table 2

Overview of the Member States (MS) lake macrophyte assessment methods (only intercalibrated methods). For detailed description of metrics see supplementary material S2, Table 3.

| MS | Metrics included in national systems | | | | Pressures addressed and pressure indicators |
|-------------|---|---|---|---|---|
| | Abundance | Composition | Sensitivity/tolerance | Functional and richness/diversity | |
| Austria | Colonisation depth Vegetation density | Species composition index | Trophic index | Type-specific zonation | EUTR and GD: TP, CHL-a, SD |
| Belgium | Abundance of submerged vegetation | Type-specific species composition index | Disturbance index | Macrophyte growth forms | EUTR and others: TP, TN, CHL-a |
| Denmark | Colonisation depth Total coverage | | Presence of indicator species | | EUTR: TP, TN, CHL-a |
| Estonia | Colonisation depth (deep lakes) | Abundance of different taxonomic groups | Abundance of sensitive/tolerant taxa | | EUTR and HM: TP, TN, CHL-a |
| Finland | | PMA index | Trophic index PTST index IBML index | | EUTR and HM: TP |
| France | | | Reference Index | | EUTR and GD: TP, chl-a, SD |
| Germany | Colonisation depth | Dominance of selected taxa | Reference Index | | EUTR and GD: TP, TN, CHL-a, SD |
| Ireland | Colonisation depth Average depth of presence | % RF Chara % RF elodeids | Plant trophic score % RF tolerant taxa | | EUTR: TP, TN, CHL-a |
| Italy | Colonisation depth | Dissimilarity index, Invasive species | Trophic score | | EUTR and GD: TP, chl-a, SD |
| Latvia | Colonisation depth | Abundance of different taxonomic groups | Abundance of sensitive/tolerant taxa | | EUTR: TP, TN, CHL-a |
| Lithuania | Colonisation depth | Dominance of selected taxa | Reference Index | | EUTR: TP, TN, CHL-a |
| Netherlands | | | Indicator species metrics | Growth form metrics | EUTR and HM: TP, TN, CHL-a |
| Norway | | | Trophic index | | EUTR: TP |
| Poland | Colonisation index | | | Pielou's index (syntax level) | EUTR and others: TP, TN, CHL-a |
| Slovenia | Colonisation depth | Depth limit of charophytes | Trophic index | | EUTR and GD: TP, CHL-a, SD |
| Sweden | | | Trophic index | | EUTR: TP |
| UK | Mean cover | Relative cover of filamentous algae | LMNI index | Number of functional groups Number of taxa | EUTR: TP, TN, CHL-a |

EUTR, eutrophication; HM, hydromorphological modifications; GD, general degradation; RF, relative frequency; TP, total phosphorus; TN, total nitrogen; CHL-a, chlorophyll-a concentration; SD, Secchi depth.

met these criteria could be intercalibrated. For example, assessment schemes must establish “biological reference conditions” from which the degree of human impact is measured using an Ecological Quality Ratio (EQR) – the ratio of the observed assessment value to the expected value under reference conditions. Additionally, the assessment method must include all the biological attributes included in the Directive, e.g., for phytoplankton: average

phytoplankton biomass, composition and abundance of planktonic taxa, frequency and intensity of planktonic blooms.

Secondly, the assessment methods were checked to ensure that intercalibration was feasible, with analyses restricted to methods that address similar water body types and anthropogenic pressure and which are based on similar concepts. For example, some assessment schemes for lake benthic invertebrates were

Table 3
Overview of the Member States (MS) lake phyto-benthos assessment methods (only intercalibrated methods). For detailed description of metrics see supplementary material S2, Table 2.

| MS | Sensitivity/tolerance metrics | Pressures addressed and pressure indicators |
|----------|--|---|
| Belgium | PISIAD index | EUTR: TP, chl-a |
| Germany | Trophic-index TI _{Nord} Quotient of reference species RAQ | EUTR: TP |
| Finland | IPS index | EUTR: TP |
| Hungary | IBD index; EPI-D index; TDIL index | EUTR: TP |
| Ireland | Lake Trophic Diatom Index | EUTR: TP |
| Poland | Trophic index TJ Index of reference species GR _j | EUTR: TP |
| Slovenia | Trophic index (Lake – TI) | EUTR: TP |
| Sweden | IPS index | EUTR: TP |
| UK | Diatom Assessment Of Lake Ecological Quality (DARLEQ) | EUTR: TP |

EUTR, eutrophication; TP, total phosphorus; CHL-a, chlorophyll-a concentration.

Table 4
Overview of the Member State (MS) lake benthic invertebrate assessment methods (only intercalibrated methods). For detailed description of metrics see supplementary material S2, Table 4.

| Metrics included in the national assessment systems | | | | | Pressures addressed and pressure indicators |
|---|-----------------------------|--|--|-------------------------------------|---|
| MS | Composition metrics | Sensitivity/tolerance metrics | Richness/diversity metrics | Functional metrics | |
| Belgium | | Number of sensitive taxa Mean tolerance score | EPT taxa richness Total taxa richness Shannon-Wiener diversity | | EUTR, HM: not tested |
| Estonia | | ASPT index Swedish Acidity index | EPT and total taxa richness Shannon-Wiener diversity | | EUTR: TP, land use |
| Finland | | BQ _{FI} index | | | EUTR: TP, TN, CHL-a, SD HM: Morphoindex |
| Germany-ALP ^a | RA of Odonata | Fauna index | Shannon-Wiener diversity | RA of gatherers <i>r/k</i> ratio | HM: Morphological Index |
| Germany-CB ^a | RA of Odonata | Fauna index | ETO taxa richness | RA of habitat lithal | EUTR, HM: Combined stressor index |
| Lithuania | RA of COP | ASPT index | CEP taxa richness Hill's number | | EUTR: TP |
| Netherlands | | DN%, DP%, KM% | KM% (taxa) | | HM: shore characteristics |
| Norway | | AWIC index; Acidity index | Ephemeroptera taxa richness Gastropoda taxa richness | | ACID: pH, ANC, Lal |
| Slovenia | | Littoral fauna index | Total taxa richness Margalef diversity | | HYMO: Lake Modification Index |
| Sweden-BQI ^b | | BQI _{SE} index | | | EUTR: TP |
| Sweden-MILA ^b | RA Ephemeroptera RA Diptera | AWIC index | Ephemeroptera taxa richness Gastropoda taxa richness | RA of predators | ACID: pH |
| UK-CPET ^c | | CPET index | | | EUTR: TP |
| UK-LAMM ^c | | LAMM index | | | ACID: pH, ANC |

RA, relative abundance; EUTR, eutrophication; HM, hydromorphological modifications; ACID, acidification; TP, total phosphorus; TN, total nitrogen; CHL-a, chlorophyll-a concentration; SD, Secchi depth; ANC, acid neutralising capacity; LAI, labile aluminium.

^a Germany intercalibrated two benthic fauna assessment methods: for Alpine and for Central Baltic lake types.

^b Sweden intercalibrated two methods: MILA for acidification am BQI for eutrophication.

^c UK intercalibrated two methods: LAMM for acidification am CPET for eutrophication.

developed to measure the impact from acidification (McFarland et al., 2010), whereas other national schemes measured the impact of eutrophication (Ruse, 2010). These two types of schemes are not comparable. Similarly, it was not possible to compare methods based on littoral and profundal benthic invertebrate communities, even if they address the same eutrophication pressure (Sandin et al., 2014). In several cases, assessment systems developed for deep lakes include specific metrics (e.g. macrophyte colonisation depth) and cannot be applied to shallow lakes (Søndergaard et al., 2010).

Thirdly, the biological quality element (BQE) and pressure data (e.g., nutrient levels for eutrophication, pH or acid neutralising capacity (ANC) levels for acidification) were collected from the countries involved and options for intercalibration were evaluated.

Intercalibration can be carried out in three ways, depending on the availability and similarity of national assessment methods (Fig. 1):

- If Member States have common sampling methods and thus similar kinds of data (e.g., the number of individuals of all taxa at identical determination level), then national methods were compared directly by applying each national method to the data of other countries (direct comparison; Sandin et al., 2014; Gassner et al., 2014);
- If Member States did not have common sampling methods, the results of national assessment methods were compared using common metrics, a metric developed that was sufficiently comparable with other countries to enable a comparison to be made (indirect comparison; Lyche-Solheim et al., 2013; Kelly et al., 2014);
- If assessment methods had not yet been developed, the Member State could choose to set boundaries based on a common database using the same assessment metrics (common boundary setting; Wolfram et al., 2009; Poikane et al., 2010).

Table 5

Overview of the Member States (MS) lake fish fauna assessment methods (only intercalibrated methods). For detailed description of metrics see supplementary material S2, Table 5.

| Metrics included in the national assessment systems | | | | | Pressures addressed and pressure indicators |
|---|--|--|--|---|---|
| MS | Abundance/composition | Sensitivity/tolerance | Functional metrics Richness/diversity | Age structure metrics | |
| Austria | Fish biomass (hydroacoustics) Abundance index of alien species | Abundance index of type specific species/sensitive species | Abundance index of migrating spawners/of spawning guilds/of small-bodied species | Length frequency of sentinel species | EUTR, HM, fisheries, alien species: index based on expert judgement |
| Finland | Total biomass of fish Total number of fish Biomass proportion of cyprinid fish | Indicator species | | | EUTR: TP, landuse |
| Germany | | Sentinel/type/side species number Sentinel species abundance | Habitat/Spawning preferences Abundance of habitat/spawning preferences | Reproduction of potentially stocked species | EUTR, HM, fisheries, alien species: index based on expert judgement |
| Ireland | Total biomass Native fish biomass Perch/roach biomass Rel. abundance of bream/rudd Rel. biomass of cyprinid species/non-native species | | Relative abundance of rheophilic/lithophilic/phytophilic species Species evenness/dominance index | Maximum length of dominant species (based on BPUE) | EUTR: TP, CHL-a |
| Italy | | NPUE of the guiding species Reduction % of guiding + accompanying species | Relative richness of alien species | Population structure of the guiding species Reproductive success % of guiding + accompanying species | EUTR, HM, fisheries, alien species: index based on expert judgement |

EUTR, eutrophication; HM, hydromorphological modifications; TP, total phosphorus; CHL-a, chlorophyll-a concentration.




Fig. 1. Flow chart of the main steps of the intercalibration process.

Due to biogeographical and typological reasons, as well as differences in data acquisition, the biological data of different countries cannot be compared without concern (Cao and Hawkins, 2011; Birk et al., 2013). As an example, the number of benthic invertebrate taxa might be generally higher in one country than in others, because the sampling covers much more area per site (Böhmer et al., 2014). The richness and diversity of fish species are related mainly to geographical and climatic variables (Bruce et al., 2013a). Hence, a “benchmarking” procedure was applied with the aim of correcting any biogeographical and methodological differences within a common dataset that can cause incomparability. Three different approaches were applied: “reference benchmarking” based on near-natural reference sites (Pardo et al., 2012), “alternative benchmarking” using sites at similar impairment levels (Birk et al., 2012b), and “continuous benchmarking” using pressure–response gradients (Kelly et al., 2014; for a detailed description of this procedure see Birk et al., 2013).

The final objective of intercalibration was to compare and harmonise national boundaries. This was done using a standardised analytical procedure and harmonised comparability criteria. The main criteria used for evaluating comparability was *boundary bias*: the deviation of a class boundary relative to the common view of the Member States (defined by the common metrics or by the global mean of all of the methods).

The most that any national boundary could deviate from the global mean view of all countries was ± 0.25 classes and therefore the most widely divergent national methods could not differ from each other by more than 0.5 classes. National methods that did not comply with these criteria were required to adjust their boundaries until acceptable boundary bias and class agreement were achieved. It is important to stress that intercalibration checks whether the results are comparable, irrespective of method.

For definition of the main terms used in the intercalibration of ecological assessment methods see supplementary material S3.

3. Lake assessment methods

3.1. Lake assessment methods: phytoplankton

The use of phytoplankton for water quality assessment of lakes has a long history, and considerable knowledge of this subject has accumulated over the past century (Willén, 2000). Phytoplankton has traditionally been one of the dominant elements in lake assessment and thus has played an important role in the development of WFD-compliant assessment systems. Nearly all of the Member States have developed phytoplankton assessment methods. Out of 24 methods participating in the intercalibration exercise, 18 methods were included in the final intercalibration results. All of the methods address eutrophication pressure and follow similar assessment principles (Table 1):

- All include measures of phytoplankton abundance – all national methods consider chlorophyll-a, and most also consider total phytoplankton biovolume (Carvalho et al., 2013b);
- Most include measures of cyanobacteria (as proxy measures of the risk of toxic algal blooms) as cyanobacteria biovolume or cyanobacteria as a percentage of the total phytoplankton biovolume (Carvalho et al., 2013b);
- All include metrics based on the trophic preferences of species or algal groups, e.g., Brettum index (Austria, Slovenia; Brettum, 1989) and PTI index (Finland, Sweden; Willén, 2007). However, functional and diversity indices are not used (except Estonia where Pielou's evenness index is used).

Despite some variation between countries in sampling methods and sampling periods (Nõges et al., 2010), all of the intercalibrated methods exhibited significant relationships with total phosphorus (TP), and some of the methods also exhibited significant relationships with total nitrogen (TN) and catchment characteristics. However, it has been difficult to establish relationships for Eastern Continental lakes and Mediterranean reservoirs due to both a lack of data (Padisák et al., 2006) and peculiarities of the water bodies. For example, Mediterranean reservoirs are affected by extreme changes in their water level due to seasonal climate variation (Marchetto et al., 2009) while Eastern Continental lakes are considered naturally eutrophic with nutrient concentrations that are considerably higher than limiting thresholds (Borics et al., 2013).

3.2. Lake assessment methods: macrophytes

Macrophytes have been used in lake assessment for many years, and the macrophyte community is generally regarded as a key indicator of ecological status both in deep (Pall and Moser, 2009) and in shallow lakes (Søndergaard et al., 2010). However, in contrast to phytoplankton assessment systems, existing macrophyte assessment was confined to relatively small geographic regions, mainly in Scandinavia and Central Europe, and was based primarily on indicator species analyses (Melzer, 1999), while the WFD requires lake assessment based on both macrophyte abundance and composition. Most European countries have undertaken significant efforts to develop WFD-compliant macrophyte assessment tools. Twenty systems have been developed recently, and 17 of these have been intercalibrated (Table 2).

All macrophyte assessment systems (except that in Poland) include sensitivity/tolerance metrics expressed as indices that are based on species indicator values, e.g., a trophic index (Pall and Moser, 2009) or relative abundance of sensitive versus tolerant taxa (Schaumburg et al., 2004; Free et al., 2006). Most macrophyte assessment systems include an assessment of abundance (except the methods of Norway, Sweden and Finland). The most widely

used abundance measures are the macrophyte colonisation depth and the abundance of submerged macrophytes.

Most macrophyte assessment systems claim to address not only eutrophication but also other pressures, such as general degradation and hydromorphological changes. However, significant pressure response relationships have only been shown for eutrophication parameters (nutrient and chlorophyll-a concentrations and water transparency) (Table 2). Even in these cases there remain some difficulties with the response of macrophyte metrics to pressure, especially the delayed response to both increasing and decreasing eutrophication (Pall and Moser, 2009) and the non-linear reaction to eutrophication following the alternative stable state theory (Penning et al., 2008). Further challenges are establishing pressure–response relationships for pressures that are themselves quite hard to quantify (hydromorphological modifications, i.e. shoreline developments and water level fluctuations), as well as for multiple stressors.

3.3. Lake assessment methods: phytobenthos

Although phytobenthos is only one part of the BQE “macrophytes and phytobenthos” (Annex V, WFD), most Member States have developed separate assessment methods for macrophytes and phytobenthos. Moreover, only 11 of the 27 Member States of the EU took part in the intercalibration of phytobenthos methods (Tables 3 and 7). Nevertheless, the collective view of the phytobenthos expert group (Kelly et al., 2014) was that a Member state cannot be considered to be fully compliant with the WFD requirements if it possesses only a macrophyte (or only phytobenthos) method. There are situations (e.g. where the lake is subject to hydromorphological stress, navigation, etc.) where macrophytes have completely disappeared and will not give any reliable indication of the impact of nutrients on littoral flora, as well as situations when the two elements react at different rates to changes in their environment (Schaumburg et al., 2004; Pall and Moser, 2009).

All national phytobenthos methods assess the composition and relative abundance of diatoms, assumed to be proxies for the phytobenthos community as a whole (Table 3). All the methods address eutrophication pressure and have demonstrated significant relationships to total phosphorus concentrations with the exception of some methods in low alkalinity lakes, where a combination of short gradients and the confounding effects of low pH led to very weak relationships with TP.

3.4. Lake assessment methods: benthic invertebrates

The lack of WFD-compliant macroinvertebrate assessment tools was identified as one of the major knowledge gaps impeding the full assessment of the ecological quality of lakes (Solimini et al., 2006). Macroinvertebrates have been recognised as one of the most difficult biological groups to consider in the assessment of lake quality for three main reasons: their complex biotic structure, their high spatial and temporal variability (Solimini et al., 2006; Free et al., 2006; Solimini and Sandin, 2012). In spite of these factors, 20 systems have been developed, of which 13 have been successfully intercalibrated (Table 4). Of the 44 metrics, almost half (42%) are sensitivity/tolerance metrics included in all national systems. Some countries have used traditional indices, such as the ASPT index (Lithuania and Estonia; Armitage et al., 1983), Benthic Quality Index (Sweden and Finland; Wiederholm, 1980) and Acidity index (Norway and Estonia; Henrikson and Medin, 1986). Most of the Member States have developed new sensitivity indices, such as the Littoral Fauna Index (Slovenia; Urbanic, 2014), Mean Tolerance Score (Belgium; Gabriels et al., 2010) and the LAMM index (United Kingdom; McFarland et al.,

2010). Eight methods also contain richness/diversity methods, the most frequently used being metrics of total taxa richness, EPT (Ephemeroptera, Plecoptera and Trichoptera) taxa richness (Lenat, 1988) and Shannon–Wiener diversity. Only four methods contain composition metrics, while functional metrics are rarely used.

Most of the current benthic fauna assessment systems are based on significant pressure–response relationships. Nonetheless, the variation explained by the models was found to be low in many cases and was considered unsatisfactory (Free et al., 2006). The low explanatory power of the models has several explanations, i.e. complex biotic structure, impact of various environmental factors and multiple pressures, and habitat impact (Solimini et al., 2006; Free et al., 2009). In short, while the development of benthic assessment methods can be considered a partial success story, there is still a need to further understand the structure of lake benthic macroinvertebrate communities and their response to anthropogenic pressure to reduce the uncertainty of the metrics developed. Recently, a new harmonized multimetric assessment system has been proposed for Western, Northern, Central and Southern Europe (including natural Mediterranean lakes; Miler et al., 2013) and its implementation in national assessment systems may represent the way forward to overcoming those limitations.

3.5. Lake assessment methods: fish fauna

It is well established that fish are sensitive indicators of environmental degradation and offer the major advantage of integrating the direct and indirect effects of stress over large scales of space and time (Minns et al., 1994). Fish exhibit reactions to eutrophication (Mehner et al., 2007), habitat destruction, shoreline degradation, lake use intensity (Belpaire et al., 2000), hydromorphological degradation, connectivity (Degerman et al., 2001), acidification (Henriksen et al., 1989) and combined degradation (Whittier, 1999). Nevertheless, the fish community is often an overlooked and neglected aspect of lake monitoring. So far, only eight Member States have finalised fish assessment systems (Kelly et al., 2012; Olin et al., 2013), and only five of these have shown significant pressure–response relationships (Table 5).

The use of fish communities as indicators of environmental quality is potentially challenging (Kelly et al., 2012), with several problems: (i) a wide variety of sampling methods are used by the Member States, including multi-mesh gill nets, electro-fishing, trawling and hydro-acoustics; (ii) the activities of fishing, stocking and the introduction of exotic species, all of which can have a large impact on the natural fish fauna; (iii) lakes are subjected to multiple pressures (shoreline degradation, eutrophication and water level regulation), and fish, at the topmost level of the trophic cascade, indirectly integrate the effects of these on lower trophic levels; (iv) high natural variability in fish metrics, which may be related to lake size, depth and water chemistry; (v) fish are mobile and can avoid areas of environmental stress, resulting in this BQE being less sensitive to pressures than others. Taking all these factors together, it is not surprising that there are few significant relationships between fish metrics and specific pressure indicators (Olin et al., 2013). Fish are, however, important indicators: they are at the top of the food chain, have significant economic and social importance and their assessment is an important part of an integrated approach to water management. One solution to this problem is to base assessment of pressure on expert judgement (e.g., Aubry and Elliott, 2006); Austria, Germany, and Italy have demonstrated a response of their overall fish assessment result to a combined pressure index (all common pressures scored and summed up to create an overall pressure index).

4. Lake intercalibration

4.1. Intercalibration groups

Fourteen Lake Intercalibration groups were formed, each focusing on a specific geographical region/BQE/anthropogenic pressure combination (Table 7, only finalised results included):

- For phytoplankton, macrophytes and phytobenthos, all geographic groups just addressed eutrophication pressure;
- For phytobenthos, all methods were intercalibrated within a single pan-European group due to the low number of methods available;
- For benthic invertebrates, the Alpine group addressed hydromorphological alterations whilst the Central Baltic group addressed combined pressures, including hydromorphological alteration and eutrophication. In the Northern region, two separate groups were created, one that addressed eutrophication using profundal communities whilst another addressed acidification using littoral communities.
- For fish fauna, the Alpine group addressed diverse pressures, but the Northern group only addressed eutrophication.

In several cases, it was not possible to intercalibrate the methods:

- Due to the low number of methods in the group (the Mediterranean benthic fauna group and the Central Baltic fish group);
- Due to different pressure/assessment concepts on which the methods were based (e.g. the Norwegian fish method which addressed acidification could not be included in the Northern fish group dealing with eutrophication).

4.2. Common metrics and benchmarking

In the next step, datasets were collected and compared. In most cases (nine groups), the assessment methods and data were sufficiently similar to enable direct comparison, where each national method was applied to all data in a common dataset. For five groups, specific common metrics were developed for the intercalibration exercise (Table 7, for details see Lyche-Solheim et al., 2013; Phillips et al., 2013). For the lake phytobenthos intercalibration exercise, the trophic index (TI: Rott et al., 1999), a widely used phytobenthos metric, was used as the common metric of intercalibration.

Half of the groups applied reference-site benchmarking. This was true primarily in the Northern and Alpine regions, where near-natural reference sites were available in sufficient numbers. Six Intercalibration groups applied continuous benchmarking, which uses the available pressure gradient (e.g. from good to bad status) to identify country-specific differences. This approach was used in the Central Baltic region, where reference sites are rare, or no longer exist, and the range of pressures was high across the group but not always adequately represented within each country. It was also used in several cases in the Northern and Alpine regions, as this approach was independent of national views of reference and is, therefore, more robust.

4.3. Boundary harmonisation

In the final step, national boundaries on the EQR scale were compared and, if necessary, adjusted if the agreed comparability criteria were exceeded.

Only three groups (Alpine phytoplankton, Alpine benthic fauna, and Northern Benthic fauna acidification methods) found that no boundary adjustment was needed, as the comparability analysis

Table 6
Overview of the Member States (MS) lake assessment methods.

| MS | Phytoplankton | Macrophytes | Phytobenthos | Benthic fauna | Fish fauna |
|-------------|-----------------------------------|-----------------------------------|----------------------------------|-----------------------------------|----------------------------------|
| Austria | Intercalibrated | Intercalibrated | – | – | Intercalibrated |
| Belgium | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated | – |
| Bulgaria | – | Submitted | – | – | – |
| Cyprus | Intercalibrated | n.a. | n.a. | n.a. | n.a. |
| Denmark | Intercalibrated | Intercalibrated | – | – | – |
| Estonia | Intercalibrated | Intercalibrated | – | Intercalibrated | – |
| Finland | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated |
| France | Submitted | Intercalibrated | Submitted | Submitted | Submitted |
| Germany | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated (2) Submitted | Intercalibrated |
| Greece | – | – | – | – | – |
| Hungary | Submitted | Submitted | Intercalibrated | Submitted | – |
| Ireland | Intercalibrated | Intercalibrated | Intercalibrated | – | Intercalibrated |
| Italy | Intercalibrated (2) | Intercalibrated | Submitted | Submitted | Intercalibrated |
| Latvia | Submitted | Intercalibrated | – | – | – |
| Lithuania | Submitted | Intercalibrated | – | Intercalibrated | – |
| Netherlands | Intercalibrated | Intercalibrated | – | Intercalibrated | Submitted |
| Norway | Intercalibrated | Intercalibrated | – | Intercalibrated | Submitted |
| Poland | Intercalibrated | Intercalibrated | Intercalibrated | – | – |
| Portugal | Intercalibrated | n.a. | n.a. | n.a. | n.a. |
| Romania | Submitted | Submitted | – | Submitted | – |
| Slovenia | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated | – |
| Spain | Intercalibrated | Submitted | – | Submitted | – |
| Sweden | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated (2) Submitted | Submitted |
| UK | Intercalibrated | Intercalibrated | Intercalibrated | Intercalibrated (2) | – |
| In total | 18 intercalibrated 5 submitted | 17 intercalibrated 3 submitted | 9 intercalibrated 2 submitted | 13 intercalibrated 6 submitted | 5 intercalibrated 4 submitted |

Intercalibrated, method intercalibrated and included in the final results; Submitted, method participated in the intercalibration but not included in the final results; n.a., not applicable, method development and intercalibration not feasible due to lack of natural lakes. (2), MS has intercalibrated 2 assessment methods (e.g., for different human impacts).

showed that the methods yielded very similar assessments (in agreement with the comparability criteria defined in the Intercalibration Guidance). These were groups with just a few (2–4) similar assessment methods (for example, Alpine phytoplankton methods all include biomass metrics and composition metrics each with harmonised boundary values agreed upon during development (Wolfram et al., 2009).

In all other groups, boundary adjustments were needed. In some cases, only the class boundaries of a few methods were adjusted. In several cases, the assessment methods were so dissimilar that more profound changes were needed, e.g., in the values associated with reference status or in the way data was used and combined. For example, reference values of the Belgian benthic fauna system were revised, and the combination rules of the specific metrics used in the Norwegian and British phytoplankton systems were adjusted.

For detailed description of one intercalibration exercise see supplementary material S3.

5. Intercalibration gaps

5.1. Intercalibration gaps: fish fauna, phytobenthos and benthic invertebrates

The situation with respect to different BQEs is highly variable. Nearly all Member States have developed and intercalibrated assessment methods for phytoplankton and macrophytes. The only exception is Greece (which has not developed methods for either BQE) and Bulgaria (which has not developed a method for phytoplankton). In contrast, only five countries have intercalibrated lake fish-based assessment methods, nine have phytobenthos methods and 10 have benthic fauna assessment methods.

5.2. Eastern Continental and Mediterranean regions

There have been particular difficulties in the development and intercalibration of ecological assessment methods in the Eastern Continental and Mediterranean regions (Table 6). Due to high evaporation/precipitation ratios and low geographic relief, lakes in the Eastern Continental region are often endorheic and naturally eutrophic. In this region, there are relatively few examples of lakes with catchments in a near natural state, and as the least impacted also have relatively high nutrient concentration, e.g. TP > 100 µg/l, pressure–response models show asymptotic behaviour and high variation (Borics et al., 2013). Thus, there have been considerable difficulties in establishing sound pressure–response relationships and setting meaningful ecological class boundaries (Poikane et al., 2014). Further research is, therefore, needed to develop and intercalibrate ecological assessment tools for these lakes.

The main difficulties encountered in the Mediterranean region are the small number and high diversity of natural lakes. In the entire Mediterranean region, only 257 lakes are reported, mostly (84%) in Spain, which includes many lakes with water surface areas of less than 50 ha. Lakes are highly diverse: for example, in the Mediterranean part of Italy, there are 13 lakes, including the paleosaline Lago di Pergusa – a unique saltwater lake in Europe, the Lago di Trasimeno – a very large shallow lake, and several volcanic lakes of different depths. Therefore, in spite of a common effort within the Mediterranean intercalibration group, it has not been possible to intercalibrate natural Mediterranean lakes, due to the lack of a sufficient number of lakes within common types. Nevertheless, Mediterranean countries must develop tools for evaluating the ecological quality of their lakes, even though this development is hindered by the limited amount of data available and the high inter- and intra-annual variability of the biological communities and water characteristics (Boix et al., 2005). In both regions, the

Table 7
Overview of the Member States (MS) lake intercalibration exercises (only finalised exercises).

| BQE | Region | Pressure addressed | Intercalibration option | Common metrics used | Benchmarking applied | MS methods intercalibrated | MS participated but not intercalibrated |
|-----------------------|--------|--------------------|--|--|------------------------------|---------------------------------------|---|
| Phytoplankton | ALP | EUTR | Direct comparison | Average of each national methods EQRs | Reference sites | AT, DE, IT _{lakes} , SI | FR |
| | CB | EUTR | Comparison via common metrics | Multimetrics of Chlorophyll-a and PTI EQR (Phillips et al., 2013) | Continuous | BE, DE, DK, EE, IE, NL, PL, UK | LV, LT |
| | MED | EUTR | Direct comparison | Average of each national methods EQRs | Reference sites + continuous | CY, ES, IT _{reservoirs} , PT | RO, FR |
| | NOR | EUTR | Direct comparison Comparison via common metrics | Multimetrics of Chlorophyll-a and PTI EQR (Phillips et al., 2013) | Continuous | FI, IE, NO, SE, UK | |
| Macro phytes | ALP | EUTR | Direct comparison | Average of each national methods EQRs | Reference sites | AT, DE, FR, IT, SI | |
| | CB | EUTR | Direct comparison | Average of each national methods EQRs | Continuous | BE, DE, DK, EE, LT, LV, NL, PL, UK | |
| | NOR | EUTR | Comparison via common metrics | Lake Macrophyte Intercalibration common metrics (Hellsten et al., 2014) | Reference sites | FI, IE, NO, SE, UK | |
| Phytobenthos | All | EUTR | Comparison via common metrics | Trophic index (Rott, 1999) | Continuous | BE, FI, DE, HU, IE, PL, SI, SE, UK | IT, FR |
| Benthic invertebrates | ALP | HM | Comparison via common metrics | Multimetrics of Fauna index, number of taxa, <i>r/k</i> ratio and RA of feeding type gatherer (Böhmer et al., 2014) | Continuous | DE _{ALP-eulit} , SI | DE _{ALP-sublit} , FR, IT |
| | CB | HM and EUTR | Comparison via common metrics | Multimetrics of EPTCBO taxa number, ASPT index, relative abundance of ETO and microhabitat type lithal (Böhmer et al., 2014) | Continuous | BE, DE, EE, LT, NL, UK | |
| | NOR | EUTR | Direct comparison | Average of each national methods EQRs | Reference sites | FI, SE _{MILA} | SE _{ASPT} , UK _{CPET} |
| | NOR | ACID | Direct comparison | Average of each national methods EQRs | Reference sites | NO, SE, UK _{LAMM} | |
| Fish fauna | ALP | Diverse impacts | Direct comparison | Global mean of all the methods | Site-specific references | AT, DE, IT | FR |
| | NOR | EUTR | Direct comparison | Global mean of all the methods | Reference sites | FI, IE | SE, NO |

EUTR, eutrophication; HM, hydromorphological modifications; ACID, acidification; BQE, biological quality element; ALP, Alpine; CB, Central Baltic; MED, Mediterranean; NOR, Northern.

use of palaeolimnological data (e.g. Bennion et al., 2004) may be more suitable for defining site-specific reference conditions.

5.3. Anthropogenic pressures addressed by lake assessment methods

Lakes in Europe are subject to manifold anthropogenic pressures (eutrophication, acidification, hydromorphological alterations, alien species and climate change). A recent analysis showed that approximately 30% of European lakes are impacted by hydromorphological modifications and 20% are impacted by acidification (EEA, 2012). Despite this, most of the methods developed focus on eutrophication impacts, while only three methods address acidification (benthic invertebrate methods of Norway, Sweden and the UK) and three – hydromorphological alterations (benthic invertebrate methods of Germany, the Netherlands and Slovenia). Some of the methods claim to address multiple pressures (mainly eutrophication and hydromorphological alterations together), but not all the

necessary pressure–response relationships have been established, or they have been established only for eutrophication parameters. These gaps in lake assessment highlight the need for further research, particularly on the impact of combined pressures (e.g. eutrophication and hydrological pressures, such as changes to natural water level regimes or flushing rates).

5.4. Heavily modified and artificial water bodies

Many lakes in Europe are either artificial (e.g. fish ponds) or heavily modified (e.g. reservoirs for hydropower or water storage). For such water bodies the environmental objective is the good ecological potential instead of the good ecological status. Overall, 15.8% of the lake water bodies are designated by the Member States as either heavily modified water bodies or artificial water bodies (EEA, 2012). For ecological potential the same biological quality elements are used as for the ecological status, with a crucial difference in how reference conditions are used. Ecological status always has the

undisturbed type-specific reference conditions as a starting point; ecological potential refers to a situation where the negative effects of a physical modification (e.g. a dam) are mitigated as much as possible, taking into account the costs and benefits of these measures (Borja and Elliott, 2007).

The WFD requires intercalibration of biological methods for all water categories including heavily modified water bodies, but so far the focus has been almost exclusively on natural water bodies. Only recently the intercalibration of ecological potential methods has been started, with a focus that is more on comparing and harmonising the ways different countries apply mitigation measures for different water uses, and how existing biological methods are used in their classification.

5.5. Detection of cyanobacteria blooms

Despite successful European policies to reduce nutrient emissions (EEA, 2012), Cyanobacterial blooms are still one of the most widespread effects of eutrophication in Europe (Dolman et al., 2012; Carvalho et al., 2013b) which are further aggravated by climate change (Paerl and Paul, 2012). MS have to measure plankton bloom intensity and frequency and to ensure that persistent summer blooms do not occur. Bloom definition has been developed (Mischke et al., 2011) and a sufficiently strong metric for phytoplankton blooms, based on cyanobacteria biovolume has been demonstrated (Carvalho et al., 2013a). All phytoplankton assessment methods reflect bloom intensity as they either include a cyanobacterial bloom metric or show a strong correlation with a bloom metric (Lyche-Solheim et al., 2014). Bloom frequency is, however, still not tackled in an appropriate way, as the current sampling frequencies are not adequate to capture the temporal dynamics of phytoplankton over days and weeks, especially short-lived blooms (Dubelaar et al., 2004; Søndergaard et al., 2011). With the development of satellite technology in the near future, high frequency and high resolution satellite imagery may enable improved temporal representation of the open water of lakes for parameters such as phytoplankton chlorophyll-a and cyanobacteria biovolume (Hunter et al., 2010), allowing missing metrics such as bloom frequency to be addressed.

5.6. Uncertainty

The intercalibration exercise was concerned solely with harmonizing the position of the high/good and good/moderate boundaries for individual BQEs. In practice, ecological status assessments will be based on the simultaneous assessment of several BQEs and, as these estimates will each differ in their precision, it would be interesting to compare the sensitivity of each method (taking into account uncertainties associated with sampling regimes and analytical procedures) and, indeed, to compare national capabilities for classifying sites when using all BQEs simultaneously. Progress has been made with understanding uncertainty associated with individual BQEs (Kelly et al., 2009; Thackeray et al., 2013), though there is still much to learn about classification procedures under the rules prescribed in the WFD (Caroni et al., 2013; Moe et al., 2015).

6. Conclusions

In total, 82 lake assessment methods were reported for the intercalibration, and, following final adjustments, 62 were successfully intercalibrated. The results include a high proportion of phytoplankton (18), macrophyte (17) and benthic invertebrate assessment methods (13), but few phytobenthos (9) and fish assessment methods (5). Most of the methods were developed to detect the impact of eutrophication (50 methods), although in addition, some methods have demonstrated a significant response to

hydromorphological pressures (3), acidification (3) or combined pressures (5). Most of the Central and Northern European countries have developed assessment methods while significant gaps exist for Eastern Continental and Mediterranean countries. The intercalibrated methods are now included in the EC Decision (EC, 2013) setting a legal obligation on Member States to use these harmonised boundaries to assess the ecological quality of their lakes. This is a major step forward in setting consistent management objectives for European water bodies but much remains to be done, mainly developing assessment systems (i) for phytobenthos and fish fauna, (ii) for other pressures and pressure combination, except eutrophication, and (iii) for Eastern Continental and Mediterranean lakes.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ecolind.2015.01.005>.

References

- Adler, R.W., 2003. The two lost books in the water quality trilogy: the elusive objectives of physical and biological integrity. *Environ. Law* 33, 29–78.
- Adler, R.W., 2013. The decline and (possible) renewal of aspiration in the Clean Water Act. *Wash. Law Rev.* 88, 759–812.
- [ANZECC] Australian and New Zealand Environment and Conservation Council, 2000. National Water Quality Management Strategy: Australian and New Zealand Guidelines for Fresh and Marine Water Quality. Canberra, Australia.
- Armitage, P.D., Moss, D., Wright, J.F., Furse, M.T., 1983. The performance of a new biological water quality score system based on a wide range of unpolluted running-water sites. *Water Res.* 17, 333–347.
- Aubry, A., Elliott, M., 2006. The use of environmental integrative indicators to assess seabed disturbance in estuaries and coasts: application to the Humber Estuary, UK. *Mar. Pollut. Bull.* 53, 175–185.
- Belpaire, C., Smolders, R., Vanden Auweele, I., Ercken, D., Breine, J., Van Thuyne, G., Ollevier, F., 2000. An index of biotic integrity characterizing fish populations and the ecological quality of Flandrian water bodies. *Hydrobiologia* 434, 17–33.
- Bennion, H., Fluin, J., Simpson, G.L., 2004. Assessing eutrophication and reference conditions for Scottish freshwater lochs using subfossil diatoms. *J. Appl. Ecol.* 41, 24–138.
- Birk, S., Bonne, W., Borja, A., Brucet, S., Courrat, A., Poikane, S., van de Bund, W., Zampoukas, N., Hering, D., 2012a. Three hundred ways to assess Europe's surface waters: an almost complete overview of biological methods to implement the Water Framework Directive. *Ecol. Indic.* 18, 31–41.
- Birk, S., van Kouwen, L., Willby, N., 2012b. Harmonising the bioassessment of large rivers in the absence of near-natural reference conditions – a case study of the Danube River. *Freshw. Biol.* 57, 1716–1732.
- Birk, S., Willby, N., Kelly, M.G., Bonne, W., Borja, A., Poikane, S., van de Bund, W., 2013. Intercalibrating classifications of ecological status: Europe's quest for common management. *Sci. Total Environ.* 454–455, 490–499.
- Boix, D., Gascón, S., Sala, J., Martinoy, M., Gifre, J., Quintana, X.D., 2005. A new index of water quality assessment in Mediterranean wetlands based on crustacean and insect assemblages. *Aquat. Conserv.* 15, 635–651.
- Borics, G., Nagy, L., Miron, S., Grigorszky, I., László-Nagy, Z., Lukács, B.A., Tóth, L., Vár-bíró, G., 2013. Which factors affect phytoplankton biomass in shallow eutrophic lakes? *Hydrobiologia* 714, 93–104.
- Borja, A., Elliott, M., 2007. What does 'good ecological potential' mean, within the European Water Framework Directive? *Mar. Pollut. Bull.* 54, 159–1564.
- Brettum, P., 1989. Algen als Indikatoren für die Gewässerqualität in norwegischen Binnenseen. Norsk institutt for vannforskning (NIVA), Oslo, Norwegian.
- Brucet, S., Pédrón, S., Mehner, T., Lauridsen, T.L., Argillier, C., Winfield, I.J., Volta, P., Emmrich, M., Hesthagen, T., Holmgren, K., et al., 2013a. Fish diversity in European lakes: geographical predictors dominate over anthropogenic pressures. *Freshw. Biol.* 58, 1779–1793.
- Brucet, S., Poikane, S., Birk, S., Lyche-Solheim, A., 2013b. Biological assessment of European lakes: ecological rationale and human impacts. *Freshw. Biol.* 58, 1106–1115.
- Böhmer, J., Arbacauskas, K., Benstead, R., Gabriels, W., Porst, G., Reeze, B., Timm, H., 2014. Water Framework Directive Intercalibration Technical Report: Central Baltic Lake Benthic invertebrate ecological assessment methods. Report EUR 26504 EN. Publications Office of the European Union, Luxembourg.
- Cao, Y., Hawkins, C.P., 2011. The comparability of bioassessments: a review of conceptual and methodological issues. *J. N. Am. Benthol. Soc.* 30, 680–701.
- Caroni, R., van de Bund, W., Clarke, R.T., Johnson, R.K., 2013. Combination of multiple biological quality elements into waterbody assessment of surface waters. *Hydrobiologia* 704, 437–451.
- Carvalho, L., McDonald, C., de Hoyos, C., Mischke, U., Phillips, G., Borics, G., Poikane, S., Skjelbred, B., Lyche-Solheim, A., Van Wichelen, J., Cardoso, A.C., 2013a. Sustaining recreational quality of European lakes: minimizing the health risks from algal blooms through phosphorus control. *J. Appl. Ecol.* 50, 315–323.

- Carvalho, L., Poikane, S., Lyche-Solheim, A., Phillips, G., Borics, G., Catalan, J., de Hoyos, C., Drakare, S., Dudley, B., Järvinen, M., et al., 2013b. Strength and uncertainty of lake phytoplankton metrics for assessing eutrophication impacts in lakes. *Hydrobiologia* 704, 127–140.
- Davies, S.P., Jackson, S.K., 2006. The Biological Condition Gradient: a conceptual model for interpreting detrimental change in aquatic ecosystems. *Ecol. Appl.* 16, 1251–1266.
- Degerman, E., Hammar, J., Nyberg, P., Svårdson, G., 2001. Human impact on the fish diversity in the four largest lakes of Sweden. *Ambio* 30, 522–528.
- de Groot, R.S., Alkemade, R., Braat, L., Hein, L., Willemens, L., 2010. Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecol. Complex* 7, 260–272.
- Diamond, J., Stribling, J.R., Huff, L., Gilliam, J., 2012. An approach for determining bioassessment performance and comparability. *Environ. Monit. Assess.* 184, 2247–2260.
- Dolman, A.M., Rucker, J., Pick, F.R., Fastner, J., Rohrlack, T., Mischke, U., Wiedner, C., 2012. Cyanobacteria and cyanotoxins: the influence of nitrogen versus phosphorus. *PLoS ONE* 7 (6), 1–14.
- Doremus, H., Dan Tarlock, A., 2013. Can the clean water act succeed as an ecosystem protection law? *J. Energy Environ. Law* 4, 46–66.
- Dubelaar, G.B.J., Geerders, P.J.F., Jonker, R.R., 2004. High frequency monitoring reveals phytoplankton dynamics. *J. Environ. Monit.* 6, 946–952.
- [EC] European Commission, 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23rd October 2000 Establishing a Framework for Community Action in the Field of Water Policy. Official Journal of the European Communities, L327/1. European Commission, Brussels.
- [EC] European Commission, 2011. Common Implementation Strategy for the Water Framework Directive (2000/60/EC). Guidance Document 14. Guidance on the Intercalibration process 2008–2011. Office for Official publications of the European Communities, Luxembourg.
- [EC] European Commission, 2013. Commission Decision of 20 September 2013 Establishing, Pursuant to Directive 2000/60/EC of the European Parliament and of the Council, the Values of the Member State Monitoring System Classifications as a Result of the Intercalibration Exercise. Official Journal of the European Communities, L266/1. European Commission, Brussels.
- [EEA] European Environment Agency, 2012. European Waters – Assessment of Status and Pressures. European Environment Agency, Copenhagen.
- Free, G., Little, R., Tierney, D., Donnelly, K., Caroni, R., 2006. A Reference Based Typology and Ecological Assessment System for Irish Lakes. Preliminary Investigations. EPA, Wexford, Ireland.
- Free, G., Solimini, A., Rossaro, B., Marziali, L., Giacchini, R., Paracchini, B., Ghiani, M., Vaccaro, S., Gawlik, B.M., Fresner, R., et al., 2009. Modelling lake macroinvertebrate species in the shallow sublittoral: relative roles of habitat, lake morphology, aquatic chemistry and sediment composition. *Hydrobiologia* 633, 123–136.
- Gabriels, W., Lock, K., De Pauw, N., Goethals, P.L.M., 2010. Multimetric Macroinvertebrate Index Flanders (MMIF) for biological assessment of rivers and lakes in Flanders (Belgium). *Limnologica* 40, 199–207.
- Gassner, H., Achleitner, D., Luger, M., Ritterbusch, D., Schubert, M., Volta, P., 2014. Water Framework Directive Intercalibration Technical Report: Alpine Lake Fish Fauna Ecological Assessment Methods. EUR 26506 EN. Publications Office of the European Union, Luxembourg.
- Hellsten, S., Willby, N., Ecke, F., Mjelde, M., Phillips, G., Tierney, D., 2014. Water Framework Directive Intercalibration Technical Report: Northern Lake Macrophyte Ecological Assessment Methods. Report EUR 26513 EN. Publications Office of the European Union, Luxembourg.
- Henriksen, A., Lien, L., Rosseland, B.O., Traaen, T.S., Sevaldrud, I.S., 1989. Lake acidification in Norway: present and predicted fish status. *Ambio* 18, 314–321.
- Henrikson, L., Medin, M., 1986. Biologisk bedömning av försurningspåverkan på Lelångens tillflöden och grundområden. *Aquaekologerna, Älvsborgs län, Swedish*.
- Hering, D., Borja, A., Carvalho, L., Feld, C.K., 2013. Assessment and recovery of European water bodies: key messages from the WISER project. *Hydrobiologia* 704, 1–9.
- Hunter, P.D., Tyler, A.N., Carvalho, L., Codd, G.A., Maberly, S.C., 2010. Hyperspectral remote sensing of cyanobacterial pigments as indicators for cell populations and toxins in eutrophic lakes. *Remote Sens. Environ.* 114, 2705–2718.
- Kelly, F.L., Harrison, A.J., Allen, M., Connor, L., Rosell, R., 2012. Development and application of an ecological classification tool for fish in lakes in Ireland. *Ecol. Indic.* 18, 608–619.
- Kelly, M., Bennion, H., Burgess, A., Ellis, J., Juggins, S., Guthrie, R., Jamieson, J., Adriessens, V., Yallop, M., 2009. Uncertainty in ecological status assessments of lakes and rivers using diatoms. *Hydrobiologia* 633, 5–15.
- Kelly, M., Urbanic, G., Acs, E., Bertrin, V., Burgess, A., Denys, L., Gottschalk, S., Kahlert, M., Karjalainen, S.M., Kennedy, B., et al., 2014. Comparing aspirations: intercalibration of ecological status concepts across European lakes using littoral diatoms. *Hydrobiologia* 734, 125–141.
- Lenat, D.R., 1988. Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. *J. N. Am. Benthol. Soc.* 7, 222–233.
- Lyche-Solheim, A., Feld, C.K., Birk, S., Phillips, G., Carvalho, L., Morabito, G., Mischke, U., Willby, N., Søndergaard, M., Hellsten, S., et al., 2013. Ecological status assessment of European lakes: a comparison of metrics for phytoplankton, macrophytes, benthic invertebrates and fish. *Hydrobiologia* 704, 57–74.
- Lyche-Solheim, A., Phillips, G., Drakare, S., Free, G., Järvinen, M., Skjelbred, B., Tierney, D., Trodd, W., 2014. Water Framework Directive Intercalibration Technical Report: Northern Lake Phytoplankton Ecological Assessment Methods. EUR 26503. Publications Office of the European Union, Luxembourg.
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being: Synthesis*. Island Press/World Resources Institute, Washington, DC.
- Marchetto, A., Padedda, B.M., Mariani, M.A., Lugliè, A., Sechi, N., 2009. A numerical index for evaluating phytoplankton response to changes in nutrient levels in deep Mediterranean reservoirs. *J. Limnol.* 68, 106–121.
- McFarland, B., Carse, F., Sandin, L., 2010. Littoral macroinvertebrates as indicators of lake acidification within the UK. *Aquat. Conserv.* 20, S105–S116.
- Mehner, T., Holmgren, K., Lauridsen, T.L., Jeppesen, E., Diekmann, M., 2007. Lake depth and geographical position modify lake fish assemblages of the European 'Central Plains' ecoregion. *Freshw. Biol.* 52, 2285–2297.
- Melzer, A., 1999. Aquatic macrophytes as tools for lake management. *Hydrobiologia* 395/396, 181–190.
- Miler, O., Porst, G., McGoff, E., Pilotto, F., Donohue, L., Jurca, T., Solimini, A.G., Sandin, L., Irvin, K., Aroviita, J., et al., 2013. Morphological alterations of lake shores in Europe: a multimetric ecological assessment approach using benthic macroinvertebrates. *Ecol. Indic.* 34, 398–410.
- Minns, C.K., Cairns, V.W., Randall, R.G., Moore, J.E., 1994. An index of biotic integrity (IBI) for fish assemblages in the littoral zone of Great Lakes' areas of concern. *Can. J. Fish. Aquat. Sci.* 51, 1804–1822.
- Mischke, U., Carvalho, L., McDonald, C., Skjelbred, B., Lyche-Solheim, A., Phillips, G., de Hoyos, C., Borics, G., Moe, J., Pahissa, J., 2011. WISER Deliverable D3. 1–2: Report on Phytoplankton Bloom Metrics. <http://www.wiser.eu/results/deliverables>
- Moe, S.J., Lyche Solheim, A., Soszka, H., Gołub, M., Hutorowicz, A., Kolada, A., Picińska-Fałtynowicz, J., Białożoz, W., 2015. Integrated assessment of ecological status and misclassification of lakes: the role of uncertainty and index combination rules. *Ecol. Indic.* 48, 605–615.
- Nöges, P., Poikane, S., Köiv, T., Nöges, T., 2010. Effect of chlorophyll sampling design on water quality assessment in thermally stratified lakes. *Hydrobiologia* 649, 157–170.
- Nöges, P., van de Bund, W., Cardoso, A.C., Solimini, A.G., Heiskanen, A.S., 2009. Assessment of the ecological status of European surface waters: a work in progress. *Hydrobiologia* 633, 197–211.
- Olin, M., Rask, M., Ruuhijärvi, J., Tammi, J., 2013. Development and evaluation of the Finnish fish-based lake classification method. *Hydrobiologia* 713, 149–166.
- Padisák, J., Borics, G., Grigorszky, I., Soróczki-Pintér, É., 2006. Use of phytoplankton assemblages for monitoring ecological status of lakes within the Water Framework Directive: the assemblage index. *Hydrobiologia* 553, 1–14.
- Paerl, H.W., Paul, V., 2012. Climate change: links to global expansion of harmful cyanobacteria. *Water Res.* 46, 1349–1363.
- Pall, K., Moser, V., 2009. Austrian index macrophytes (AIM-Module 1) for lakes: a Water Framework Directive compliant assessment system for lakes using aquatic macrophytes. *Hydrobiologia* 633, 83–104.
- Palmer, M.A., Bernhardt, E.S., Allan, J.D., Lake, P.S., Alexander, G., Brooks, S., Carr, J., Clayton, S., Dahm, C.N., Follstad-Shah, J., et al., 2005. Standards for ecologically successful river restoration. *J. Appl. Ecol.* 42, 208–217.
- Pardo, I., Gómez-Rodríguez, C., Wasson, J.G., Owen, R., van de Bund, W., Kelly, M., Bennett, C., Birk, S., Buffagni, A., Erba, S., et al., 2012. The European reference condition concept: a scientific and technical approach to identify minimally-impacted river ecosystems. *Sci. Total Environ.* 420, 33–42.
- Penning, E.W., Dudley, B., Mjelde, M., Hellsten, S., Hanganu, J., Kolada, A., van den Berg, M., Poikane, S., Phillips, G., Willby, N., et al., 2008. Using aquatic macrophyte community indices to define the ecological status of European lakes. *Aquat. Ecol.* 42, 253–264.
- Phillips, G., Lyche-Solheim, A., Skjelbred, B., Mischke, U., Drakare, S., Free, G., Järvinen, M., de Hoyos, C., Morabito, G., Poikane, S., et al., 2013. A phytoplankton trophic index to assess the status for the Water Framework Directive. *Hydrobiologia* 704, 75–95.
- Poikane, S., Alves, M., Argillier, C., van den Berg, M., Buzzi, F., Hoehn, E., de Hoyos, C., Karottki, I., Laplace-Treytore, C., Solheim, A., et al., 2010. Defining chlorophyll-a reference conditions in European lakes. *Environ. Manag.* 45 (6), 1286–1298.
- Poikane, S., Portielje, R., van den Berg, M., Phillips, G., Brucet, S., Carvalho, L., Ott, I., Soszka, H., Van Wichelen, J., 2014. Defining ecologically relevant water quality targets for lakes in Central Europe. *J. Appl. Ecol.* 51, 592–602.
- Rott, E., Pipp, E., Pfister, P., Van Dam, H., Ortler, K., Binder, N., Pall, K., 1999. Indikationslisten für Aufwuchsalgen in österreichischen Fließgewässern.
- Ruse, L., 2010. Classification of nutrient impact on lakes using the chironomid pupal exuvial technique. *Ecol. Indic.* 10, 594–601.
- Sandin, L., Schartau, A.K., Aroviita, J., Carse, F., Colvill, D., Fozzard, I., Goedkoop, W., Göthe, E., Little, R., McFarland, B., et al., 2014. Water Framework Directive Intercalibration Technical Report: Northern Lake Benthic Fauna Ecological Assessment Methods. Report EUR 26510 EN. Publications Office of the European Union, Luxembourg.
- Schaumburg, J., Schranz, C., Hofmann, G., Stelzer, D., Schneider, S., Schmedtje, U., 2004. Macrophytes and phytobenthos as indicators of ecological status in German lakes – a contribution to the implementation of the Water Framework Directive. *Limnologica* 34, 302–314.
- Solimini, A.G., Free, G., Donohue, I., Irvine, K., Pusch, M., Rossaro, B., Sandin, L., Cardoso, A.C., 2006. Using Benthic Macroinvertebrates to Assess Ecological Status of Lakes. EUR 22347 EN. Office for Official Publications of the European Communities, Luxembourg.
- Solimini, A.G., Sandin, L., 2012. The importance of spatial variation of benthic invertebrates for the ecological assessment of European lakes. *Fundam. Appl. Limnol.* 180, 85–89.

- Søndergaard, M., Johansson, L.S., Lauridsen, T.L., Jørgensen, T.B., Liboriussen, L., Jeppesen, E., 2010. Submerged macrophytes as indicators of the ecological quality of lakes. *Freshw. Biol.* 55, 893–908.
- Søndergaard, M., Larsen, S.E., Jørgensen, T.B., Jeppesen, E., 2011. Using chlorophyll-a and cyanobacteria in the ecological classification of lakes. *Ecol. Indic.* 11, 1403–1412.
- Thackeray, S.J., Nöges, P., Dunbar, M.J., Dudley, B.J., Skjelbred, B., Morabito, G., Carvalho, L., Phillips, G., Mischke, U., Catalan, J., et al., 2013. Quantifying uncertainties in biologically-based water quality assessment: a pan-European analysis of lake phytoplankton community metrics. *Ecol. Indic.* 29, 34–47.
- Urbanic, G., 2014. A Littoral Fauna Index for assessing the impact of lakeshore alterations in Alpine lakes. *Ecohydrology* 7 (2), 703–716.
- Whittier, T.R., 1999. Development of IBI metrics for lakes in southern New England. In: Simon, T.P. (Ed.), *Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities*. CRC-Press, pp. 563–582.
- Wiederholm, T., 1980. Use of zoobenthos in lake monitoring. *J. Water Pollut. Control Fed.* 52, 537–547.
- Willén, E., 2007. Växtplankton i sjöar – bedömningsgrunder. Institutionen för Miljöanalys (SLU). Rapport 2007:5. Swedish.
- Willén, E., 2000. Phytoplankton in water quality assessment - An indicator concept. In: Heinonen, P., Giuliano, Z., Van der Beken, A. (Eds.), *Hydrological and Limnological Aspects of Lake Monitoring*. Wiley and Sons Ltd, West Sussex, England, pp. 57–80.
- Wolfram, G., Argillier, C., de Bortoli, J., Buzzi, F., Dalmiglio, A., Dokulil, M.T., Hoehn, E., Marchetto, A., Martinez, P.J., Morabito, G., et al., 2009. Reference conditions and WFD compliant class boundaries for phytoplankton biomass and chlorophyll-a in Alpine lakes. *Hydrobiologia* 633, 45–58.
- Yoder, C.O., Barbour, M.T., 2009. Critical technical elements of state bioassessment programs: a process to evaluate program rigor and comparability. *Environ. Monit. Assess.* 150, 31–42.