

Landscape history, time lags and drivers of change: urban natural grassland remnants in
Potchefstroom, South Africa

Marié J. du Toit · D. Johan Kotze · Sarel S. Cilliers

Marié J. du Toit (Corresponding author) · Sarel S. Cilliers

Unit of Environmental Sciences and Management,

North West University, Potchefstroom Campus,

Private Bag X 6001, Potchefstroom 2520, South Africa

e-mail: 13062638@nwu.ac.za

phone: +2718 299 2518/ fax: +2718 299 2509

D. Johan Kotze

Department of Environmental Sciences,

Niemenkatu 73, FIN-15140,

University of Helsinki, Lahti, Finland

Abstract

Context: The history of the landscape directly affects biotic assemblages, resulting in time lags in species response to disturbances. In highly fragmented environments, this phenomenon often
5 causes extinction debts. However, few studies have been carried out in urban settings.

Objectives: To determine if there are time lags in the response of temperate natural grasslands to urbanization. Does it differ for indigenous species and for species indicative of disturbance and between woody and open grasslands? Do these time lags change over time? What are the
10 potential landscape factors driving these changes? What are the corresponding vegetation changes?

Methods: In 1995 and 2012 vegetation sampling was carried out in 43 urban grassland sites. We calculated six urbanization and landscape measures in a 500 m buffer area surrounding
15 each site for 1938, 1961, 1970, 1994, 1999, 2006, and 2010. We used generalized linear models and model selection to determine which time period best predicted the contemporary species richness patterns.

Results: Woody grasslands showed time lags of 20 - 40 years. Contemporary open grassland
20 communities were, generally, associated with more contemporary landscapes. Altitude and road network density of natural areas were the most frequent predictors of species richness. The importance of the predictors changed between the different models. Species richness, specifically, indigenous herbaceous species, declined from 1995 to 2012.

25 **Conclusions:** The history of urbanization affects contemporary urban vegetation assemblages.
This indicates potential extinction debts, which have important consequences for biodiversity
conservation planning and sustainable future scenarios.

Keywords: urban ecology, urban landscape measures, temperate grasslands, indigenous
30 species, exotic species, disturbance indicator species, extinction debt.

Introduction

The study of patterns of biodiversity in the landscape is of considerable interest in ecology. Theories have been proposed on the drivers of these patterns and numerous studies compare current land-use effects with biotic assemblages (e.g. [Godefroid and Koedam 2003](#); [Hansen and Clevenger 2005](#); [Duguay et al. 2006](#); [Vakhlamova et al. 2014](#)). However, an often ignored perspective is that of the impact of the history of the landscape. A study in France revealed evidence of the irreversible impacts of ancient farming practices on soil nutrient availability and plant diversity of contemporary forested landscapes ([Dupouey et al. 2002](#)). Other studies also provided clear evidence, indicating that many current landscape patterns are directly influenced by past anthropogenic activities (e.g. [Foster 1992](#); [Koerner et al. 1997](#); [Gustavsson et al. 2007](#); [Rhemtulla et al. 2009](#)). Such impacts are called legacy effects and are defined as "*the impacts that previous conditions have on current processes or properties*" ([Monger et al. 2015](#)). Therefore, in order to understand current observed patterns and processes of ecological systems it is critical to incorporate information on past human-mediated disturbances to determine the influence of potential anthropogenic legacy effects ([Hamburg and Sanford 1986](#)).

Not only does history affect ecosystems, but communities are often slow in reacting to disturbances, and time lags occur between causes and their effects (e.g. [Magnuson 1990](#); [Burel 1993](#); [Tilman et al. 1994](#); [Foster et al. 2003](#); [Gustavsson et al. 2007](#); [Koyanagi et al. 2012](#)). Time lags occur because processes operating in landscapes take time, relict species can remain in altered landscapes, movements across the landscape may be slow, some biological events require certain rare concurrent conditions to take place, and time lags can be accumulated by a sequence of events ([Magnuson 1990](#)). Species respond in different ways to disturbances based on their life history characteristics ([Ewers and Didham 2006](#); [Chiba et al. 2009](#); [Kuussaari et al. 2009](#)), therefore, observed time lags vary. In vegetation, longer living trees show longer time

lags than grassland and herbaceous plants ([Davis 1986](#)). Time lags of 150 – 250 years were estimated for Alaskan forests in response to climate change ([Chapin and Starfield 1997](#); [Lloyd 2005](#)), while semi-natural grasslands in Sweden displayed a 50 -100 year time lag in species diversity changes due to changes in habitat connectivity ([Lindborg and Eriksson 2004](#)). Similarly, [Helm et al. \(2006\)](#) showed that in alvar grasslands of Estonia, current habitat specialist species were best described by habitat configurations 70 years earlier. In Japanese semi-natural grasslands, [Koyanagi et al. \(2012\)](#) found that time lags differed between 30 – 50 years depending on the specific plant species functional traits.

In studying time lags, it is important to distinguish between research on exotic and indigenous species. Studies of time lags in exotic species usually pertain to the time lag between introduction and eventual establishment and invasion; the focus is generally on the increase of species richness with time ([Sakai et al. 2001](#); [Crooks 2005](#); [Daehler 2009](#)). Studies focusing on indigenous species, however, invariably investigate time lags in the decrease of species richness, i.e. declines in populations or even species extinctions ([Brooks et al. 1999](#); [Hanski and Ovaskainen 2002](#); [Kuussaari et al. 2009](#)). Disturbances not only influence vegetation composition but severe disturbances, such as habitat destruction or fragmentation, can push species to extinction ([Tilman et al. 2001](#)). However, due to the existence of species time lags in response to these disturbances, delayed extinctions can occur (i.e. extinction debts, [Tilman et al. 1994](#); [Kuussaari et al. 2009](#)). Therefore, a sobering picture is revealed of current observed extant vegetation potentially already doomed by past disturbances; a situation calling for critical reflection in fragmented contextual settings.

In urban ecological studies, there is a paucity of studies on anthropogenic legacy effects and time lags. The limited number of studies documenting time lags in vegetation patterns in urban areas almost exclusively pertains to the relationship between neighbourhood social characteristics and woody vegetation cover ([Pickett et al. 2008](#); [Boone et al. 2009](#); [Luck et al. 2009](#); [Grove et al. 2014](#); [Locke and Baine 2014](#)). However, these do not document responses

of vegetation to a disturbance, but rather identify corresponding social contextual plant
85 establishment periods. Nonetheless, there are a few studies on extinction debts in urban areas.
[Hahs et al. \(2009\)](#) studied plant extinction rates in 22 cities worldwide and determined that
contemporary urban areas potentially have large extinction debts. A subsequent study of
European countries determined that “*irrespective of recent conservation actions, large scale
risks to biodiversity lag considerably behind contemporary levels of socioeconomic pressures.*
90 *The negative impact of human activities on current biodiversity will not become fully realized
until several decades into the future*” ([Dullinger et al. 2013](#)).

This study aims to (i) enhance research on time lags in the response of urban remnant
natural vegetation to anthropogenic disturbances; (ii) address the paucity of urban re-visitation
studies ([Williams et al. 2005](#)), (iii) add to long-term vegetation change studies in new locations
95 in order to enhance global knowledge; and (iv) enhance urban research in temperate natural
grasslands. Moreover, no such study has been carried out in Africa. We answer the call in
urban ecological literature to include temporal dynamics ([Ramalho and Hobbs 2012](#)) and to
more fully understand mechanisms driving current vegetation patterns ([McDonnell and Hahs
2013](#)). The main objective is to determine if there are time lags in the response of temperate
100 natural grasslands to urbanization. More specifically, we asked (i) does it differ for indigenous
species and disturbance indicator species and between woody grassland communities and open
grasslands? (ii) Do these time lags change over time? (iii) What are the potential landscape
factors driving these changes? Additionally, (iv) what are the corresponding vegetation
changes over a 17-year time period?

105 We hypothesize that (a) indigenous species patterns will correspond to past landscape
patterns, (b) disturbance indicator species richness will correspond to periods of rapid urban
growth and will have longer time lags, (c) woody grassland communities will have longer time
lags than open grasslands, (d) intensification of disturbances (such as rapid urban growth) will

shorten observed time lags, and (e) indigenous species richness will decline while exotics and
110 species indicative of disturbances will increase with urbanization.

Methods

Study area and history

115
The study was conducted in the city of Potchefstroom (26° 42' 53'' S; 27° 05' 49'' E) and its
immediate natural surroundings, situated in the North-West Province of South Africa (Fig. 1a).
The city covers a 55 km² area at an elevation of 1350 m, with a population of approximately
250 000 ([Tlokwe City Council 2015](#)). The city experiences average temperatures of 0°C in
120 winter with frequent frost and 30°C in summer months. The mean annual rainfall is about 550
mm received mainly in summer months (Supplementary Information Fig. S1). Potchefstroom
was established in its current location in 1841 ([van den Bergh 1992](#)). Simultaneous to town
settlement was the delineation of farms around the town boundary ([van den Bergh 1990](#)).
However, along the western boundary of the town, a town commonage area was demarcated
125 for use as grazing (Fig. 1b). The commonage reached a maximum area of 25 434 ha by 1889
due to the continuous demand for more grazing ([van den Bergh 1992](#)). The rapid advance of
the motorcar from 1929 to 1945 signalled the end of the large-scale use of the town
commonage for grazing purposes ([Smit 1989](#)). Therefore, the commonage area was continually
grazed for about 100 years. Thereafter, urbanization and its associated disturbances became the
130 major influence. In subsequent years, most of the urban development expanded into the town
commonage area, as the most readily available municipal land (Fig. 1c). A crucial factor in the
urban development of the town commonage area was the Group Areas Act of 1950, mandating
the compulsory zoning of separate racial residential areas. Development of these residential
areas of mostly poor residents started in 1956 in the town commonage area ([Neser 1967](#)). The

135 democratic elections of 1994 and the lift of urban restriction laws resulted in an immense
growth of urban residential areas of the town commonage area (Fig. 1c).

#Figure 1 approximately here#

140 *Vegetation sampling*

Potchefstroom is situated in temperate natural grasslands at the confluence of three vegetation types: the Rand Highveld Grassland, Carletonville Dolomite Grassland and the Andesite Mountain Bushveld ([Mucina and Rutherford 2006](#)), the latter being Savanna elements found on
145 hills and ridges in the area (Supplementary Information Fig. S2), described in this study as woody grassland communities. Vegetation sampling sites are all situated in the town commonage area (Fig. 1b) with most on the urban-rural fringe and some existing as remnant fragments within residential areas. Extensive vegetation sampling of the Potchefstroom area in 1995-6 ([Cilliers et al. 1999](#)) allowed us to compare vegetation composition before and after the
150 rapid urban expansion period (Fig. 1d and e). The original surveys were phytosociological relevés sampled using Braun-Blanquet cover abundance values ([Mueller-Dombois and Ellenberg 1974](#)). Seventy-eight sample plots were surveyed in 1995-6 of which 43 were re-surveyed in 2012 comprising of 26 woody grassland sites and 17 open grassland sites. Eight of the 78 sites were lost to urban development, the precise location of 15 sites are unknown and of
155 the remaining sites we only chose those that were historically untransformed, i.e. we excluded old fields and any other direct anthropogenically-disturbed sites. Most of the sites are situated in the western part of the study area in an unbroken extensive grassland area, some of which later became fragmented. The rest of the sites are located on small remnant areas fragmented due to urbanization (Supplementary Information Fig. S3). Resampled site locations were as
160 close as possible to the original sites, using datasheet site descriptions, aerial photographs on

which some of the site locations were marked, and one of the author's (Cilliers) own recollections. Woody grasslands were sampled in 10 x 10 m sample plots and open grasslands in 4 x 4 m sample plots ([Bredenkamp and Theron 1978](#)), with one plot per site. Woody communities are naturally occurring habitat 'islands' in the grassland matrix, mainly situated on hills and ridges forming part of the Andesite Mountain Bushveld vegetation type ([Mucina and Rutherford 2006](#)) with the dominant taxa represented by trees and large shrubs. However, historically open grasslands can also be transformed by bush encroachment due to higher rainfall frequencies ([Ward 2005](#)), which allows tree seedlings to survive in the grasslands or as a result of over-grazing and unnatural fire regimes that stimulate tree growth when competition with grassland species is lower ([Kraaij and Ward 2006](#); [Van Auken 2009](#)).

The percentage cover of all species within the sample plots was recorded. The 1995 Braun-Blanquet cover abundance data were converted to mean percentage values ([Eckhardt et al. 1993](#)) to allow direct comparison between the resampled and original data. For this paper, only the perennial species data were used as per [Williams et al. \(2005\)](#) to reduce the potential for recording false absences of species. We calculated indigenous species richness (ISR), exotic species richness (ESR) and species richness of plants that indicate disturbance (DSR) per sample plot, as well as total species richness, percentage indigenous species and percentage indigenous species cover. However, the last three measures were excluded from our analyses because they all correlated with either ISR or ESR. For instance, ISR correlated highly and significantly with total species richness in both years and for both woody and open grassland communities (Person's $r \geq 0.98$). Furthermore, ESR correlated highly with percentage indigenous species ($-0.86 < r < -0.97$) and percentage indigenous species cover ($-0.68 < r < -0.83$). To calculate DSR, we identified 31 (indigenous and exotic) species from the recorded species occurrences that indicate grassland vegetation in poor condition, specifically overgrazing and trampling, pioneer species, weeds, invaders, and bush encroachment (e.g. [Louw 1951](#); [Bredenkamp et al. 1989](#); [Bezuidenhout et al. 1994a](#); [Cilliers and Bredenkamp](#)

[1999](#); [Cilliers et al. 1999](#)). Because humans disturb natural areas in various ways, through trampling, plant removal, rubbish dumping etc., we wanted to determine whether these species that flourish in disturbed environments increase with increasing urbanization and if their richness patterns correspond to urban landscape measures. These disturbance indicator species were identified from published studies of Potchefstroom and its surrounding regional grassland setting (Supplementary Information Table 1).

Urban landscape measures

We digitized 500 m buffer areas around each site to quantify the landscape matrix. Optimal matrix scales have not been determined to study the effects of urbanization on vegetation ([Dauber et al. 2003](#)), therefore we decided on 500 m as some authors state this a suitable buffer area (e.g. [Tscharntke et al. 2005](#); [Duguay et al. 2006](#); [Kleijn et al. 2009](#); [Vakhlamova et al. 2014](#)). Additionally, since we are interested in the anthropogenic effects on vegetation dynamics and the area of interest is mostly inhabited by poor residents, we analysed firewood collection distances. The field data sheets of Cilliers in 1995-6 revealed that sites where firewood was gathered were all situated within 200-500 m from residential areas. The limited literature on grassland species dispersal also indicate that most dispersal of grassland species, excluding long-distance dispersal by animals, occur well within the 500 m limit of our buffer zone ([O'Connor 1991](#); [Morgan 1995](#)).

Measures were calculated for the time periods 1938, 1961, 1970, 1994, 1999, 2006, and 2010 as these were the dates for which we could readily obtain aerial or satellite imagery. For 1938-1994, we georeferenced digital aerial photographs, procured from government archives (www.ngi.gov.za), in ArcMap 10 ([ESRI 2010](#)) using digital topographical maps of periods nearest to the imagery dates as reference material. For the periods 1999-2010 we obtained orthorectified satellite imagery. The chosen measures for this study included: age of

urbanization (Age), altitude, road network density of natural areas (RNDN), percentage natural area (PN), density of dwellings (CD) and landscape diversity (H'). These measures were
215 calculated in ArcMap 10, additionally using the Hawth's Analysis Tools version 3.27 ([Beyer 2006](#)) add-in to calculate RNDN, CD and H. To determine the age of urbanization we used the aerial and satellite images to determine when any form of urban buildings first occurred in the 500 m buffer, for example if urban infrastructure occurred in 1938, the age of urbanization in the 2012 models was 74 years. Because we could not accurately state in which year
220 urbanization occurred before 1938, all the urbanized sites in the 1938 GLM models were given the age of 1 year (in a sense presence/absence data). We included altitude (recorded with a GPS device: 1332 - 1432 m) since the sampling site locations varies from plains to foot slopes and small hills and ridges.

The effects of roads and trampling on vegetation are well documented ([Cilliers and Bredenkamp 2000](#); [Gelbard and Belnap 2003](#); [Hansen and Clewenger 2005](#); [Kissling et al. 2009](#); [Deng et al. 2011](#)). We specifically wanted to determine the effect of roads (tar and dirt)
225 and footpaths on the vegetation dynamics of natural areas as an index of fragmentation and disturbance, i.e. a proxy for habitat quality. Therefore, all roads and footpaths inside and traversing the natural areas were digitized. We modified the standard road network density measure (e.g. [Hahs and McDonnell 2006](#)) by dividing only the digitized roads inside the
230 natural area by the size of the natural area inside each 500 m buffer area. Thus, road network density of natural areas (RNDN) was measured as the length of roads per m² of natural areas. Percentage natural areas was simply the proportion of natural areas inside the 500 m buffer area. We used the measure density of dwellings as a surrogate for population density for two
235 reasons; first we could accurately digitize it for each time period as there are no reciprocal fine scale census data available for any of the time periods. Second, comparison in the literature showed that in PCA and FA analyses where both measures were used, it had similar loadings as the population density measure ([Hahs and McDonnell 2006](#); [Andersson et al. 2009](#); [du Toit](#)

[and Cilliers 2011](#)). Thus, the density of dwellings (CD) was calculated as the count of all
240 buildings per buffer area. Landscape diversity (H') was calculated using the proportion of
different habitats analysed with the Shannon Index ([Shannon 1948](#)) (after [Romme and Knight
1982](#); [Purschke et al. 2012](#); [Janišová et al. 2014](#)). The buffer areas were classified by manually
digitizing it from aerial and satellite images into the following classes: mine dumps and
excavations, natural, urban, cultivated land, plantations and water. The percentage area per
245 class was then used as input into Primer 6 ([Clarke and Gorley 2006](#)) to calculate the Shannon
index.

Data analysis

250 All analyses were performed with the statistical package R version 3.0.2 ([R Core Team 2013](#)).
Four sets of analyses were performed per vegetation type (woody and open grassland); two for
the 1995 vegetation data (ISR, DSR) and two for the 2012 vegetation data (ISR, DSR). We
were unable to analyse ESR due to very low numbers of exotic species in the dataset. For each
analysis, the following steps were followed: In the first step, spatial autocorrelation between
255 the sample plots were tested using Moran's I in the ape version 3.1-4 package in R. Site
longitude and latitude were used to perform this test. The presence of spatial structure was
further evaluated by comparing the AIC value of a null model with no covariance structure, to
AIC values of models including various covariate structures (including exponential, Gaussian,
spherical, linear and rational quadratics that are based on site longitude and latitude values)
260 (the procedure is explained in full here:
http://statistics.ats.ucla.edu/stat/r/faq/variogram_lmme.htm). The lme command in the nlme
version 3.1-111 package in R was used to perform this analysis. No indication of spatial
structure was detected in any of the datasets.

In the second step, generalized linear models were performed for each of the eight
265 vegetation parameters. The response variables were modelled following a Poisson error
distribution. Predictor variables for all models included altitude, age of urbanization (Age),
percentage natural area (PN), road network density of natural areas (RNDN), density of
dwellings (CD) and landscape diversity (H'). For the data collected in 1995, each response
variable (e.g., ISR of woody grasslands in 1995) was modelled against all predictor variables
270 for 1938, then for 1961, then 1970 and finally for 1994. Model selection was performed for
each year (e.g., ISR woody grasslands for 1995 against the 1938 landscape, ISR woody
grasslands for 1995 against the 1961 landscape, etc.) by removing less significant variables one
at a time until only variables of $p < 0.2$ were left in the models. AIC values of these final
models for, say ISR woody grasslands for 1995, were compared between the four years (1938,
275 1961, 1970, 1994) and the one with the lowest AIC was selected as the optimal model for this
particular vegetation parameter. For the data collected in 2012, the same procedure was
followed as described above, except that the response variables here were modelled against all
predictor variables for 1938, 1961, 1970, 1994, 1999, 2006, and 2010.

All variables were on a continuous scale, but a few had to be converted to categorical
280 variables due to highly irregular values in the variable. This was particularly true for variables
measured in 1938 and 1961, and a few variables in 1970.

Results

285 *Landscape matrix change*

The percentage natural area declined steadily with time; however woody grasslands situated,
on average, at higher altitudes, had consistently higher average percentage natural areas left
(Fig. 2a). The road network density of natural areas (RNDN) indicated that open grasslands

290 had consistently higher densities of footpaths and roads, except in 2006. RNDN in both habitats increased steadily with a noticeable increase in 1994 and then a drastic increase in 2010, consistent with rapid urban development in the area (Fig. 2b). The density of dwellings measure showed a marked increase in population density since 1994 (Fig. 2c). Open grassland landscape diversity was consistently more complex than woody grassland landscape diversity and both increased from 1938 to 2010 (Fig. 2d).

#Figure 2 approximately here#

Vegetation community time lags

300 Results of the GLM models indicate that in 1995 and 2012 the open grassland vegetation (ISR) was best predicted by the 1994 and 2010 landscape variables as they returned the lowest AIC value (Table 1). Since we did not have landscape data for 1995 and 2012 this can either be interpreted as a 1 and 2 year time lag, respectively, or as no time lag as those are the most current landscape variables available. The 1995 DSR, on the other hand, was best explained by 305 the 1970s landscape, indicating a 25 year lag. In 2012, this changed to a six year lag, best predicted by the 2006 landscape. The woody grassland richness patterns (ISR and DSR), however, showed longer time lags (Table 1). In 1995, ISR was best predicted by the 1970 landscape, indicating a 25 year time lag. In 2012, the best model was of the 1994 landscape, 310 indicating an 18 year time lag. The 1995 DSR was best predicted by the 1961 landscape, a 34 year lag. This changed to 42 years in 2012 as the 1970 landscape represented the best fit model.

Altitude and road network density of natural areas (Table 1, Fig. 3) were the most frequent predictors of species richness. Five of the eight vegetation models had a statistically significant response to both altitude and RNDN. Open grassland species richness increased 315

with altitude and RNDN, while woody grassland species richness decreased with altitude and RNDN (Fig. 3). Percentage natural area was statistically significant in four models with open grassland communities decreasing in species richness with increasing PN, while the woody grassland DSR in 2012 increased with PN (Table 1, Fig. 4). The density of dwellings and landscape diversity were each significant in three models. The age of urbanization was the least important variable with only one statistically significant contribution in the 2012 open grassland ISR model (Table 1).

#Table 1 approximately here#

325

#Figure 3 approximately here#

#Figure 4 approximately here#

330 *Vegetation change between 1995 and 2012*

The open grassland sites gained 19 species between 1995 and 2012, but lost 58 species, with a net loss of 39. The woody grasslands gained 26, but lost 51, with a net loss of 25. Collectively, 24 of the species (3 grasses, 10 herbs, 10 shrubs—1 exotic, 1 tree) lost in the open and woody grasslands did not occur in any habitats surveyed in the recent literature ([du Toit 2009](#); [Jansen van Rensburg 2010](#); [van der Walt 2013](#)), and have a potential local extinction risk (Supplementary Information Table 2). These 24 species were all rare, occurring only in 3 or fewer sites. None of the species gained are new additions to the area. Species were either recorded in the alternate habitat in 1995 (e.g. if recorded in the 2012 woody grasslands it was present in the 1995 open grasslands) or listed in older literature of the Potchefstroom area ([Bezuidenhout and Bredenkamp 1990, 1991](#); [Bezuidenhout et al. 1994b](#); [Bredenkamp et al.](#)

340

[1994](#); [Retief and Herman 1997](#); [Cilliers et al. 1998](#); [Cilliers and Bredekamp 1998, 1999](#); [Cilliers et al. 1999](#); [Cilliers and Bredekamp 2000](#)). Both habitats in 1995 had 20% rare species (recorded in a single site and habitat only). This increased to 24 % for the open
345 grasslands and 22 % in the woody grasslands in 2012. Growth forms between years remained essentially the same, except that herbaceous species declined drastically (26 in open and 19 in woody grasslands, respectively) (Table 2). Exotic species declined in the open grasslands and increased in the woody grasslands, with very low percentage cover throughout the study periods. Sites that displayed high species richness in the open grasslands in 1995, also did so in
350 2012 (Supplementary Information Fig. S4a). Woody sites, however, had distinct variation with time (Supplementary Information Fig. S4b) For the disturbance indicator species (Table 2), the species richness declined in the open grasslands and stayed the same in the woody grasslands, with very low percentage covers throughout the study periods. All 14 DSR species recorded in the 2012 open grasslands were also present in 1995. The woody sites shared 22 DSR species
355 and the two newly recorded species in 2012 were both present in the 1995 and 2012 open grasslands.

Table 2 approximately here#

360 **Discussion**

Time lags were observed for both the open and woody grassland vegetation. Time lags observed for the woody grasslands preceded major urban development in the study area. The township of Ikageng was established in 1957-1963 and in 1969 resettlement started in Promosa
365 ([Neser 1967](#)). In the period between 1961 and 1970 the township of Mohadin was established, and 1994 preceded the start of rapid urban growth (see Fig. 1). The woody grassland ISR lags remained constant at an average of 21 years, moving 'forward' with urban development. The

woody DSR averaged 38 years, corresponding to the first major urban development in the town commonage area.

370 The open grassland ISR patterns can either be interpreted as a 1 and 2 year lag, respectively, or as no time lag because of an absence of data for the specific years in which the sampling was done. A study by [Adriaens et al. \(2006\)](#) on semi-natural grasslands in Belgium found no time lags in connection with past habitat area and connectivity, however they also had a 19 year gap between their current landscape variables and the nearest historical one.

375 [O'Connor \(1991\)](#) researched local extinctions in perennial grasslands and reported that many palatable grass species are obligate seed producers that produce low numbers of seeds that are poorly dispersed. The seed bank of these 'extinction-prone' species need annual seed input as the seeds only survive for 2-3 years in the soil. If the extant plant is removed and the soil seed bank depleted or seedlings fail to establish due to disturbances, predation or low rainfall the

380 species can quickly become locally extinct ([O'Connor 1991](#)). Moreover, a soil seed bank experiment in temperate natural *Themeda triandra* grasslands in Australia found that perennial indigenous species (specifically herbs) were almost completely absent from the soil seed bank ([Morgan 1998](#)). This implies quick species turnover times in short-lived grassland species, which in combination with described hyperdynamism of highly fragmented areas ([Laurance](#)

385 [2002](#)) most probably accounts for the short or absent time lags observed in the open grassland ISR patterns.

 It is important to note that the accuracy of the time lag calculation is restricted by the time periods used. This is especially pertinent for the evaluation of the open grassland models. After the 1970 landscape there is a 24 year gap wherein we could not obtain accurate aerial

390 imagery before the 1994 dataset. All the models indicate comparative time lags between the two sampling periods, except the open grassland DSR models. The 1995 DSR corresponded to the 1970 landscape indicating a 25 year lag and the 2012 DSR corresponded to 2006 indicating a 6 year lag. However, close scrutiny of our results indicates that both of the time lags of the

DSR models in the open grasslands are indicative of the last time period before the most
395 current landscape variables, i.e. in 1995 it was the 1970 landscape and in 2012 the 2006
landscape. Therefore, the 25 year gap displayed by the 1995 model could be inaccurate as it is
the closest available historical landscape prior to the current landscape. Therefore, the 1995
open grassland DSR patterns could probably have better corresponded to the observed 6 year
lag of the 2012 landscape if we had more landscape datasets in the 24 year interval.

400 Time lags observed for the open and woody grassland DSR patterns both exceeded
that observed for the ISR patterns. Ecosystem response lags behind disturbances ([Magnuson
1990](#); [Hayashida 2005](#)). For instance, the disturbance indicator species used in this study are
pioneers or bush encroachers and their presence are the first indication that a site experienced
disturbances. The resulting community changes and direct disturbance impacts in turn
405 influence the rest of the community. It is therefore expected that disturbance indicator species
patterns will always lag behind extant indigenous species patterns. The presence of disturbance
indicator species can subsequently potentially serve as an early warning sign that other species
losses will occur. There are no equivalent time lag studies carried out in temperate natural
grasslands with which to compare our findings. However, the observed time lags for the woody
410 community are relatively close to the 30-50 year time lags of Japanese semi-natural grasslands
([Koyanagi et al. 2012](#)). Other studies in semi-natural grasslands document 50-100 years
([Lindborg and Eriksson 2004](#)) and 70 years for Estonian semi-natural grasslands ([Helm et al.
2006](#)).

The 1995 open grassland indigenous species richness patterns were best predicted by
415 the 1994 landscape variables where higher altitudes, older urbanization influences with smaller
percentage natural areas in the buffer, higher road network densities, and more heterogeneous
landscapes had the highest ISR (Fig. 5a and b). The 2012 ISR patterns were best predicted by
2010 landscape variables; higher altitudes, older urbanization influences, higher road network
densities, population densities and landscape diversity predicted higher ISR. The increase in

420 species richness with a decrease in percentage natural areas might be explained by the fact that
species richness was positively related to urbanization age. Therefore, older sites closer to the
urban area had higher quality habitats (as indicated by higher ISR) (Fig. 5b). The town
commonage area in which the sites further from the urban area are situated was continuously
grazed for approximately 100 years, thus the sites close to the urban area would have been
425 largely protected from grazers. This then could explain the counter-intuitive negative
relationship with percentage natural areas in the buffer. Moreover, of the two variables only
urbanization age remained significant in the 2012 ISR model. This relationship with
urbanization age also explains the positive relationship with RNDN, as the vegetation
fragments are smaller and many footpaths traverse them between the residential areas (Fig. 5b).
430 The same argument holds for the positive relationship of the 2012 ISR with the density of
dwellings and landscape diversity. Further evidence is also provided when examining the ISR
pattern changes per site from 1995 to 2012 (Supplementary Information Fig. S4a). In the open
grassland DSR models, only a single variable best predicted the patterns in 1995 and 2012. The
1995 model indicated that lower percentage natural area sites of 1970 had higher DSR (Fig. 5c)
435 whereas sites with higher road network densities in 2006 predicted sites with higher DSR in
2012 (Fig. 5d).

For the indigenous species richness of woody grasslands, on the other hand, lower
altitudes with smaller percentage natural areas in the buffer, but more homogenous and low
dwelling densities in 1970 predicted sites with higher ISR in 1995 (Fig. 6a). In 2012, however,
440 lower altitudes, lower road network densities but higher dwelling densities in 1994 predicted
sites with higher ISR (Fig 6b). The negative relationship with RNDN for both years can be
explained by the fact that the footpaths in woody sites are indicative not only of thoroughfares
but of direct disturbances i.e. firewood harvesting, whereas most of the footpaths in open
grassland sites are only used as thoroughfares. The negative relationship with altitude might
445 also partly be explained by the ecotone concept ([Risser 1995](#); [Kark and van Rensburg 2006](#)),

where lower lying woody sites have more open grassland species as the community changes from woody to open grasslands enhancing species richness of these sites. Urban expansion mostly took place on lower lying areas, which in combination with higher species richness of ecotones might explain the negative relationship with PN in 1995 and the subsequent positive
450 relationship with urbanization age and density of dwellings in 2012 (Fig. 6b). The percentage natural area in the buffer, as with the open grasslands, was again not important in the 2012 model.

#Figure 5 approximately here#

455

The woody grassland DSR in 1995 was best predicted by the 1961 landscape, indicating that lower altitude, lower road network density, and more homogenous sites had higher species richness (Fig. 6c). In 2012, lower altitudes with less natural area in the buffer had higher DSR values (Fig. 6d). The sites nearest to urban areas are on average situated at lower altitudes, and
460 the contradictory lower road network densities can be ascribed to the fact that RNDN values for 1961 was highest in sites situated in small grassland fragments on the opposite side of the densely populated informal residential areas (Fig. 6c and Supplementary Information Fig. S3). Many footpaths traversed this area, due to the vicinity of military grounds and a water purification plant. In contrast, sites in the non-fragmented homogenous western part of the
465 study area—despite the nearness of many to urban areas— still had fewer roads and footpaths in relation to the size of natural areas.

#Figure 6 approximately here#

470 The loss of indigenous species in both the woody and open grassland vegetation is in line with documented loss of species in urban areas with time ([Chocholoušková and Pyšek 2003](#); [Tait et](#)

[al. 2005](#); [Williams et al. 2006](#); [Stehlik et al. 2007](#)). Long time-lags in the response of woody grassland communities might be the reason why these communities lost fewer species and shared more species between the time periods than the open grasslands. The decline in exotic species and disturbance species richness of the open grasslands seem to indicate that these sites do not receive as much direct disturbances as that of woody sites, which are particularly targeted for firewood collection ([Madubansi and Shackleton 2007](#)). Of specific concern is the major loss of herbaceous species in both habitat types. Moreover, despite the presence of time lags to indicate possible extinction debts, the high proportion of rare species and its slight increase from 1995 to 2012 for both the open and woody grasslands might also be an extinction debt signature ([Hanski and Ovaskainen 2002](#)).

Natural stressors in these grasslands are grazing, climate and fire. In our study area, fire is largely suppressed at sites in the city (however mowing can simulate the effects of fire due to biomass removal ([Williams et al. 2006](#); [Cilliers et al. 2008](#))) and those on the fringe experience irregular burning. Anthropogenically-caused and suppressed fires together with natural fires cause changed disturbance regimes, which impact indigenous species richness ([Williams et al. 2006](#)) and bush encroachment ([Friedel 1987](#)). In their study, [Williams et al. \(2006\)](#) linked changed fire regimes (as a proxy of habitat quality) to declines in indigenous species and indicated that habitat quality rather than patch size or isolation determined local extinction risks of grassland species. Similarly, in explaining the distribution of herbaceous species in woodlands, [Dupré and Ehrlén \(2002\)](#) found that habitat quality (measured by soil properties) was more important than spatial configuration. However, fragmentation effects can be masked by confounding factors ([Ewers and Didham 2006](#)). In the current study, some small fragmented areas harboured higher diversity than sites situated in the extensive grassland area in the western part of the study area, emphasizing the conservation value of smaller natural remnants ([Lawson et al. 2008](#)). However, the overall loss of species from 1995 to 2012 in almost every site indicates the negative effects of urban anthropogenic disturbances and subsequent

fragmentation on vegetation communities, as also indicated by [Krauss et al. \(2010\)](#). Other important factors influencing vegetation patterns are precipitation and climate, not explicitly incorporated in this study. However, in their study on semi-arid sagebrush steppe [Anderson and Inouye \(2001\)](#) found that vegetation cover and density lagged 2-4 years in response to precipitation. The rainfall patterns prior to the sampling periods in this study indicate that both periods received rainfall close to the overall mean of the study area (Supplementary Information Fig. S1). Therefore, despite above average rainfall during the 1995-6 sampling period this would probably not have profoundly influenced cover abundance at the time of sampling (Supplementary Information Fig. S1). Results indicate that urban areas are complex and that different components of plant communities react differently to urbanization-related disturbances. This study is unique in the sense that we had two sampling periods with which to test time lags and as a direct result thereof we highlighted the importance of the selection of landscape measures. The relevance of the measures changed in explaining the observed vegetation patterns for different sampling periods. Moreover, between the different years, some of the relationships between species richness and the variables changed (e.g. from negative to positive). The urban landscape measures were determined for a 500 m radius buffer and we acknowledge that the scale we used to determine the landscape measures can have an impact on the reliability of our results. The nature of the sites that were chosen, i.e. relevés (representative sample plots) instead of entire fragmented areas and the specific landscape measures used can also influence our results; the possibility arises that grasslands might respond better to other variables not measured or at different scales.

520 **Conclusions**

Our first hypothesis that indigenous species richness patterns correspond to past landscape patterns was partly confirmed, as indigenous woody grassland species richness showed long

time lags, while open grassland had very short or even no time lags. The second hypothesis
525 that disturbance indicator species correspond to periods of rapid growth and exhibit longer time
lags was partly supported, as woody DSR was linked to initial urban expansion periods (the
establishment of Ikageng and Promosa) but not to rapid urbanization after 1994. However,
DSR time lags were consistently longer than ISR time lags for all habitats. We believe that
DSR time lags can act as an early warning system and proxy for changes to local vegetation
530 communities. The third hypothesis that woody grassland will have longer time lags was
confirmed. The fourth hypothesis the time lags will shorten as urbanization intensifies was not
confirmed, except in the case of grassland DSR, however this could be as a result of the
available data. Of specific significance is that despite rapid urbanization in the area, all other
models showed relatively constant time lags. The last hypothesis was partly supported as
535 indigenous species richness declined, however exotic species richness declined in open
grassland sites but increased in woody grasslands, whereas disturbance species richness also
declined in open grasslands but remained the same in woody grasslands.

Our study is the first to specifically test for time lags in the response of remnant natural
vegetation to urbanization. Additionally, we tested two different vegetation sampling periods
540 indicating potentially constant time lag periods in the response of remnant natural vegetation to
urbanization influences, despite rapid urban expansion in the intervening years. The modified
measure of road network density of natural areas shows potential as it was one of the most
frequent predictors in the models. However, it will have more ecological utility if equally sized
sites are compared with one another. The results of this study highlight the importance of
545 legacy effects on patterns observed in contemporary communities ([Burel 1993](#); [Ewers and
Didham 2006](#)). Indigenous species losses over the 17-year period and the presence of clear
time lags in the woody grasslands indicate that these communities at least have a probable
extinction debt ([Kuussaari et al. 2009](#); [Krauss et al. 2010](#)) to pay in the future.

550 **Acknowledgements**

We would like to thank the South African Weather Service for long-term climate data of Potchefstroom and the National Research Foundation (NRF) for financial assistance towards the studies of MJDT and incentive funding for SSC.

555 **Tables**

Table 1 GLM results. Model coefficients (\pm SE) are presented, as well as p-values. Int = model intercept, Alt = altitude, Age = age of urbanisation, PN = percentage natural area, RNDN = road network density of natural areas, CD = density (the number) of dwellings, H' = Shannon's diversity index of landscape diversity. AIC = Akaike's Information Criterion. AIC values represent the model with the highest AIC value (and year), and the model with the lowest AIC (and year, in bold). GLM model coefficients and SE and p values presented here are from models with the lowest AIC values. L2 and L3 represent the second and third levels of a variable classified as a factor. See Data analysis.

| | Int | Alt | Age | PN | RNDN | CD | H' | AIC (model year) |
|-----------------------|------------|------------|------------|-----------|-------------|-----------|-----------|-------------------------|
| Open grassland | | | | | | | | |
| 1995 | | | | | | | | |
| ISR | -19.992 | 0.017 | 0.006 | -0.007 | 39.058 | | | 153.40 (1938) -> |
| | (5.030) | (0.004) | (0.004) | (0.004) | (14.328) | | | 135.38 (1994) |
| | <0.001 | <0.001 | 0.133 | 0.045 | 0.006 | | | |
| DSR | 3.388 | | | -0.026 | | | | 70.91 (1994) -> |
| | (0.387) | | | (0.005) | | | | 67.51 (1970) |
| | <0.001 | | | <0.001 | | | | |

2012

| | | | | | | | | |
|-----|---------|---------|---------|--|----------|----------|---------|------------------------|
| ISR | -20.550 | 0.016 | 0.011 | | 18.750 | 0.001 | 0.752 | 130.06 (1938) -> |
| | (5.139) | (0.004) | (0.003) | | (8.713) | (0.0003) | (0.376) | 116.21 (2010) |
| | <0.001 | <0.001 | 0.002 | | 0.031 | 0.003 | 0.046 | |
| DSR | -0.935 | | | | 107.187 | | | 69.58 (1938) -> |
| | (0.507) | | | | (25.946) | | | 59.40 (2006) |
| | 0.065 | | | | <0.001 | | | |

Woody grassland

1995

| | | | | | | | | |
|-----|---------|---------|--|------------|----------|------------|---------|------------------------|
| ISR | 20.647 | -0.012 | | -0.285 | -43.493 | -0.460 | -1.429 | 211.05 (1938) -> |
| | (3.585) | (0.003) | | (0.211) | (24.447) | (0.144) | (0.355) | 178.76 (1970) |
| | <0.001 | <0.001 | | 0.175 (L2) | 0.075 | 0.001 (L2) | <0.001 | |
| | | | | -0.553 | | -0.126 | | |
| | | | | (0.228) | | (0.239) | | |
| | | | | 0.015 (L3) | | 0.600 (L3) | | |
| DSR | 53.572 | -0.037 | | | -68.388 | | -0.731 | 157.33 (1970) -> |

| | | | | | |
|---------|---------|--|----------|------------|------------------------|
| (9.026) | (0.006) | | (34.550) | (0.255) | 125.66 (1961) |
| <0.001 | <0.001 | | 0.048 | 0.004 (L2) | |

2012

| | | | | | | |
|-----|---------|---------|---------|------------|----------|------------------------|
| ISR | 8.980 | -0.004 | 0.005 | -39.360 | 0.001 | 197.46 (2006) -> |
| | (3.196) | (0.002) | (0.003) | (15.070) | (0.0003) | 187.98 (1994) |
| | 0.005 | 0.080 | 0.116 | 0.009 | <0.001 | |
| DSR | 21.426 | -0.015 | | -0.036 | | 113.04 (2010) -> |
| | (7.815) | (0.006) | | (0.271) | | 109.21 (1970) |
| | 0.006 | 0.010 | | 0.894 (L2) | | |
| | | | | 0.612 | | |
| | | | | (0.294) | | |
| | | | | 0.037 (L3) | | |

Table 2 Species composition of each habitat type for all species and the disturbance indicator species specifically in 1995-6 and 2012. Percentage values are given in brackets.

| All species | SR | Indigenous | Exotic | Grass | Herb | Shrub | Tree |
|----------------------------|-----------|-------------------|---------------|--------------|-------------|--------------|-------------|
| Grasslands 1995 | 153 | 142 (92.8%) | 11 (7.2%) | 38 (24.8%) | 95 (62.1%) | 19 (12.4%) | 1 (0.7%) |
| Grasslands 2012 | 114 | 107 (93.9%) | 7 (6.1%) | 31 (27.2%) | 69 (60.5%) | 13 (11.4%) | 1 (0.9%) |
| Woody 1995 | 164 | 150 (91.5%) | 14 (8.5%) | 29 (17.7%) | 80 (48.8%) | 38 (23.2%) | 17 (10.4%) |
| Woody 2012 | 139 | 120 (86.3%) | 19 (13.7%) | 29 (20.9%) | 61 (43.9%) | 32 (23%) | 17 (12.2%) |
| Disturbance species | | | | | | | |
| Grasslands 1995 | 26 | 21 (81%) | 5 (19%) | 8 (31%) | 12 (46%) | 6 (23%) | 0 |
| Grasslands 2012 | 14 | 11 (79%) | 3 (21%) | 4 (29%) | 8 (57%) | 2 (14%) | 0 |
| Woody 1995 | 24 | 17 (71%) | 7 (29%) | 7 (29%) | 13 (54%) | 4 (17%) | 0 |
| Woody 2012 | 24 | 17 (71%) | 7 (29%) | 7 (29%) | 13 (54%) | 4 (17%) | 0 |

565

Figure Captions

Fig. 1 The location of Potchefstroom in South Africa (a), the town commonage area (black outline) and the location of the sites (see legend of map c) (b), urban development from 1863 to 570 2010 (1863 obtained from an early hand drawn map, 1944-2006 were digitized from government topographical maps and 2010 from a satellite image) (c), sites and urban area in 1995 (d), sites and urban area in 2010, indicating the 1995 urban area (hatched) (e).

Fig. 2 Plots indicating change in a) the percentage natural (PN) area, b) the road network 575 density of the natural areas (RNDN), c) the density of dwellings (CD), and d) landscape diversity (H') for the open grassland (G) (open boxes) and woody (W) (black boxes) communities spanning the period 1938-2010. Means \pm SE's are shown.

Fig. 3 The relationships between species richness and altitude (top plot) and the road network 580 density of natural areas (bottom plot) as indicated in Table 1. Codes indicate open grassland (G) or woody community (W) patterns in 1995 (95) or 2012 (12) for indigenous (ISR) and disturbance (DSR) species richness.

Fig. 4 The relationships between species richness and the percentage natural area as indicated 585 in Table 1. Codes indicate open grassland (G) or woody community (W) patterns in 1995 (95) or 2012 (12) for indigenous (ISR) and disturbance (DSR) species richness.

Fig. 5 Open grassland indigenous (a, b) and disturbance (c, d) species richness per site for 1995 590 and 2012. Indicated in each map are the extent of the urban area and the roads and footpaths in the natural areas (RN Natural) corresponding to the best predicted model.

Fig. 6 Woody grassland indigenous (a, b) and disturbance (c, d) species richness per site for 1995 and 2012. Indicated in each map are the extent of the urban area and the roads and footpaths in the natural areas (RN Natural) corresponding to the best predicted model.




Fig. 1






Fig. 2



600

Fig. 3





Fig. 4

605





Fig. 5



610 Fig. 6

References

- 615 Adriaens D, Honnay O, Hermy M (2006) No evidence of a plant extinction debt in highly fragmented calcareous grasslands in Belgium. *Biol Conserv* 133(2):212-224
- Anderson JE, Inouye RS (2001) Landscape-scale changes in plant species abundance and biodiversity of a sagebrush steppe over 45 years. *Ecol Monogr* 71(4):531-556
- Andersson E, Ahrné K, Pyykönen M, Elmqvist T (2009) Patterns and scale relations among urbanization measures in Stockholm, Sweden. *Landscape Ecol* 24(10):1331-1339
- 620 Beyer HL (2006) Hawth's Analysis Tools Version 3.27.
- Bezuidenhout H, Bredenkamp GJ (1990) A reconnaissance survey of the vegetation of the dolomitic region in the Potchefstroom-Ventersdorp-Randfontein area, South Africa. *Phytocoenologia* 18:387-403
- 625 Bezuidenhout H, Bredenkamp GJ (1991) The vegetation of the Bc land type in the western Transvaal Grassland, South Africa. *Phytocoenologia* 19:497-518
- Bezuidenhout H, Bredenkamp GJ, Theron GK (1994a) A classification of the vegetation of the western Transvaal dolomite and chert grassland, South Africa. *S Afr J Bot* 60(3):152-161
- Bezuidenhout H, Bredenkamp GJ, Theron GK (1994b) Syntaxonomy of the vegetation of the Fb land type in the western Transvaal grassland, South Africa. *S Afr J Bot* 60(1):72-81

- 630 Boone CG, Cadenasso ML, Grove JM, Schwarz K, Buckley GL (2009) Landscape, vegetation characteristics, and group identity in an urban and suburban watershed: why the 60s matter. *Urban Ecosystems* 13(3):255-271
- Bredenkamp GJ, Bezuidenhout H, Naude C, Joubert H (1994) The vegetation of the Boskop Dam Nature Reserve, Potchefstroom. *Koedoe* 37(1):19-33
- 635 Bredenkamp GJ, Joubert AF, Bezuidenhout H (1989) A reconnaissance survey of the plains of the Potchefstroom-Parys-Fochville area. *S Afr J Bot* 55:199-206
- Bredenkamp GJ, Theron GK (1978) A synecological account of the Suikerbosrand Nature Reserve 1. The phytosociology of the Witwatersrand geological system. *Bothalia* 12:513-529
- Brooks TM, Pimm SL, Oyugi JO (1999) Time lag between deforestation and bird extinction in
640 tropical forest fragments. *Conserv Biol* 13(5):1140-1150
- Burel F (1993) Time lags between spatial pattern changes and species distribution changes in dynamic landscapes. *Landscape Urban Plann* 24:161-166
- Chapin FS, III, Starfield AM (1997) Time lags and novel ecosystems in response to transient climatic change in arctic Alaska. *Clim Change* 35:449-461
- 645 Chiba S, Okochi I, Ohbayashi T et al (2009) Effects of habitat history and extinction selectivity on species-richness patterns of an island land snail fauna. *J Biogeogr* 36(10):1913-1922

- Chocholoušková Z, Pyšek P (2003) Changes in composition and structure of urban flora over 120 years: a case study of the city of Plzeň. *Flora - Morphology, Distribution, Functional Ecology of Plants* 198(5):366-376
- 650 Cilliers S, Schoeman L, Bredenkamp G (1998) Wetland plant communities in the Potchefstroom Municipal area, North-west, South Africa. *Bothalia* 28(2):213-229
- Cilliers SS, Bredenkamp GJ (1998) Vegetation analysis of railway reserves in the Potchefstroom municipal area, North West Province, South Africa. *S Afr J Bot* 64(5):271-280
- 655 Cilliers SS, Bredenkamp GJ (1999) Ruderal and degraded natural vegetation on vacant lots in the Potchefstroom Municipal Area, North West Province, South Africa. *S Afr J Bot* 65(2):163-173
- Cilliers SS, Bredenkamp GJ (2000) Vegetation of road verges on an urbanisation gradient in Potchefstroom, South Africa. *Landscape Urban Plann* 46:217-239
- 660 Cilliers SS, Van Wyk E, Bredenkamp GJ (1999) Urban nature conservation: vegetation of natural areas in the Potchefstroom municipal area, North West Province, South Africa. *Koedoe* 42(1):1-30
- Cilliers SS, Williams NSG, Barnard FJ (2008) Patterns of exotic plant invasions in fragmented urban and rural grasslands across continents. *Landscape Ecol* 23(10):1243-1256
- Clarke KR, Gorley RN (2006) *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth,

- 665 Crooks JA (2005) Lag times and exotic species: the ecology and management of biological
invasions in slow-motion. *Ecoscience* 12(3):316-329
- Daehler CC (2009) Short lag times for invasive tropical plants: evidence from experimental
plantings in Hawai'i. *PLoS ONE* 4(2):e4462
- Dauber J, Hirsch M, Simmering D, Waldhardt R, Otte A, Wolters V (2003) Landscape
670 structure as an indicator of biodiversity: matrix effects on species richness. *Agric, Ecosyst
Environ* 98(1-3):321-329
- Davis MB (1986) Climatic instability, time lags, and community disequilibrium. In: Diamond
J. and Case T. J. (eds), *Community Ecology*. Harper & Row, pp. 269-284
- Deng X, Huang J, Huang Q, Rozelle S, Gibson J (2011) Do roads lead to grassland degradation
675 or restoration? A case study in Inner Mongolia, China. *Environment and Development
Economics* 16(06):751-773
- du Toit MJ (2009) Grassland ecology along an urban-rural gradient using GIS techniques in
Klerksdorp, South Africa., North-West University <http://hdl.handle.net/10394/4197>
- du Toit MJ, Cilliers SS (2011) Aspects influencing the selection of representative urbanization
680 measures to quantify urban-rural gradients. *Landscape Ecol* 26(2):169-181
- Duguay S, Eigenbrod F, Fahrig L (2006) Effects of surrounding urbanization on non-native
flora in small forest patches. *Landscape Ecol* 22(4):589-599

- Dullinger S, Essl F, Rabitsch W et al (2013) Europe's other debt crisis caused by the long legacy of future extinctions. *Proceedings of the National Academy of Sciences* 110(18):7342-685 7347
- Dupouey JL, Dambrine E, Laffite JD, Moares C (2002) Irreversible Impact of past Land Use on Forest Soils and Biodiversity. *Ecology* 83(11):2978-2984
- Dupré C, Ehrlén J (2002) Habitat configuration, species traits and plant distributions. *J Ecol* 90(5):796-805
- 690 Eckhardt H, Van Rooyen N, Bredenkamp G (1993) Use of Braun-Blanquet data for the assessment of veld condition and grazing capacity in grassland. *African Journal of Range & Forage Science* 10(1):41-46
- ESRI (2010) ArcGIS, version 10.0. Environmental Systems Research Institute, Redlands, CA,
- Ewers RM, Didham RK (2006) Confounding factors in the detection of species responses to 695 habitat fragmentation. *Biol Rev Camb Philos Soc* 81(1):117-42
- Foster D, Swanson F, Aber J et al (2003) The Importance of Land-Use Legacies to Ecology and Conservation. *Bioscience* 53(1):77-88
- Foster DR (1992) Land-Use History (1730-1990) and Vegetation Dynamics in Central New England, USA. *J Ecol* 80(4):753-771

- 700 Friedel MH (1987) A preliminary investigation of woody plant increase in the western Transvaal and implications for veld assessment. *Journal of the Grassland Society of Southern Africa* 4(1):25-30
- Gelbard JL, Belnap J (2003) Roads as conduits for exotic plant invasions in a semiarid landscape. *Conserv Biol* 17(2):420-432
- 705 Godefroid S, Koedam N (2003) Distribution pattern of the flora in a peri-urban forest: an effect of the city–forest ecotone. *Landscape Urban Plann* 65(4):169-185
- Grove JM, Locke DH, O'Neil-Dunne JP (2014) An ecology of prestige in New York City: examining the relationships among population density, socio-economic status, group identity, and residential canopy cover. *Environ Manage* 54(3):402-19
- 710 Gustavsson E, Lennartsson T, Emanuelsson M (2007) Land use more than 200 years ago explains current grassland plant diversity in a Swedish agricultural landscape. *Biol Conserv* 138(1-2):47-59
- Hahs AK, McDonnell MJ (2006) Selecting independent measures to quantify Melbourne's urban–rural gradient. *Landscape Urban Plann* 78(4):435-448
- 715 Hahs AK, McDonnell MJ, McCarthy MA et al (2009) A global synthesis of plant extinction rates in urban areas. *Ecol Lett* 12(11):1165-73
- Hamburg SP, Sanford RL, Jr. (1986) Disturbance, "Homo Sapiens," and Ecology. *Bulletin of the Ecological Society of America* 67(2):169-171

- Hansen MJ, Cleverger AP (2005) The influence of disturbance and habitat on the presence of
720 non-native plant species along transport corridors. *Biol Conserv* 125(2):249-259
- Hanski I, Ovaskainen O (2002) Extinction debt at extinction threshold. *Conserv Biol*
16(3):666-673
- Hayashida FM (2005) Archaeology, Ecological History, and Conservation. *Annual Review of
Anthropology* 34:43-65
- 725 Helm A, Hanski I, Partel M (2006) Slow response of plant species richness to habitat loss and
fragmentation. *Ecol Lett* 9(1):72-7
- Janišová M, Michalcová D, Bacaro G, Ghisla A (2014) Landscape effects on diversity of semi-
natural grasslands. *Agric, Ecosyst Environ* 182:47-58
- Jansen van Rensburg JP (2010) Investigation of the microbial diversity and functionality of soil
730 in fragmented South African grasslands along an urbanization gradient. North-West University
- Kark S, van Rensburg BJ (2006) Ecotones: Marginal or Central Areas of Transition? *Israel
Journal of Ecology & Evolution* 52(1):29-53
- Kissling M, Hegetschweiler KT, Rusterholz H-P, Baur B (2009) Short-term and long-term
effects of human trampling on above-ground vegetation, soil density, soil organic matter and
735 soil microbial processes in suburban beech forests. *Applied Soil Ecology* 42(3):303-314
- Kleijn D, Kohler F, Baldi A et al (2009) On the relationship between farmland biodiversity and
land-use intensity in Europe. *Proc Biol Sci* 276(1658):903-9

- 740 Koerner W, Dupouey JL, Dambrine E, Benoit M (1997) Influence of Past Land Use on the
Vegetation and Soils of Present Day Forest in the Vosges Mountains, France. *J Ecol* 85(3):351-
358
- Koyanagi T, Kusumoto Y, Yamamoto S, Okubo S, Iwasaki N, Takeuchi K (2012) Grassland
plant functional groups exhibit distinct time-lags in response to historical landscape change.
Plant Ecol 213(2):327-338
- 745 Kraaij T, Ward D (2006) Effects of rain, nitrogen, fire and grazing on tree recruitment and
early survival in bush-encroached savanna, South Africa. *Plant Ecol* 186(2):235-246
- Krauss J, Bommarco R, Guardiola M et al (2010) Habitat fragmentation causes immediate and
time-delayed biodiversity loss at different trophic levels. *Ecol Lett* 13(5):597-605
- Kuussaari M, Bommarco R, Heikkinen RK et al (2009) Extinction debt: a challenge for
biodiversity conservation. *Trends Ecol Evol* 24(10):564-71
- 750 Laurance WF (2002) Hyperdynamism in fragmented habitats. *Journal of Vegetation Science*
13(4):595
- Lawson DM, Lamar CK, Schwartz MW (2008) Quantifying Plant Population Persistence in
Human-Dominated Landscapes
- 755 Cuantificación de la Persistencia de Poblaciones de Plantas en Paisajes Dominados por
Humanos. *Conserv Biol* 22(4):922-928

- Lindborg R, Eriksson O (2004) Historical landscape connectivity affects present plant species diversity. *Ecology* 85(7):1840-1845
- Lloyd AH (2005) Ecological histories from Alaskan tree lines provide insight into future change. *Ecology* 86(7):1687-1695
- 760 Locke DH, Baine G (2014) The good, the bad, and the interested: how historical demographics explain present-day tree canopy, vacant lot and tree request spatial variability in New Haven, CT. *Urban Ecosystems*
- Louw WJ (1951) An ecological account of the vegetation of the Potchefstroom area. *Memoirs of the Botanical Survey of South Africa*, No. 24
- 765 Luck GW, Smallbone LT, O'Brien R (2009) Socio-Economics and Vegetation Change in Urban Ecosystems: Patterns in Space and Time. *Ecosystems* 12(4):604-620
- Madubansi M, Shackleton CM (2007) Changes in fuelwood use and selection following electrification in the Bushbuckridge lowveld, South Africa. *J Environ Manage* 83(4):416-26
- Magnuson JJ (1990) Long-Term Ecological Research and the Invisible Present. *Bioscience*
770 40(7):495-501
- McDonnell MJ, Hahs AK (2013) The future of urban biodiversity research: Moving beyond the 'low-hanging fruit'. *Urban Ecosystems* 16(3):397-409
- Monger C, Sala OE, Duniway MC et al (2015) Legacy effects in linked ecological–soil–geomorphic systems of drylands. *Front Ecol Environ* 13(1):13-19

- 775 Morgan J (1995) Ecological studies of the endangered *Rutidosia leptorrhynchoides*. II. Patterns of seedling emergence and survival in a native grassland. *Aust J Bot* 43(1):13-24
- Morgan JW (1998) Composition and seasonal flux of the soil seed bank of species-rich *Themeda triandra* grasslands in relation to burning history. *Journal of Vegetation Science* 9:145-156
- 780 Mucina L, Rutherford MC (eds) (2006) The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia* 19. South African National Biodiversity Institute, Pretoria
- Mueller-Dombois D, Ellenberg H (1974) Aims and methods of plant ecology. John Wiley and Sons, New York,
- Neser L (1967) Die Kleurlinge van Potchefstroom: 'n algemene kultuurbeeld met besondere
785 verwysing na gesinsgebruike. Potchefstroomse Universiteit vir Christelike Hoër Onderwys
- O'Connor TG (1991) Local extinction in perennial grasslands: A life-history approach. *The American Naturalist* 137(6):753-773
- Pickett ST, Cadenasso ML, Grove JM et al (2008) Beyond urban legends: an emerging
framework of urban ecology, as illustrated by the Baltimore Ecosystem Study. *Bioscience*
790 58(2):139-150
- Purschke O, Sykes MT, Reitalu T, Poschlod P, Prentice HC (2012) Linking landscape history and dispersal traits in grassland plant communities. *Oecologia* 168(3):773-83

- R Core Team (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://R-project.org/>.
- 795 Ramalho CE, Hobbs RJ (2012) Time for a change: dynamic urban ecology. *Trends Ecol Evol* 27(3):179-88
- Retief E, Herman PPJ (1997) Plants of the northern provinces of South Africa: keys and diagnostic characters. *Strelitzia* 6
- Rhemtulla JM, Mladenoff DJ, Clayton MK (2009) Legacies of Historical Land Use on
800 Regional Forest Composition and Structure in Wisconsin, USA (Mid-1800s-1930s-2000s). *Ecol Appl* 19(4):1061-1078
- Risser PG (1995) The Status of the Science Examining Ecotones. *Bioscience* 45(5):318-325
- Romme WH, Knight DH (1982) Landscape diversity: the concept applied to Yellowstone Park. *Bioscience* 32(8):664-670
- 805 Sakai AK, Allendorf FW, Holt JS et al (2001) The population biology of invasive species. *Annu Rev Ecol Syst* 32:305-332
- Shannon CE (1948) A mathematical theory of communication. *Bell System Technical Journal* 27:379-423, 623-656
- Smit PA (1989) Die ekonomiese ontwikkeling van Potchefstroom vanaf 1902 tot 1945.
810 Potchefstroom University for CHE

- Stehlik I, Caspersen JP, Wirth LEA, Holderegger R (2007) Floral free fall in the Swiss lowlands: environmental determinants of local plant extinction in a peri-urban landscape. *J Ecol* 95(4):734-744
- 815 Tait CJ, Daniels CB, Hill RS (2005) Changes in species assemblages within the Adelaide Metropolitan Area, Australia, 1836-2002. *Ecol Appl* 15(1):346-359
- Tilman D, Fargione J, Wolff B et al (2001) Forecasting agriculturally driven global environmental change. *Science* 292(5515):281-4
- Tilman D, May RM, Lehman CL, Nowak MA (1994) Habitat destruction and the extinction debt. *Nature* 371:65-66
- 820 Tlokwe City Council (2015) Business Overview. Available from <http://www.potch.co.za/cdepstatment.htm> accessed 25 April 2015)
- Tscharntke T, Klein AM, Kruess A, Steffan-Dewenter I, Thies C (2005) Landscape perspectives on agricultural intensification and biodiversity — ecosystem service management. *Ecol Lett* 8(8):857-874
- 825 Vakhlamova T, Rusterholz H-P, Kanibolotskaya Y, Baur B (2014) Changes in plant diversity along an urban–rural gradient in an expanding city in Kazakhstan, Western Siberia. *Landscape Urban Plann* 132:111-120
- Van Auken OW (2009) Causes and consequences of woody plant encroachment into western North American grasslands. *J Environ Manage* 90(10):2931-42

830 van den Bergh G (1990) Voortrekker plaasbesetting op die Transvaalse Hoëveld: 'n versteurde beeld. *South African Journal of Surveying and Mapping* 20(Part 7):301-311

van den Bergh G (1992) Die tweede Potchefstroom, opmeting en besetting 1841-60. *South African Journal of Surveying and Mapping* 21(Part 4):167-178

van der Walt L (2013) Landscape functionality and plant diversity of grassland fragments
835 along an urban-rural gradient in the Tlokwe Municipal area, South Africa. North-West University <http://hdl.handle.net/10394/9732>

Ward D (2005) Do we understand the causes of bush encroachment in African savannas? *African Journal of Range & Forage Science* 22(2):101-105

Williams NSG, Morgan JW, McCarthy MA, McDonnell MJ (2006) Local extinction of
840 grassland plants: The landscape matrix is more important than patch attributes. *Ecology* 87(12):3000-3006

Williams NSG, Morgan JW, McDonnell MJ, McCarthy MA (2005) Plant traits and local extinctions in natural grasslands along an urban-rural gradient. *J Ecol* 93(6):1203-1213