

<https://helda.helsinki.fi>

---

## Early gestational ethanol exposure in mice : Effects on brain structure, energy metabolism and adiposity in adult offspring

Zhang, Christine R.

2019-03

---

Zhang , C R , Kurniawan , N D , Yamada , L , Fleming , W , Kaminen-Ahola , N , Ahola , A , Galloway , G & Chong , S 2019 , ' Early gestational ethanol exposure in mice : Effects on brain structure, energy metabolism and adiposity in adult offspring ' , Alcohol (New York) , vol. 75 , pp. 1-10 . <https://doi.org/10.1016/j.alcohol.2018.04.008>

---

<http://hdl.handle.net/10138/306943>

<https://doi.org/10.1016/j.alcohol.2018.04.008>

---

cc\_by\_nc\_nd

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*


*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Accepted Manuscript

Early gestational ethanol exposure in mice: effects on brain structure, energy metabolism and adiposity in adult offspring

Christine R. Zhang, Nyoman D. Kurniawan, Lisa Yamada, Whitney Fleming, Nina Kaminen-Ahola, Arttu Ahola, Graham Galloway, Suyinn Chong



PII: S0741-8329(18)30022-3

DOI: [10.1016/j.alcohol.2018.04.008](https://doi.org/10.1016/j.alcohol.2018.04.008)

Reference: ALC 6800

To appear in: *Alcohol*

Received Date: 31 January 2018

Revised Date: 20 April 2018

Accepted Date: 20 April 2018

Please cite this article as: Zhang C.R, Kurniawan N.D, Yamada L., Fleming W., Kaminen-Ahola N., Ahola A., Galloway G. & Chong S., Early gestational ethanol exposure in mice: effects on brain structure, energy metabolism and adiposity in adult offspring, *Alcohol* (2018), doi: 10.1016/j.alcohol.2018.04.008.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Early gestational ethanol exposure in mice: effects on brain structure, energy**  
2 **metabolism and adiposity in adult offspring**

3

4 Christine R Zhang<sup>a</sup>, Nyoman D Kurniawan<sup>b</sup>, Lisa Yamada<sup>a</sup>, Whitney Fleming<sup>a</sup>, Nina  
5 Kaminen-Ahola<sup>c</sup>, Arttu Ahola<sup>d</sup>, Graham Galloway<sup>b</sup> and Suyinn Chong<sup>a</sup>

6

7 <sup>a</sup>Mater Research Institute – The University of Queensland, Translational Research Institute,  
8 Woolloongabba, Queensland, Australia

9 <sup>b</sup>Centre for Advanced Imaging, The University of Queensland, St Lucia, Queensland,  
10 Australia

11 <sup>c</sup>Department of Medical and Clinical Genetics, University of Helsinki, Helsinki, Finland

12 <sup>d</sup>Queensland Institute of Medical Research, Herston, Queensland, Australia

13

14 Corresponding author: Suyinn Chong, Epigenetics Group, Mothers', Babies' and Women's  
15 Health Program, Mater Research Institute – The University of Queensland, Translational  
16 Research Institute, Level 4, 37 Kent St, Woolloongabba, Queensland 4102, Australia. E-mail:  
17 [suyinn.chong@mater.uq.edu.au](mailto:suyinn.chong@mater.uq.edu.au)

18

19 Declarations of interest: none

**20 Abstract**

21 We examined whether an early-life event — ethanol exposure in the initial stages of  
22 pregnancy — affected offspring brain structure, energy metabolism and body composition in  
23 later life. Consumption of 10% (v/v) ethanol by inbred C57BL/6J female mice from 0.5 to 8.5  
24 days post coitum was used to model alcohol exposure during the first 3-4 weeks of gestation  
25 in humans, when pregnancy is not typically recognized. At adolescence (postnatal day (P) 28)  
26 and adulthood (P64), the brains of male offspring were scanned *ex vivo* using ultra-high field  
27 (16.4 Tesla) magnetic resonance imaging and diffusion tensor imaging. Energy metabolism  
28 and body composition were measured in adulthood by indirect calorimetry and dual-energy  
29 X-ray absorptiometry (DXA), respectively. Ethanol exposure had no substantial impact on  
30 white matter organization in the anterior commissure, corpus callosum, hippocampal  
31 commissure, internal capsule, optic tract or thalamus. Whole brain volume and the volumes  
32 of the neocortex, cerebellum and caudate putamen were also unaffected. Subtle, but non-  
33 significant, effects were observed on the hippocampus and the hypothalamus in adult ethanol-  
34 exposed male offspring. Ethanol exposure was additionally associated with a trend towards  
35 decreased oxygen consumption, carbon dioxide production and reduced daily energy  
36 expenditure as well as significantly increased adiposity, albeit with normal body weight and  
37 food intake, in adult male offspring. In summary, ethanol exposure restricted to early  
38 gestation had subtle long-term effects on the structure of specific brain regions in male  
39 offspring. The sensitivity of the hippocampus to ethanol-induced damage is reminiscent of  
40 that reported by other studies — despite differences in the level, timing and duration of  
41 exposure — and likely contributes to the cognitive impairment which characteristically  
42 results from prenatal ethanol exposure. The hypothalamus plays an important role in  
43 regulating metabolism and energy homeostasis. Our finding of altered daily energy  
44 expenditure and adiposity in adult ethanol-exposed males is consistent the idea that central

45 nervous system abnormalities also underpin some of the metabolic phenotypes associated  
46 with ethanol exposure in pregnancy.

47

#### 48 **Keywords**

49 prenatal, alcohol, brain structure, energy metabolism, body composition

50

#### 51 **Introduction**

52 Adverse early-life environmental exposures such as gestational undernutrition, overnutrition  
53 and diabetes mellitus have been shown to increase susceptibility to obesity and its related  
54 diseases in later life (Curhan et al., 1996; Fraser et al., 2010; Law, Barker, Osmond, Fall, &  
55 Simmonds, 1992; Lunde et al., 2016; Ravelli, van Der Meulen, Osmond, Barker, & Bleker,  
56 1999; Zhao et al., 2016). The underlying mechanisms are not fully understood; however,  
57 affected pathways include those involved in the control of appetite (Bellinger, Lilley, &  
58 Langley-Evans, 2004; Franke et al., 2005), glucose metabolism (Phillips, Barker, Hales,  
59 Hirst, & Osmond, 1994), circadian rhythms (Borengasser et al., 2014; Sutton, Centanni, &  
60 Butler, 2010) and adipogenesis (Borengasser et al., 2013; Lukaszewski et al., 2011).

61 There is growing evidence that prenatal ethanol exposure has similar consequences, with  
62 links to increased adiposity, insulin resistance and glucose intolerance in animal models  
63 (Chen & Nyomba, 2003; Dobson et al., 2012; Gardebjer, Anderson, Pantaleon, Wlodek, &  
64 Moritz, 2015; Gardebjer et al., 2017) and reports of increased rates of obesity in children and  
65 adolescents with fetal alcohol spectrum disorders (FASDs) (Fuglestad et al., 2014; Werts,  
66 Van Calcar, Wargowski, & Smith, 2014). Structural and functional abnormalities of the  
67 central nervous system are common in FASDs (Astley, 2010) and could potentially underpin  
68 other phenotypes associated with prenatal ethanol exposure.

69 In this study, we examined the consequences of prenatal ethanol exposure on brain structure  
70 at adolescence and adulthood using ultra-high field magnetic resonance imaging (MRI) and  
71 diffusion tensor imaging (DTI). Long-term effects on energy metabolism, body weight and  
72 body composition were also explored. Notably, ethanol exposure was restricted to early  
73 gestation — equivalent to the period between conception and pregnancy recognition in  
74 humans — when the chances of alcohol consumption by pregnant women are high  
75 (McCormack et al., 2017), but information on possible adverse outcomes is limited.

## 76 **Materials and Methods**

### 77 *Prenatal ethanol exposure*

78 Animal work was conducted in accordance with the Australian code for the care and use of  
79 animals for scientific purposes, and was approved by Animal Ethics Committees at the  
80 Queensland Institute of Medical Research (P986, A0606-609M) and The University of  
81 Queensland (MRI-UQ/TRI/430/13). Prenatal ethanol exposure involved consumption of 10%  
82 (v/v) ethanol by pregnant C57BL/6J dams from 0.5 to 8.5 days post coitum (dpc) and was  
83 performed as described previously (Kaminen-Ahola, Ahola, Maga, et al., 2010). C57BL/6J  
84 mice show a significant preference for 10% (v/v) ethanol when given *ad libitum* two-bottle  
85 choice between ethanol and water (Belknap, Crabbe, & Young, 1993). In this study, single-  
86 bottle administration of 10% (v/v) ethanol was used to reduce variation in ethanol exposure  
87 and, consequently, the number of animals needed for the experiment. This approach is  
88 potentially more stressful than a two-bottle choice paradigm. Animal welfare during the  
89 exposure period was monitored by measurements of fluid intake, body weight and daily  
90 observations of mouse appearance and behaviour. Adult (6-8 week old) C57BL/6J mice were  
91 obtained from the Animal Resources Centre (Canning Vale, WA, Australia) and acclimatized  
92 to a 12-hour light/dark cycle for up to one week. Males were then caged with a single female  
93 overnight and detection of a vaginal plug the next morning indicated that mating had taken

94 place (designated 0.5 dpc). Males were then removed and females were provided with a drink  
95 bottle containing either 10% (v/v) ethanol (ethanol-exposed) or water (control) *ad libitum* for  
96 the following eight days. Liquid consumption (to the nearest 0.2 ml) was measured every 24  
97 hours. At 8.5 dpc, the ethanol-exposed females were placed back on water. All mice had free  
98 access to standard mouse chow (Irradiated Rat and Mouse Diet, Specialty Feeds, Glen  
99 Forrest, WA, Australia) at all times. Body weight was measured at 0.5 dpc and 8.5 dpc for  
100 ethanol-exposed and control dams, and at 12 weeks of age for their offspring.

### 101 ***MRI and DTI studies***

102 At P28 or P64, male mice were transcardially perfused with 4% paraformaldehyde according  
103 to standard protocols. The brains were then removed and scanned *ex vivo* as described  
104 previously (Kurniawan et al., 2014). Brain samples were incubated in 0.2% (v/v) Magnevist  
105 (Bayer AG, Leverkusen, Germany) for 4 days prior to imaging, and MRI scans were  
106 performed using a Bruker 16.4 Tesla widebore Avance II NMR spectrometer (Bruker  
107 Biospin, Karlsruhe, Germany) and 15 mm SAW coil (M2M Imaging, Brisbane, Australia).  
108 Whole brain scans were performed using: (1) 3D diffusion-weighted spin-echo sequence at  
109 100  $\mu\text{m}$  isotropic resolution, with 30 diffusion directions,  $b=5000 \text{ s/mm}^2$ ; and (2)  $T_1/T_2^*$ -  
110 weighted 3D gradient echo sequence at 50  $\mu\text{m}$  isotropic resolution. The total acquisition time  
111 was ~16 hours. Using the C57BL/6 MRI brain atlas (Ma et al., 2005), regions of interest  
112 (ROI) were registered to each gradient echo dataset using FSL linear (FLIRT) and non-linear  
113 (FNIRT) registration protocols ([www.fmrib.ox.ac.uk/fsl](http://www.fmrib.ox.ac.uk/fsl)). Registered ROIs were then  
114 examined and manually corrected using a histology-based C57BL/6 mouse brain atlas  
115 (Franklin & Paxinos, 2008). Volumes were calculated using ITK-SNAP (Yushkevich et al.,  
116 2006). DiffusionToolkit (Ruopeng Wang, Van J. Wedeen, TrackVis.org, Martinos Center for  
117 Biomedical Imaging, Massachusetts General Hospital) was used to process the diffusion data  
118 and calculate the DTI parametric maps. Fibretracking was performed using Q-ball and fibre

119 assignment for continuous tract (FACT). Reconstructed DTI data was visualized and  
120 analysed using TrackVis software (Ruopeng Wang, Van J. Wedeen, TrackVis.org, Martinos  
121 Center for Biomedical Imaging, Massachusetts General Hospital). All MRI and DTI analyses  
122 were conducted blind to treatment group.

### 123 *Dual-energy X-ray absorptiometry (DXA)*

124 Body composition was analysed using a PIXImus2 densitometer and software version 2.10  
125 (Lunar, Madison, WI, USA). The head was excluded from all analyses. Measurements  
126 automatically supplied by the software included lean tissue (g), fat tissue (g), % fat, bone  
127 mineral content (g), bone area (cm<sup>2</sup>) and bone mineral density (g/cm<sup>2</sup>).

### 128 *Indirect calorimetry*

129 PhenoMaster metabolic cages (TSE Systems, Chesterfield, MO, USA) were used to monitor  
130 food (g) and water (ml) intake, body weight (g), oxygen consumption (VO<sub>2</sub>; ml/h/kg) and  
131 carbon dioxide production (VCO<sub>2</sub>; ml/h/kg) in a home cage environment. Mice were housed  
132 individually at 22 °C on a 12-hour light/12-hour dark cycle, with paper bedding and free  
133 access to water and standard chow (Irradiated Rat and Mouse Diet, Specialty Feeds, Glen  
134 Forrest, WA, Australia). Data was collected at 20 minute intervals for 3 days following an  
135 initial acclimatization period of 2-3 days. Respiratory exchange ratio (RER) was calculated as  
136 the ratio between VCO<sub>2</sub> and VO<sub>2</sub>. Daily energy expenditure was calculated as described by  
137 Meyer and colleagues (Meyer, Reitmeir, & Tschop, 2015) using the Heldmaier conversion  
138 equation.

### 139 *Statistics*

140 Previous work with this model has shown that early gestational ethanol exposure affects  
141 offspring outcomes in a stochastic manner, producing substantial intra-litter variation even in  
142 inbred mice (Kaminen-Ahola, Ahola, Maga, et al., 2010; Zhang, Ho, Vega, Burne, & Chong,  
143 2015). Therefore each animal was considered an independent unit of analysis (Elswick,



144 Welsch, & Janszen, 2000). Statistical analyses were conducted using either *R* (R  
145 Development Core Team, 2010) or GraphPad Prism 6. The Student's *t*-test was used to  
146 analyse differences between treatment groups (ethanol-exposed and control) for maternal  
147 liquid consumption and weight gain over the 8-day exposure period, litter size at weaning and  
148 offspring brain structure. Adjustment for multiple testing used the Holm-Sidak method. Two-  
149 way ANOVAs with a Tukey's multiple comparison *post hoc* test were used to analyse the  
150 effects of sex, treatment and the interaction of sex and treatment on offspring body weight,  
151 body composition, food and water intake and energy metabolism.

## 152 **Results**

### 153 ***Consumption of 10% (v/v) ethanol by C57BL/6J female mice***

154 As in our previous studies (Kaminen-Ahola, Ahola, Maga, et al., 2010; Zhang et al., 2015),  
155 there was no effect of ethanol treatment on the average volume of liquid consumed per day  
156 by pregnant dams, maternal body weight gain during the exposure period (0.5 to 8.5 dpc) or  
157 litter size at weaning (Figure 1), indicating that the exposure was not detrimental to maternal  
158 health or offspring viability.

### 159 ***Structural MRI in adolescent and adult male offspring exposed to ethanol in utero***

160 The volumes of the whole brain, neocortex, cerebellum, caudate putamen, hippocampus and  
161 hypothalamus were compared between ethanol-exposed and control male offspring both at  
162 adolescence (P28) and adulthood (P64). There was no effect of prenatal ethanol exposure on  
163 whole brain volume at either age (P28: control =  $366.30 \pm 7.48 \text{ mm}^3$ , ethanol-exposed =  
164  $344.92 \pm 11.07 \text{ mm}^3$ , unadjusted  $P=0.15$ ; P64: control =  $369.46 \pm 5.69 \text{ mm}^3$ , ethanol-exposed  
165 =  $370.32 \pm 4.70 \text{ mm}^3$ , unadjusted  $P=0.91$ ). At P28, all of the other brain regions that were  
166 analysed were similarly unaffected by ethanol exposure (Table 1). At P64, there was a trend  
167 towards a smaller hippocampus (3.8%, adjusted  $P=0.053$ ) and larger hypothalamus (4.3%,  
168 adjusted  $P=0.17$ ) in ethanol-exposed offspring, but the differences were not statistically

169 significant (Table 1). Volumetric analysis of hippocampal subfields in adult males, including  
170 the dentate gyrus and cornu ammonis (CA) 1-2 and 3 regions, found no significant  
171 differences between treatment groups, suggesting that the reduction in overall hippocampal  
172 volume was not caused by a change in any one specific subregion (Supplementary Figure S1  
173 and Table 2).

174 ***DTI analysis of white matter microstructure in adolescent and adult male offspring***  
175 ***exposed to ethanol in utero***

176 The anterior commissure, corpus callosum, hippocampal commissure, internal capsule, optic  
177 tract, and thalamus were analysed by DTI. Comparisons were made between treatment  
178 groups for fractional anisotropy as well as fibre tract number, volume and length. At P28 a  
179 reduction in fibre tract length was observed in the hippocampal commissure of ethanol-  
180 exposed mice (unadjusted  $P < 0.05$ ) however this resolved by adulthood (Tables 3 and 4).  
181 None of the other regions analysed were altered at either P28 or P64 indicating that white  
182 matter integrity and connectivity were not substantially affected by ethanol exposure early in  
183 pregnancy.

184 ***Ethanol exposure early in pregnancy is associated with changes in energy metabolism and***  
185 ***body composition in adulthood***

186 Body weight and body composition were measured in both male and female offspring of  
187 ethanol-exposed and control dams at 12 weeks of age (Figure 2 and Figure 3). Energy intake  
188 and expenditure were measured in littermates at 21-26 weeks of age (Figure 4 and Figure 5).  
189 There was an effect of sex on body weight ( $F(1,92)=547.8$ ,  $P < 0.0001$ ), lean mass  
190 ( $F(1,49)=196.2$ ,  $P < 0.0001$ ) and fat mass ( $F(1,49)=23.3$ ,  $P < 0.0001$ ), with adult male  
191 offspring being significantly heavier than female offspring in the same treatment group  
192 (Figure 2a-c). Ethanol exposure had no significant effect on body weight or lean mass in  
193 either sex (Figure 2a-b), but ethanol-exposed males exhibited increased (18.7%; control = 2.9

194  $\pm 0.5$  g, ethanol-exposed =  $3.5 \pm 0.4$  g) fat mass compared to control males (treatment  
195  $F(1,49)=4.7$ ,  $P<0.05$ ; treatment x sex interaction  $F(1,49)=5.1$ ,  $P<0.05$ , Figure 2c).  
196 Furthermore, the increased adiposity in ethanol-exposed males resulted in percentage fat  
197 levels (~14%) similar to that of female offspring (Figure 2d). Prenatal ethanol exposure had  
198 no effect on bone mineral content, bone area or bone mineral density in either sex, although  
199 there was a significant effect of sex on bone mineral content and bone area (Figure 3).  
200 Metabolic phenotyping by indirect calorimetry revealed significant differences in oxygen  
201 consumption ( $VO_2$ ; sex  $F(1,37)=18.9$ ,  $P<0.001$ , treatment  $F(1,37)=9.6$ ,  $P<0.01$ ), carbon  
202 dioxide production ( $VCO_2$ ; sex  $F(1,37)=12.9$ ,  $P<0.001$ , treatment  $F(1,37)=6.4$ ,  $P<0.05$ ) and  
203 daily energy expenditure (sex  $F(1,37)=11.08$ ,  $P<0.01$ , treatment  $F(1,37)=4.9$ ,  $P<0.05$ ) by sex  
204 and by treatment group (Figure 4a, b and d, respectively). Moreover, ethanol-exposed males  
205 tended to have lower oxygen consumption (9.1%, adjusted  $P=0.10$ ), carbon dioxide  
206 production (8.8%, adjusted  $P=0.16$ ) and daily energy expenditure (7.9%, adjusted  $P=0.11$ )  
207 compared to sex- and age-matched controls (Figure 4a, b and d, respectively). There was no  
208 effect of sex or treatment on offspring respiratory exchange ratio (RER, Figure 4c). In  
209 addition, ethanol exposure did not significantly affect food or water intake in offspring of  
210 either sex (Figure 5) although there was an effect of light cycle on these measures (male  
211 water intake:  $F(1,34)=410.9$ ,  $P<0.0001$ ; female water intake:  $F(1,40)=98.7$ ,  $P<0.0001$ ; male  
212 food intake:  $F(1,34)=391.7$ ,  $P<0.0001$ ; female food intake:  $F(1,40)=131.0$ ,  $P<0.0001$ ).

## 213 Discussion

214 Recognition of pregnancy by women often results in the reduction of risky behaviours such  
215 as alcohol consumption; however, this does not usually occur before the fourth week of  
216 gestation. Thus, there is a critical window of early development when inadvertent alcohol  
217 exposure is possible, warranting investigation of its impact on offspring health. We have an  
218 established inbred C57BL/6J mouse model of gestational ethanol exposure which

219 encompasses implantation, gastrulation and early organogenesis, and is developmentally  
220 equivalent to the first 3-4 weeks of a human pregnancy (Kaminen-Ahola, Ahola, Maga, et al.,  
221 2010). Prior work has shown that this type of exposure is capable of producing changes in  
222 adolescent body weight (Kaminen-Ahola, Ahola, Flatscher-Bader, et al., 2010; Kaminen-  
223 Ahola, Ahola, Maga, et al., 2010) and craniofacial structure (Kaminen-Ahola, Ahola, Maga,  
224 et al., 2010) as well as adult behaviour (Sanchez Vega, Chong, & Burne, 2013). In this report,  
225 we extend our studies to show that early gestational ethanol exposure can also have long-term  
226 consequences on offspring brain structure, energy metabolism and body composition.

227 This study has several limitations. First, the blood alcohol concentrations in female mice  
228 consuming the ethanol are unknown. Second, imaging at P28 involved a small number of  
229 samples and was likely underpowered to detect subtle changes in brain structure. Third, MRI  
230 and DTI were not performed on female offspring. Therefore it remains unclear whether early  
231 gestational ethanol exposure affects brain structure in females in a similar manner to that  
232 observed in males.

233 Previous brain imaging studies in mouse models of prenatal ethanol exposure have focused  
234 on the impact of acute high dose exposures (blood alcohol concentrations of 350–420 mg/dl)  
235 and have shown alterations in both grey and white matter, although the negative effect of  
236 these exposures on offspring viability has largely prevented the examination of long-term  
237 consequences (O'Leary-Moore, Parnell, Lipinski, & Sulik, 2011). High dose ethanol  
238 exposures on gestational day (GD) 7, 8 or 10 affected overall brain volume, the forebrain,  
239 lateral ventricles, olfactory bulbs, hippocampus and cerebellum in fetuses at GD17 (Godin et  
240 al., 2009; O'Leary-Moore et al., 2010; Parnell et al., 2009). Furthermore, different structures  
241 were affected depending on the day of the exposure (O'Leary-Moore et al., 2011). A DTI  
242 study following high dose ethanol exposure on GD7 revealed effects on the fetal internal and  
243 external capsule, fimbria/fornix and corpus callosum at GD17 (O'Leary-Moore et al., 2011).

244 Cao and colleagues used quantitative susceptibility mapping to identify abnormalities in the  
245 anterior commissure, hippocampal commissure and corpus callosum in mice at P45  
246 (equivalent to 10–12 years in humans) after high dose exposure at GD7; however, analysis of  
247 the same midline structures using DTI failed to identify any differences between the ethanol-  
248 exposed and control mice (Cao et al., 2014).

249 Maternal consumption of 10% (v/v) ethanol likely results in lower blood alcohol  
250 concentrations than two intraperitoneal injections (4 hours apart) of 23-25% (v/v) ethanol at  
251 2.8–2.9 g/kg, as used in the acute high dose studies, and is expected to be closer to a  
252 moderate exposure (Allan, Chynoweth, Tyler, & Caldwell, 2003). In contrast to studies using  
253 acute high dose exposures, we found that most grey and white matter structures were either  
254 consistently unaffected by early gestational ethanol exposure, or exhibited transient changes  
255 in adolescence which were resolved by adulthood. At P64 (9 weeks of age), when brain  
256 development is complete, we identified disproportionate volumetric changes in both the  
257 hippocampus and the hypothalamus; however, the differences were not statistically  
258 significant following correction for multiple comparisons, possibly due to the small  
259 magnitude of the changes involved (~4%). The lack of significant white matter changes in  
260 adult mice suggests that these structures are not susceptible to the type of exposure used in  
261 this study; however, we cannot exclude the possibility that DTI may not be sensitive enough  
262 to detect subtle changes (Cao et al., 2014).

263 The hippocampus is known to be particularly sensitive to intrauterine alcohol exposure  
264 (Autti-Ramo et al., 2002). Neuroimaging studies have revealed changes in hippocampal  
265 volume, shape and neurometabolites (Moore, Migliorini, Infante, & Riley, 2014; Wang &  
266 Kroenke, 2015). Moreover, the type and severity of brain structural changes vary with the  
267 timing, dosage and duration of ethanol exposure. Our finding of limited changes in brain  
268 structure overall following early gestational ethanol exposure supports the idea that the

269 hippocampus and hypothalamus may be more vulnerable to ethanol-induced damage. Altered  
270 hippocampal structure in individuals with FASD has been correlated with performance in  
271 learning and memory tests (Coles et al., 2011; Willoughby, Sheard, Nash, & Rovet, 2008).  
272 Behavioural profiling of adult mice subjected to the same type of ethanol exposure as used in  
273 this study identified alterations in performance in the Morris water maze (Sanchez Vega et  
274 al., 2013); however, further work is necessary to determine the extent to which the changes in  
275 hippocampal structure influence learning and memory in this model.

276 Changes in hypothalamic structure have previously been documented in humans and animals  
277 following binge-like prenatal ethanol exposure (Coulter, Leech, Schaefer, Scheithauer, &  
278 Brumback, 1993; Fish et al., 2016). Our results show that ethanol exposure restricted to early  
279 pregnancy is sufficient to influence the volume of the hypothalamus in adult male offspring.  
280 The basis of this hypothalamic enlargement is unknown, but could involve increased  
281 neurogenesis similar to that reported in rats after low-moderate ethanol exposure in mid-late  
282 pregnancy (GD9-21) (Chang, Karatayev, Liang, Barson, & Leibowitz, 2012). Furthermore,  
283 deficits in hypothalamic function have been reported in rats after low-moderate ethanol  
284 exposures either late in gestation (Abate, Hernandez-Fonseca, Reyes-Guzman, Barbosa-Luna,  
285 & Mendez, 2014) or throughout gestation (Dembele, Yao, Chen, & Nyomba, 2006; Glavas,  
286 Ellis, Yu, & Weinberg, 2007).

287 The hypothalamus is an important regulator of metabolism and energy homeostasis and  
288 structural damage in this region has been linked with long-term weight gain and obesity in  
289 humans (Pinkney, Wilding, Williams, & MacFarlane, 2002). We found a significant increase  
290 in adiposity (18.7% or an average of 0.6 g) specifically in adult male offspring as a  
291 consequence of early pregnancy ethanol exposure; however, this did not translate into a  
292 substantial change in body weight, possibly due to a subtle (0.3 g) but non-significant  
293 decrease in lean tissue in the same group. A recent study using a similar, periconceptual,

294 ethanol exposure combined with a postnatal high-fat diet in rats reported similar, male-  
295 specific, effects on fat mass, fat-free mass and body weight (Gardebjer et al., 2017). It  
296 remains unclear whether the change in fat mass represents a uniform increase across all fat  
297 depots, or whether some depots are affected to a greater extent than others. Metabolic  
298 phenotyping revealed reduced energy expenditure, but no change in energy (food) intake, in  
299 adult ethanol-exposed offspring of both sexes, but tending to be stronger in males. The  
300 decrease in energy expenditure could be due to primary perturbations in basal metabolic rate,  
301 thermoregulation and/or physical activity; however, further work is required to determine  
302 whether one or all of these factors are involved. Further work is also necessary to discern  
303 whether the increased adiposity in ethanol-exposed males is a cause or consequence of their  
304 altered energy balance.

305 There is increasing evidence for sex-specific responses to a variety of adverse environmental  
306 exposures *in utero* (Bolton, Auten, & Bilbo, 2014; Giesbrecht, Letourneau, Campbell,  
307 Alberta Pregnancy, & Nutrition Study, 2016; Paolozza, Munn, Munoz, & Reynolds, 2015) as  
308 well as for intrinsic sex differences in energy metabolism and the development of obesity  
309 (Fried, Lee, & Karastergiou, 2015; Mauvais-Jarvis, 2015). In this study, early gestational  
310 ethanol exposure significantly affected fat mass specifically in adult male offspring. The  
311 mechanisms underpinning this difference between male and female offspring are not known,  
312 but could involve sex hormones and/or sexually dimorphic gene expression. We have  
313 previously identified male-specific changes in both gene expression and epigenetic state in  
314 the adult hippocampus following early gestational ethanol exposure (Zhang et al., 2015),  
315 lending support to the idea that similar modifications could occur in tissues (e.g.  
316 hypothalamus, adipose tissue) relevant to the phenotype described here.

317 In summary, our ultra-high field MRI study indicates that ethanol exposure early in  
318 pregnancy has limited effects on long-term brain structure in male mice, with only subtle

319 changes in the hippocampus and hypothalamus. Our results are also consistent with the idea  
320 that ethanol-induced changes in hypothalamic structure contribute to perturbations in energy  
321 metabolism and altered body composition in later life.

### 322 **Acknowledgements**

323 We thank Dr Brian Tse, Preclinical Imaging Facility, Translational Research Institute and Dr  
324 Johanna Barclay, Metabolic Medicine Group, Mater Research for technical assistance with  
325 the PIXImus2 and PhenoMaster, respectively. We thank Professor Emma Whitelaw for  
326 assistance with transcardial perfusions.

327 Funding: This work was supported by an Australian National Health and Medical Research  
328 Council grant (1003038) to S Chong, N Kaminen-Ahola, N D Kurniawan and G Galloway;  
329 and funding from the Mater Foundation to S Chong. C R Zhang was supported by  
330 scholarships from The University of Queensland and Mater Research. S Chong was  
331 supported by an Australian Research Council Future Fellowship (FT100100333). The 16.4T  
332 is supported by the Queensland State Government through the Queensland NMR Network,  
333 and the Australian Government through NCRIS and the National Imaging Facility. The  
334 Translational Research Institute is supported by a grant from the Australian Government.



335 **References**

- 336 Abate, P., Hernandez-Fonseca, K., Reyes-Guzman, A. C., Barbosa-Luna, I. G., & Mendez, M. (2014).  
337 Prenatal ethanol exposure alters met-enkephalin expression in brain regions related with  
338 reinforcement: possible mechanism for ethanol consumption in offspring. *Behav Brain Res*,  
339 274, 194-204. doi:10.1016/j.bbr.2014.08.022
- 340 Allan, A. M., Chynoweth, J., Tyler, L. A., & Caldwell, K. K. (2003). A mouse model of prenatal ethanol  
341 exposure using a voluntary drinking paradigm. *Alcohol Clin Exp Res*, 27(12), 2009-2016.  
342 doi:10.1097/01.ALC.0000100940.95053.72
- 343 Astley, S. J. (2010). Profile of the first 1,400 patients receiving diagnostic evaluations for fetal alcohol  
344 spectrum disorder at the Washington State Fetal Alcohol Syndrome Diagnostic & Prevention  
345 Network. *Can J Clin Pharmacol*, 17(1), e132-164.
- 346 Autti-Ramo, I., Autti, T., Korkman, M., Kettunen, S., Salonen, O., & Valanne, L. (2002). MRI findings in  
347 children with school problems who had been exposed prenatally to alcohol. *Dev Med Child*  
348 *Neurol*, 44(2), 98-106.
- 349 Belknap, J. K., Crabbe, J. C., & Young, E. R. (1993). Voluntary consumption of ethanol in 15 inbred  
350 mouse strains. *Psychopharmacology (Berl)*, 112(4), 503-510.
- 351 Bellinger, L., Lilley, C., & Langley-Evans, S. C. (2004). Prenatal exposure to a maternal low-protein  
352 diet programmes a preference for high-fat foods in the young adult rat. *Br J Nutr*, 92(3), 513-  
353 520.
- 354 Bolton, J. L., Auten, R. L., & Bilbo, S. D. (2014). Prenatal air pollution exposure induces sexually  
355 dimorphic fetal programming of metabolic and neuroinflammatory outcomes in adult  
356 offspring. *Brain Behav Immun*, 37, 30-44. doi:10.1016/j.bbi.2013.10.029
- 357 Borengasser, S. J., Kang, P., Fasje, J., Gomez-Acevedo, H., Blackburn, M. L., Badger, T. M., & Shankar,  
358 K. (2014). High fat diet and in utero exposure to maternal obesity disrupts circadian rhythm  
359 and leads to metabolic programming of liver in rat offspring. *PLoS One*, 9(1), e84209.  
360 doi:10.1371/journal.pone.0084209
- 361 Borengasser, S. J., Zhong, Y., Kang, P., Lindsey, F., Ronis, M. J., Badger, T. M., . . . Shankar, K. (2013).  
362 Maternal obesity enhances white adipose tissue differentiation and alters genome-scale  
363 DNA methylation in male rat offspring. *Endocrinology*, 154(11), 4113-4125.  
364 doi:10.1210/en.2012-2255
- 365 Cao, W., Li, W., Han, H., O'Leary-Moore, S. K., Sulik, K. K., Allan Johnson, G., & Liu, C. (2014). Prenatal  
366 alcohol exposure reduces magnetic susceptibility contrast and anisotropy in the white  
367 matter of mouse brains. *Neuroimage*, 102P2, 748-755.  
368 doi:10.1016/j.neuroimage.2014.08.035
- 369 Chang, G. Q., Karatayev, O., Liang, S. C., Barson, J. R., & Leibowitz, S. F. (2012). Prenatal ethanol  
370 exposure stimulates neurogenesis in hypothalamic and limbic peptide systems: possible  
371 mechanism for offspring ethanol overconsumption. *Neuroscience*, 222, 417-428.  
372 doi:10.1016/j.neuroscience.2012.05.066
- 373 Chen, L., & Nyomba, B. L. (2003). Glucose intolerance and resistin expression in rat offspring exposed  
374 to ethanol in utero: modulation by postnatal high-fat diet. *Endocrinology*, 144(2), 500-508.  
375 doi:10.1210/en.2002-220623
- 376 Coles, C. D., Goldstein, F. C., Lynch, M. E., Chen, X., Kable, J. A., Johnson, K. C., & Hu, X. (2011).  
377 Memory and brain volume in adults prenatally exposed to alcohol. *Brain Cogn*, 75(1), 67-77.  
378 doi:10.1016/j.bandc.2010.08.013
- 379 Coulter, C. L., Leech, R. W., Schaefer, G. B., Scheithauer, B. W., & Brumback, R. A. (1993). Midline  
380 cerebral dysgenesis, dysfunction of the hypothalamic-pituitary axis, and fetal alcohol effects.  
381 *Arch Neurol*, 50(7), 771-775.
- 382 Curhan, G. C., Willett, W. C., Rimm, E. B., Spiegelman, D., Ascherio, A. L., & Stampfer, M. J. (1996).  
383 Birth weight and adult hypertension, diabetes mellitus, and obesity in US men. *Circulation*,  
384 94(12), 3246-3250.

- 385 Dembele, K., Yao, X. H., Chen, L., & Nyomba, B. L. (2006). Intrauterine ethanol exposure results in  
386 hypothalamic oxidative stress and neuroendocrine alterations in adult rat offspring. *Am J*  
387 *Physiol Regul Integr Comp Physiol*, 291(3), R796-802. doi:10.1152/ajpregu.00633.2005
- 388 Dobson, C. C., Mongillo, D. L., Brien, D. C., Stepita, R., Poklewska-Koziell, M., Winterborn, A., . . .  
389 Reynolds, J. N. (2012). Chronic prenatal ethanol exposure increases adiposity and disrupts  
390 pancreatic morphology in adult guinea pig offspring. *Nutr Diabetes*, 2, e57.  
391 doi:10.1038/nutd.2012.31
- 392 Elswick, B. A., Welsch, F., & Janszen, D. B. (2000). Effect of different sampling designs on outcome of  
393 endocrine disruptor studies. *Reprod Toxicol*, 14(4), 359-367.
- 394 Fish, E. W., Holloway, H. T., Rumble, A., Baker, L. K., Wieczorek, L. A., Moy, S. S., . . . Parnell, S. E.  
395 (2016). Acute alcohol exposure during neurulation: Behavioral and brain structural  
396 consequences in adolescent C57BL/6J mice. *Behav Brain Res*, 311, 70-80.  
397 doi:10.1016/j.bbr.2016.05.004
- 398 Franke, K., Harder, T., Aerts, L., Melchior, K., Fahrenkrog, S., Rodekamp, E., . . . Plagemann, A. (2005).  
399 'Programming' of orexigenic and anorexigenic hypothalamic neurons in offspring of treated  
400 and untreated diabetic mother rats. *Brain Res*, 1031(2), 276-283.  
401 doi:10.1016/j.brainres.2004.11.006
- 402 Franklin, K. B. J., & Paxinos, G. (2008). *The mouse brain in stereotaxic coordinates* (3rd ed.).  
403 Amsterdam: Elsevier.
- 404 Fraser, A., Tilling, K., Macdonald-Wallis, C., Sattar, N., Brion, M. J., Benfield, L., . . . Lawlor, D. A.  
405 (2010). Association of maternal weight gain in pregnancy with offspring obesity and  
406 metabolic and vascular traits in childhood. *Circulation*, 121(23), 2557-2564.  
407 doi:10.1161/CIRCULATIONAHA.109.906081
- 408 Fried, S. K., Lee, M. J., & Karastergiou, K. (2015). Shaping fat distribution: New insights into the  
409 molecular determinants of depot- and sex-dependent adipose biology. *Obesity (Silver*  
410 *Spring)*, 23(7), 1345-1352. doi:10.1002/oby.21133
- 411 Fuglestad, A. J., Boys, C. J., Chang, P. N., Miller, B. S., Eckerle, J. K., Deling, L., . . . Wozniak, J. R.  
412 (2014). Overweight and obesity among children and adolescents with fetal alcohol spectrum  
413 disorders. *Alcohol Clin Exp Res*, 38(9), 2502-2508. doi:10.1111/acer.12516
- 414 Gardebjer, E. M., Anderson, S. T., Pantaleon, M., Wlodek, M. E., & Moritz, K. M. (2015). Maternal  
415 alcohol intake around the time of conception causes glucose intolerance and insulin  
416 insensitivity in rat offspring, which is exacerbated by a postnatal high-fat diet. *FASEB J*, 29(7),  
417 2690-2701. doi:10.1096/fj.14-268979
- 418 Gardebjer, E. M., Cuffe, J. S. M., Ward, L. C., Steane, S., Anderson, S. T., Dorey, E. S., . . . Moritz, K. M.  
419 (2017). The effects of periconceptional maternal alcohol intake and a postnatal high-fat diet  
420 on obesity and liver disease in male and female rat offspring. *Am J Physiol Endocrinol Metab*,  
421 ajpendo 00251 02017. doi:10.1152/ajpendo.00251.2017
- 422 Giesbrecht, G. F., Letourneau, N., Campbell, T. S., Alberta Pregnancy, O., & Nutrition Study, T. (2016).  
423 Sexually dimorphic and interactive effects of prenatal maternal cortisol and psychological  
424 distress on infant cortisol reactivity. *Dev Psychopathol*, 1-14.  
425 doi:10.1017/S0954579416000493
- 426 Glavas, M. M., Ellis, L., Yu, W. K., & Weinberg, J. (2007). Effects of prenatal ethanol exposure on basal  
427 limbic-hypothalamic-pituitary-adrenal regulation: role of corticosterone. *Alcohol Clin Exp*  
428 *Res*, 31(9), 1598-1610. doi:10.1111/j.1530-0277.2007.00460.x
- 429 Godin, E. A., O'Leary-Moore, S. K., Khan, A. A., Parnell, S. E., Ament, J. J., Dehart, D. B., . . . Sulik, K. K.  
430 (2009). Magnetic resonance microscopy defines ethanol-induced brain abnormalities in  
431 prenatal mice: effects of acute insult on gestational day 7. *Alcohol Clin Exp Res*, 34(1), 98-  
432 111. doi:10.1111/j.1530-0277.2009.01071.x
- 433 Kaminen-Ahola, N., Ahola, A., Flatscher-Bader, T., Wilkins, S. J., Anderson, G. J., Whitelaw, E., &  
434 Chong, S. (2010). Postnatal growth restriction and gene expression changes in a mouse

- 435 model of fetal alcohol syndrome. *Birth Defects Res A Clin Mol Teratol*, 88(10), 818-826.  
436 doi:10.1002/bdra.20729
- 437 Kaminen-Ahola, N., Ahola, A., Maga, M., Mallitt, K. A., Fahey, P., Cox, T. C., . . . Chong, S. (2010).  
438 Maternal ethanol consumption alters the epigenotype and the phenotype of offspring in a  
439 mouse model. *PLoS Genet*, 6(1), e1000811. doi:10.1371/journal.pgen.1000811
- 440 Kurniawan, N. D., Richards, K. L., Yang, Z., She, D., Ullmann, J. F., Moldrich, R. X., . . . Reutens, D. C.  
441 (2014). Visualization of mouse barrel cortex using ex-vivo track density imaging.  
442 *Neuroimage*, 87, 465-475. doi:10.1016/j.neuroimage.2013.09.030
- 443 Law, C. M., Barker, D. J., Osmond, C., Fall, C. H., & Simmonds, S. J. (1992). Early growth and  
444 abdominal fatness in adult life. *J Epidemiol Community Health*, 46(3), 184-186.
- 445 Lukaszewski, M. A., Mayeur, S., Fajardy, I., Delahaye, F., Dutriez-Casteloot, I., Montel, V., . . . Breton,  
446 C. (2011). Maternal prenatal undernutrition programs adipose tissue gene expression in  
447 adult male rat offspring under high-fat diet. *Am J Physiol Endocrinol Metab*, 301(3), E548-  
448 559. doi:10.1152/ajpendo.00011.2011
- 449 Lunde, E. R., Washburn, S. E., Golding, M. C., Bake, S., Miranda, R. C., & Ramadoss, J. (2016). Alcohol-  
450 Induced Developmental Origins of Adult-Onset Diseases. *Alcohol Clin Exp Res*, 40(7), 1403-  
451 1414. doi:10.1111/acer.13114
- 452 Ma, Y., Hof, P. R., Grant, S. C., Blackband, S. J., Bennett, R., Slate, L., . . . Benveniste, H. (2005). A  
453 three-dimensional digital atlas database of the adult C57BL/6J mouse brain by magnetic  
454 resonance microscopy. *Neuroscience*, 135(4), 1203-1215.  
455 doi:10.1016/j.neuroscience.2005.07.014
- 456 Mauvais-Jarvis, F. (2015). Sex differences in metabolic homeostasis, diabetes, and obesity. *Biol Sex*  
457 *Differ*, 6, 14. doi:10.1186/s13293-015-0033-y
- 458 McCormack, C., Hutchinson, D., Burns, L., Wilson, J., Elliott, E., Allsop, S., . . . Mattick, R. (2017).  
459 Prenatal Alcohol Consumption Between Conception and Recognition of Pregnancy. *Alcohol*  
460 *Clin Exp Res*, 41(2), 369-378. doi:10.1111/acer.13305
- 461 Meyer, C. W., Reitmeir, P., & Tschop, M. H. (2015). Exploration of Energy Metabolism in the Mouse  
462 Using Indirect Calorimetry: Measurement of Daily Energy Expenditure (DEE) and Basal  
463 Metabolic Rate (BMR). *Curr Protoc Mouse Biol*, 5(3), 205-222.  
464 doi:10.1002/9780470942390.mo140216
- 465 Moore, E. M., Migliorini, R., Infante, M. A., & Riley, E. P. (2014). Fetal Alcohol Spectrum Disorders:  
466 Recent Neuroimaging Findings. *Curr Dev Disord Rep*, 1(3), 161-172. doi:10.1007/s40474-014-  
467 0020-8
- 468 O'Leary-Moore, S. K., Parnell, S. E., Godin, E. A., Dehart, D. B., Ament, J. J., Khan, A. A., . . . Sulik, K. K.  
469 (2010). Magnetic resonance microscopy-based analyses of the brains of normal and ethanol-  
470 exposed fetal mice. *Birth Defects Res A Clin Mol Teratol*, 88(11), 953-964.  
471 doi:10.1002/bdra.20719
- 472 O'Leary-Moore, S. K., Parnell, S. E., Lipinski, R. J., & Sulik, K. K. (2011). Magnetic resonance-based  
473 imaging in animal models of fetal alcohol spectrum disorder. *Neuropsychol Rev*, 21(2), 167-  
474 185. doi:10.1007/s11065-011-9164-z
- 475 Paolozza, A., Munn, R., Munoz, D. P., & Reynolds, J. N. (2015). Eye movements reveal sexually  
476 dimorphic deficits in children with fetal alcohol spectrum disorder. *Front Neurosci*, 9, 76.  
477 doi:10.3389/fnins.2015.00076
- 478 Parnell, S. E., O'Leary-Moore, S. K., Godin, E. A., Dehart, D. B., Johnson, B. W., Allan Johnson, G., . . .  
479 Sulik, K. K. (2009). Magnetic resonance microscopy defines ethanol-induced brain  
480 abnormalities in prenatal mice: effects of acute insult on gestational day 8. *Alcohol Clin Exp*  
481 *Res*, 33(6), 1001-1011. doi:10.1111/j.1530-0277.2009.00921.x
- 482 Phillips, D. I., Barker, D. J., Hales, C. N., Hirst, S., & Osmond, C. (1994). Thinness at birth and insulin  
483 resistance in adult life. *Diabetologia*, 37(2), 150-154.
- 484 Pinkney, J., Wilding, J., Williams, G., & MacFarlane, I. (2002). Hypothalamic obesity in humans: what  
485 do we know and what can be done? *Obes Rev*, 3(1), 27-34.

- 486 R Development Core Team. (2010). R: A language and environment for statistical computing. Vienna,  
487 Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org>
- 488 Ravelli, A. C., van Der Meulen, J. H., Osmond, C., Barker, D. J., & Bleker, O. P. (1999). Obesity at the  
489 age of 50 y in men and women exposed to famine prenatally. *Am J Clin Nutr*, *70*(5), 811-816.
- 490 Sanchez Vega, M. C., Chong, S., & Burne, T. H. (2013). Early gestational exposure to moderate  
491 concentrations of ethanol alters adult behaviour in C57BL/6J mice. *Behav Brain Res*, *252*,  
492 326-333. doi:10.1016/j.bbr.2013.06.003
- 493 Sutton, G. M., Centanni, A. V., & Butler, A. A. (2010). Protein malnutrition during pregnancy in  
494 C57BL/6J mice results in offspring with altered circadian physiology before obesity.  
495 *Endocrinology*, *151*(4), 1570-1580. doi:10.1210/en.2009-1133
- 496 Wang, X., & Kroenke, C. D. (2015). Utilization of Magnetic Resonance Imaging in Research Involving  
497 Animal Models of Fetal Alcohol Spectrum Disorders. *Alcohol Res*, *37*(1), 39-51.
- 498 Werts, R. L., Van Calcar, S. C., Wargowski, D. S., & Smith, S. M. (2014). Inappropriate feeding  
499 behaviors and dietary intakes in children with fetal alcohol spectrum disorder or probable  
500 prenatal alcohol exposure. *Alcohol Clin Exp Res*, *38*(3), 871-878. doi:10.1111/acer.12284
- 501 Willoughby, K. A., Sheard, E. D., Nash, K., & Rovet, J. (2008). Effects of prenatal alcohol exposure on  
502 hippocampal volume, verbal learning, and verbal and spatial recall in late childhood. *J Int  
503 Neuropsychol Soc*, *14*(6), 1022-1033. doi:10.1017/S1355617708081368
- 504 Yushkevich, P. A., Piven, J., Hazlett, H. C., Smith, R. G., Ho, S., Gee, J. C., & Gerig, G. (2006). User-  
505 guided 3D active contour segmentation of anatomical structures: significantly improved  
506 efficiency and reliability. *Neuroimage*, *31*(3), 1116-1128.  
507 doi:10.1016/j.neuroimage.2006.01.015
- 508 Zhang, C. R., Ho, M. F., Vega, M. C., Burne, T. H., & Chong, S. (2015). Prenatal ethanol exposure alters  
509 adult hippocampal VGLUT2 expression with concomitant changes in promoter DNA  
510 methylation, H3K4 trimethylation and miR-467b-5p levels. *Epigenetics Chromatin*, *8*, 40.  
511 doi:10.1186/s13072-015-0032-6
- 512 Zhao, P., Liu, E., Qiao, Y., Katzmarzyk, P. T., Chaput, J. P., Fogelholm, M., . . . Group, I. R. (2016).  
513 Maternal gestational diabetes and childhood obesity at age 9-11: results of a multinational  
514 study. *Diabetologia*. doi:10.1007/s00125-016-4062-9

515

516

517 **Figure Legends**

518 **Figure 1.** Maternal liquid consumption per day **(a)** and % weight gain **(b)** over the 8-day  
519 ethanol exposure period (0.5-8.5 dpc). **(c)** Litter size at weaning (P21). EtOH: ethanol-  
520 exposed mice. All data points are overlaid on box and whisker graphs where the box extends  
521 from the 25th to 75th percentile, the line in the middle is the median and the whiskers show  
522 the maximum and minimum values obtained.

523 **Figure 2.** Body weight and body composition of adult (12 week old) male and female  
524 offspring. Control offspring are indicated by open circles ( $\circ$ ) and ethanol-exposed (EtOH)  
525 offspring are indicated by filled circles ( $\bullet$ ). **(a)** Body weight of control males (n=21 from 6  
526 litters), EtOH males (n=23 from 7 litters), control females (n=23 from 6 litters) and EtOH  
527 females (n=29 from 7 litters). Lean mass **(b)**, fat mass **(c)** and % fat **(d)** are also shown for a  
528 subset of control males (n=11 from 6 litters), EtOH males (n=14 from 7 litters), control  
529 females (n=12 from 6 litters) and EtOH females (n=16 from 7 litters). All data points are  
530 overlaid on box and whisker graphs where the box extends from the 25th to 75th percentile,  
531 the line in the middle is the median and the whiskers show the maximum and minimum  
532 values obtained; Tukey's multiple comparisons test  $*P<0.05$ ,  $****P<0.0001$ .

533 **Figure 3.** Bone mineral content (BMC), bone area and bone mineral density (BMD) of adult  
534 (12 week old) male and female offspring. Control offspring are indicated by open circles ( $\circ$ )  
535 and ethanol-exposed (EtOH) offspring are indicated by filled circles ( $\bullet$ ). BMC **(a)**, bone area  
536 **(b)** and BMD **(c)** are plotted for four groups of mice consisting of control males (n=11 from 6  
537 litters), EtOH males (n=14 from 7 litters), control females (n=12 from 6 litters) and EtOH  
538 females (n=16 from 7 litters). All data points are overlaid on box and whisker graphs where  
539 the box extends from the 25th to 75th percentile, the line in the middle is the median and the  
540 whiskers show the maximum and minimum values obtained; Tukey's multiple comparisons  
541 test  $***P<0.001$ ,  $****P<0.0001$ .

542 **Figure 4.** Energy metabolism in adult (21-26 week old) male and female offspring. Control  
543 offspring are indicated by open circles ( $\circ$ ) and ethanol-exposed (EtOH) offspring are  
544 indicated by filled circles ( $\bullet$ ). **(a)** Oxygen consumption ( $\text{VO}_2$ ), **(b)** carbon dioxide production  
545 ( $\text{VCO}_2$ ), **(c)** respiratory exchange ratio (RER) and **(d)** daily energy expenditure. Control  
546 male (n=10 from 6 litters), EtOH male (n=9 from 7 litters), control female (n=10 from 6  
547 litters) and EtOH female (n=12 from 7 litters) data points are overlaid on box and whisker  
548 graphs where the box extends from the 25th to 75th percentile, the line in the middle is the  
549 median and the whiskers show the maximum and minimum values obtained; Tukey's  
550 multiple comparisons test \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , \*\*\*\* $P < 0.0001$ .

551 **Figure 5.** Water and food intake by adult (21-26 week old) male and female offspring.  
552 Control offspring are indicated by open circles ( $\circ$ ) and ethanol-exposed (EtOH) offspring are  
553 indicated by filled circles ( $\bullet$ ). **(a-b)** water intake and **(c-d)** food intake were analysed  
554 separately for the light and dark phases of the light cycle. Control male (n=10 from 6 litters),  
555 EtOH male (n=9 from 7 litters), control female (n=10 from 6 litters) and EtOH female (n=12  
556 from 7 litters) data points are overlaid on box and whisker graphs where the box extends from  
557 the 25th to 75th percentile, the line in the middle is the median and the whiskers show the  
558 maximum and minimum values obtained; Tukey's multiple comparisons test \*\*\*\* $P < 0.0001$ .



**Table 2.** Volumetric analysis of hippocampal subfields in male offspring at P64 (n=11 per group from 9 litters)

Hippocampal subfield	Relative Volume <sup>a</sup>		P-value
	Control	Ethanol-exposed	
Dentate gyrus	0.0119 ± 0.00121	0.0116 ± 0.000506	0.47
CA1 and CA2	0.0265 ± 0.00206	0.0265 ± 0.00159	1.00
CA3	0.00362 ± 0.00118	0.00382 ± 0.000929	0.67

<sup>a</sup>Mean±SD. All volumes were normalised to whole brain volume.

**Table 3.** DTI analysis at P28 (n=5 per group from 2-3 litters)

Region	Parameter	Mean $\pm$ SD		P-value
		Control	Ethanol-exposed	
AC	FA	0.36 $\pm$ 0.06	0.37 $\pm$ 0.01	0.55
	FT number	762 $\pm$ 305	547 $\pm$ 222	0.24
	FT Volume <sup>a</sup>	3 109 $\pm$ 1 056	2 258 $\pm$ 679	0.17
	FT Length (mm)	7.99 $\pm$ 1.98	6.57 $\pm$ 1.72	0.26
CC	FA	0.36 $\pm$ 0.03	0.35 $\pm$ 0.03	0.40
	FT number	6 920 $\pm$ 1 829	6 197 $\pm$ 2 823	0.65
	FT Volume <sup>a</sup>	19 663 $\pm$ 4 261	18 341 $\pm$ 6 902	0.73
	FT Length (mm)	3.77 $\pm$ 0.88	3.64 $\pm$ 1.04	0.84
HC	FA	0.42 $\pm$ 0.03	0.40 $\pm$ 0.03	0.29
	FT number	3 293 $\pm$ 548	2 777 $\pm$ 731	0.25
	FT Volume <sup>a</sup>	8 776 $\pm$ 2 235	7 236 $\pm$ 1 588	0.25
	FT Length (mm)	5.00 $\pm$ 0.32	4.50 $\pm$ 0.30	0.03*
IC	FA	0.30 $\pm$ 0.01	0.30 $\pm$ 0.01	1.00
	FT number	5 825 $\pm$ 518	4 562 $\pm$ 1 035	0.05
	FT Volume <sup>a</sup>	14 522 $\pm$ 1 235	12 164 $\pm$ 2 357	0.09
	FT Length (mm)	5.22 $\pm$ 0.37	4.76 $\pm$ 0.30	0.07
OT	FA	0.44 $\pm$ 0.02	0.42 $\pm$ 0.02	0.29
	FT number	429 $\pm$ 105	415 $\pm$ 132	0.37
	FT Volume <sup>a</sup>	1 588 $\pm$ 153	1 421 $\pm$ 267	0.74
	FT Length (mm)	3.62 $\pm$ 0.23	3.71 $\pm$ 0.41	0.78
TH	FA	0.21 $\pm$ 0.02	0.19 $\pm$ 0.02	0.97
	FT number	1 251 $\pm$ 324	1 128 $\pm$ 362	0.54
	FT Volume <sup>a</sup>	3 148 $\pm$ 681	2 989 $\pm$ 813	0.87
	FT Length (mm)	1.48 $\pm$ 0.23	1.34 $\pm$ 0.16	0.82

SD: standard deviation, AC: anterior commissure, CC: corpus callosum, HC: hippocampal commissure, IC: internal capsule, OT: optic tract, TH: thalamus, FA: fractional anisotropy and FT: fibre tract. <sup>a</sup>Volume is shown as voxel numbers. Each voxel is 10<sup>-6</sup> mm<sup>3</sup>. \*P<0.05.



**Table 4.** DTI analysis at P64 (n=11 per group from 9 litters)


Region	Parameter	Mean $\pm$ SD		P-value
		Control	Ethanol-exposed	
AC	FA	0.45 $\pm$ 0.05	0.45 $\pm$ 0.04	0.79
	FT number	457 $\pm$ 83	534 $\pm$ 160	0.38
	FT Volume <sup>a</sup>	2 334 $\pm$ 331	2 600 $\pm$ 839	0.54
	FT Length (mm)	7.89 $\pm$ 0.76	6.25 $\pm$ 1.45	0.07
CC	FA	0.43 $\pm$ 0.01	0.44 $\pm$ 0.01	0.62
	FT number	7 008 $\pm$ 612	6 919 $\pm$ 621	0.82
	FT Volume <sup>a</sup>	19 904 $\pm$ 1 469	19 454 $\pm$ 1 801	0.68
	FT Length (mm)	6.23 $\pm$ 0.81	5.57 $\pm$ 0.28	0.15
HC	FA	0.42 $\pm$ 0.01	0.42 $\pm$ 0.02	0.73
	FT number	4 613 $\pm$ 884	4 447 $\pm$ 848	0.77
	FT Volume <sup>a</sup>	10 702 $\pm$ 1 733	10 562 $\pm$ 2 153	0.91
	FT Length (mm)	5.96 $\pm$ 0.51	5.93 $\pm$ 0.55	0.93
IC	FA	0.32 $\pm$ 0.02	0.32 $\pm$ 0.02	0.89
	FT number	3 157 $\pm$ 888	2 550 $\pm$ 741	0.28
	FT Volume <sup>a</sup>	8 625 $\pm$ 2 049	7 438 $\pm$ 1 350	0.32
	FT Length (mm)	4.31 $\pm$ 0.43	4.60 $\pm$ 0.54	0.38
OT	FA	0.45 $\pm$ 0.02	0.44 $\pm$ 0.02	0.33
	FT number	498 $\pm$ 109	428 $\pm$ 134	0.39
	FT Volume <sup>a</sup>	1 944 $\pm$ 211	1 864 $\pm$ 327	0.66
	FT Length (mm)	3.68 $\pm$ 0.33	4.08 $\pm$ 0.57	0.22
TH	FA	0.20 $\pm$ 0.02	0.19 $\pm$ 0.03	0.65
	FT number	1 354 $\pm$ 378	1 136 $\pm$ 388	0.39
	FT Volume <sup>a</sup>	4 449 $\pm$ 789	3 789 $\pm$ 960	0.27
	FT Length (mm)	2.88 $\pm$ 0.39	2.54 $\pm$ 0.30	0.17


SD: standard deviation, AC: anterior commissure, CC: corpus callosum, HC: hippocampal commissure, IC: internal capsule, OT: optic tract, TH: thalamus, FA: fractional anisotropy and FT: fibre tract. <sup>a</sup>Volume is shown as voxel numbers. Each voxel is 10<sup>-6</sup> mm<sup>3</sup>.


**Table 1.** Relative volumes of selected brain regions in adolescent (P28) and adult (P64) male offspring


Brain region	Relative volume <sup>a</sup>		Raw <i>P</i> -value	Adjusted <i>P</i> -value
	Control	Ethanol-exposed		
P28 (n=5/group from 2-3 litters)				
Neocortex	0.343 ± 0.00910	0.342 ± 0.00600	0.79	0.99
Cerebellum	0.111 ± 0.00616	0.109 ± 0.00926	0.71	0.99
Caudate putamen	0.0536 ± 0.000673	0.0535 ± 0.00170	0.90	0.99
Hippocampus	0.0503 ± 0.00160	0.0505 ± 0.00360	0.94	0.99
Hypothalamus	0.0211 ± 0.00163	0.0217 ± 0.00223	0.65	0.99
P64 (n=11/group from 9 litters)				
Neocortex	0.314 ± 0.00563	0.310 ± 0.0108	0.40	0.68
Cerebellum	0.123 ± 0.00559	0.122 ± 0.00393	0.59	0.68
Caudate putamen	0.0534 ± 0.00258	0.0523 ± 0.00257	0.32	0.68
Hippocampus	0.0533 ± 0.00141	0.0515 ± 0.00159	0.011*	0.053
Hypothalamus	0.0228 ± 0.000830	0.0235 ± 0.000808	0.045*	0.17


<sup>a</sup>Mean±SD. All volumes were normalized to whole brain volume to minimize the effects of subtle inter-individual variation. \**P*<0.05











**1 Highlights**

- 2 • The long-term effects of early pregnancy alcohol exposure were examined in mice.
- 3 • Hippocampal and hypothalamic structure were altered in adult male offspring.
- 4 • Energy metabolism and body composition were also changed in adult males.
- 5 • CNS abnormalities may underpin other outcomes linked to prenatal alcohol exposure.