

<https://helda.helsinki.fi>

Weak effects of geolocators on small birds : a meta-analysis controlled for phylogeny and publication bias

Brlík, Vojtěch

2020-01

Brlík, V., Koleček, J., Burgess, M., Hahn, S., Humple, D., Krist, M., Weiser, E. L., Adamík, P., Alves, J. A., Arlt, D., Bariaei, S., Becker, V., Both, C., Bravo, S. P., Briedis, M., Chutný, B., Čikovi, D., Cueto, V. R., Emmenegger, T., Fraser, K., Gilg, O., Guerrero, M., Hallworth, M. T., Hewson, C., Jiguet, F., Johnson, J. A., Kelly, T., Kishkinev, D., Leconte, M., Lislevand, T., Lisovski, S., López, C., McFarland, K. P., Marra, P. P., Matsuoka, S. M., Matyjasiak, P., Metzger, B., Monrós, J. S., Neumann, R., Newman, A., Norris, R., Pärt, T., Pavel, V., Perlut, N., Piha, M., Reneerkens, J., Rimmer, C. C., Roberto-Charron, A., Scandolara, C., Sokolova, N., Takenaka, M., Tolkmitt, D., van Oosten, H., Wellbrock, A. H. J., Wheeler, H., van der Winden, J., Witte, K., Woodworth, B. & Procházka, P. 2020, 'Weak effects of geolocators on small birds : a meta-analysis controlled for phylogeny and publication bias', *Journal of Animal Ecology*, vol. 89, no. 1, pp. 207-220. <https://doi.org/10.1111/1365-2656.12962>

<http://hdl.handle.net/10138/311721>

<https://doi.org/10.1111/1365-2656.12962>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Journal of Animal Ecology

MR VOJTĚCH BRLÍK (Orcid ID : 0000-0002-7902-8123)

DR JAROSLAV KOLEČEK (Orcid ID : 0000-0003-1069-6593)

DR MILOŠ KRIST (Orcid ID : 0000-0002-6183-686X)

MISS JANNE OUWEHAND (Orcid ID : 0000-0003-2573-6287)

DR JOSE A ALVES (Orcid ID : 0000-0001-7182-0936)

MR NATHAN WANDS COOPER (Orcid ID : 0000-0002-4667-1542)

DR MICHAEL T. HALLWORTH (Orcid ID : 0000-0002-6385-3815)

DR CHRIS M. HEWSON (Orcid ID : 0000-0002-8493-5203)

DR SIMEON LIISOVSKI (Orcid ID : 0000-0002-6399-0035)

Article type : Research Article

Handling Editor: Jenny Dunn

Weak effects of geolocators on small birds: a meta-analysis controlled for phylogeny and publication bias

Vojtěch Brlík^{1,2}, Jaroslav Koleček¹, Malcolm Burgess³, Steffen Hahn⁴, Diana Humple⁵, Miloš Krist⁶,

Janne Ouwehand⁷, Emily L. Weiser^{8,9}, Peter Adamík^{6,10}, José A. Alves^{11,12}, Debora Arlt¹³, Sanja

Barišić¹⁴, Detlef Becker¹⁵, Eduardo J. Belda¹⁶, Václav Beran^{6,17,18}, Christiaan Both⁷, Susana P. Bravo¹⁹,

Martins Briedis⁴, Bohumír Chutný²⁰, Davor Ćiković¹⁴, Nathan W. Cooper²¹, Joana S. Costa¹¹, Víctor R.

Cueto¹⁹, Tamara Emmenegger⁴, Kevin Fraser²², Olivier Gilg^{23,24}, Marina Guerrero²⁵, Michael T.

Hallworth²⁶, Chris Hewson²⁷, Frédéric Jiguet²⁸, James A. Johnson²⁹, Tosha Kelly³⁰, Dmitry

Kishkinev^{31,32}, Michel Leconte³³, Terje Lislevand³⁴, Simeon Lisovski⁴, Cosme López³⁵, Kent P.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1365-2656.12962

This article is protected by copyright. All rights reserved.

McFarland³⁶, Peter P. Marra²⁶, Steven M. Matsuoka^{29,37}, Piotr Matyjasiak³⁸, Christoph M. Meier⁴, Benjamin Metzger³⁹, Juan S. Monrós⁴⁰, Roland Neumann⁴¹, Amy Newman⁴², Ryan Norris⁴², Tomas Pärt¹³, Václav Pavel^{6,43}, Noah Perlut⁴⁴, Markus Piha⁴⁵, Jeroen Reneerkens⁷, Christopher C. Rimmer³⁶, Amélie Roberto-Charron²², Chiara Scandolaro⁴, Natalia Sokolova^{46,47}, Makiko Takenaka⁴⁸, Dirk Tolkmitt⁴⁹, Herman van Oosten^{50,51}, Arndt H. J. Wellbrock⁵², Hazel Wheeler⁵³, Jan van der Winden⁵⁴, Klaudia Witte⁵², Brad Woodworth⁵⁵, Petr Procházka¹

Author for correspondence: Vojtěch Brlík, The Czech Academy of Sciences, Institute of Vertebrate Biology, Květná 8, CZ-603 65 Brno, Czech Republic. E-mail: vojtech.brlik@gmail.com

Affiliations

¹ The Czech Academy of Sciences, Institute of Vertebrate Biology, Květná 8, 603 65 Brno, Czech Republic

² Department of Ecology, Faculty of Science, Charles University in Prague, Viničná 7, 128 44 Prague 2, Czech Republic

³ Royal Society for the Protection of Birds – Centre for Conservation Science, The Lodge, Sandy, SG19 2DL Beds, UK

⁴ Bird Migration Department, Swiss Ornithological Institute, Seerose 1, 6204 Sempach, Switzerland

⁵ Point Blue Conservation Science, 3820 Cypress Drive 11, Petaluma, California 94954, USA

⁶ Department of Zoology, Faculty of Science, Palacký University, tř. 17. listopadu 50, 771 46 Olomouc, Czech Republic

⁷ Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands

⁸ Kansas State University, Division of Biology, 116 Ackert Hall, Manhattan, Kansas 66506, USA

⁹ U.S. Geological Survey, Upper Midwest Environmental Sciences Center, 2630 Fanta Reed Rd, La Crosse, Wisconsin 54603, USA

¹⁰ Museum of Natural History, nám. Republiky 5, 771 73 Olomouc, Czech Republic

¹¹ Department of Biology & Centre for Environmental and Marine Studies, University of Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

¹² University of Iceland, South Iceland Research Centre, Lindarbraut 4, IS-840 Laugarvatn, Iceland

¹³ Department of Ecology, Swedish University of Agricultural Sciences, PO Box 7044, 75007 Uppsala, Sweden

¹⁴ Institute of Ornithology, Croatian Academy of Sciences and Arts, Gundulićeva 24, 10000 Zagreb, Croatia

¹⁵ Museum Heineanum, Domplatz 36, 38820 Halberstadt, Germany

¹⁶ Universitat Politècnica de València, C/ Paranimfo, 1, 46730 Gandia, Valencia, Spain

¹⁷ Municipal Museum of Ústí nad Labem, Masarykova 1000/3, 40001 Ústí nad Labem, Czech Republic

¹⁸ ALKA Wildlife o.p.s., Liděřovice 62, 38001 Dačice, Czech Republic

¹⁹ CIEMEP, CONICET/UNPSJB, Roca 780, Esquel, CP 9200, Chubut, Argentina

²⁰ Malinová 1650/27, 10600 Prague 10, Czech Republic

²¹ Migratory Bird Center, Smithsonian Conservation Biology Institute, National Zoological Park, PO Box 37012 MRC 5503, Washington, D.C. 20013, USA

²² Avian Behaviour and Conservation Lab, Department of Biological Sciences, University of Manitoba, 50 Sifton Road, Winnipeg, Manitoba R3T 2N2, Canada

²³ UMR 6249 Chrono-environnement, Université de Bourgogne Franche-Comté, 16 route de Gray,
25000 Besançon, France

²⁴ Groupe de recherche en Ecologie Arctique, 16 rue de Vernot, 21440 Francheville, France

²⁵ Servicio de Jardines, Bosques y Huertas, Patronato de la Alhambra y el Generalife.C/ Real de la
Alhambra, 18009 Granada, Spain

²⁶ Migratory Bird Center – Smithsonian Conservation Biology Institute, National Zoological Park,
Washington DC 20013, USA

²⁷ British Trust for Ornithology, The Nunnery, Thetford, Norfolk IP24 2PU, UK

²⁸ UMR7204 CESCO, MNHN-CNRS-Sorbonne Université, CP135, 43 Rue Buffon, 75005 Paris, France

²⁹ U.S. Fish and Wildlife Service, Migratory Bird Management, 1011 East Tudor Road, Anchorage,
Alaska 99503, USA

³⁰ Advanced Facility for Avian Research, Western University, 32 Wellington Dr, N6G 4W4, London,
Ontario, Canada

³¹ School of Natural Sciences, Bangor University, Deiniol Road, Bangor, LL57 2UW, Gwynedd, UK

³² Biological station Rybachy, Zoological Institute of Russian Academy of Sciences, Rybachy,
Kaliningrad region 238535, Russia

³³ Quartier du Caü, F-64260 Arudy, France

³⁴ University Museum of Bergen, Department of Natural History, University of Bergen, PO Box 7800,
5020 Bergen, Norway

³⁵ Department of Zoology, Faculty of Biology, Green Building, Avenue Reina Mercedes, 41012 Seville,
Spain

³⁶ Vermont Center for Ecostudies, PO Box 420, Norwich, 05055 Vermont, USA

³⁷ U.S. Geological Survey Alaska Science Center, 4210 University Drive, Anchorage, Alaska 99508, USA

³⁸ Department of Evolutionary Biology, Faculty of Biology and Environmental Sciences, Cardinal Stefan Wyszyński University in Warsaw, Wóycickiego 1/3, PL-01-938 Warsaw, Poland

³⁹ Rua da Esperanca 43/3D, 1200-655 Lisbon, Portugal

⁴⁰ Cavanilles Institute of Biodiversity and Evolutionary Biology, University of Valencia, C/ Catedrático José Beltrán 2, E-46980 Paterna, València, Spain

⁴¹ Kritzmower Weg 1, 18198 Stäbelow, Germany

⁴² Department of Integrative Biology, University of Guelph, 50 Stone Rd E, Guelph, ON N1G 2W1, Canada

⁴³ Centre for Polar Ecology, University of South Bohemia, Branišovská 31, 370 05 České Budějovice, Czech Republic

⁴⁴ University of New England, Department of Environmental Studies, 11 Hills Beach Rd, Biddeford, Maine, USA

⁴⁵ Finnish Museum of Natural History LUOMUS, University of Helsinki, PO Box 17, 00014 Helsinki, Finland

⁴⁶ Arctic Research Station of Institute of Plant and Animal Ecology, Ural Branch Russian Academy of Sciences, Zelenaya Gorka Str. 21, 629400 Labytnangi, Russia

⁴⁷ Arctic Research Center of Yamal-Nenets Autonomous District, Respublika str. 73, 629008 Salekhard, Russia

⁴⁸ Tokai University Sapporo Campus, Minamisawa 5-1-1-1, Minami-ku, Sapporo, Hokkaido 005-8601, Japan

⁴⁹ Menckestraße 34, 04155 Leipzig, Germany

⁵⁰ Oenanthe Ecologie, Hollandseweg 42, 6706 KR Wageningen, The Netherlands

⁵¹ Institute for Water and Wetland Research, Animal Ecology, Physiology & Experimental Plant Ecology, Radboud University, PO Box 9100, 6500 GL Nijmegen, The Netherlands

⁵² Institute of Biology, Department of Chemistry – Biology, Faculty of Science and Technology, University of Siegen, Adolf-Reichwein-Str. 2, 57076 Siegen, Germany

⁵³ Wildlife Preservation Canada, 5420 Highway 6 North, Guelph, Ontario N1H 6J2, Canada

⁵⁴ Ecology Research & Consultancy, Dantelaan 115, 3533 VC Utrecht, The Netherlands

⁵⁵ School of Biological Sciences, The University of Queensland, Brisbane, Queensland, Australia

ORCID

Vojtěch Brlík: <https://orcid.org/0000-0002-7902-8123>

Jaroslav Koleček: <https://orcid.org/0000-0003-1069-6593>

Malcolm Burgess: <https://orcid.org/0000-0003-1288-1231>

Steffen Hahn: <https://orcid.org/0000-0002-4924-495X>

Miloš Krist: <https://orcid.org/0000-0002-6183-686X>

Janne Ouwehand: <https://orcid.org/0000-0003-2573-6287>

Emily L. Weiser: <https://orcid.org/0000-0003-1598-659X>

Peter Adamík: <https://orcid.org/0000-0003-1566-1234>

José A. Alves: <https://orcid.org/0000-0001-7182-0936>

Debora Arlt: <https://orcid.org/0000-0003-0874-4250>

Sanja Barišić: <https://orcid.org/0000-0003-3472-3285>

Eduardo J. Belda: <https://orcid.org/0000-0003-1995-1271>

Christiaan Both: <https://orcid.org/0000-0001-7099-9831>

Martins Briedis: <https://orcid.org/0000-0002-9434-9056>

Davor Ćiković: <https://orcid.org/0000-0002-3234-0574>

Joana S. Costa: <https://orcid.org/0000-0002-1532-8936>

Tamara Emmenegger: <https://orcid.org/0000-0002-2839-6129>

Olivier Gilg: <https://orcid.org/0000-0002-9083-4492>

Chris Hewson: <https://orcid.org/0000-0002-8493-5203>

Frédéric Jiguet: orcid.org/0000-0002-0606-7332

Dmitry Kishkinev: <https://orcid.org/0000-0002-2619-1197>

Terje Lislevand: <https://orcid.org/0000-0003-1281-7061>

Piotr Matyjasiak: <https://orcid.org/0000-0003-0384-2935>

Kent McFarland: <https://orcid.org/0000-0001-7809-5503>

Christoph M. Meier: <https://orcid.org/0000-0001-9584-2339>

Tomas Pärt: <https://orcid.org/0000-0001-7388-6672>

Markus Piha: <https://orcid.org/0000-0002-8482-6162>

Jeroen Reneerkens: <https://orcid.org/0000-0003-0674-8143>

Natalia Sokolova: <https://orcid.org/0000-0002-6692-4375>

Arndt H. J. Wellbrock: <https://orcid.org/0000-0001-9929-7091>

Klaudia Witte: <https://orcid.org/0000-0002-2812-9936>

Running head: Geolocator effects on small birds

Abstract

1. Currently, the deployment of tracking devices is one of the most frequently used approaches to study movement ecology of birds. Recent miniaturisation of light-level geolocators enabled studying small bird species whose migratory patterns were widely unknown. However, geolocators may reduce vital rates in tagged birds and may bias obtained movement data.
2. There is a need for a thorough assessment of the potential tag effects on small birds, as previous meta-analyses did not evaluate unpublished data and impact of multiple life-history traits, focused mainly on large species and the number of published studies tagging small birds has increased substantially.
3. We quantitatively reviewed 549 records extracted from 74 published and 48 unpublished studies on over 7,800 tagged and 17,800 control individuals to examine the effects of geolocator tagging on small bird species (body mass <100 g). We calculated the effect of tagging on apparent survival, condition, phenology and breeding performance and identified the most important predictors of the magnitude of effect sizes.
4. Even though the effects were not statistically significant in phylogenetically controlled models, we found a weak negative impact of geolocators on apparent survival. The negative effect on apparent survival was stronger with increasing relative load of the device and with geolocators attached using elastic harnesses. Moreover, tagging effects were stronger in smaller species.
5. In conclusion, we found a weak effect on apparent survival of tagged birds and managed to pinpoint key aspects and drivers of tagging effects. We provide recommendations for

establishing matched control group for proper effect size assessment in future studies and outline various aspects of tagging that need further investigation. Finally, our results encourage further use of geolocators on small bird species but the ethical aspects and scientific benefits should always be considered.

Keywords: condition, migration, phenology, reproduction, return rate, survival, tracking device, tag effect

Introduction

Tracking devices have brought undisputed insights into the ecology of birds. Use of these tags has enabled researchers to gather valuable information about the timing of life events across annual cycles, the year-round geographic distribution of populations and other important ecological patterns in many species whose movement ecology was widely unknown (e.g. Patchett, Finch, & Cresswell, 2018; Stanley, MacPherson, Fraser, McKinnon, & Stutchbury, 2012; Weimerskirch et al., 2002). A significant proportion of recently published tracking studies use light-level geolocators on small bird species (body mass up to 100 g; Bridge et al., 2013; McKinnon & Love, 2018); however, the increasing use of these tags on small birds raises questions about ethics of tagging and how representative the behaviour of tagged individuals is (Jewell, 2013; Wilson & McMahon, 2006).

Studies using tracking devices such as archival light-level geolocators (hereafter 'geolocators') frequently report the effect of tagging. The published results on the effects of geocator tagging are equivocal: some found reduced apparent survival, breeding success and parental care (Arlt, Low, & Pärt, 2013; Pakanen, Rönkä, Thomson, & Koivula, 2015; Scandolara et al., 2014; Weiser et al., 2016) while others report no obvious effects (Bell, Harouchi, Hewson, & Burgess, 2017; Fairhurst et al., 2015; Peterson et al., 2015; van Wijk, Souchay, Jenni-Eiermann, Bauer, &

Schaub, 2015). Recent meta-analyses evaluating the effects of geolocators (Costantini & Møller, 2013) and other tracking devices (Barron, Brawn, & Weatherhead, 2010; Bodey et al., 2018) showed slightly negative effects on apparent survival, breeding success and parental care. These studies also discussed relative load as an aspect affecting the tagged birds (Costantini & Møller, 2013), or suggested multiple threshold values of relative load on birds (Barron et al., 2010; Bodey et al., 2018). However, these studies involved mainly large bird species where the same additional relative load will more negatively affect surplus power and thus the flight performance than in smaller species (Caccamise & Hedin, 1985). Moreover, previous studies did not control for the effect of small-sample studies, or phylogenetic non-independence and its uncertainty. There is thus a lack of systematic and complex evaluation of geocator effects on small birds including species' life-history and ecological traits, geocator design, and type of attachment.

Almost all prior meta-analyses reporting effects of tagging relied only on published sources and could thus be affected by publication bias (Koricheva, Gurevitch, & Mengersen, 2013), as omitting unpublished sources in meta-analyses may obscure the result (see e.g. Sánchez-Tójar et al. 2018). The main source of publication bias in movement ecology could be a lower probability of publishing studies based on a small sample size, including studies where no or only few tagged birds were successfully recovered due to a strong tagging effect. Additionally, geocator effects most frequently rely on comparisons between tagged and control birds and a biased choice of control individuals may directly lead to the misestimation of the tagging effect sizes. The bias in the control groups can be due to selection of smaller birds, birds being caught in different spatio-temporal conditions, including non-territorial individuals, or different effort put into recapturing control and tagged individuals.

The number of studies tagging small birds is rapidly increasing each year even though our understanding of tag effects is incomplete. In this study, we evaluated the effects of tagging on apparent survival, condition, phenology, and breeding performance for small bird species (<100 g) in

a robust dataset of both published and unpublished studies to minimize the impact of publication bias. Moreover, we assess whether the tagging effects are related to species' ecological and life-history traits, type of control treatment as well as geolocator and attachment designs. We build on the most recent advances in meta-analytical statistical modelling to get unbiased estimates of the geolocator deployment effects controlled for phylogenetic non-independence and its uncertainty (Doncaster & Spake, 2017; Guillerme & Healy 2017; Hadfield, 2010; Viechtbauer, 2010).

Predictions

- i) Geolocators will negatively affect apparent survival, condition, phenology and breeding performance of small birds.
- ii) Negative effects will be stronger in unpublished studies than in published studies.
- iii) Deleterious effects will be most prominent in studies establishing matched control groups compared to studies with potentially-biased control groups.
- iv) Geolocators which constitute a higher relative load will imply stronger negative effects.
- v) Geolocators with a longer light stalk/pipe will cause stronger negative effects because of increased drag in flight and thus increased energetic expenditure (Bowlin et al., 2010; Pennycuik, Fast, Ballerstädt, & Rattenborg, 2012). These effects will be stronger in aerial foragers than in other foraging guilds (Costantini & Møller, 2013).
- vi) Non-elastic harnesses will cause stronger negative effects than elastic harnesses, which better adjust to intra-annual body mass changes and avoid flight restriction (Blackburn et al., 2016).

Material and Methods

Data search

We conducted a comprehensive search for both published and unpublished studies deploying geolocators on bird species with body mass up to 100 g. We searched the Web of Science Core Collection (search terms: TS = (geoloc* AND (bird* OR avian OR migra*) OR geologg*)) and Scopus databases (search terms: TITLE-ABS-KEY (geoloc* AND (bird* OR migra*) OR geologg*)), to find published studies listed to 18 February 2018. Moreover, we searched reference lists of studies using geolocators on small birds and included studies from previous comparative studies (Bridge et al., 2013; Costantini & Møller, 2013; Weiser et al., 2016). In order to obtain information from unpublished studies, we inquired geocator producers and the Migrant Landbird Study Group to disseminate our request for unpublished study details among their customers and members, respectively. In addition, we asked the corresponding authors of the published studies to share any unpublished data. The major geocator producers – Biotrack, Lotek, Migrate Technology and the Swiss Ornithological Institute – sent our request to their customers. To find whether the originally unpublished studies were published over the course of this study, we inspected their status on 1 December 2018. The entire process of search and selection of studies and records (described below) is presented in a flow-chart (Fig. S1).

Inclusion criteria; additional data requesting

We included studies that met the following criteria:

1. The study reported response variables (e.g. return rates, body masses) necessary for effect size calculation.
2. The study included a control group of birds alongside the geocator-tagged individuals or reported a pairwise comparison of tagged birds during geocator deployment and recovery.

3. As a control group, the study considered birds marked on the same site, of the same sex and age class without any indication of a difference in recapture effort between tagged and control groups.
4. For pairwise comparisons, the study presented correlation coefficients or raw data.
5. The variable of interest was presented outside the interaction with another variable.

In order to obtain robust and unbiased results, we asked the corresponding authors for missing data or clarification when the criteria were not met or when it was not clear whether the study complied with the criteria (70% response rate [n = 115]). In addition, we excluded birds that had lost geolocators before subsequent recapture as we did not know when the bird lost the geolocator, and excluded all individuals tagged repeatedly over years because of possible inter-annual carry-over effects of the devices. VBr assessed all studies for eligibility and extracted data, the final dataset was cross-checked by JK and PP. A list of all published studies included in the meta-analysis is provided in the Published Data Sources section.

Trait categories; effect size calculation; explanatory variables

We divided all collected data into four trait categories: apparent survival, condition, phenology and breeding performance based on the response variables reported (e.g. inter-annual recapture rates, body mass changes, arrival dates, or clutch sizes; Table S2). These categories represent the main traits possibly affected in the geolocator-tagged individuals. Subsequently, analyses were run separately for each trait category. We calculated the effect sizes for groups of tagged birds from the same study site and year of attachment, of the same sex (if applicable) and specific geolocator and attachment type accompanied with the corresponding control groups. For simplicity, we call these units *records* throughout the text. For each record, we extracted a contingency table with the treatment arm continuity correction (Schwarzer, Carpenter, & Rucker, 2014) or mean, variance, and sample size, to calculate the unbiased standardised mean difference – Hedges' *g* (Borenstein,

Hedges, Higgins, & Rothstein, 2009) – and its variance with correction for the effect of small sample sizes (Doncaster & Spake, 2018). We used the equation from Sweeting et al. (2004) to calculate variance in pairwise comparisons. When raw data were not provided, we used the reported test statistics (F , t or χ^2) and sample sizes to calculate the effect size using the R package `compute.es` (Del Re 2013). Besides the effect size measures, we extracted additional variables of potential interest – ecological and life-history traits per species, methodological aspects of the study, geolocator and attachment designs and harness material elasticity (Table 1).

Accounting for dependency

We accounted for data non-independence on several levels. When multiple records shared one control group (e.g. several geolocator types and attachment designs used in one year), we split the sample size in the shared control group by the number of records to avoid a false increase in record precision. When multiple measures were available for the same individuals, we randomly chose one effect size measure in each trait category ($n = 8$). If the study provided both recapture and re-encounter rates, we chose the re-encounter rate as a more objective measure of apparent survival. Re-encounters included captures and observations of tagged birds and thus the bias towards the tagged birds caused by the potentially higher recapture effort to retrieve the geolocators should be lower. Finally, we accounted for phylogenetic non-independence between the species and the uncertainty of these relationships using 100 phylogenetic trees (Jetz, Thomas, Joy, Hartmann, & Mooers, 2012) downloaded from the BirdTree.org (www.birdtree.org) using the backbone of Hackett et al. (2008). Moreover, we used the random intercepts of species and study sites in all models, the latter to account for possible site-specific differences (such as different netting effort or other field methods used by particular research teams).

Overall effect sizes and heterogeneity

We calculated the overall effect size for each trait category from all available records using meta-analytical null models. We employed the *MCMCglmm* function from the *MCMCglmm* package (Hadfield, 2010) to estimate overall effect sizes not controlled for phylogeny (model 1, Table S3). We then used the *mulTree* function from the *mulTree* package (Guillerme & Healy, 2017) to automatically fit a *MCMCglmm* model on each phylogenetic tree and summarized the results from all these models to obtain phylogenetically controlled overall effect size estimates (model 2, Table S3). We used weakly informative inverse-Gamma priors ($V = 1$, $\nu = 0.002$) in all models. All fitted *MCMCglmm* models converged and Gelman-Rubin statistic was always <1.1 for all parameters. As our data contained many effect sizes based on small sample sizes, which could lead to a biased estimate of the overall effect size variance, all effect sizes were weighted by their mean-adjusted sampling variance (Doncaster & Spake, 2018). We considered effect sizes (Hedge's g) of 0.2, 0.5 and 0.8 weak, moderate and large effects, respectively. Moreover, we calculated the amount of between-study heterogeneity in all null models using the equation described in Nakagawa and Santos (2012). Phylogenetic heritability (H^2) expressing the phylogenetic signal was estimated as the ratio of phylogenetic variance ($\sigma^2_{phylogeny}$) against the sum of phylogenetic and species variance ($\sigma^2_{species}$) from the models (Table S3; Hadfield & Nakagawa, 2010):

$$H^2 = \sigma^2_{phylogeny} / (\sigma^2_{phylogeny} + \sigma^2_{species})$$

Multivariate meta-analysis

To unveil the most important dependencies of the geolocator effects, we calculated three types of multivariate models: a full trait model (model 3), an ecological model (model 4) and models of publication bias (model 5, Table S3). In the full trait model, we used methodological, species, geolocator specification and attachment variables (Table 1) to estimate their impact on apparent

Accepted Article

survival (model 3). We did not compare the tagging effects of different attachment types due to their use in specific groups of species (e.g. the leg-flagged attachment in shorebirds or the full-body harnesses in nightjars and swifts only). Prior to fitting the ecological model, we employed a principal component analysis of the inter-correlated log continuous life-history traits and extracted the two most important ordination axes – PC1 and PC2 (Table 1). The PC1 explained 54.4% of the variability and expressed a gradient of species characterised mainly by increasing body mass, egg mass and clutch mass (Fig. S4). The PC2 explained 18.7% of variance and was characterised mainly by increasing clutch sizes, number of broods and decreasing migration distances (Fig. S4). These axes together with the categorical ecological traits (Table 1) were then entered into the ecological model to estimate their effect on apparent survival (model 4). Finally, we tested for differences in effect sizes between published and unpublished results in each trait category using all available records (model 5). In these models, we employed the *rma.mv* function from the R package *metafor* (Viechtbauer, 2010) weighted by the mean-adjusted sampling error (Doncaster & Spake, 2018). Continuous predictors were scaled and centred. None of the model residuals violated the assumptions of normal distribution. Because the phylogenetic relatedness of the species explained only a small amount of variation and the phylogenetic relatedness correlates with the life-history and ecological traits, we did not control for phylogeny in the multivariate models but incorporated the random intercepts of species and study site. We calculated R^2 for the full trait and ecological models using the residual between-study variability (τ^2_{residual}) and the total between-study variability (τ^2_{total}) according to the equation (López-López, Marín-Martínez, Sánchez-Meca, Van den Noortgate, & Viechtbauer, 2014):

$$R^2 = (1 - \tau^2_{\text{residual}} / \tau^2_{\text{total}}) \times 100$$

Publication bias; body mass manipulation

We used funnel plots to visually check for potential asymmetry caused by publication bias in each trait category (Fig. S5). To quantify the level of asymmetry in each trait category, we applied the Egger's regression tests of the meta-analytical residuals from all null models of the trait categories (calculated using the *rma.mv* function) against effect size precision ($1 / \text{mean-adjusted standard error}$; Nakagawa & Santos, 2012). An intercept significantly differing from zero suggests the presence of publication bias. In order to find differences in log body mass between the tagged and control individuals during the tagging and marking, we applied a linear mixed-effect model with species and study site as a random intercept weighted by the sample sizes. We considered all effect sizes significant when the 95% credible interval (CrI; using *MCMCglmm* function) or confidence interval (CI; using *rma.mv* function) did not overlap zero. All analyses were conducted in R version 3.3.1 (R Core Team, 2016).

Results

We assessed 854 records for eligibility of effect size calculation and excluded 36% of these records mainly due to a missing control group (59% of ineligible records) or missing essential values for effect size calculation (21%; Fig. S1). Finally, a total of 122 studies containing 549 effect sizes were included in our meta-analysis wherein 35% effect sizes originated from unpublished sources (Table 2). The vast majority of the analysed effect sizes originated from Europe or North America (94%; Fig. S6) and the data contained information about 7,829 tagged and 17,834 control individuals of 69 species from 27 families and 7 orders (Table S7).

We found a weak overall negative effect (Hedges' g : -0.2 ; 95% CrI -0.29 , -0.11 ; $P < 0.001$) only on apparent survival in the model not controlled for phylogeny (model 1). Although we found no statistically significant overall tagging effects in any trait category when controlling for

phylogenetic relatedness, the estimates were similar to those not controlled for phylogeny (model 2, Fig. 1). The phylogenetic signal ($H^2 = 59\%$) was statistically significant only for apparent survival, suggesting that closely related species have more similar response to tagging than less related species, but the variances explained by phylogeny and species were very low for all models (Table S8).

The full trait model of apparent survival revealed that tagging effects were stronger with increasing load on tagged individuals and that geolocators with elastic harnesses affected birds more negatively than geolocators with non-elastic harnesses (Table 3, Fig. 2). However, we found no statistically significant effect on apparent survival for control group type, sex, stalk length, foraging strategy or the interaction between stalk length and foraging strategy (model 3, Table 3). The ecological model suggested a relationship of apparent survival with the PC1, with negative effects being stronger with decreasing body, egg and clutch mass (model 4, Table 3). The full trait model explained 21.1% and the ecological model 11.8% of the between-study variance.

We did not find any evidence for publication bias in any of the trait categories, either visually in the funnel plots (Fig. S5), or using Egger's regression tests (Table 2). Moreover, there were no statistically significant differences in tagging effects between published and unpublished studies (model 5, Table S9). The geolocator-tagged birds were on average 3.8% heavier than control individuals prior to the geolocator deployment and marking (LMM: estimate 0.008 ± 0.003 , $t = 2.47$, $P = 0.014$).

Discussion

Geolocator deployment has a potential to reduce a bird's apparent survival, condition, breeding performance, or may delay events of the annual cycle leading to biases in movement data. By conducting a quantitative review of published studies deploying geolocators on small bird species

and incorporating unpublished data, we revealed only a weak overall effect of geolocators on apparent survival of tagged birds while we found no clear overall effect on condition, phenology and breeding performance. Moreover, we found no statistically significant effects of tagging in any of trait categories when accounting for phylogenetic relationships. Tagging effects on apparent survival were stronger with a higher relative load, when the geolocators were attached with elastic harnesses and in small-bodied species.

Overall tag effects

A negative overall effect of geocator tagging on apparent survival found in this study seems to be prevalent across previous comparative studies of tagging effects (Barron et al., 2010; Bodey et al., 2018; Costantini & Møller, 2013; Trefry, Diamond, & Jesson, 2012; Weiser et al., 2016). However, unlike previous comparative (Barron et al., 2010; Bodey et al., 2018) and primary studies (e.g. Adams et al., 2009; Arlt et al., 2013; Snijders et al., 2017), we found no overall negative effects of tagging on variables associated with breeding performance in our analysis. We also did not find evidence for overall effects of tagging on body condition and phenology, which was consistent with equivocal results of previous studies: some found reduced body condition (Adams et al. 2009, Elliott et al., 2012) or delayed timing of annual cycle events (Arlt et al., 2013, Scandolara et al., 2014), while others found no evidence for tagging effects on these traits (Bell et al., 2017; Fairhurst et al., 2015; Peterson et al., 2015; van Wijk et al., 2015).

Tagged individuals that returned to the study site are potentially in better condition than the tagged individuals that did not return – this potentially contributes to the weak tagging effects on condition, phenology and breeding performance. However, the lack of effect we found on phenology and breeding performance could also be an artefact of the small sample sizes, as collecting these data is probably more challenging in small avian species, which are more difficult to re-sight and recapture and have shorter life-spans than the relatively heavier species included in the previous

Accepted Article

studies. Similarly, effects of tagging on condition could be underestimated in our analysis due to the initial differences we found between the body mass of tagged and control birds. Additionally, the intra-annual body mass changes could be biased in studies where timing of geolocator deployment and geolocator recovery differs. Unfortunately, the timing of captures and recaptures was rarely reported and could not be analysed in our study. Overall, the weak effects of tagging we found support several primary studies (e.g. Bell et al., 2017; Fairhurst et al., 2015; Peterson et al., 2015; van Wijk et al., 2015), indicating that geolocator tagging is both ethical and provides credible information on bird movements. On the other hand, care should be taken as the tagging effect may be specific to populations or species. For example, Weiser et al. (2016) found a negligible overall effect but significant reduction of return rates in the smallest species in their meta-analysis. The negative effect of geolocators can also vary between years (Bell et al., 2017, Scandolara et al., 2014), or be induced by occasional bad weather conditions (Snijders et al., 2017), or food shortages (Saraux et al., 2011; Wilson et al., 2015).

Inferring unbiased overall effect sizes

We minimised publication bias in our estimates of overall effects by including substantial amount of unpublished results (192 records of 38 species) and contacting authors of published studies for additional data. Still, some of these studies might get published in the future despite the delay between our data collation and the final analysis. We did not find any evidence that tagging effects differed between published and unpublished studies, suggesting that the tagging effect may not be a critical consideration for publishing a study.

Moreover, we found no support for stronger tag effects in studies with matched control individuals compared to studies with less strict control treatments. However, this result is potentially confounded by the fact that tagged birds were on average larger and in potentially better condition than control birds, which would underestimate the negative effects of tagging. We thus suggest

establishing carefully matched control groups in all future studies to enable a more reliable estimation of tagging effects. Such a control group should include: i) randomly selected individuals of the same species, sex and age class; ii) individuals caught at the same time of the season and year; iii) at the same time of the day; iv) of similar size and condition as tagged individuals, and v) exclude non-territorial birds or individuals passing through the site.

Influence of relative load and species' life-histories

Our results support the current evidence (Bodey et al., 2018; Weiser et al., 2016) for reduced apparent survival in studies with a relatively higher tag load on treated individuals. Moreover, we found an increasing negative effect in studies tagging smaller species with smaller eggs and clutch masses. The lower body mass in these species is likely accompanied with a higher relative tag load due to technical constraints of lower tag weights. Although recent miniaturisation has led to the development of smaller tags, these tags have been predominantly applied to smaller species instead of reducing tag load in larger species (Portugal & White, 2018). The various relative loads used without observed tagging effects (e.g. Bell et al., 2017, Peterson et al., 2015; van Wijk et al., 2015) indicate the absence of a generally applicable rule for all small bird species (Schacter & Jones, 2017) and we thus recommend the use of reasonably small tags despite potential disadvantages (e.g. reduced battery lifespan or light sensor quality).

Harness material

Contrary to our prediction, we found higher apparent survival in birds tagged with harnesses made of non-elastic materials. Non-elastic harnesses are usually individually adjusted on each individual, whereas elastic harnesses are often prepared before attachment to fit the expected body size of the tagged individuals according to allometric equations (e.g. Naef-Daenzer, 2007). As pre-sized elastic

harnesses cannot match perfectly the size of every captured individual, they may be in the end more frequently tightly fitted as some researches might tend to tag larger individuals or avoid too loose harnesses to prevent geolocator loss. Non-elastic harnesses may also be more frequently looser than elastic harnesses as researchers try to reduce the possibility of non-elastic harness getting tight when birds accumulate fat. Tight harnesses significantly reduced the return rates in whinchat (*Saxicola rubetra*; Blackburn et al. 2016), and it may be difficult to register whether elastic harnesses are restricting physical movement of birds when deploying tags. In contrast, non-elastic harnesses, which are more commonly tailored according to the actual size, are often made sufficiently loose to account for body mass changes of each individual. Prepared elastic harnesses are usually used to reduce the handling time during the geolocator deployment (Streby et al. 2015) but this advantage may be outweighed by the reduced apparent survival of geolocators with tied elastic harnesses. We thus suggest to consider stress during geolocator deployment together with the potentially reduced apparent survival and the risk of tag loss when choosing harness material.

Variables without statistically significant impact on tagging effect

Migratory distance did not affect the magnitude of the effect sizes, contrasting with some previous findings (Bodey et al., 2018; Costantini & Møller, 2013). However, none of these studies used population-specific distances travelled; instead, they used latitudinal spans between ranges of occurrence (Costantini & Møller, 2013) or travelled distance categorised into three distances groups (Bodey et al., 2018). These types of distance measurements could greatly affect the results especially in species that migrate mainly in an east-west direction (Lislevand et al., 2015; Stach, Kullberg, Jakobsson, Ström, & Fransson, 2016) or in species whose populations largely differ in their travel distances (Bairlein et al., 2012; Schmaljohann, Buchmann, Fox, & Bairlein, 2012). Moreover, light-level geolocators were most frequently deployed to the long-distance migrants in our study and the result can be thus applicable to these species only.

Additionally, we found no overall effect of species' foraging strategy, contrary to the strong overall negative effect found for aerial foraging species (Costantini and Møller 2013). Despite the tag shape altering the drag and thus energy expenditure during flight (Bowlin et al., 2010; Pennycuik et al., 2012), apparent survival tended to be better in individuals fitted with stalked geolocators and we found no interaction between stalk length and foraging strategy on the tagging effect size.

Geolocators with longer stalks have been more frequently used in heavier birds with low relative load where the expected tag effect is weak. Moreover, previous results of strong negative effects in aerial foragers led to a preferential use of stalkless geolocators in these species and probably minimised the tagging effect in this foraging guild (Morganti et al., 2018; Scandolara et al., 2015).

However, the evidence for the negative effects in non-aerial foragers is low as there is only one field study focusing on stalk length effects on the return rates (Blackburn et al., 2016).

Future considerations

Future studies evaluating the use of geolocators on birds should focus on assessing inter-annual differences in tagging effects, effects of varying relative loads, different stalk lengths or different attachment methods to minimise the negative effects of tagging. We also suggest to focus on the impact of various movement strategies such as fattening and moulting schedules on the tagging effect. All future studies should carefully set matched controls and transparently report on tagging effects. Finally, our results encourage use of geolocators on small bird species but the ethical and scientific benefits should always be considered.

Authors' contributions

VBr, JK and PP conceived the idea and designed the methodology. VBr reviewed the literature and collected data, JK and PP checked the data extracted for analysis. VBr and PP analysed the data. VBr led the writing of the manuscript with significant contributions from JK and PP. MB, SH, DH, MK, JO and EW contributed with unpublished data and their comments and suggestions significantly improved the manuscript. PA, JA, DA, SB, DB, EB, VBe, CB, SB, MBr, BC, DC, NC, JC, VC, TE, KF, OG, MG, MH, CH, FJ, JJ, TK, DK, ML, TL, SL, CL, KM, PMar, SM, PMat, CM, BM, JM, RNe, AN, RNo, TP, VP, NP, MP, JR, CR, AR, CS, NS, MT, DT, HO, AW, HW, JW, KW and BW contributed unpublished data and critically revised the manuscript. All authors gave final approval for publication.

Acknowledgements

We thank James W. Fox (Migrate Technology), the Swiss Ornithological Institute, Biotrack/Lotek employees for circulating the call for sharing the unpublished study results among their customers and Rien van Wijk for sharing our inquiry for unpublished data among the Migrant Landbird Study Group members. We are grateful to Carlos Camacho, Vladimir G. Grinkov, Helene M. Lampe, Ken Otter, Jaime Potti, Milica Požgayová, Scott M. Ramsay and Helmut Sternberg for providing unpublished data and to Marie Hánová for extracting part of the species-specific life-history data.

We thank Martin Sládeček, anonymous reviewers and editors for valuable comments on the earlier version of the manuscript and Adéla Stupková for the graphics. The fieldwork in Greenland and Russia (Yamal Peninsula) was supported by the RFBR through grant Arctic-18-05-60261, Yamal-LNG company (Sabetta) and the French Polar Institute (IPEV, program 1036 "Interactions"). DK was supported by the Russian Science Foundation grant (project no. 17-14-01147) and by a Leverhulme Trust research grant to Richard Holland (RPG-2013288). The study was funded by the Czech Science Foundation (project no. 13-06451S) and by the Institutional Research Plan (RVO: 68081766). We are grateful to the funders, supporters and researchers of the many studies included herein. Any use of

trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Data accessibility

Data described in this article are available at <https://doi.org/10.5281/zenodo.1886530> (Brlík et al., 2018).

References

- Adams, J., Scott, D., McKechnie, S., Blackwell, G., Shaffer, S. A., & Moller, H. (2009). Effects of geolocation archival tags on reproduction and adult body mass of sooty shearwaters (*Puffinus griseus*). *New Zealand Journal of Zoology*, *36*, 355–366.
<https://doi.org/10.1080/03014220909510160>
- Arlt, D., Low, M., & Pärt, T. (2013). Effect of geolocators on migration and subsequent breeding performance of a long-distance passerine migrant. *PLoS ONE*, *8*, e82316.
<https://doi.org/10.1371/journal.pone.0082316>
- Bairlein, F., Norris, D. R., Nagel, R., Bulte, M., Voigt, C. C., Fox, J., ... Schmalljohann, H. (2012). Cross-hemisphere migration of a 25 g songbird. *Biology Letters*, *8*, 505–507.
<https://doi.org/10.1098/rsbl.2011.1223>
- Barron, D. G., Brawn, J. D., & Weatherhead, P. J. (2010). Meta-analysis of transmitter effects on avian behaviour and ecology. *Methods in Ecology and Evolution*, *1*, 180–187.
<https://doi.org/10.1111/j.2041-210X.2010.00013.x>
- Bell, S. C., Harouchi, M. E. L., Hewson, C. M., & Burgess, M. D. (2017). No short- or long-term effects of geolocator attachment detected in Pied Flycatchers *Ficedula hypoleuca*. *Ibis*, *159*, 734–743.

<https://doi.org/10.1111/ibi.12493>

Blackburn, E., Burgess, M., Freeman, B., Risely, A., Izang, A., Ivande, S., ... Cresswell, W. (2016). An experimental evaluation of the effects of geolocator design and attachment method on between-year survival on Whinchats *Saxicola rubetra*. *Journal of Avian Biology*, *47*, 530–539. <https://doi.org/10.1111/jav.00871>

Bodey, T. W., Cleasby, I. R., Bell, F., Parr, N., Schultz, A., Votier, S. C., & Bearhop, S. (2018). A phylogenetically controlled meta-analysis of biologging device effects on birds: Deleterious effects and a call for more standardized reporting of study data. *Methods in Ecology and Evolution*, *9*, 946–955. <https://doi.org/10.1111/2041-210X.12934>

Bodey, T. W., Cleasby, I. R., Bell, F., Parr, N., Schultz, A., Votier, S. C., & Bearhop, S. (2018). A phylogenetically controlled meta-analysis of biologging device effects on birds: Deleterious effects and a call for more standardized reporting of study data. *Dryad Digital Depository*, <https://doi.org/10.5061/dryad.0rp52>

Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to meta-analysis*. John Wiley & Sons. <https://doi.org/10.1002/9780470743386>

Bowlin, M. S., Henningson, P., Muijres, F. T., Vleugels, R. H. E., Liechti, F., & Hedenström, A. (2010). The effects of geolocator drag and weight on the flight ranges of small migrants. *Methods in Ecology and Evolution*, *1*, 398–402. <https://doi.org/10.1111/j.2041-210X.2010.00043.x>

Bridge, E. S., Kelly, J. F., Contina, A., Gabrielson, R. M., MacCurdy, R. B., & Winkler, D. W. (2013). Advances in tracking small migratory birds: A technical review of light-level geolocation. *Journal of Field Ornithology*, *84*, 121–137. <https://doi.org/10.1111/jfo.12011>

Brlík, V., Koleček, J., Burgess, M. D., Hahn, S., Humple, D., Krist, M., ... Procházka, P. (2018). Weak effect of geolocators on small birds: a meta-analysis controlled for phylogeny and potential publication bias [Dataset]. *Zenodo*, <https://doi.org/10.5281/zenodo.1886530>

Caccamise, D. F., & Hedin, R. S. (1985). An aerodynamic basis for selecting transmitter loads in birds.

The Wilson Bulletin, 97, 306–318.

Costantini, D., & Møller, A. P. (2013). A meta-analysis of the effects of geolocator application on

birds. *Current Zoology*, 59, 697–706. <https://doi.org/10.1093/czoolo/59.6.697>

Cramp, S., Perrins, C. M., (1977–1994). The birds of the Western Palearctic. Volumes 1–9. Oxford,

UK: Oxford University Press.

Del Re, A. C. (2013). compute.es: Compute effect sizes. R package version 0.2-2. URL: [https://cran.r-](https://cran.r-project.org/web/packages/compute.es/index.html)

[project.org/web/packages/compute.es/index.html](https://cran.r-project.org/web/packages/compute.es/index.html)

Doncaster, C. P., & Spake, R. (2018). Correction for bias in meta-analysis of little-replicated studies.

Methods in Ecology and Evolution, 9, 634–644. <https://doi.org/10.1111/2041-210X.12927>

Elliott, K. H., McFarlane, L., Burke, C. M., Hedd, A., Montevecchi, W. A., & Anderson, W. G. (2012).

Year-long deployments of small geolocators increase corticosterone levels in murres. *Marine*

Ecology Progress Series, 466, 1–7. <https://doi.org/10.3354/meps09975>

Fairhurst, G. D., Berzins, L. L., David, W., Laughlin, A. J., Romano, A., Romano, M., ... Clark, R. G.

(2015). Assessing costs of carrying geolocators using feather corticosterone in two species of

aerial insectivore. *Royal Society Open Science*, 2, 150004. <https://doi.org/10.1098/rsos.150004>

Guillerme, T., & Healy, K. (2017). mulTree: Performs MCMCgImm on multiple phylogenetic trees. R

package version 1.3.1. <https://github.com/TGuillerme/mulTree>

Hackett, S., Kimball, R., Reddy, S., Bowie, R., Braun, E., Braun, M., ... Yuri, T. (2008). A phylogenomic

study of birds reveals their evolutionary history. *Science*, 320, 1763–1768.

<https://doi.org/10.1126/science.1157704>

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: The

MCMCgImm R package. *Journal of Statistical Software*, 33, 1–22.

- Hadfield, J. D., & Nakagawa, S. (2010). General quantitative genetic methods for comparative biology: Phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of Evolutionary Biology*, 23, 494–508. <https://doi.org/10.1111/j.1420-9101.2009.01915.x>
- Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*, 491, 444–448. <https://doi.org/10.1038/nature11631>
- Jewell, Z. (2013). Effect of monitoring technique on quality of conservation science. *Conservation Biology*, 27(3), 501–508. <https://doi.org/10.1111/cobi.12066>
- Koricheva, J., Gurevitch, J., & Mengersen, K. (2013). *Handbook of meta-analysis in ecology and evolution*. Princeton University Press.
- Lislevand, T., Chutný, B., Byrkjedal, I., Pavel, V., Briedis, M., Adamík, P., & Hahn, S. (2015). Red-spotted Bluethroats *Luscinia s. svecica* migrate along the Indo-European flyway: a geolocator study. *Bird Study*, 62, 508–515. <https://doi.org/10.1080/00063657.2015.1077781>
- López-López, J. A., Marín-Martínez, F., Sánchez-Meca, J., Van den Noortgate, W., & Viechtbauer, W. (2014). Estimation of the predictive power of the model in mixed-effects meta-regression: A simulation study. *British Journal of Mathematical and Statistical Psychology*, 67, 30–48. <https://doi.org/10.1111/bmsp.12002>
- McKinnon, E. A., & Love, O. P. (2018). Ten years tracking the migrations of small landbirds: Lessons learned in the golden age of bio-logging. *The Auk*, 135, 834–856. <https://doi.org/10.1642/AUK-17-202.1>
- Morganti, M., Rubolini, D., Åkesson, S., Bermejo, A., de la Puente, J., Lardelli, R., ... Ambrosini, R. (2018). Effect of light-level geolocators on apparent survival of two highly aerial swift species. *Journal of Avian Biology*, 49, jav-01521. <https://doi.org/10.1111/jav.01521>
- Naef-Daenzer, B. (2007). An allometric function to fit leg-loop harnesses to terrestrial birds. *Journal*

of Avian Biology, 38, 404–407. <https://doi.org/10.1111/j.2007.0908-8857.03863.x>

Nakagawa, S., & Santos, E. S. A. (2012). Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology*, 26, 1253–1274. <https://doi.org/10.1007/s10682-012-9555-5>

Pakanen, V. M., Rönkä, N., Thomson, R. L., & Koivula, K. (2015). No strong effects of leg-flagged geolocators on return rates or reproduction of a small long-distance migratory shorebird. *Ornis Fennica*, 92, 101–111.

Patchett, R., Finch, T., & Cresswell, W. (2018). Population consequences of migratory variability differ between flyways. *Current Biology*, 28, R340–R341. <https://doi.org/10.1016/j.cub.2018.03.018>

Pennycuik, C. J., Fast, P. L. F., Ballerstädt, N., & Rattenborg, N. (2012). The effect of an external transmitter on the drag coefficient of a bird's body, and hence on migration range, and energy reserves after migration. *Journal of Ornithology*, 153, 633–644. <https://doi.org/10.1007/s10336-011-0781-3>

Peterson, S. M., Streby, H. M., Kramer, G. R., Lehman, J. a., Buehler, D. a., & Andersen, D. E. (2015). Geolocators on Golden-winged Warblers do not affect migratory ecology. *The Condor*, 117, 256–261. <https://doi.org/10.1650/CONDOR-14-200.1>

Portugal, S. J., & White, C. R. (2018). Miniaturisation of biologgers is not alleviating the 5% rule. *Methods in Ecology and Evolution*, 9, 1662–1666. <https://doi.org/10.1111/2041-210X.13013>

R Core Team 2018. R: a language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. URL: <https://www.R-project.org/>

Rodewald, P. (2015). The birds of North America. Cornell Laboratory of Ornithology, Ithaca, NY. URL: <https://birdsna.org>

Sánchez-Tójar, A., Nakagawa, S., Sánchez-Fortún, M., Martín, D. A., Ramani, S., Girndt, A., ...

Schroeder, J. (2018). Meta-analysis challenges a textbook example of status signalling and demonstrates publication bias. *eLife*, 7, e37385. <https://doi.org/10.7554/eLife.37385>

Saraux, C., Le Bohec, C., Durant, J. M., Viblanc, V. A., Gauthier-Clerc, M., Beaune, D., ... Le Maho, Y. (2011). Reliability of flipper-banded penguins as indicators of climate change. *Nature*, 469, 203–206. <https://doi.org/10.1038/nature09630>

Scandolaro, C., Rubolini, D., Ambrosini, R., Caprioli, M., Hahn, S., Liechti, F., ... Saino, N. (2014). Impact of miniaturized geolocators on barn swallow *Hirundo rustica* fitness traits. *Journal of Avian Biology*, 45, 417–423. <https://doi.org/10.1111/jav.00412>

Schacter, C. R., & Jones, I. L. (2017). Effects of geolocation tracking devices on behavior, reproductive success, and return rate of *Aethia* auklets: An evaluation of tag mass guidelines. *The Wilson Journal of Ornithology*, 129, 459–468. <https://doi.org/10.1676/16-084.1>

Schmaljohann, H., Buchmann, M., Fox, J. W., & Bairlein, F. (2012). Tracking migration routes and the annual cycle of a trans-Saharan songbird migrant. *Behavioral Ecology and Sociobiology*, 66, 915–922. <https://doi.org/10.1007/s00265-012-1340-5>

Schönwetter, M. (1960–1992). *Handbuch der Oologie*. Akademie Verlag, Berlin.

Schwarzer, G., Carpenter, J. R., & Rücker, G. (2014). *Meta-analysis with R*. Springer. <https://doi.org/10.1007/978-3-319-21416-0>

Snijders, L., Nieuwe Weme, L. E., De Goede, P., Savage, J. L., Van Oers, K., & Naguib, M. (2017). Context-dependent effects of radio transmitter attachment on a small passerine. *Journal of Avian Biology*, 48, 650–659. <https://doi.org/10.1111/jav.01148>

Stach, R., Kullberg, C., Jakobsson, S., Ström, K., & Fransson, T. (2016). Migration routes and timing in a bird wintering in South Asia, the Common Rosefinch *Carpodacus erythrinus*. *Journal of Ornithology*, 157, 756–767. <https://doi.org/10.1007/s10336-016-1329-3>

Stanley, C. Q., MacPherson, M., Fraser, K. C., McKinnon, E. A., & Stutchbury, B. J. M. (2012). Repeat tracking of individual songbirds reveals consistent migration timing but flexibility in route. *PLoS ONE*, 7, e40688. <https://doi.org/10.1371/journal.pone.0040688>

Streby, H. M., McAllister, T. L., Peterson, S. M., Kramer, G. R., Lehman, J. a., & Andersen, D. E. (2015). Minimizing marker mass and handling time when attaching radio-transmitters and geolocators to small songbirds. *The Condor*, 117, 249–255. <https://doi.org/10.1650/CONDOR-14-182.1>

Sweeting, M. J., Sutton, A. J., & Lambert, P. C. (2004). What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data. *Statistics in Medicine*, 23, 1351–1375. <https://doi.org/10.1002/sim.1761>

Trefry, S. A., Diamond, A. W., & Jesson, L. K. (2012). Wing marker woes: a case study and meta-analysis of the impacts of wing and patagial tags. *Journal of Ornithology*, 154, 1–11. <https://doi.org/10.1007/s10336-012-0862-y>

van Wijk, R. E., Souchay, G., Jenni-Eiermann, S., Bauer, S., & Schaub, M. (2015). No detectable effects of lightweight geolocators on a Palaearctic-African long-distance migrant. *Journal of Ornithology*, 157, 255–264. <https://doi.org/10.1007/s10336-015-1274-6>

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal Of Statistical Software*, 36, 1–48.

Weimerskirch, H., Bonadonna, F., Bailleul, F., Mabile, G., Dell’Omo, G., & Lipp, H.-P. (2002). GPS tracking of foraging albatrosses. *Science*, 295, 1259. <https://doi.org/10.1126/science.1068034>

Weiser, E. L., Lanctot, R. B., Brown, S. C., Alves, J. A., Battley, P. F., Bentzen, R., ... Sandercock, B. K. (2016). Effects of geolocators on hatching success, return rates, breeding movements, and change in body mass in 16 species of Arctic-breeding shorebirds. *Movement Ecology*, 4, 12. <https://doi.org/10.1186/s40462-016-0077-6>

Wilson, R. P., & McMahon, C. R. (2006). Measuring devices on wild animals: what constitutes

acceptable practice? *Frontiers in Ecology and the Environment*, 4, 147–154.

[https://doi.org/10.1890/1540-9295\(2006\)004\[0147:MDOWAW\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)004[0147:MDOWAW]2.0.CO;2)

Wilson, R. P., Sala, J. E., Gómez-Laich, A., Ciancio, J., & Quintana, F. (2015). Pushed to the limit: Food abundance determines tag-induced harm in penguins. *Animal Welfare*, 24, 37–44.

<https://doi.org/10.7120/09627286.24.1.037>

Published Data Sources

Alonso, D., Arizaga, J., Meier, C. M., & Liechti, F. (2017). Light-level geolocators confirm resident status of a Southern European Common Crossbill population. *Journal of Ornithology*, 158, 75–81. <https://doi.org/10.1007/s10336-016-1388-5>

Arbeiter, S., Schulze, M., Todte, I., & Hahn, S. (2012). Das Zugverhalten und die Ausbreitung von in Sachsen-Anhalt brütenden Bienenfressern (*Merops apiaster*). *Berichte der Vogelwarte Hiddensee*, 21, 33–41.

Arlt, D., Low, M., & Pärt, T. (2013). Effect of geolocators on migration and subsequent breeding performance of a long-distance passerine migrant. *PLoS ONE*, 8, e82316.

<https://doi.org/10.1371/journal.pone.0082316>

Arlt, D., Olsson, P., Fox, J. W., Low, M., & Pärt, T. (2015). Prolonged stopover duration characterises migration strategy and constraints of a long-distance migrant songbird. *Animal Migration*, 2, 47–62. <https://doi.org/10.1515/ami-2015-0002>

<https://doi.org/10.1515/ami-2015-0002>

Bächler, E., Hahn, S., Schaub, M., Arlettaz, R., Jenni, L., Fox, J. W., ... Liechti, F. (2010). Year-round tracking of small trans-Saharan migrants using light-level geolocators. *PLoS ONE*, 5, e9566.

<https://doi.org/10.1371/journal.pone.0009566>

Bairlein, F., Norris, D. R., Nagel, R., Bulte, M., Voigt, C. C., Fox, J., ... Schmalljohann, H. (2012). Cross-hemisphere migration of a 25 g songbird. *Biology Letters*, *8*, 505–507.

<https://doi.org/10.1098/rsbl.2011.1223>

Bell, S. C., Harouchi, M. E. L., Hewson, C. M., & Burgess, M. D. (2017). No short- or long-term effects of geolocator attachment detected in Pied Flycatchers *Ficedula hypoleuca*. *Ibis*, *159*, 734–743.

<https://doi.org/10.1111/ibi.12493>

Blackburn, E., Burgess, M., Freeman, B., Risely, A., Izang, A., Ivande, S., ... Cresswell, W. (2016). An experimental evaluation of the effects of geolocator design and attachment method on between-year survival on Whinchats *Saxicola rubetra*. *Journal of Avian Biology*, *47*, 530–539.

<https://doi.org/10.1111/jav.00871>

Bravo, S. P., Cueto, V. R., & Andre, C. (2017). Migratory timing, rate, routes and wintering areas of White-crested Elaenia (*Elaenia albiceps chilensis*), a key seed disperser for Patagonian forest regeneration. *PLoS ONE*, *12*, e0170188. <https://doi.org/10.1371/journal.pone.0170188>

Briedis, M., Beran, V., Hahn, S., & Adamík, P. (2016). Annual cycle and migration strategies of a habitat specialist, the Tawny Pipit *Anthus campestris*, revealed by geolocators. *Journal of Ornithology*, *157*, 619–626. <https://doi.org/10.1007/s10336-015-1313-3>

Briedis, M., Hahn, S., Gustafsson, L., Henshaw, I., Träff, J., Král, M., & Adamík, P. (2016). Breeding latitude leads to different temporal but not spatial organization of the annual cycle in a long-distance migrant. *Journal of Avian Biology*, *47*, 743–748. <https://doi.org/10.1111/jav.01002>

Briedis, M., Träff, J., Hahn, S., Ilieva, M., Král, M., Peev, S., & Adamík, P. (2016). Year-round spatiotemporal distribution of the enigmatic Semi-collared Flycatcher *Ficedula semitorquata*. *Journal of Ornithology*, *157*, 895–900. <https://doi.org/10.1007/s10336-016-1334-6>

Brlík, V., Ilieva, M., Lisovski, S., Voigt, C. C., & Procházka, P. (2018). First insights into the migration route and migratory connectivity of the Paddyfield Warbler using geolocator tagging and stable

isotope analysis. *Journal of Ornithology*, 159, 879–882. <https://doi.org/10.1007/s10336-018-1557-9>

Callo, P. A., Morton, E. S., & Stutchbury, B. J. M. (2013). Prolonged spring migration in the Red-eyed Vireo (*Vireo olivaceus*). *The Auk*, 130, 240–246. <https://doi.org/10.1525/auk.2013.12213>

Cooper, N. W., Hallworth, M. T., & Marra, P. P. (2017). Light-level geolocation reveals wintering distribution, migration routes, and primary stopover locations of an endangered long-distance migratory songbird. *Journal of Avian Biology*, 48, 209–2019. <https://doi.org/10.1111/jav.01096>

Cormier, R. L., Humple, D. L., Gardali, T., & Seavy, N. E. (2013). Light-level geolocators reveal strong migratory connectivity and within-winter movements for a coastal California Swainson's thrush (*Catharus ustulatus*) population. *The Auk*, 130, 283–290. <https://doi.org/10.1525/auk.2013.12228>

Cormier, R. L., Humple, D. L., Gardali, T., & Seavy, N. E. (2016). Migratory connectivity of Golden-crowned Sparrows from two wintering regions in California. *Animal Migration*, 3, 48–56. <https://doi.org/10.1515/ami-2016-0005>

Cresswell, B., & Edwards, D. (2013). Geolocators reveal wintering areas of European Nightjar (*Caprimulgus europaeus*). *Bird Study*, 60, 77–86. <https://doi.org/10.1080/00063657.2012.748714>

DeLuca, W. V., Woodworth, B. K., Rimmer, C. C., Marra, P. P., Taylor, P. D., McFarland, K. P., ... Norris, D. R. (2015). Transoceanic migration by a 12 g songbird. *Biology Letters*, 11, 20141045. <https://doi.org/10.1098/rsbl.2014.1045>

Evens, R., Convey, G. J., Henderson, I. G., Cresswell, W., Jiguet, F., Moussy, C., ... Artois, T. (2017). Migratory pathways, stopover zones and wintering destinations of Western European Nightjars *Caprimulgus europaeus*. *Ibis*, 159, 680–686. <https://doi.org/10.1111/ijlh.12426>

Fairhurst, G. D., Berzins, L. L., Bradley, D. W., Laughlin, A. J., Romano, A., Romano, M., ... Clark, R. G. (2015). Assessing costs of carrying geolocators using feather corticosterone in two species of aerial insectivore. *Royal Society Open Science*, 2, 150004. <https://doi.org/10.1098/rsos.150004>

Fairhurst, G. D., Berzins, L. L., Bradley, D. W., Laughlin, A. J., Romano, A., Romano, M., ... Clark, R. G. (2015). Data from: Assessing costs of carrying geolocators using feather corticosterone in two species of aerial insectivore. *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.sq184>

Fraser, K. C., Cousens, B., Simmons, M., Nightingale, A., Cormier, L., Humple, D. L., & Shave, A. C. (2018). Classic pattern of leapfrog migration in Sooty Fox Sparrow (*Passerella iliaca unalaschcensis*) is not supported by direct migration tracking of individual birds. *Auk*, 135, 572–582. <https://doi.org/10.1642/AUK-17-224.1>

Fraser, K. C., Stutchbury, B. J. M., Silverio, C., Kramer, P. M., Barrow, J., Newstead, D., ... Tautin, J. (2012). Continent-wide tracking to determine migratory connectivity and tropical habitat associations of a declining aerial insectivore. *Proceedings of the Royal Society B-Biological Sciences*, 279, 4901–4906. <https://doi.org/10.1098/rspb.2012.2207>

Gersten, A., & Hahn, S. (2016). Timing of migration in Common Redstarts (*Phoenicurus phoenicurus*) in relation to the vegetation phenology at residence sites. *Journal of Ornithology*, 157, 1029–1036. <https://doi.org/10.1007/s10336-016-1359-x>

Gómez, J., Michelson, C. I., Bradley, D. W., Ryan Norris, D., Berzins, L. L., Dawson, R. D., & Clark, R. G. (2014). Effects of geolocators on reproductive performance and annual return rates of a migratory songbird. *Journal of Ornithology*, 155, 37–44. <https://doi.org/10.1007/s10336-013-0984-x>

Hallworth, M. T., Sillett, T. S., Van Wilgenburg, S. L., Hobson, K. A., & Marra, P. P. (2015). Migratory connectivity of a neotropical migratory songbird revealed by archival light-level geolocators. *Ecological Applications*, 25, 336–347. <https://doi.org/10.1890/14-0195.1>

Heckscher, C. M., Taylor, S. M., Fox, J. W., & Afanasyev, V. (2011). Veery (*Catharus fuscescens*) wintering locations, migratory connectivity, and a revision of its winter range using geolocator technology. *The Auk*, *128*, 531–542. <https://doi.org/10.1525/auk.2011.10280>

Horns, J., Buechley, E., Chynoweth, M., Aktay, L., Çoban, E., Kırpık, M., ... Şekercioğlu, Ç. H. (2016). Geolocator tracking of great reed warbler (*Acrocephalus arundinaceus*) identifies key regions of importance to migratory wetland specialist throughout the Middle East and Sub-Saharan Africa. *The Condor*, *118*, 835–849. <https://doi.org/10.1650/CONDOR-16-63.1>

Jiménez, J. E., Jahn, A. E., Rozzi, R., & Seavy, N. E. (2016). First documented migration of individual White-Crested Elaenias (*Elaenia albiceps chilensis*) in South America. *The Wilson Journal of Ornithology*, *128*, 419–425. <https://doi.org/10.1163/187529271X00756>

Johnson, J. A., Matsuoka, S. M., Tessler, D. F., Greenberg, R., & Fox, J. W. (2012). Identifying migratory pathways used by Rusty Blackbirds breeding in southcentral Alaska. *The Wilson Journal of Ornithology*, *124*, 698–703. <https://doi.org/10.1676/1559-4491-124.4.698>

Koleček, J., Procházka, P., El-Arabany, N., Tarka, M., Ilieva, M., Hahn, S., ... Hansson, B. (2016). Cross-continental migratory connectivity and spatiotemporal migratory patterns in the great reed warbler. *Journal of Avian Biology*, *47*, 756–767. <https://doi.org/10.1111/jav.00929>

Laughlin, A. J., Taylor, C. M., Bradley, D. W., LeClair, D., Clark, R. G., Dawson, R. D., ... Norris, D. R. (2013). Integrating information from geolocators, weather radar, and citizen science to uncover a key stopover area of an aerial insectivore. *The Auk*, *130*, 230–239. <https://doi.org/10.1525/auk.2013.12229>

Lemke, H. W., Tarka, M., Klaassen, R. H. G., Åkesson, M., Bensch, S., Hasselquist, D., & Hansson, B. (2013). Annual cycle and migration strategies of a trans-Saharan migratory songbird: A geolocator study in the great reed warbler. *PLoS ONE*, *8*, e79209. <https://doi.org/10.1371/journal.pone.0079209>

Liechti, F., Scandola, C., Rubolini, D., Ambrosini, R., Korner-Nievergelt, F., Hahn, S., ... Saino, N.

(2015). Timing of migration and residence areas during the non-breeding period of barn swallows *Hirundo rustica* in relation to sex and population. *Journal of Avian Biology*, *46*, 254–265. <https://doi.org/10.1111/jav.00485>

Liechti, F., Witvliet, W., Weber, R., & Bächler, E. (2013). First evidence of a 200-day non-stop flight in a bird. *Nature Communications*, *4*, 2554. <https://doi.org/10.1038/ncomms3554>

Lislevand, T., Briedis, M., Heggøy, O., & Hahn, S. (2016). Seasonal migration strategies of Common Ringed Plovers *Charadrius hiaticula*. *Ibis*, *159*, 225–229. <https://doi.org/10.1111/ibi.12424>

Lislevand, T., Chutný, B., Byrkjedal, I., Pavel, V., Briedis, M., Adamík, P., & Hahn, S. (2015). Red-spotted Bluethroats *Luscinia s. svecica* migrate along the Indo-European flyway: a geolocator study. *Bird Study*, *62*, 508–515. <https://doi.org/10.1080/00063657.2015.1077781>

Lislevand, T., & Hahn, S. (2013). Effects of geolocator deployment by using flexible leg-loop harnesses in a small wader. *Wader Study Group Bulletin*, *120*, 108–113.

Macdonald, C. A., Mckinnon, E. A., Gilchrist, H. G., & Love, O. P. (2016). Cold tolerance, and not earlier arrival on breeding grounds, explains why males winter further north in an Arctic-breeding songbird. *Journal of Avian Biology*, *47*, 7–15. <https://doi.org/10.1111/jav.00689>

Matyjasiak, P., Rubolini, D., Romano, M., & Saino, N. (2016). No short-term effects of geolocators on flight performance of an aerial insectivorous bird, the Barn Swallow (*Hirundo rustica*). *Journal of Ornithology*, *157*, 653–661. <https://doi.org/10.1007/s10336-015-1314-2>

McNeil, S. E. M., Tracy, D., & Cappello, C. D. (2015). Loop migration by a Western Yellow-billed Cuckoo wintering in the gran chaco. *Western Birds*, *46*, 244–255.

Meier, C. M., Karaard, H., Aymí, R., Peev, S. G., Bächler, E., Weber, R., ... Liechti, F. (2018). What makes Alpine swift ascend at twilight? Novel geolocators reveal year-round flight behaviour. *Behavioral Ecology and Sociobiology*, *72*, 45. <https://doi.org/10.1007/s00265-017-2438-6>

Minton, C., Gosbell, K., Johns, P., Christie, M., Klaassen, M., Hassell, C., ... Fox, J. (2013). New insights from geolocators deployed on waders in Australia. *Wader Study Group Bulletin*, 120, 37–46.

Minton, C., Gosbell, K., Johns, P., Christie, M., Klaassen, M., Hassell, C., ... Fox, J. W. (2011). Geocator studies on Ruddy Turnstones *Arenaria interpres* and Greater Sandplovers *Charadrius leschenaultii* in the East Asian- Australasia Flyway reveal widely different migration strategies. *Wader Study Group Bulletin*, 118, 87–96.

Nelson, A. R., Cormier, R. L., Humple, D. L., Scullen, J. C., Sehgal, R., & Seavy, N. E. (2016). Migration patterns of San Francisco Bay Area Hermit Thrushes differ across a fine spatial scale. *Animal Migration*, 3, 1–13. <https://doi.org/10.1515/ami-2016-0001>

Norevik, G., Åkesson, S., & Hedenström, A. (2017). Migration strategies and annual space-use in an Afro- Palaearctic aerial insectivore – the European nightjar. *Journal of Avian Biology*, 48, 738–747. <https://doi.org/10.1111/jav.01071>

Ouwehand, J., Ahola, M. P., Ausems, A. N. M. A., Bridge, E. S., Burgess, M., Hahn, S., ... Both, C. (2016). Light-level geolocators reveal migratory connectivity in European populations of pied flycatchers *Ficedula hypoleuca*. *Journal of Avian Biology*, 47, 69–83. <https://doi.org/10.1111/jav.00721>

Ouwehand, J., & Both, C. (2017). African departure rather than migration speed determines variation in spring arrival in pied flycatchers. *Journal of Animal Ecology*, 86, 88–97. <https://doi.org/10.1111/1365-2656.12599>

Ouwehand, J., & Both, C. (2017). Data from: African departure rather than migration speed determines variation in spring arrival in pied flycatchers. *Dryad Digital Depository*, <https://doi.org/10.5061/dryad.k6q68>

Pakanen, V. M., Rönkä, N., Thomson, R. L., & Koivula, K. (2015). No strong effects of leg-flagged geolocators on return rates or reproduction of a small long-distance migratory shorebird. *Ornis Fennica*, *92*, 101–111.

Perlut, N. G. (2018). Prevalent transoceanic fall migration by a 30-gram songbird, the Bobolink. *The Auk*, *135*, 992–997. <https://doi.org/10.1642/AUK-18-56.1>

Peterson, S. M., Streby, H. M., Kramer, G. R., Lehman, J. a., Buehler, D. a., & Andersen, D. E. (2015). Geolocators on Golden-winged Warblers do not affect migratory ecology. *The Condor*, *117*, 256–261. <https://doi.org/10.1650/CONDOR-14-200.1>

Pillar, A. G., Marra, P. P., Flood, N. J., & Reudink, M. W. (2016). Molt migration in Bullock's orioles (*Icterus bullockii*) confirmed by geolocators and stable isotope analysis. *Journal of Ornithology*, *157*, 265–275. <https://doi.org/10.1007/s10336-015-1275-5>

Procházka, P., Brlík, V., Yohannes, E., Meister, B., Auerswald, J., Ilieva, M., & Hahn, S. (2018). Across a migratory divide: divergent migration directions and non-breeding grounds of Eurasian reed warblers revealed by geolocators and stable isotopes. *Journal of Avian Biology*, *49*, jav-012516. <https://doi.org/10.1111/jav.01769>

Renfrew, R. B., Kim, D., Perlut, N., Smith, J., Fox, J., & Marra, P. P. (2013). Phenological matching across hemispheres in a long-distance migratory bird. *Diversity and Distributions*, *19*, 1008–1019. <https://doi.org/10.1111/ddi.12080>

Ross, J. D., Bridge, E. S., Rozmarynowycz, M. J., & Bingman, V. P. (2014). Individual variation in migratory path and behaviour among Eastern Lark Sparrows. *Animal Migration*, *2*, 29–33. <https://doi.org/10.2478/ami-2014-0003>

Ryder, T. B., Fox, J. W., & Marra, P. P. (2011). Estimating migratory connectivity of Gray Catbirds (*Dumetella carolinensis*) using geocator and mark—recapture data. *The Auk*, *128*, 448–453. <https://doi.org/10.1525/auk.2011.11091>

- Salewski, V., Flade, M., Poluda, A., Kiljan, G., Liechti, F., Lisovski, S., & Hahn, S. (2013). An unknown migration route of the “globally threatened” Aquatic Warbler revealed by geolocators. *Journal of Ornithology*, *154*, 549–552. <https://doi.org/10.1007/s10336-012-0912-5>
- Scandolaro, C., Rubolini, D., Ambrosini, R., Caprioli, M., Hahn, S., Liechti, F., ... Saino, N. (2014). Impact of miniaturized geolocators on barn swallow *Hirundo rustica* fitness traits. *Journal of Avian Biology*, *45*, 417–423. <https://doi.org/10.1111/jav.00412>
- Schmaljohann, H., Buchmann, M., Fox, J. W., & Bairlein, F. (2012). Tracking migration routes and the annual cycle of a trans-Sahara songbird migrant. *Behavioral Ecology and Sociobiology*, *66*, 915–922. <https://doi.org/10.1007/s00265-012-1340-5>
- Schmaljohann, H., Meier, C., Arlt, D., Bairlein, F., van Oosten, H., Morbey, Y. E., ... Eikenaar, C. (2016). Proximate causes of avian protandry differ between subspecies with contrasting migration challenges. *Behavioral Ecology*, *27*, 321–331. <https://doi.org/10.1093/beheco/arv160>
- Seavy, N. E., Humple, D. L., Cormier, R. L., & Gardali, T. (2012). Establishing the breeding provenance of a temperate-wintering north american passerine, the golden-crowned sparrow, using light-level geolocation. *PLoS ONE*, *7*, e34886. <https://doi.org/10.1371/journal.pone.0034886>
- Sechrist, J., Paxton, E., Ahlers, D., Doster, R., & Ryan, V. M. (2012). One year of migration data for a western yellow-billed cuckoo. *Western Birds*, *43*, 2–11.
- Smith, M., Bolton, M., David, J., Summers, R. W., Ellis, P., & Wilson, J. D. (2014). Short communication Geolocator tagging reveals Pacific migration of Red-necked Phalarope *Phalaropus lobatus* breeding in Scotland. *Ibis*, *156*, 870–873. <https://doi.org/10.1111/ibi.12196>
- Stutchbury, B. J. M., Gow, E. A., Done, T., MacPherson, M., Fox, J. W., & Stutchbury, B. J. M. (2010). Effects of post-breeding moult and energetic condition on timing of songbird migration into the tropics. *Proceedings of the Royal Society B: Biological Sciences*, *278*, 131–137. <https://doi.org/10.1098/rspb.2010.1220>

Stutchbury, B. J. M., Tarof, S. A., Done, T., Gow, E., Kramer, P. M., Tautin, J., ... Afanasyev, V. (2009).

Tracking long-distance songbird migration by using geolocators. *Science*, 323, 896.

<https://doi.org/10.1126/science.1166664>

Szép, T., Liechti, F., Nagy, K., Nagy, Z., & Hahn, S. (2017). Discovering the migration and non-breeding

areas of sand martins and house martins breeding in the Pannonian basin (central-eastern

Europe). *Journal of Avian Biology*, 48, 114–122. <https://doi.org/10.1111/jav.01339>

Tøttrup, A. P., Klaassen, H. G., Strandberg, R., Thorup, K., Kristensen, M. W., Jørgensen, P. S., ...

Alerstam, T. (2012). The annual cycle of a trans-equatorial Eurasian–African passerine migrant:

different spatio-temporal strategies for autumn and spring migration. *Proceedings of the Royal*

Society B: Biological Sciences, 279, 1009–1016. <https://doi.org/10.1098/rspb.2011.1323>

van Oosten, H. H., Versluijs, R., & van Wijk, R. (2014). Twee Nederlandse Tapuiten in de Sahel:

trekroutes en winterlocaties ontrafeld. *Limosa*, 87, 168–172.

van Wijk, R. E., Schaub, M., Tolkmitt, D., Becker, D., & Hahn, S. (2013). Short-distance migration of

Wrynecks *Jynx torquilla* from Central European populations. *Ibis*, 155, 886–890.

<https://doi.org/10.1111/ibi.12083>

van Wijk, R. E., Souchay, G., Jenni-Eiermann, S., Bauer, S., & Schaub, M. (2015). No detectable effects

of lightweight geolocators on a Palaearctic-African long-distance migrant. *Journal of*

Ornithology, 157, 255–264. <https://doi.org/10.1007/s10336-015-1274-6>

Weiser, E. L., Lanctot, R. B., Brown, S. C., Alves, J. A., Battley, P. F., Bentzen, R., ... Sandercock, B. K.

(2016). Effects of geolocators on hatching success, return rates, breeding movements, and

change in body mass in 16 species of Arctic-breeding shorebirds. *Movement Ecology*, 4, 12.

<https://doi.org/10.1186/s40462-016-0077-6>

Wellbrock, A. H. J., Bauch, C., Rozman, J., & Witte, K. (2017). "Same procedure as last year?" – Repeatedly tracked swifts show individual consistency in migration pattern in successive years. *Journal of Avian Biology*, 48, 897–903. <https://doi.org/10.1111/jav.01251>

Woodworth, B. K., Newman, A. E. M., Turbek, S. P., Dossman, B. C., Hobson, K. A., Wassenaar, L. I., ...

Norris, D. R. (2016). Differential migration and the link between winter latitude, timing of migration, and breeding in a songbird. *Oecologia*, 181, 413–422.

<https://doi.org/10.1007/s00442-015-3527-8>

Xenophontos, M., Blackburn, E., & Cresswell, W. (2017). Cyprus Wheatears *Oenanthe cypriaca* likely reach sub-Saharan African wintering grounds in a single migratory flight. *Journal of Avian Biology*, 48, 529–535. <https://doi.org/10.1111/jav.01119>

Table 1. Explanatory variables used in the multivariate meta-analysis of apparent survival extracted from published and unpublished geolocator studies or from the literature. *N* presents the number of records specified as the groups of tagged birds from the same study site, year of attachment, of the same sex, and the specific geolocator and the attachment type accompanied with the corresponding control groups.

Methodological aspect	<i>Description</i>	<i>N</i>
Published data	Published – data from published studies (for details see Methods), data from unpublished sources from years following an already published study, or data initially collected as unpublished but published by 31 August 2018	303
	Unpublished – data from unpublished studies	123
Control group	Matched – birds handled in the exactly same way as geolocator-tagged birds except for geolocator deployment	102
	Marked only – birds of the same sex, age, from the same year and study site or birds from the same site, from different years	324
Species trait		
Foraging strategy ^{1,2}	Aerial forager	122
	Non-aerial forager	304
Sex	Males	195
	Females	120
Geolocator specification		
Relative load	% of geolocator mass (including the harness) of the body mass of the tagged birds	418
Stalk/pipe length*	Length (mm) of the stalk/pipe holding the light sensor or guiding the light towards the sensor (0 mm for stalkless models)	371

Attachment specification

Attachment type	Leg-loop harness	304
	Full-body harness	80
	Leg-flag attachment	42
Material elasticity*	Elastic – elastan, ethylpropylen, neoprene, rubber, silicone, silastic, or Stretch Magic	235
	Non-elastic – cord, kevlar, nylon, plastic, polyester, or teflon	146

Ecological trait

Life-histories	Great circle distance between geolocator deployment site and population-specific centroid of the non-breeding (or breeding) range	426
	Male body mass (g)	426
	Female body mass (g)	426
	Nest type – open/close	426
	Clutch size (number of eggs)	426
	Number of broods per year	426
	Dense habitat preference (species occurs especially in dense habitats e.g. reeds or scrub) – yes/no	426
	Egg mass (g) – mean fresh mass ³	426
	Clutch mass (g) – egg mass × clutch size	426

* only used for harness attachments

¹ Cramp & Perrins, 1977–1994

² Rodewald, 2015

³ Schönwetter, 1960–1992

Table 2. Number of unpublished effect sizes included in the analysis and Egger's regression tests of the null model residuals against their precision to assess the presence of publication bias.

<i>Trait category</i>	<i>Unpublished (%)</i>		<i>Egger's regression</i>			
	<i>Effect sizes</i>	<i>N</i>	<i>Intercept</i>	<i>t</i>	<i>SE</i>	<i>P</i>
Apparent survival	28.9	426	0.12	1.53	0.08	0.121
Condition	63.3	79	-0.36	-1.70	0.21	0.088
Phenology	59.1	22	-0.26	-1.28	0.21	0.217
Breeding performance	27.3	22	-0.01	-0.01	0.61	0.993

Table 3. Summary of the full trait model (n = 281; model 3) and the ecological model (n = 426; model 4) of the geolocator effects on apparent survival. Levels contrasted against the reference level are given in parentheses.

Full trait model					
<i>Trait</i>	<i>Estimate</i>	<i>SE</i>	<i>Z</i>	<i>95% CI</i>	<i>P</i>
Intercept	-0.25	0.10	-2.59	(-0.44; -0.06)	0.010
Published (published)	0.14	0.10	1.39	(-0.06; 0.34)	0.164
Control type (matched)	-0.05	0.09	-0.61	(-0.23; 0.12)	0.542
Foraging strategy (aerial)	-0.09	0.14	-0.61	(-0.36; 0.19)	0.540
Sex (males)	-0.07	0.05	-1.30	(-0.17; 0.03)	0.192
Relative load	-0.12	0.05	-2.36	(-0.23; -0.02)	0.018
Stalk/pipe length	0.07	0.04	1.77	(-0.01; 0.15)	0.077
Material elasticity (non-elastic)	0.19	0.08	2.21	(0.03; 0.35)	0.026
Foraging strategy (aerial) × stalk length	-0.10	0.07	-1.40	(-0.25; 0.04)	0.161
Ecological model					
<i>Trait</i>	<i>Estimate</i>	<i>SE</i>	<i>Z</i>	<i>95% CI</i>	<i>P</i>
Intercept	-0.26	0.08	-3.20	(-0.42; -0.10)	0.001
PC1	0.06	0.03	2.32	(0.01; 0.11)	0.026
PC2	0.02	0.03	0.47	(-0.05; 0.08)	0.638
Dense habitat (yes)	0.03	0.13	0.21	(-0.22; 0.27)	0.834
Nest type (open)	0.14	0.11	1.27	(-0.08; 0.36)	0.205

