

<https://helda.helsinki.fi>

---

## Dermatitis herpetiformis arising within vitiligo in a patient with autoimmune polyendocrine syndrome type 3

Köykkä, Hanna

2019-06

---

Köykkä , H , Lappalainen , K & Kluger , N 2019 , ' Dermatitis herpetiformis arising within vitiligo in a patient with autoimmune polyendocrine syndrome type 3 ' , International Journal of Dermatology , vol. 58 , no. 6 , pp. E123-E125 . <https://doi.org/10.1111/ijd.14446>

---

<http://hdl.handle.net/10138/312893>

<https://doi.org/10.1111/ijd.14446>

---

draft

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

## Correspondence

### Dermatitis herpetiformis arising within vitiligo in a patient with autoimmune polyendocrine syndrome type 3

The association between dermatitis herpetiformis (DH) and vitiligo is infrequent. It concerns 1.3% of the patients with DH<sup>1</sup> and 1% of the patients with vitiligo.<sup>2</sup> Case reports showed a peculiar affinity of DH lesions for vitiligo patches.<sup>3-9</sup> We report here an illustrative case of DH that displayed affinity phenomenon on vitiligo patches in a patient with autoimmune polyendocrine syndrome (APS) type 3.

A 47-year-old woman presented with a 2-week history of pruritic papulovesicular rash. Her past medical history included type 1 diabetes, hypothyroidism, vitiligo, asthma, hypertension, atopic eczema, and mood disorder. Her two children had celiac disease. The patient had a previous blood screening test with positive tissue transglutaminase IgA antibodies (19; normal <7). She denied any intestinal symptoms, and past gastroscopy and

biopsies of the duodenum were normal. She had been followed in the past in our clinic for an itchy rash. Despite recurrent suspicions of DH, direct immunofluorescence (DIF) remained negative on several occasions 20 years ago. The rash had been treated since as atopic eczema with topical corticosteroids and moisturizers. Upon examination, the patient had symmetrical pruritic erythematous papules, vesicles, and excoriations on extensor surfaces of elbows, knees, and buttocks. She also displayed lesions on wrists, ankles, pubic area, and hairline (Fig. 1). The lesions were mainly localized within vitiligo. Vitiligo patches on the inner thighs and abdomen were mainly spared. DH disease was again highly suspected. A punch skin biopsy of a vesicle showed upper dermal inflammation with lymphocytes, macrophages and eosinophils, and subepidermal cleavage. Two DIF from an erythematous area and a normal appearing perilesional area revealed granular IgA deposition on the basal



**Figure 1** Vesicles of DH restricted to vitiligo of the elbows and forearm (a), knees (b), and dorsum of the hand (c). Complete regression after 3 months with dapsone 100 mg/day and gluten-free diet (d)

**Table 1** Reported cases of dermatitis herpetiformis associated with vitiligo


Case	Author	Gender	Age (years) at the time of diagnosis of DH	Co-localization	First pathology	Other autoimmune conditions
1	Allende <i>et al.</i> , 1964 <sup>3</sup>	M	35	No	Vitiligo	No
2	Ortonne <i>et al.</i> , 1976 <sup>2</sup>	–	–	–	–	–
3	Olholm-Larsen <i>et al.</i> , 1980 <sup>4</sup>	W	33	Yes	DH	No
4	Combs <i>et al.</i> , 1980 <sup>5</sup>	M	25	Yes	Vitiligo	MGN
5	Hogan <i>et al.</i> , 1986 <sup>6</sup>	W	46	Yes	Vitiligo	Antithyroid antibodies without hypothyroidism
6	Reunala <i>et al.</i> , 1997 <sup>1</sup>	W	–	–	–	–
7	Reunala <i>et al.</i> , 1997 <sup>1</sup>	M	–	–	–	–
8	Reunala <i>et al.</i> , 1997 <sup>1</sup>	M	–	–	–	–
9	Reunala <i>et al.</i> , 1997 <sup>1</sup>	M	–	–	DH	–
10	Amato <i>et al.</i> , 2000 <sup>7</sup>	W	53	Yes	Vitiligo	Antithyroid antibodies without hypothyroidism
11	Karabudak <i>et al.</i> , 2007 <sup>8</sup>	M	21	Yes	Vitiligo	No
12	Macbeth <i>et al.</i> , 2013 <sup>9</sup>	W	37	Yes	Vitiligo	APS-2 (thyroiditis, adrenal insufficiency, DT1, CD)
13	Present case, 2018	W	47	Yes	Vitiligo	APS-3 (DT1, thyroiditis, vitiligo)

APS, autoimmune polyendocrine syndrome; CD, celiac disease; DT1, type 1 diabetes; M, man; MGN, membranous glomerulonephritis; NA, not available; W, woman.

<sup>a</sup>Cases extracted from large series. No specific clinical data given.

membrane. She started oral disulone 100 mg/day along with gluten-free diet. No gastroscopy was performed. At 3-month follow-up, the patient was clear of any lesion.

We report here a new case of association between DH and vitiligo. Our case is striking by the very strong autoimmune personal and familial context. The patient had APS type 3, which is defined by autoimmune thyroiditis with other organ-specific immunity, here type 1 diabetes and vitiligo, but without adrenal insufficiency.<sup>10</sup> A previous case was reported with APS type 2.<sup>9</sup> The localization and affinity of DH lesions within vitiligo is striking. We found 12 other cases of DH coexisting with vitiligo in the literature (Table 1).<sup>1–9</sup> Half were women, and the median age at diagnosis of DH was 36 years. In most of the cases (7/9), vitiligo preceded DH. DH was co-localized totally or at least partly on vitiligo in 7/8 patients. Only Allende *et al.* reported a case of DH sparing vitiligo patches.<sup>3</sup> Additional autoimmune features were found in 5/8 patients with either APS (2/8), circulating antithyroid antibodies (2/8), or in one case membranous glomerulonephritis during follow-up.<sup>5</sup> The mechanism of DH affinity for vitiligo patches is unknown. Hypotheses of the literature include: (i) Köebner phenomenon,<sup>4</sup> (ii) genetic mosaicism in case of segmental vitiligo,<sup>9</sup> (iii) fortuitous as DH lesions may be located on other areas without vitiligo.<sup>8</sup> Besides, DH antigens could be unmasked in vitiliginous areas under the action of UV exposure, trauma, or modified local immunity associated with melanocyte destruction.

Hanna Köykkä, MD  
Katriina Lappalainen, MD, PhD  
Nicolas Kluger \*, MD, PhD 

Department of Dermatology, Allergology and Venereology,  
Helsinki University Central Hospital, Helsinki, Finland  
\*E-mail: nicolas.kluger@hus.fi

Funding: none.

Conflict of interest: none declared.

doi: 10.1111/ijd.14446

## References

- Reunala T, Collin P. Diseases associated with dermatitis herpetiformis. *Br J Dermatol* 1997; **136**: 315–318.
- Ortonne JP, Perrot H, Thivolet J. Clinical and statistical study of 100 cases of vitiligo. II. – Associated lesions. *Sem Hop* 1976; **52**: 679–686.
- Allende MF, Reed E. Dermatitis herpetiformis and vitiligo. *Arch Dermatol* 1964; **89**: 156.
- Olholm-Larsen P, Kavli G. Dermatitis herpetiformis and vitiligo. *Dermatologica* 1980; **160**: 41–44.
- Combs R, Hazelrigg DE. Dermatitis herpetiformis and Membranous Glomerulonephritis. *Cutis* 1980; **25**: 660–661.
- Hogan DJ, Lane PR. Dermatitis herpetiformis and vitiligo. *Cutis* 1986; **38**: 195–197.
- Amato L, Gallerani I, Fuligni A, *et al.* Dermatitis herpetiformis and vitiligo: report of a case and review of the literature. *J Dermatol* 2000; **27**: 462–466.
- Karabudak O, Dogan B, Yildirim S, *et al.* Case report. Dermatitis herpetiformis and vitiligo. *J Chin Med Assoc* 2007; **70**: 504–506.
- Macbeth AE, Lee KYC, Igali L, *et al.* Dermatitis herpetiformis arising within vitiligo in the presence of multiple autoimmune conditions. *J Dermatol Case Rep* 2013; **7**: 101–102.
- Warzée E, Legros JJ, Geenen V. Autoimmune polyendocrine syndrome (APS). *Rev Med Liege* 2002; **57**: 710–714.