




Contents lists available at ScienceDirect

European Journal of Obstetrics & Gynecology and Reproductive Biology

journal homepage: www.elsevier.com/locate/ejogrb

Maternal hemoglobin associates with preterm delivery and small for gestational age in two Finnish birth cohorts



Justiina Ronkainen^{a,b,*}, Estelle Lowry^{a,b}, Anni Heiskala^{a,b}, Iida Uusitalo^{a,b},
Peppi Koivunen^{b,c,d}, Eero Kajantie^{e,f,g,h}, Marja Väärasmäki^{e,f}, Marjo-Riitta Järvelin^{a,b,i,j},
Sylvain Sebert^{a,b,k}

^a Center for Life Course Health Research, University of Oulu, Oulu, Finland

^b Biocenter Oulu, University of Oulu, Oulu, Finland

^c Faculty of Biochemistry and Molecular Medicine, University of Oulu, Oulu, Finland

^d Oulu Center for Cell-Matrix Research, University of Oulu, Oulu, Finland

^e PEDEGO Research Unit, MRC Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland

^f Public Health Promotion Unit, Department of Public Health Solutions, National Institute for Health and Welfare, Helsinki, Finland

^g Hospital for Children and Adolescents, Helsinki University Central Hospital and University of Helsinki, Helsinki, Finland

^h Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology, Trondheim, Norway

ⁱ Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom

^j MRC-PHE Centre for Environment and Health, School of Public Health, Imperial College London, London, United Kingdom

^k Department for Genomics of Common Diseases, School of Medicine, Imperial College London, London, United Kingdom

ARTICLE INFO

Article history:

Received 20 February 2019

Received in revised form 27 March 2019

Accepted 29 April 2019

Keywords:

Intrauterine growth restriction

Large for gestational age

Maternal hemoglobin

Preterm delivery

Small for gestational age

ABSTRACT

Objective: To test whether maternal hemoglobin during pregnancy associates with offspring perinatal outcomes in a developed country. Changes in maternal hemoglobin concentration during pregnancy are partly physiological phenomena reflecting alterations of maternal blood volume. Especially hemoglobin measures outside the physiological range may influence maternal health and fetal growth with long-lasting consequences.

Study design: We studied an unselected sample drawn from two regional birth cohorts born 20 years apart: The Northern Finland Birth Cohorts 1966 and 1986. These are two mother-and-child population-based birth cohorts together comprising 21,710 mothers and their children. After exclusions, the sample size of the current study was 20,554. Concentrations of maternal hemoglobin at first and last antenatal visits were categorized as low (lowest 10%), medium (reference) or high (highest 10%). Multinomial logistic regression analyses for categories of maternal hemoglobin and perinatal outcomes such as preterm delivery and full-term small and large for gestational age were conducted with adjustments for maternal cofactors.

Results: Low maternal hemoglobin at early pregnancy associated with decreased risk of full-term small for gestational age (adjusted OR 0.73, 95% CI [0.58, 0.93], $p=0.010$). At late pregnancy, low maternal hemoglobin associated with increased risk of preterm delivery (adjusted OR 1.60, 95% CI [1.26, 2.02], $p<0.0005$) whereas high maternal hemoglobin associated with increased risk of full-term small for gestational age (adjusted OR 1.29, 95% CI [1.07, 1.56], $p=0.009$). Maternal hemoglobin did not show constant association with risk of large for gestational age.

Conclusion: The results from this study support evidence that both low and high maternal hemoglobin associate with adverse perinatal outcomes. Low maternal hemoglobin associated with preterm delivery and high with full-term small for gestational age. Association was mainly present when maternal hemoglobin was measured during the third trimester. These results indicate that it is important to monitor both extremes of maternal hemoglobin throughout the pregnancy.

© 2019 Elsevier B.V. All rights reserved.

Introduction

Both low and high maternal hemoglobin (mHb) concentration has been consistently associated with adverse perinatal outcomes

* Corresponding author at: PO BOX 5000, FI-90014, Oulu, Finland.
E-mail address: justiina.ronkainen@oulu.fi (J. Ronkainen).

such as preterm birth and fetal growth restriction [1–7] and in our own data with later developmental indices [8]. Low hemoglobin and anemia (generally defined as mHb < 110 g/l) are global problems affecting both the mother and the developing child. In 2011, the global prevalence of anemia among pregnant women was 38%, which translated to 32 million mothers [9]. Regionally, the prevalence of maternal anemia vary substantially from 22% in European and North American countries to 56% in Central and West Africa [10].

Several environmental and behavioral maternal factors affect mHb and the developing child. Smoking during pregnancy has been associated with placental defects, prematurity and growth restriction (reviewed in [11]). Furthermore, smoking associates with increased mHb levels, which may contribute to increased risk of adverse perinatal outcomes [7,12]. Low maternal BMI is associated with increased risk of spontaneous preterm birth whereas some studies report obesity to have a protective effect [13]. However, obese mothers are in risk for hypertensive disorders, which associate with increased risk of medically indicated preterm birth [13]. Mother's low education was associated with prematurity and small for gestational age (SGA) in recent meta-analysis comprising of 12 European countries, although there were differences between countries [14]. Nulliparous mothers are at increased risk of preterm delivery and SGA

specifically if the mother is under 18 years old and maternal age 35 or more is associated with increased risk of SGA but not prematurity [15].

In the current study, we study the association between mHb and adverse perinatal outcomes within two study populations located in Finland. These populations are highly homogenous due to standardized antenatal care and pregnancy counselling, which contributes to fewer environmental confounders than in more diverse settings. Association between mHb at different time points of pregnancy and such outcomes as prematurity and small and large for gestational age (SGA and LGA) are studied.

Material and methods

Inclusion and exclusion criteria

The study population is the mothers and children of Northern Finland Birth Cohorts (NFBC) 1966 and 1986 which are population-based birth cohorts comprising 98% (n=21,710) of all deliveries occurred in 1966 and 1985–86 in Oulu and Lapland provinces of Finland. Multiple pregnancies, individuals without both early and late mHb measurements and individuals with mHb over 5 SD from mean (n = 1,156) were excluded from the analyses resulting in final

Table 1

Characteristics of mothers and singleton newborns in the pooled sample and stratified by the cohort.

	Pooled	NFBC1966	NFBC1986	p value [*]
Early maternal hemoglobin				< 0.0005 ^A
Reference, n (%)	15248 (80.4)	8348 (83.7)	6900 (76.8)	
Low, n (%)	1888 (10.0)	827 (8.3)	1061 (11.8)	
High, n (%)	1817 (9.6)	798 (8.0)	1019 (11.3)	
Late maternal hemoglobin				< 0.0005 ^A
Reference, n (%)	16096 (80.3)	9118 (81.4)	6978 (76.0)	
Low, n (%)	1978 (9.9)	926 (8.3)	1052 (11.9)	
High, n (%)	1961 (9.8)	1153 (10.3)	808 (9.1)	
Mean maternal age, years (SD)	27.8 (6.2)	27.8 (6.7)	27.8 (5.5)	0.911
Mean pre-pregnancy BMI, kg/m² (SD)	22.8 (3.4)	23.1 (3.2)	22.4 (3.5)	< 0.0005
Maternal smoking				< 0.0005 ^A
No, n (%)	15581 (76.9)	8827 (78.1)	6754 (75.4)	
Yes, but stopped, n (%)	986 (4.9)	753 (6.7)	233 (2.6)	
Yes, n (%)	3701 (18.3)	1728 (15.3)	1973 (22)	
Maternal SES				< 0.0005 ^B
Professional, n (%)	3697 (18.4)	1435 (12.6)	2262 (25.9)	
Skilled worker, n (%)	5981 (29.7)	2469 (21.7)	3512 (40.2)	
Unskilled worker, n (%)	3539 (17.6)	985 (8.7)	2554 (29.2)	
Housewife, n (%)	6906 (34.3)	6497 (57.1)	409 (4.7)	
Parity				0.076
First child, n (%)	6813 (33.2)	3770 (32.7)	3043 (33.9)	
Second or later child, n (%)	13712 (66.8)	7768 (67.3)	5944 (66.1)	
Maternal hypertensive disorders				< 0.0005 ^C
Normotensive, n (%)	13535 (83)	6460 (76.3)	7075 (90.3)	
Gestational hypertension, n (%)	1253 (7.7)	983 (11.6)	270 (3.4)	
Pre-eclampsia, n (%)	423 (2.6)	241 (2.8)	182 (2.3)	
Chronic hypertension, n (%)	881 (5.4)	668 (7.9)	213 (2.7)	
Superimposed pre-eclampsia, n (%)	208 (1.3)	116 (1.4)	92 (1.2)	
Mean birth weight, grams (SD)	3518 (556)	3478 (560)	3570 (546)	< 0.0005
Mean gestational age, weeks (SD)	39.9 (1.9)	40.0 (2.1)	39.8 (1.7)	< 0.0005
Boys, n (%)	10529 (51.2)	5917 (51.2)	4612 (51.3)	0.978
Preterm delivery, n (%)	1003 (5.0)	610 (5.5)	393 (4.5)	0.002
Full-term SGA, n (%)	1746 (9.2)	1172 (11.1)	574 (6.9)	< 0.0005
Full-term LGA, n (%)	1773 (9.4)	908 (8.6)	865 (10.3)	< 0.0005

^{*} Comparison between NFBC1966 and NFBC1986.

^A p < 0.05 between all mHb categories.

^B p < 0.05 between all SES categories except professional vs. skilled worker.

^C p < 0.05 between all hypertensive disorder categories except normotensive vs. superimposed pre-eclampsia, gestational hypertension vs. chronic hypertension and pre-eclampsia vs. superimposed pre-eclampsia.

sample size of 20,554. Concentration of mHb at early pregnancy was available from 18,953 mothers and at late pregnancy from 20,035 mothers. Informed consent was obtained from study subjects for the use of their data in the study. Approval for the studies was granted by the ethics committee of the Northern Ostrobothnia Hospital District in Oulu, Finland in accordance with the declaration of Helsinki.

Predictor variables

Mean gestational week for early mHb measurement was 13.3 (SD 4.8) and for late mHb 38.2 (SD 3.4). In NFBC1966, mHb was measured with Tallquist chart, hemometer or photometer and in NFBC1986, with standard photometer. In NFBC1966, mHb was first standardized for measurement method and then mHb measures were transformed to z values [$z(\text{mHb}) = (\text{mHb} - \text{mean}(\text{mHb})) / \text{SD}(\text{mHb})$] separately for each cohort and pregnancy time point. Z value of mHb below 10th percentile was defined as low and above 90th percentile as high mHb while the middle 80% was used as reference.

Perinatal outcomes

Children born before 37 weeks of gestation were defined as preterm. SGA and LGA were analyzed from full-term children (gestational age 37 weeks or more) only. SGA was defined as below 10th percentile and LGA as above 90th percentile of the Finnish national reference according to [16].

Covariates

Following factors were used as covariates in the statistical models: maternal smoking, maternal hypertensive disorders, maternal pre-pregnancy body mass index (BMI), maternal socioeconomic status (SES), parity and maternal age. Maternal smoking was categorized as not smoked during pregnancy (reference, $n = 15,581$), smoked, but not after second month of pregnancy ($n = 986$) and smoked after second month of pregnancy ($n = 3,701$). Maternal hypertensive disorders were defined as described in [17]: Normotensive (reference, $n = 13,535$), gestational hypertension ($n = 1,253$), pre-eclampsia ($n = 423$), chronic hypertension ($n = 881$) and superimposed pre-eclampsia ($n = 208$). Maternal SES was categorized as professionals (reference, $n = 3,697$), skilled workers ($n = 5,981$), unskilled workers ($n = 3,539$) and housewives, farmers and farmer's wives ($n = 6,906$). Parity was dichotomized as nulliparous (reference, $n = 6,813$) and multiparous ($n = 13,712$). Maternal pre-pregnancy BMI (weight in kg/(height in m)²) and age were used as continuous variables. Cohort was included in the models, NFBC1966 as a reference ($n = 11,554$) and NFBC1986 ($n = 9,000$). Sex of the child was not included in the models because it did not associate with prematurity in the current

population and SGA and LGA were determined from sex specific growth curves [16].

Statistics

Data from two cohorts were pooled. Differences in maternal and fetal characteristics between the cohorts were analyzed with independent samples *t*-test for continuous variables and chi squared test of independence with subsequent post hoc analyses for categorical variables. Association between mHb at different time points of pregnancy and perinatal outcomes was analyzed using multinomial logistic regression (MLR). Odds ratios (OR) and 95% confidence intervals (95% CI) were recorded and p-values according to Wald's test that were below 0.0125 (0.05/4 tests) were considered statistically significant. Association was analyzed without adjustments (base model), with adjustment for cohort, maternal smoking and maternal hypertensive disorders (model 1) and with adjustment for cohort, maternal smoking, maternal hypertensive disorders, pre-pregnancy BMI, SES, parity and maternal age at delivery (model 2). Statistical analyses were conducted using R software package version 3.5.0 [18].

Results

Characteristics of the sample population

Distribution of offspring sex, maternal age and parity were similar between the cohorts (Table 1). Other maternal and fetal characteristics changed within the 20 years between the cohort initiations. Number of mothers with low mHb at any point of pregnancy was lower in NFBC1966 than in NFBC1986 whereas number of housewives was considerably higher in NFBC1966. Furthermore, number of mothers with either gestational or chronic hypertension was higher in NFBC1966.

Multinomial logistic regression for perinatal outcomes

As shown in Table 2, low mHb at late pregnancy associated with increased risk of prematurity (fully adjusted OR (aOR) 1.60, 95% CI [1.26, 2.02], $p < 0.0005$). Associations between mHb and SGA and LGA are indicated in Table 3. Low mHb at early pregnancy associated with decreased risk of SGA (aOR 0.73, 95% CI [0.58, 0.93], $p = 0.010$) and high mHb at late pregnancy with increased risk of SGA (aOR 1.29, 95% CI [1.07, 1.56], $p = 0.009$, Table 2). Low mHb at early pregnancy associated with increased risk of LGA; however, the association was not statistically significant after adjustment for maternal cofactors. High mHb at early pregnancy associated with decreased risk of LGA in fully adjusted model but not in other models. Results are shown separately for each cohort in Supplementary Table 1. To characterize spontaneous preterm delivery and normotensive SGA, we conducted additional MLR analyses for pregnancies without maternal hypertensive disorders

Table 2

Association between maternal hemoglobin during pregnancy and preterm delivery. OR, odds ratio; 95% CI, 95% confidence interval; p, p-value according to Wald's test, significant when $p < 0.0125$.

Outcome	mHb	Base model			Model 1 ^a			Model 2 ^b		
		OR	95% CI	p	OR	95% CI	p	OR	95% CI	p
Preterm delivery	Low early	1.17	[0.94, 1.44]	0.157	1.23	[0.97, 1.56]	0.081	1.24	[0.97, 1.59]	0.079
	High early	1.12	[0.90, 1.39]	0.323	1.10	[0.86, 1.41]	0.437	1.18	[0.91, 1.52]	0.209
	Low late	1.58	[1.29, 1.93]	< 0.0005	1.64	[1.31, 2.05]	< 0.0005	1.60	[1.26, 2.02]	< 0.0005
	High late	0.87	[0.67, 1.12]	0.275	0.81	[0.61, 1.09]	0.166	0.83	[0.62, 1.13]	0.236

^a Model 1 adjusted for cohort, maternal smoking and maternal hypertensive disorders.

^b Model 2 adjusted for cohort, maternal smoking, maternal hypertensive disorders, pre-pregnancy BMI, maternal SES, parity and maternal age.

Table 3

Association between maternal hemoglobin during pregnancy and full-term small and large for gestational age (SGA and LGA, respectively). OR, odds ratio; 95% CI, 95% confidence interval; p, p-value according to Wald's test, significant when $p < 0.0125$.

Outcome	mHb	Base model			Model 1 ^a			Model 2 ^b		
		OR	95% CI	p	OR	95% CI	p	OR	95% CI	p
SGA	Low early	0.74	[0.61, 0.90]	0.003	0.75	[0.59, 0.94]	0.011	0.73	[0.58, 0.93]	0.010
	High early	1.10	[0.93, 1.30]	0.280	1.07	[0.87, 1.31]	0.518	1.15	[0.93, 1.42]	0.199
	Low late	0.88	[0.74, 1.06]	0.173	0.82	[0.67, 1.02]	0.073	0.84	[0.67, 1.05]	0.129
	High late	1.29	[1.10, 1.50]	0.001	1.30	[1.09, 1.56]	0.004	1.29	[1.07, 1.56]	0.009
LGA	Low early	1.23	[1.05, 1.44]	0.010	1.25	[1.05, 1.50]	0.013	1.23	[1.02, 1.49]	0.030
	High early	0.95	[0.80, 1.14]	0.598	0.83	[0.68, 1.03]	0.086	0.72	[0.57, 0.91]	0.005
	Low late	1.22	[1.04, 1.43]	0.014	1.25	[1.04, 1.50]	0.015	1.22	[1.01, 1.48]	0.039
	High late	0.96	[0.81, 1.13]	0.605	0.93	[0.76, 1.12]	0.436	0.99	[0.81, 1.21]	0.915

^a Model 1 adjusted for cohort, maternal smoking and maternal hypertensive disorders.

^b Model 2 adjusted for cohort, maternal smoking, maternal hypertensive disorders, pre-pregnancy BMI, maternal SES, parity and maternal age.

as these have been shown to associate with preterm delivery as well as SGA (reviewed in [19]). As shown in Supplementary Table 2, similar to main analyses, low mHb at late pregnancy was associated with increased risk of spontaneous preterm delivery (aOR 1.65, 95% CI [1.27, 2.15], $p < 0.0005$). High mHb at late pregnancy was associated with increased risk of normotensive full-term SGA in model 1 (OR 1.34, 95% CI [1.08, 1.66], $p = 0.009$) and borderline significant in other models.

Sensitivity analyses

As we do not have the exact gestational age at the last antenatal visit for all mothers, we were not able to estimate the effect of exact mHb measurement timing on the perinatal outcome. This is critical specifically with the case of prematurity, because during normal pregnancy, mHb decreases until 20 weeks of gestation and then begins to increase around 30 weeks of gestation [20]. To confirm that our results are not due to normal mHb change during pregnancy, we conducted sensitivity analyses such as i) MLR of individuals with available information about gestational age at last antenatal visit ($n = 7,281$), ii) MLR of pregnancies with birth at 32 weeks of gestation or more and iii) MLR of gestational age-adjusted late pregnancy mHb. These analyses showed similar results with the main analyses (Supplementary Tables 3–5). Association between prematurity and low mHb at late pregnancy was not statistically significant when gestational age at last antenatal visit was added to the models (Supplementary Table 3). This is most likely due to lack of power because only 45 preterm deliveries had the information about gestational age at last antenatal visit and were included in the gestational age at last antenatal visit-adjusted model.

Comment

The current study indicates that low mHb at late pregnancy is associated with increased risk of preterm delivery and high mHb with increased risk of full-term SGA. These results support the previously published concept of U-shaped association between mHb concentration and the risk of adverse perinatal outcomes [7,21]. However, there are discrepancies between previous publications with respect of this association. One problem related to the generalization of the results from mHb studies concerns the differences between study populations [22]. In low-income countries, the mHb level often relates to the poor nutritional status of the mother contributing to adverse perinatal outcomes. In addition, the prevalence of infectious diseases such as HIV and malaria is higher in the developing countries, which adds further burden to maternal health [5]. This socioeconomic and nutritional disparity may lead to severe residual confounding in the analyses. Current study population is located in Finland where standardized antenatal care system and relatively high living standards induce smaller differences in

environmental factors affecting the development of the fetus and yet still, abnormal mHb was associated with prematurity and SGA.

Another issue varying between studies relates to the timing of the mHb measurement during pregnancy. Level of mHb changes naturally throughout the pregnancy and it may be critical for the results whether the measurement occurs at early or late pregnancy [20]. In the current study, we wanted to evaluate mHb at early and late pregnancy to find out how they contribute to overall risk of adverse perinatal outcomes. Our results indicate that abnormal mHb measured at late pregnancy associates with increased risk of prematurity and SGA. Fetus grows rapidly during the third trimester when iron and other micronutrient demands are specifically high [23]. Abnormal mHb may reflect insufficient availability and/or delivery of nutrients to the fetus and specifically at the end of pregnancy alter the rapid growth phase. It is safe to assume that due to systematic antenatal care in Finland, if low mHb was measured at the beginning of the pregnancy, it had been attempted to increase by administration of iron supplements or other medical actions. Interestingly, low mHb at the beginning of pregnancy was in fact associated with decreased risk of SGA in current study. In high-income countries, approximately 60% of the pregnant women with low hemoglobin are estimated to be amenable to iron supplementation [9]. Thus, the mothers with low mHb at late pregnancy may be the ones that do not respond to iron supplementation and are at increased risk of preterm delivery in the current study. We know that iron supplementation was recommended already in 1960s for pregnant women to prevent gestational anemia [8,24]; however, we do not have information about its administration or iron status of the mother in the current study to research this issue in more detail.

Further variation within mHb studies is added by varying cutoffs for categorical variables of low and high mHb. WHO defines anemia during pregnancy as mHb less than 110 g/l, which is widely used cutoff for low mHb [9]. However, discussion about an appropriate cutoff during pregnancy is ongoing as some studies found association only with mHb levels below 100 g/l [25,26]. Furthermore, similar to previous studies located in Finland and Sweden [22,27], the prevalence of anemia with the 110 g/l cutoff was low in the current study, only approximately 4%. The earlier Finnish study with more stringent cutoff of 100 g/l did not find association between mHb and preterm delivery and SGA, which may result from decreased statistical power due to low number of mothers with “true anemia” [22]. To overcome this issue, we decided to define lowest 10% of the mHb z values as low mHb and highest 10% of the mHb z values as high mHb. These correspond to cutoff values for low mHb of approximately 110 g/l in NFBC1966 and 117 g/l in NFBC1986. For high mHb, cutoff values were 135 g/l and 140 g/l in NFBC1966 and NFBC1986, respectively. Z values were calculated for each cohort and pregnancy time point separately, thus the extremes of mHb are relative for the reference, not for the

absolute mHb concentration that may differ throughout the pregnancy.

The mechanism behind the association is unclear. Iron deficiency may contribute to the restricted fetal growth and prematurity through such mechanisms as increased maternal and fetal stress, increased serum norepinephrine production and oxidative damage to erythrocytes and fetoplacental unit (reviewed in [28]). Mothers with low mHb show increased placental vascularization suggesting earlier placental maturity, which may contribute to increased risk of preterm delivery [29,30]. Low mHb may cause chronic hypoxia which in turn may induce a stress-response with increased placental corticotropin-releasing hormone (CRH) production. During normal pregnancy, a rapid increase in bioavailable CRH induces labor and if this occurs too early, it may induce preterm delivery (28). High mHb, specifically during late pregnancy may indicate incomplete hemodilution and may lead to fetal hypoxia due to increased blood viscosity and impairment of maternal-fetal exchange and contribute to the risk of SGA [1]. Further studies located in high-income countries as well as deeper mechanistic understanding are needed with respect of association between mHb and fetal growth.

Strengths and weaknesses of the study

Large sample size in a homogenous population is major strength of the current study. Universal antenatal care available in Finland and high standard of living offer less variation in the environmental factors affecting the intrauterine development of the child. First limitation is that we do not have reliable information about exact gestational age at mHb measurements from all study subjects in the current study. The measurement timing is specifically important when prematurity is assessed because mHb begins to naturally increase after 30 weeks of gestation. To overcome this, we conducted series of sensitivity analyses that showed similar results as our main models indicating that the association between low mHb at late pregnancy and prematurity is not mediated by the timing of mHb measurement. Other limitations relate to the lack of clinical measures of fetal hypoxia and iron metabolism. Our data suggests that there may be distinct outcomes depending on the mother's responsiveness on iron supplementation. The current study shows that both low and high mHb may indicate increased risk for distress in the developing child. Mechanistic studies are warranted to reveal the possible causal influence of abnormal mHb on perinatal outcomes.

Conflict of interest

The authors report no conflict of interest.

Source of funding

This work was supported by the European Union's Horizon 2020 research and innovation program under grant agreement No. 633595 (DynaHEALTH), and grant agreement No. 733206 (Life-Cycle); the academy of Finland EGEA-project (285547) and the Biocenter Oulu.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.ejogrb.2019.04.045>.

References

- [1] Cordina M, Bhatti S, Fernandez M, Syngelaki A, Nicolaidis KH, Kametas NA. Association between maternal haemoglobin at 27-29 weeks gestation and intrauterine growth restriction. *Pregnancy Hypertens* 2015;5(October(4)):339–45.
- [2] Gonzales GF, Steenland K, Tapia V. Maternal hemoglobin level and fetal outcome at low and high altitudes. *Am J Physiol Regul Integr Comp Physiol* 2009;297(November (5)):R1477–85.
- [3] Jwa SC, Fujiwara T, Yamanobe Y, Kozuka K, Sago H. Changes in maternal hemoglobin during pregnancy and birth outcomes. *BMC Pregnancy Childbirth* 2015;2(April (15)):80 015–0516–1.
- [4] Tandu-Umba B, Mbangama AM. Association of maternal anemia with other risk factors in occurrence of Great obstetrical syndromes at university clinics, Kinshasa, DR Congo. *BMC Pregnancy Childbirth* 2015;21(August (15)):183 015–0623–z.
- [5] Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, et al. Maternal anemia and risk of adverse birth and health outcomes in low- and middle-income countries: systematic review and meta-analysis. *Am J Clin Nutr* 2016;103(February (2)):495–504.
- [6] Levy A, Fraser D, Katz M, Mazor M, Sheiner E. Maternal anemia during pregnancy is an independent risk factor for low birthweight and preterm delivery. *Eur J Obstet Gynecol Reprod Biol* 2005;122(October (2)):182–6.
- [7] Scanlon KS, Yip R, Schieve LA, Cogswell ME. High and low hemoglobin levels during pregnancy: differential risks for preterm birth and small for gestational age. *Obstet Gynecol* 2000;96(November (5 Pt 1)):741–8.
- [8] Fararouei M, Robertson C, Whittaker J, Sovio U, Ruokonen A, Pouta A, et al. Maternal Hb during pregnancy and offspring's educational achievement: a prospective cohort study over 30 years. *Br J Nutr* 2010;104(November (9)):1363–8.
- [9] WHO. The global prevalence of anaemia in 2011. Geneva: World Health Organization; 2015.
- [10] Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. *Lancet Glob Health* 2013;1(July (1)):e16–25.
- [11] Salihu HM, Wilson RE. Epidemiology of prenatal smoking and perinatal outcomes. *Early Hum Dev* 2007;83(November (11)):713–20.
- [12] Milman N, Pedersen AN. Blood haemoglobin concentrations are higher in smokers and heavy alcohol consumers than in non-smokers and abstainers: should we adjust the reference range? *Ann Hematol* 2009;88(July (7)):687–94.
- [13] Hendler I, Goldenberg RL, Mercer BM, Iams JD, Meis PJ, Moawad AH, et al. The Preterm Prediction Study: association between maternal body mass index and spontaneous and indicated preterm birth. *Am J Obstet Gynecol* 2005;192(March (3)):882–6.
- [14] Ruiz M, Goldblatt P, Morrison J, Kukla L, Svancara J, Riitta-Jarvelin M, et al. Mother's education and the risk of preterm and small for gestational age birth: a DRIVERS meta-analysis of 12 European cohorts. *J Epidemiol Community Health* 2015;69(September (9)):826–33.
- [15] Kozuki N, Lee AC, Silveira MF, Sania A, Vogel JP, Adair L, et al. The associations of parity and maternal age with small-for-gestational-age, preterm, and neonatal and infant mortality: a meta-analysis. *BMC Public Health* 2013;13(Suppl. 3):S2 2458–13–S3–S2. Epub 2013 Sep 17.
- [16] Pihkala J, Hakala T, Voutilainen P, Raivio K. Characteristic of recent fetal growth curves in Finland. *Duodecim* 1989;105(18):1540–6.
- [17] Mannisto T, Karumanchi SA, Pouta A, Vaarasmaki M, Mendola P, Miettola S, et al. Preeclampsia, gestational hypertension and subsequent hypothyroidism. *Pregnancy Hypertens* 2013;3(January (1)):21–7.
- [18] Team RC. R: a language and environment for statistical computing. 2018.
- [19] Vest AR, Cho LS. Hypertension in pregnancy. *Curr Atheroscler Rep* 2014;16(March (3)):395 013–0395–8.
- [20] Steer PJ. Maternal hemoglobin concentration and birth weight. *Am J Clin Nutr* 2000;71(May (Suppl. 5)):1285S–7S.
- [21] Dewey KG, Oaks BM. U-shaped curve for risk associated with maternal hemoglobin, iron status, or iron supplementation. *Am J Clin Nutr* 2017;(October):25.
- [22] Hamalainen H, Hakkarainen K, Heinonen S. Anaemia in the first but not in the second or third trimester is a risk factor for low birth weight. *Clin Nutr* 2003;22(June (3)):271–5.
- [23] Kumar KJ, Asha N, Murthy DS, Sujatha M, Manjunath V. Maternal anemia in various trimesters and its effect on newborn weight and maturity: an observational study. *Int J Prev Med* 2013;4(February (2)):193–9.
- [24] Nevanlinna HR. Therapy of anemia. *Duodecim* 1965;81:141–5.
- [25] Kozuki N, Lee AC, Katz J, Child Health Epidemiology Reference Group. Moderate to severe, but not mild, maternal anemia is associated with increased risk of small-for-gestational-age outcomes. *J Nutr* 2012;142(February (2)):358–62.
- [26] Gonzales GF, Tapia V, Gasco M, Carrillo CE. Maternal hemoglobin concentration and adverse pregnancy outcomes at low and moderate altitudes in Peru. *J Matern Fetal Neonatal Med* 2012;25(July (7)):1105–10.
- [27] Stephansson O, Dickman PW, Johansson A, Cnattingius S. Maternal hemoglobin concentration during pregnancy and risk of stillbirth. *JAMA* 2000;284(November (20)):2611–7.
- [28] Allen LH. Biological mechanisms that might underlie iron's effects on fetal growth and preterm birth. *J Nutr* 2001;131(February (2S-2)):581S–9S.
- [29] Lelic M, Bogdanovic G, Ramic S, Brkicevic E. Influence of maternal anemia during pregnancy on placenta and newborns. *Med Arch* 2014;68(June (3)):184–7.
- [30] Stangret A, Wnuk A, Szweczyk G, Pyzlak M, Szukiewicz D. Maternal hemoglobin concentration and hematocrit values may affect fetal development by influencing placental angiogenesis. *J Matern Fetal Neonatal Med* 2017;30(January (2)):199–204.