



ORIGINAL ARTICLE

Diastolic function in young patients with cryptogenic stroke: A case-control pilot study

Jani Pirinen¹ | Jouni Kuusisto² | Vesa Järvinen¹ | Nicolas Martinez-Majander³ |
Juha Sinisalo² | Pauli Pöyhönen² | Jukka Putaala³

¹HUS Medical Imaging Center, Clinical Physiology and Nuclear Medicine, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

²Heart and Lung Center, Helsinki University Hospital and Helsinki University, Helsinki, Finland

³Clinical Neurosciences, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

Correspondence

Jani Pirinen, HUS Medical Imaging Center, Clinical Physiology and Nuclear Medicine, University of Helsinki and Helsinki University Hospital, Finland, Haartmaninkatu 4, FIN-00029, Helsinki, Finland.

Email: jani.pirinen@helsinki.fi

Funding information

Finska Läkaresällskapet; Helsingin ja Uudenmaan Sairaanhoidopiiri; Maire Taposen Säätiö; Aarne Koskelon Säätiö; Academy of Finland; Stiftelsen Dorothea Olivia, Karl Walter och Jarl Walter Perkléns Minne

Abstract

Background: Ischaemic stroke in young individuals often remains cryptogenic. In this pilot study, we investigated, whether advanced echocardiography methods could find differences in the diastolic function between young cryptogenic stroke patients and stroke-free controls.

Methods: We recruited 30 cryptogenic ischaemic stroke patients aged 18–49 and 30 age- and sex-matched stroke-free controls among participants of the Searching for Explanations for Cryptogenic Stroke in the Young: Revealing the Etiology, Triggers, and Outcome (SECRETO) study (NCT01934725). We measured diastolic function parameters derived from speckle tracking strain rate, Doppler techniques and 4D volumetry. We also performed statistical analyses comparing only the highest and lowest tertile of cases and controls for each parameter.

Results: None of our patients or controls had diastolic dysfunction according to ASE/EACVI criteria. However, compared to stroke-free controls, the stroke patient group had lower E/A ratio of mitral inflow, lower lateral and mean e', lower A/a' ratio, lower strain rate in early diastole and lower speckle tracking-derived e/a ratio. When comparing the lowest tertiles, patients also had a lower peak filling rate by 4D volumetry, a lower peak early filling fraction (fraction of left ventricular filling during early diastole), and lower velocities in a series of the tissue Doppler-derived diastolic parameters and blood flow/tissue velocity ratios.

Conclusion: Our study displayed subtle differences in diastolic function between patients and stroke-free controls, which may play a role in early-onset cryptogenic stroke. The differences were clearer when the lowest tertiles were compared, suggesting that there is a subgroup of young cryptogenic stroke patients with subclinical heart disease.

KEYWORDS

4D volumetry, brain infarction, echocardiography, strain imaging, strain rate imaging

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Clinical Physiology and Functional Imaging* published by John Wiley & Sons Ltd on behalf of Scandinavian

1 | INTRODUCTION

Ischaemic stroke can occur due to cardioembolism. Some cardiac diseases are categorized as high-risk sources of cardioembolism, due to an over 2% annual ischaemic stroke risk (Ay et al., 2005). Part of ischaemic strokes remains cryptogenic, that is with undetermined aetiology despite intensive diagnostic investigations. Stroke at younger ages particularly often remains cryptogenic (Putala et al., 2012). A portion of these strokes are probably of cardiac origin, although how large this portion is, is unknown.

Cardiac findings in particularly young (below 50 years) cryptogenic stroke patients have been described very scarcely in the literature. In older patients, altered diastolic function is associated with a higher risk of left atrial appendix thrombosis in patients with atrial fibrillation (AF), and diastolic dysfunction is associated with worse outcome after ischaemic stroke (Doukky et al., 2016; Garshick et al., 2018; Park et al., 2016).

We sought to evaluate, whether advanced echocardiography methods can detect differences in left ventricular diastolic function between young cryptogenic stroke patients and stroke-free controls, and hence detect whether diastolic function could play a role in cryptogenic stroke in the young.

2 | MATERIALS AND METHODS

2.1 | Study population

Participants in this study were recruited among those enrolled into the Searching for Explanations for Cryptogenic Stroke in the Young: Revealing the Etiology, Triggers, and Outcome study (SECRETO, NCT01934725), which is an international prospective multicentre case-control study of young adults (age 18–49) with a first-ever ischaemic stroke of undetermined aetiology. The study protocol has earlier been published in more detail (Putala et al., 2017). Written informed consent was obtained from all study participants. SECRETO has been approved by the Ethics Committee of Helsinki and Uusimaa Hospital District.

Patients were included after standardized diagnostic procedures, including brain magnetic resonance imaging, imaging of intracranial and extracranial arteries with either computed tomography angiography or magnetic resonance angiography, and cardiac imaging to rule out established causes of ischaemic stroke. Cardiac imaging included standardized transthoracic and transesophageal echocardiography (Saeed, Gerdtts, Waje-Andreassen, Sinisalo, & Putala, 2019). We ruled out AF with a \geq 24-hr Holter monitoring. Patients were classified according to embolic stroke of undetermined source (ESUS) criteria as ESUS (+) and ESUS (–) (Hart et al., 2014). Body surface area (BSA) was calculated using the Mosteller formula (Mosteller, 1987). Patients included in the present study suffered their stroke between December 2013 and May 2017.

Patients were age- and sex-matched to stroke-free controls in a 1:1 fashion. Controls were searched randomly through population

registers, and in some cases patients' nonrelated proxies. Detailed clinical history was recorded from all study subjects, including arterial blood pressure, height, weight, alcohol consumption and the presence of right-to-left shunt. The definition of right-to-left shunt was a positive finding (any degree of shunt) on either transesophageal echocardiography or transcranial Doppler bubble test; each patient and control underwent at least one of these tests. Participants were dichotomized according to their right-to-left shunt status. Patients with prior PFO closure were excluded from the study.

Detailed baseline data and clinical data and basic left ventricular (LV) measurements have been published earlier (Pirinen et al. 2020). Compared with controls, patients were heavier, had a larger waist circumference, higher body-mass index (BMI) and a larger BSA. Patients also had a higher prevalence of right-to-left shunt. We found no significant differences between patients and controls regarding established cardiovascular risk factors. There were no differences between the two groups in basic LV diameter, LV wall thickness or LV ejection fraction.

2.2 | Echocardiography methods

We performed advanced echocardiography on our study subjects between November 2017 and October 2018. The same echocardiographer (J.Pi) examined all patients and control subjects blinded to the case-control status and clinical data, with a General Electric Vivid E9 version 113 cardiac ultrasound device, using M5Sc and 4V probes (General Electric).

Measurements of mitral inflow E wave and A wave were obtained from the apical 4-chamber view, as were e' and a' according to international standards (Nagueh et al., 2016). We calculated E/e' , A/a' and E/a' for all combinations of septal, lateral and mean measurements. The diastolic function was evaluated according to ASE/EACVI criteria (Nagueh et al., 2016).

Left ventricular strain rate analysis was derived from the apical 4-chamber, 2-chamber and 3-chamber views, using speckle tracking imaging. Only the longitudinal component of strain rate was analysed. A true apical angle was verified using the triplane mode, with which the echocardiographer can confirm that both the apex and the mitral valve orifice are in the midline of the ultrasound beam in all three planes simultaneously. The apical 4-chamber view was defined as a view with centre in LV apex and the mitral orifice, viewing both ventricles and both atria, with a rotation angle maximizing the area of the right ventricle (RV). The 2-chamber view was defined as a counter-clockwise virtual rotation of 60° from the 4-chamber view, hence viewing the LV and the left atrium (LA). The 3-chamber view was defined as a further counter-clockwise virtual rotation of 60°, hence viewing the LV, the ascending aorta and the LA. The typical frame rate used for this imaging was 60–80 fps. Strain rate was measured using speckle tracking from 2D images. Global strain rate was defined as a mean of the entire LV strain rate in the three apical views. Strain rate

was measured in two different time points: in early diastole and in late diastole (during atrial contraction), using EchoPAC version 113 (General Electric) (Figure 1).

Left ventricular volume data points with respective time stamps over whole cardiac cycle were exported from Echopac software to sheet data. From these data, volume change rate was calculated: volume change between all successive volume

points was calculated and divided by sampling cycle length $dV_n/dt_n = (V_n + 1 - V_n)/(t_n + 1 - t_n)$. For this average, time between these two data points was used (t_m). Volume change rate curves were then generated (Figure 2). Early positive peak represented peak early filling rate, as earlier described by Hieda, and latter positive peak late peak filling rate (Hieda et al., 2018). Time to peak early filling rate was determined both in milliseconds and relative time




FIGURE 1 Left ventricular strain rate analysis in 4-chamber view (upper), 2-chamber view (middle) and 3-chamber view (lower). The early diastolic maximum strain rates of this control person in the three views were 2.36/s, 1.94/s and 1.93/s. The late diastolic peak strain rates were 0.59/s, 0.55/s and 0.58/s, respectively. Hence, the global early diastolic strain rate was 2.08/s, global late diastolic strain rate was 0.57/s, and the speckle tracking e/a ratio 3.65. The yellow lines mark the end of diastole, and the green line marks aortic valve closure, and hence the end of systole




FIGURE 2 Analysis of left ventricular volume dynamics. (A) Volume and time data exported from EchoPac 4D LVQ software and plotted as volume–time curve. From these data, volume change rate is calculated and displayed as (B) volume change rate–time, and (C) volume change rate–volume curves. In figure B, early positive peak represents early peak filling rate. In figure C, the red arrow depicts ventricular systole (from a to b), the yellow arrow early systole (b to c) and the green arrow late diastole during atrial contraction (c to a). Point a represents end-diastolic volume, b end-systolic volume, and c mid-diastolic volume after early filling

of the whole cardiac cycle. Volume change rate–volume curves were also generated. These curves were used to detect LV volumes after early filling to calculate early filling fraction.

2.3 | Statistical methods

We used the Wilcoxon signed rank test for all continuous parameters. For dichotomous clinical parameters, we used the McNemar test. Tertile analyses were performed using Wilcoxon signed rank test, selecting only the minimum or maximum tertile of case and

control measurements of each parameter, and comparing these. All analyses used IBM SPSS 22 (IBM), or RStudio 1.2.1335 (RStudio).

3 | RESULTS

3.1 | Performance of measurements

Heart rate during echocardiography varied between 39/min and 87/min in our subjects. LV 4D volumetry, to a degree acceptable for flow analysis, was successful in 51 of 60 subjects, constituting 21 case–control pairs. Mitral inflow E and A, and the measurement of septal and lateral e' and a', could be obtained in all subjects. Speckle tracking strain rate analysis was successful in 27 case–control pairs.

3.2 | Analysis of LV diastolic function

None of the patients or controls had evident diastolic dysfunction according to American Society of Echocardiography and European Association of Cardiovascular Imaging (ASE/EACVI) criteria. Controls had higher mitral inflow E/A ratio than patients, and larger mean e'. In strain rate analysis, both early diastolic strain rate (e), and the ratio between early diastolic and late diastolic strain rate (e/a), were higher in controls than in patients (Table 1).

3.3 | Minimum and maximum tertile analysis of diastolic parameters

In minimum tertile analysis, stroke patients had a lower E/A ratio, lower E wave velocity, lower septal e' velocity, lower mean e', lower e'/a' ratio regardless of measurement point, lower septal A/a' ratio, and also lower peak early filling rate, time to peak filling rate and early filling fraction (Table 2). In maximum tertile analysis, controls had a higher E/A ratio, higher septal e', whereas stroke patients had a higher lateral a', and a higher peak filling rate in atrial systole (Table 3).

4 | DISCUSSION

To the best of our knowledge, this is the first study to investigate LV diastolic function in young cryptogenic stroke patients. Our observations support the hypothesis that diastolic function has a role in the pathogenesis of ischaemic stroke. However, the relationship between diastolic parameters and stroke has been only scarcely studied to date, with only a few reports showing positive associations, mostly in the presence of AF (Doukky et al., 2016; Garshick et al., 2018; Park et al., 2016).

Our main result is that even without actual diastolic dysfunction, there seems to be an association between poorer values of diastolic function parameters and ischaemic stroke, since the stroke patient group had a lower E/A ratio, lower mean e' velocity, and lower peak

TABLE 1 Comparison of echocardiographic findings between stroke patients and controls

Mitral inflow measurements	Patients	Controls	p-value
E wave maximum velocity, m/s	0.8 (0.3)	0.7 (0.2)	.424
E wave deceleration time, ms	160 (40)	160 (40)	.896
A wave maximum velocity, m/s	0.6 (0.2)	0.5 (0.2)	.081
E/A ratio	1.4 (0.8)	1.5 (1.1)	.039
Mitral annulus tissue Doppler imaging			
Septal e', cm/s	11.0 (3.0)	11.5 (4.0)	.074
Lateral e', cm/s	13.0 (5.0)	14.0 (5.0)	.039
Mean e', cm/s	12.0 (2.7)	13.3 (4.3)	.016
Septal a', cm/s	8.0 (3.0)	8.0 (2.0)	.680
Lateral a', cm/s	7.0 (4.0)	8.0 (2.0)	.830
Mean a', cm/s	9.5 (3.4)	9.8 (2.0)	.206
Septal e'/a'	1.3 (0.8)	1.4 (0.7)	.247
Lateral e'/a'	1.9 (1.1)	1.9 (1.0)	.136
Mean e'/a'	1.6 (1.0)	1.7 (0.8)	.102
Mitral inflow/annulus tissue Doppler ratios			
Septal E/e'	6.9 (2.1)	6.3 (2.3)	.737
Lateral E/e'	5.8 (2.0)	5.4 (2.0)	.399
Mean E/e'	6.2 (2.0)	5.8 (2.2)	.341
Septal A/a'	7.0 (3.8)	6.0 (3.3)	.073
Lateral A/a'	7.4 (2.8)	6.3 (2.6)	.094
Mean A/a'	6.8 (2.7)	6.2 (2.9)	.047
Left ventricular strain rate analysis			
Peak e strain rate, 1/s ²	1.4 (0.4)	1.5 (0.5)	.005
Peak a strain rate, 1/s ²	0.7 (0.3)	0.7 (0.2)	.755
Peak strain rate e/a	2.1 (1.4)	2.2 (1.0)	.013
Left ventricular 4D volumetry			
Peak early filling rate, ml/s	277 (148)	303 (108)	.289
Time to peak early filling rate, ms	79 (50)	88 (39)	.322
% time to peak early filling rate	89 (70)	95 (63)	.339
Peak filling rate in atrial systole, ml/s	225 (128)	207 (66)	.076
Early filling fraction, %	71 (16)	70 (8)	.658

Note: Numbers are median (interquartile range).

early diastolic strain rate and strain rate e/a ratio. The minimum tertile analysis findings of patients with lower maximum E wave velocity, higher A wave velocity, lower septal e', lower a'/e', higher A/a', lower peak early filling rate and lower early filling fraction are also suggestive that there is a subgroup of young stroke patients with poorer diastolic function than in their stroke-free counterparts. Also, the maximum tertile analysis, demonstrating higher lateral a' and peak filling rate in atrial systole in this subgroup of patients, suggests pronounced atrial contraction, possibly to compensate weakness in early filling.

Earlier, diastolic dysfunction measured by E/e' ratio has been linked to higher risk of thrombosis in patients with AF (Doukky et al., 2016; Garshick et al., 2018). Also, poorer diastolic function assessed by higher E/e' is associated with higher risk of stroke

TABLE 2 Lowest tertile analysis of echocardiographic parameters between stroke patients and controls

Mitral inflow measurements	Patients	Controls	p
E wave maximum velocity, m/s	0.5 (0.2)	0.6 (0.1)	.005
E wave deceleration time, ms	130 (20)	130 (20)	.907
A wave maximum velocity, m/s	0.4 (0.1)	0.3 (0.11)	.040
E/A ratio	0.9 (0.1)	1.1 (0.1)	.006
Mitral annulus tissue Doppler imaging			
Septal e', cm/s	8 (1)	9.5 (1)	.015
Lateral e', cm/s	10.5 (1.8)	11.5 (2.5)	.097
Mean e', cm/s	9.5 (1.5)	10 (0.9)	.031
Septal a', cm/s	6.5 (1.8)	7 (1)	.587
Lateral a', cm/s	6 (0.8)	6 (2.5)	.968
Mean a', cm/s	8 (0.8)	8.8 (0.5)	.039
Septal e'/a'	0.8 (0.14)	1.0 (0.1)	.005
Lateral e'/a'	1.0 (0.2)	1.3 (0.4)	.015
Mean e'/a'	1.0 (0.1)	1.3 (0.2)	<.001
Mitral inflow/annulus tissue Doppler ratios			
Septal E/e'	5.4 (1.2)	5.4 (0.5)	.791
Lateral E/e'	4.4 (0.6)	4.3 (0.6)	.271
Mean E/e'	4.9 (0.7)	4.6 (0.5)	.272
Septal A/a'	4.8 (0.5)	4.3 (0.5)	.009
Lateral A/a'	5.3 (0.9)	4.5 (1.2)	.130
Mean A/a'	5.2 (0.8)	4.5 (0.9)	.028
Left ventricular strain rate analysis			
Peak e strain rate, 1/s ²	0.8 (0.3)	1.1 (0.3)	.028
Peak a strain rate, 1/s ²	0.5 (0.2)	0.5 (0.2)	.448
Peak strain rate e/a	1.1 (0.3)	1.5 (0.7)	.043
Left ventricular 4D volumetry			
Peak early filling rate, ml/s	223.5 (63.3)	273.2 (51.8)	.029
Time to peak early filling rate, ms	56 (8)	73 (22)	.014
% time to peak early filling rate	53 (19)	65 (19)	.058
Peak filling rate in atrial systole, ml/s	179.4 (56.6)	143.6 (37.0)	.105
Early filling fraction, %	59.9 (14.8)	66.7 (3.1)	.004

Note: Numbers are median (interquartile range)

recurrence, suggesting that diastolic function is associated with the mechanisms of intracardiac thrombosis (Park et al., 2016). However, we did not find significant differences in the E/e' ratio between patients and controls, although we found differences in several other diastolic parameters. One reason for that may be that E/e' is a quite specific marker for diastolic dysfunction and elevated LV filling pressure, but lacks sensitivity (Sharifov et al., 2016). Furthermore, since all patients and controls had normal diastolic function according to ASE/EACVI criteria and normal E/e' values, the association between E/e' and LV filling pressure may not be accurate.

TABLE 3 Highest tertile analysis of echocardiographic parameters between stroke patients and controls

Mitral inflow measurements	Patients	Controls	p
E wave maximum velocity, m/s	1.0 (0.1)	0.9 (0.2)	.427
E wave deceleration time, ms	190 (15)	185 (28)	1.000
A wave maximum velocity, m/s	0.7 (0.1)	0.6 (0.2)	.139
E/A ratio	1.8 (0.3)	2.3 (0.4)	.003
Mitral annulus tissue Doppler imaging			
Septal e', cm/s	13.0 (12)	14.5 (1.0)	.040
Lateral e', cm/s	16.0 (2.8)	18.0 (1.8)	.079
Mean e', cm/s	14.5 (2.5)	15.8 (1.6)	.087
Septal a', cm/s	10.5 (1.8)	10.0 (0.8)	.163
Lateral a', cm/s	11.0 (3.0)	9.0 (2.0)	.046
Mean a', cm/s	11.0 (0.5)	11.3 (0.9)	.086
Septal e'/a'	2.0 (0.7)	2.1 (0.7)	.520
Lateral e'/a'	2.5 (0.8)	2.7 (0.3)	.343
Mean e'/a'	2.1 (0.6)	2.4 (0.7)	.496
Mitral inflow/annulus tissue Doppler ratios			
Septal E/e'	8.9 (1.6)	8.6 (1.2)	.423
Lateral E/e'	6.8 (1.4)	6.7 (1.1)	.239
Mean E/e'	7.8 (0.9)	7.3 (1.0)	.384
Septal A/a'	9.3 (1.2)	8.0 (1.9)	.064
Lateral A/a'	9.5 (1.5)	8.5 (1.5)	.058
Mean A/a'	9.5 (1.4)	8.3 (0.8)	.121
Left ventricular strain rate analysis			
Peak e strain rate, 1/s ²	1.6 (0.3)	1.8 (0.2)	.075
Peak a strain rate, 1/s ²	0.9 (0.2)	0.8 (0.1)	.069
Peak strain rate e/a	2.8 (0.7)	3.3 (1.9)	.272
Left ventricular 4D volumetry			
Peak filling rate, ml/s	405.1 (81.7)	441.4 (79.7)	.529
Time to peak early filling rate, ms	110 (57)	132 (36)	.623
% time to peak early filling rate	136 (45)	141 (73)	.597
Peak filling rate in atrial systole, ml/s	317.1 (46.6)	227.2 (29.7)	.002
Early filling fraction, %	75.9 (5.0)	76.9 (6.2)	.529

Note: Numbers are median (interquartile range).

In our study, the diastolic strain rate e and strain rate e/a ratio were significantly higher in controls than patients, which suggests that the patients have stiffer LV than the controls. Our method of examining LV diastolic function with speckle tracking SR was earlier used to identify increased risk of AF among general ischaemic stroke patients (Skaarup et al., 2017). In that study, early diastolic SR was the only LV speckle tracking parameter associated with AF in ischaemic stroke patients, although they did not use our method of calculating the SR-derived e/a ratio, which was also lower in our patients than in controls.

Our patients had a higher prevalence of right-to-left shunts than the controls, which might be a confounding factor regarding stroke

aetiology. However, the effect of a right-to-left shunt on diastolic function has been studied very scarcely, although one small study suggests that PFO closure, and hence the elimination of right-to-left shunt, has no effect on the diastolic E/A ratio or e' (Yalonetsky, Schwartz, & Lorber, 2007). This finding supports the theory of diastolic function being an entity independent of right-to-left shunt, and thus might play an independent role in thrombogenesis in our patients.

Based on our data, we can only hypothesize the possible underlying pathways explaining the link between diastolic dysfunction and ischaemic stroke. As our patients were more obese compared with controls, one possible explanation includes the contribution of obesity with diastolic dysfunction, hypercoagulation and occult AF.

Strengths of our study include a systematic and well-matched case-control population, a prospective design, a precise protocol with very few missing data, blinding of the echocardiographer, and the use of modern echocardiography methods such as 4D volumetry and strain rate analysis. Limitations include a relatively small sample size, which was justified with the imaging protocol being very demanding, and the purpose of the study to create new hypotheses to be tested more widely in the future. Other limitations were shortcomings inherent to case-control studies, such as selection bias.

5 | CONCLUSIONS

Young cryptogenic ischaemic stroke patients tend to have normal LV diastolic function. Aside from actual diastolic dysfunction, our pilot study found subtle differences in the apparently normal diastolic function between young cryptogenic stroke patients and stroke-free controls. Diastolic function may thus be one of the players in cryptogenic ischaemic stroke at young age by increasing susceptibility for intracardiac thrombosis through slower blood flow. Future studies should enrol larger patient populations to verify our inaugural observations.

ACKNOWLEDGMENTS

We thank our funders: Academy of Finland, Helsinki and Uusimaa Hospital District, Finska läkaresällskapet, Maire Taposen säätiö, Aarne Koskelon säätiö, and Stiftelsen Dorothea Olivia, Karl Walter och Jarl Walter Perkléns Minne. We also wish to thank Laura-Leena Kupari, RN, for all her efforts during this study.

CONFLICTS OF INTEREST

None.

ORCID

Jani Pirinen  <https://orcid.org/0000-0002-4030-9358>

Jouni Kuusisto  <https://orcid.org/0000-0003-2221-2621>

Nicolas Martinez-Majander  <https://orcid.org/0000-0001-8489-7051>

Juha Sinisalo  <https://orcid.org/0000-0002-0169-5137>

Pauli Pöyhönen  <https://orcid.org/0000-0002-3031-5391>

Jukka Putaala  <https://orcid.org/0000-0002-6630-6104>

REFERENCES

- Ay, H., Furie, K. L., Singhal, A., Smith, W. S., Sorensen, A. G., & Koroshetz, W. J. (2005). An evidence-based causative classification system for acute ischemic stroke. *Annals of Neurology*, *58*, 688–697. <https://doi.org/10.1002/ana.20617>
- Doukky, R., Garcia-Sayan, E., Patel, M., Pant, R., Wassouf, M., Shah, S., ... Kehoe, R. F. (2016). Impact of Diastolic Function Parameters on the Risk for Left Atrial Appendage Thrombus in Patients with Nonvalvular Atrial Fibrillation: A Prospective Study. *Journal of the American Society of Echocardiography*, *29*, 545–553. <https://doi.org/10.1016/j.echo.2016.01.014>
- Garshick, M. S., Mulliken, J., Schoenfeld, M., Riedy, K., Guo, Y., Zhong, J., ... Skolnick, A. H. (2018). Average e' velocity on transthoracic echocardiogram is a novel predictor of left atrial appendage sludge or thrombus in patients with atrial fibrillation. *Echocardiography*, *35*, 1939–1946. <https://doi.org/10.1111/echo.14148>
- Hart, R. G., Diener, H. C., Coutts, S. B., Easton, J. D., Granger, C. B., O'Donnell, M. J., ... Connolly, S. J. (2014). Cryptogenic Stroke/ESUS International Working Group. Embolic strokes of undetermined source: The case for a new clinical construct. *The Lancet Neurology*, *13*, 429–438. [https://doi.org/10.1016/S1474-4422\(13\)70310-7](https://doi.org/10.1016/S1474-4422(13)70310-7)
- Hieda, M., Parker, J., Rajabi, T., Fujimoto, N., Bhella, P. S., Prasad, A., ... Levine, B. D. (2018). Left ventricular volume-time relation in patients with heart failure with preserved ejection fraction. *American Journal of Cardiology*, *121*, 609–614. <https://doi.org/10.1016/j.amjcard.2017.11.033>
- Mosteller, R. D. (1987). Simplified calculation of body-surface area. *New England Journal of Medicine*, *317*, 1098. <https://doi.org/10.1056/NEJM198710223171717>
- Nagueh, S. F., Smiseth, O. A., Appleton, C. P., Byrd, B. F., Dokainish, H., Edvardsen, T., ... Waggoner, A. D. (2016). Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *European Heart Journal of Cardiovascular Imaging*, *17*, 1321–1360. <https://doi.org/10.1093/ehjci/jew082>
- Park, H. K., Kim, B. J., Yoon, C. H., Yang, M. H., Han, M. K., & Bae, H. J. (2016). Left ventricular diastolic dysfunction in ischemic stroke: functional and vascular outcomes. *J Stroke*, *18*, 195–202. <https://doi.org/10.5853/jos.2015.01697>
- Pirinen, J., Järvinen, V., Martinez-Majander, N., Sinisalo, J., Pöyhönen, P., & Putaala, J. (2020). Left atrial dynamics is altered in young adults with cryptogenic ischemic stroke: a case-control study utilizing advanced echocardiography. *Journal of the American Heart Association*, *7*(9), e014578. <https://doi.org/10.1161/JAHA.119.014578>
- Putaala, J., Martinez-Majander, N., Saeed, S., Yesilot, N., Jäkälä, P., Nerg, O., ... Tatlisumak, T. (2017). Searching for explanations for cryptogenic stroke in the young: revealing the triggers, causes, and outcome (SECRETO): Rationale and design. *Eur Stroke J*, *2*, 116–125. <https://doi.org/10.1177/2396987317703210>
- Saeed, S., Gerdtts, E., Waje-Andreassen, U., Sinisalo, J., & Putaala, J. (2019). Searching for explanations for cryptogenic stroke in the young: revealing the etiology, triggers, and outcome (SECRETO): Echocardiography performance protocol. *Echo Res Pract*, *6*, 53–61. <https://doi.org/10.1530/ERP-19-0025>
- Sharifov, O. F., Schiros, C. G., Aban, I., Denney, T. S., & Gupta, H. (2016). Diagnostic accuracy of tissue doppler index E/e' for evaluating left ventricular filling pressure and diastolic dysfunction/heart failure with preserved ejection fraction: A systematic review and meta-analysis. *J Am Heart Assoc*, *5*(1), e002530. <https://doi.org/10.1161/JAHA.115.002530>
- Skaarup, K. G., Christensen, H., Høst, N., Mahmoud, M. M., Ovesen, C., Olsen, F. J., ... Biering-Sørensen, T. (2017). Usefulness of left ventricular speckle tracking echocardiography and novel measures of left atrial structure and function in diagnosing paroxysmal atrial fibrillation in ischemic stroke and transient ischemic attack patients. *International Journal of Cardiovascular Imaging*, *33*, 1921–1929. <https://doi.org/10.1007/s10554-017-1204-1>
- Yalonetsky, S., Schwartz, Y., & Lorber, A. (2007). Analysis of left and right ventricular Doppler tissue imaging in patients undergoing percutaneous closure of patent foramen ovale. *J Invasive Cardiol*, *19*, 252–254.
- Yesilot Barlas, N., Putaala, J., Waje-Andreassen, U., Vassilopoulou, S., Nardi, K., Odier, C., ... Bahar, S. Z. (2012). Etiology of first-ever ischemic stroke in European young adults: The 15 cities young stroke study. *Stroke*, *43*, 2624–2630. <https://doi.org/10.1111/ene.12228>

How to cite this article: Pirinen J, Kuusisto J, Järvinen V, et al. Diastolic function in young patients with cryptogenic stroke: A case-control pilot study. *Clin Physiol Funct Imaging*. 2020;40:336–342. <https://doi.org/10.1111/cpf.12640>