

<https://helda.helsinki.fi>

First record of a Nathusius' pipistrelle (*Pipistrellus nathusii*) overwintering at a latitude above 60°N

Blomberg, Anna S.

2021-01

Blomberg , A S , Vasko , V , Salonen , S , Petersons , G & Lilley , T M 2021 , ' First record of a Nathusius' pipistrelle (*Pipistrellus nathusii*) overwintering at a latitude above 60°N ' , *Mammalia : international journal of the systematics, biology and ecology of mammals* , vol. 85 , no. 1 , pp. 74-78 . <https://doi.org/10.1515/mammalia-2020-0019>

<http://hdl.handle.net/10138/325873>

<https://doi.org/10.1515/mammalia-2020-0019>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Short note

Anna S. Blomberg*, Ville Vasko, Saku Salonen, Gunārs Pētersons and Thomas M. Lilley

First record of a *Nathusius'* pipistrelle (*Pipistrellus nathusii*) overwintering at a latitude above 60°N

<https://doi.org/10.1515/mammalia-2020-0019>

Received February 24, 2020; accepted June 12, 2020; published online August 19, 2020

Abstract: Highly mobile species are considered to be the first to respond to climate change by transforming their ranges of distribution. There is evidence suggesting that *Pipistrellus nathusii*, a species capable of long-distance migration, is expanding both its reproduction and overwintering ranges to the North. We recorded the echolocation calls of bats at 16 sites in South-Western Finland on two consecutive winters, and detected calls of *P. nathusii* at one of the sites throughout the second winter. To our knowledge, this is the northernmost record of an overwintering *P. nathusii*, and contributes to evidence that the species is already responding to climate change.

Keywords: chiroptera; climate change; hibernation; migration; range expansion.

Climate change is already affecting the range limits and phenology of organisms across a plethora of taxa (Hickling et al. 2006; Parmesan and Yohe 2003; Tingley et al. 2009). According to a review by Thomas (2010), it is possible that more than half of observed animal range boundaries have shown a response to climate change. In most cases, range expansion has occurred on the boundary with lower ambient temperatures, which appears to take place more rapidly than local extinctions at the boundary with higher ambient temperatures (Brommer et al. 2012; Hickling et al. 2006; Thomas et al. 2006). Highly mobile species, such as bats, are most likely to be the first to respond to a changing climate by shifting their ranges towards the poles or to

higher latitudes (Brommer et al. 2012; Warren et al. 2001). While it is difficult to differentiate changes caused by climate change from those resulting from habitat loss or other local changes, or an increase in sampling effort, there is some evidence that bats are also already responding to the changing climate through range expansion (Lundy et al. 2010; Sherwin et al. 2013).

Winter is often a critical time for animals inhabiting high latitudes, and the overwintering strategies of organisms have received considerable attention with regards to climate change and associated range shifts (Humphries et al. 2002; Maclean et al. 2008; Sorte and Iii 2007). For many relatively sedentary species, such as most hibernating mammals, climate change can have unfavorable effects (Lane et al. 2012). However, bats (order Chiroptera), as an example of hibernating mammals, are mobile and more capable of shifting their overwintering ranges as a response. Indeed, Humphries et al. (2002) suggest that the hibernation ranges of North American bats will shift towards the North because of climate change. Furthermore, according to Newson et al. (2009), climate change is likely to affect the hibernation site selection and species composition in hibernacula to an extent that the abundance of bats, and changes in the distribution and species composition at underground hibernation sites can be used as an indicator for climate change.

In Europe, *Pipistrellus kuhlii* (Ancillotto et al. 2016; Sachanowicz et al. 2006; 2017) and *Hypsugo savii* (Lehotská and Lehotský 2006; Uhrin et al. 2016) have shown a remarkable increase in their geographical range, whereas in North-America, Willis and Brigham (2003) reported the tree-roosting bat, *Lasiurus borealis*, in Southwestern Saskatchewan, Canada, 300 km from the nearest previous observation. In addition, *Pipistrellus nathusii* has expanded its range considerably during the past decades (Benda et al. 2008; Lundy et al. 2010; Sachanowicz and Ciechanowski 2006; Sachanowicz et al. 2019). Breeding colonies, and a likely increase in the amount of individuals, have been documented in the British Isles (Matthews et al. 2018; Russ et al. 2001), Northern Italy (Martinoli et al. 2000) and on the Iberian Peninsula (Flaquer et al. 2005).

*Corresponding author: Anna S. Blomberg, Department of Biology, University of Turku, Vesilinnantie 5, Turku 20014, Finland, E-mail: asblom@utu.fi. <https://orcid.org/0000-0002-6754-4948>

Ville Vasko and Thomas M. Lilley: Finnish Museum of Natural History, University of Helsinki, P. Rautatiekatu 13, PL17, 00100 Helsinki, Finland. <https://orcid.org/0000-0001-5864-4958> (T.M. Lilley)

Saku Salonen: Department of Biology, University of Turku, Vesilinnantie 5, Turku 20014, Finland

Gunārs Pētersons: Latvian University of Life Sciences and Technologies, Lielā iela 2, Jelgava, LV-3001, Latvia

Table 1: Number of minutes (Min) with recordings of *Pipistrellus nathusii* at Härmälä cleft (Finland) during the winter 2018–2019.

Oct 2018	Min	Nov 2018	Min	Dec 2018	Min	Feb 2019	Min	Mar 2019	Min	Apr 2019	Min
6	29	3	5	20	2	15	5	23	2	16	2
8	9	13	10							17	1
9	21									24	4
10	6										
11	47										
12	21										
13	7										
14	6										
15	1										


The more northerly populations of *P. nathusii* are considered long-distance migrants, with distances of up to 1905 km between its breeding and overwintering areas (Petersons 2004), whereas in central Europe the species is a short-distance migrant or even, a sedentary species (Sachanowicz et al. 2019). *P. nathusii* overwinters predominantly in Western, Central and Southern Europe, usually hibernating solitarily or in small groups above ground (Sachanowicz et al. 2019). Typical winter roosts are buildings, hollow trees, woodpiles and sometimes rock crevices (Dietz and Kiefer 2016). Therefore, hibernation surveys fail to record the species during the winter (Sachanowicz et al. 2019), and alternative methodology is needed to map the overwintering range of the species.

Bats are known to regularly arouse (Thomas et al. 1990), fly and even forage outside hibernacula during the winter (Bernard and McCracken 2017; Hope and Jones 2012). We recorded the echolocation calls of bats at 16 sites in South-Western Finland over two consecutive winters (2017–2018 and 2018–2019) by using SongMeter SM2 + Bat passive ultrasound detectors and SMX-US ultrasound microphones (Wildlife Acoustics). These sites included four outcrop formations, three ancient shores, three glacial erratic or boulder formations, three cellars and three control sites in diverse environments where we did not expect bats to be hibernating. We filtered the data using

Kaleidoscope Pro (Wildlife Acoustics) and identified the calls manually. In our data, the echolocation calls of *P. nathusii* occurred regularly at a single site, Härmälä cleft, on several occasions during the winter of 2018–2019 (Table 1, Figure 1), suggesting that the species overwintered at the site. Due to similarities in echolocation characteristics (Zsebok et al. 2012), we cannot fully disclose the possibility of our recordings belonging to *P. kuhlii*, or its subspecies, *P. kuhlii lepidus*. This species has not been observed in Finland previously, but both *P. kuhlii* and *P. kuhlii lepidus* has shown considerable range expansion during the last decades (Ancillotto et al. 2016; Sachanowicz et al. 2006, 2017). The probability of either of these taxa occurring in Finland is still small, but the observed range shifts call for increased vigilance in identification procedures.

Härmälä cleft (N60.488358, E22.006638) is an outcrop formation on the shore of the Baltic Sea in the Masku municipality, South-Western Finland. The cleft is approximately 20 meters high with multiple deep crevices. While winter activity, as evidenced by acoustic data, suggests bats hibernate in the cleft, we never observed a bat in the crevices. We cannot disclose the possibility of the bat occupying cavities in large trees for hibernation in the vicinity of the recorder.

In addition to being the coldest month of that winter with a mean temperature of -4.4 °C, January was also the only month without any recordings of *P. nathusii*. The lowest

**Figure 1:** Sonogram of *Pipistrellus nathusii* calls recorded at Härmälä cleft (Finland) on February 15, 2019.




Figure 2: Known breeding colonies (round symbols) and overwintering site (square symbol) of *Pipistrellus nathusii* in Finland.

ambient temperature with observed *P. nathusii* activity was $-2.9\text{ }^{\circ}\text{C}$ on 20th of December 2018. At the other 15 sites, no *P. nathusii* were recorded later than 5th of November.

There is an increasing number of records of hibernating individuals to the east of earlier records, mainly from Poland, Slovenia, Slovakia and Hungary (Benda and Hotový 2004; Nusová et al. 2019; Sachanowicz and Ciechanowski 2006; Sachanowicz et al. 2019). Our record is the northernmost so far. The previous northernmost record to our knowledge was from Riga, Latvia, (16.01.2014), where an individual was found hibernating in a crevice between a window frame and concrete wall (authors' own data). Sachanowicz et al. (2019) stated that *P. nathusii* might be utilizing urban heat islands to extend their overwintering range towards North-Eastern Europe. As a result, the breeding and overwintering areas of *P. nathusii* are now overlapping in Central Europe (Sachanowicz et al. 2019). Our observations in Finland are consistent with such overlapping as, in the summer of 2017, we captured two *P. nathusii* weanlings in Turku, approximately

10 km from Härmälä cleft, indicating that the species is also breeding in the area.

While climate change may lead to loss in biodiversity especially in Southern Europe, many studies predict that species richness in Northern Europe and Scandinavia will increase (Levinsky et al. 2007; Rebelo et al. 2010). According to Mikkonen et al. (2015), the mean annual temperatures in Finland have risen $2.3 \pm 0.4\text{ }^{\circ}\text{C}$ from the mid-nineteenth century. The highest increases have taken place over the winter months, most notably in December when the monthly mean temperature has risen $4.8\text{ }^{\circ}\text{C}$. The spring months have also warmed more than the annual average. Until recently, *P. nathusii* has been considered a vagrant species in Finland during migration periods in late May and late August- early October (Ijäs et al. 2017; Rydell et al. 2014). The current range of the species in Finland covers the coastal areas in Southern and South-Western Finland, with observations as north as $64^{\circ}15'\text{N}$ (Tidenberg et al. 2019). The first breeding colony was found in Southern

Finland in 2006 (Hagner-Wahlsten and Kyheröinen 2008). Since then, records of *P. nathusii* have become more frequent outside the migration period and more breeding colonies have been discovered (Hagner-Wahlsten and Karlsson 2009, Eeva-Maria Tidenberg, personal communication, authors' own data). However, until now, the species has never been recorded in Finland during the hibernation period (Figure 2). Haarsma et al. (2019) reported that *Myotis dasycneme* males have altered their overwintering areas, suggesting that hibernating closer to breeding areas extend their mating season while saving energy needed for migration. We propose that the recent findings on the range expansion of *P. nathusii* may indicate that the migratory behavior of the species could be shifting in a similar way and that the sedentary populations of *P. nathusii* may be expanding towards the North.

Our finding indicates that acoustic surveillance of hibernation sites can yield valuable information on elusive bat species. Winter activity of bats has also been studied using acoustic monitoring in similar climatic conditions in North America, revealing unexpected levels of activity at low temperatures and new information about the ecology of several species (Lemen et al. 2017). In Alberta, Canada, bats were active even when temperatures were as low as $-8\text{ }^{\circ}\text{C}$ (Lausen and Barclay 2006) and $-10.4\text{ }^{\circ}\text{C}$ (Klüg-Baerwald et al. 2016). Acoustic monitoring during the hibernation period can also give insight to sickness behavior demonstrated by bats as a result of white-nose syndrome (WNS) (Bernard and McCracken 2017; Carr et al. 2014). Therefore, acoustic monitoring can be used as a surveillance tool for WNS (Schwab and Mabee 2014).

It is likely that recent advancements in technology used for acoustic surveillance as well as an increase in the number of bat workers has contributed to the elevated number of *P. nathusii* records in Finland. However, we suggest that these factors cannot explain the increased occurrence of the species, as observations have also increased on sites where monitoring has been continuous for more than a decade (authors' own data). Bat species previously known to overwinter above 60°N in Fennoscandia include *Eptesicus nilssonii*, *Myotis daubentonii*, *Myotis brandtii*, *Myotis mystacinus*, *M. dasycneme*, *Myotis nattereri* and *Plecotus auritus* (Siivonen and Wermundsen 2003; Wermundsen and Siivonen 2010). Our results suggest that *P. nathusii* has now joined these seven species.

Acknowledgments: The study was funded by Kone foundation.

Author contribution: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: The study was funded by Kone foundation and H2020 Marie Skłodowska-Curie Actions.

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

- Ancillotto, L., Santini, L., Ranc, N., Maiorano, L., and Russo, D. (2016). Extraordinary range expansion in a common bat: the potential roles of climate change and urbanisation. *Sci. Nat.* 103: 15.
- Benda, P. and Hotový, J. (2004). Hibernation record of (*Pipistrellus nathusii*) in Southern Moravia (Czech Rep). *Vespertilio* 8: 137–139. (in Czech).
- Benda, P., Georgiakakis, P., Dietz, C., Hanák, V., Galanaki, K., Markantonatou, V., Chudárková, A., Hulva, P., and Horáček, I. (2008). Bats (Mammalia: Chiroptera) of the Eastern Mediterranean and Middle East. Part 7. The bat fauna of Crete, Greece. *Acta Soc. Zool. Bohem.* 72: 105–190.
- Bernard, R.F. and McCracken, G.F. (2017). Winter behavior of bats and the progression of white-nose syndrome in the Southeastern United States. *Ecol. Evol.* 7: 1487–1496.
- Brommer, J.E., Lehikoinen, A., and Valkama, J. (2012). The breeding ranges of Central European and Arctic bird species move poleward. *PLoS One* 7: e43648.
- Carr, J.A., Bernard, R.F., and Stiver, W.H. (2014). Unusual bat behavior during winter in Great Smoky Mountains National Park. *Southeast. Nat.* 13: N18–N21.
- Dietz, C. and Kiefer, A. (2016). *Bats of Britain and Europe*. Bloomsbury, London.
- Flaquer, C., Ruiz-Jarillo, R., Torre, I., and Arrizabalaga, A. (2005). First resident population of *Pipistrellus nathusii* (Keyserling and Blasius, 1839) in the Iberian Peninsula. *Acta Chiropterol.* 7: 183–188.
- Haarsma, A.-J., Lina, P.H.C., Voûte, A.M., and Siepel, H. (2019). Male long-distance migrant turned sedentary; the west European pond bat (*Myotis dasycneme*) alters their migration and hibernation behaviour. *PLoS One* 14: e0217810.
- Hagner-Wahlsten, N. and Karlsson, R. (2009). *Bat survey of Seurasaari 2009*. BatHouse, Kauniainen. (in Finnish).
- Hagner-Wahlsten, N. and Kyheröinen, E.-M. (2008). First observation of breeding Nathusius' pipistrelle (*Pipistrellus nathusii*) in Finland. *Memo. Soc. Fauna Flora Fenn.* 84: 36–40.
- Hickling, R., Roy, D.B., Hill, J.K., Fox, R., and Thomas, C.D. (2006). The distributions of a wide range of taxonomic groups are expanding polewards. *Global Change Biol.* 12: 450–455.
- Hope, P.R. and Jones, G. (2012). Warming up for dinner: torpor and arousal in hibernating Natterer's bats (*Myotis nattereri*) studied by radio telemetry. *J. Comp. Physiol. B.* 182: 569–578.
- Humphries, M.M., Thomas, D.W., and Speakman, J.R. (2002). Climate-mediated energetic constraints on the distribution of hibernating mammals. *Nature*. 418: 313–316.
- Ijäs, A., Kahilainen, A., Vasko, V.V., and Lilley, T.M. (2017). Evidence of the migratory bat, *Pipistrellus nathusii*, aggregating to the coastlines in the Northern Baltic Sea. *Acta Chiropterol.* 19: 127–139.
- Klüg-Baerwald, B.J., Gower, L.E., Lausen, C.L., and Brigham, R.M. (2016). Environmental correlates and energetics of winter flight by bats in southern Alberta, Canada. *Can. J. Zool.* 94: 829–836.

- Lane, J.E., Kruuk, L.E.B., Charmantier, A., Murie, J.O., and Dobson, F.S. (2012). Delayed phenology and reduced fitness associated with climate change in a wild hibernator. *Nature*. 489: 554–557.
- Lausen, C.L. and Barclay, R.M.R. (2006). Winter bat activity in the Canadian prairies. *Can. J. Zool.* 84: 1079–1086.
- Lehotská, B. and Lehotský, R. (2006). First record of *Hypsugo savii* (Chiroptera) in Slovakia. *Biologĳa (Bratisl.)* 61: 192.
- Lemen, C.A., Freeman, P.W., and White, J.A. (2017). Acoustic evidence of bats using rock crevices in winter: a call for more research on winter roosts in North America. *Trans. Neb. Acad. Sci. Affil. Soc.* 36: 9–13.
- Levinsky, I., Skov, F., Svenning, J.C., and Rahbek, C. (2007). Potential impacts of climate change on the distributions and diversity patterns of European mammals. *Biodivers. Conserv.* 16: 3803–3816.
- Lundy, M., Montgomery, I., and Russ, J. (2010). Climate change-linked range expansion of Nathusius' pipistrelle bat, *Pipistrellus nathusii* (Keyserling & Blasius, 1839). *J. Biogeogr.* 37: 2232–2242.
- Maclean, I.M.D., Austin, G.E., Rehfishch, M.M., Blew, J., Crowe, O., Delany, S., Devos, K., Deceuninck, B., Günther, K., Laursen, K., et al. (2008). Climate change causes rapid changes in the distribution and site abundance of birds in winter. *Global Change Biol.* 14: 2489–2500.
- Martinoli, A., Preatoni, D.G., and Tosi, G. (2000). Does Nathusius' pipistrelle *Pipistrellus nathusii* (Keyserling & Blasius, 1839) breed in Northern Italy? *J. Zool.* 250: 217–220.
- Matthews, F., Kubasiewicz, L.M., Gurnell, J., Harrower, C.A., McDonald, R.A., and Shore, R.F. (2018). *Review of the population and conservation status of British mammals*. Natural England, Peterborough.
- Mikkonen, S., Laine, M., Mäkelä, H.M., Gregow, H., Tuomenvirta, H., Lahtinen, M., and Laaksonen, A. (2015). Trends in the average temperature in Finland, 1847–2013. *Stoch. Environ. Res. Risk Assess.* 29: 1521–1529.
- Newson, S., Mendes, S., Crick, H., Dulvy, N., Houghton, J., Hays, G., Hutson, A., Macleod, C., Pierce, G., and Robinson, R. (2009). Indicators of the impact of climate change on migratory species. *Endanger. Species Res.* 7: 101–113.
- Nusová, G., Uhrin, M., and Kaňuch, P. (2019). Go to the city: urban invasions of four pipistrelle bat species in Eastern Slovakia. *Eur. J. Ecol.* 5: 23–26.
- Parmesan, C. and Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*. 421: 37–42.
- Petersons, G. (2004). Seasonal migrations of North-Eastern populations of Nathusius bat *Pipistrellus nathusii* (Chiroptera). *Myotis* 41–42: 29–56.
- Rebelo, H., Tarroso, P., and Jones, G. (2010). Predicted impact of climate change on European bats in relation to their biogeographic patterns. *Global Change Biol.* 16: 561–576.
- Russ, J.M., Hutson, A.M., Montgomery, W.I., Racey, P.A., and Speakman, J.R. (2001). The status of Nathusius' pipistrelle (*Pipistrellus nathusii* Keyserling & Blasius, 1839) in the British Isles. *J. Zool.* 254: 91–100.
- Rydell, J., Bach, L., Bach, P., Diaz, L.G., Furmankiewicz, J., Hagner-Wahlsten, N., Kyheröinen, E.-M., Lilley, T., Masing, M., Meyer, M.M., et al. (2014). Phenology of migratory bat activity across the Baltic Sea and the South-Eastern North Sea. *Acta Chiropterol.* 16: 139–147.
- Sachanowicz, K. and Ciechanowski, M. (2006). First winter record of the migratory bat *Pipistrellus nathusii* (Keyserling and Blasius 1839) (Chiroptera: Vespertilionidae) in Poland: yet more evidence of global warming? *Mammalia* 70: 168–169.
- Sachanowicz, K., Wower, A., and Bashta, A.-T. (2006). Further range extension of *Pipistrellus kuhlii* (Kuhl, 1817) in Central and Eastern Europe. *Acta Chiropterol.* 8: 543–548.
- Sachanowicz, K., Piskorski, M., and Tereba, A. (2017). Systematics and taxonomy of *Pipistrellus kuhlii* (Kuhl, 1817) in Central Europe and the Balkans. *Zootaxa* 4306: 53.
- Sachanowicz, K., Ciechanowski, M., Tryjanowski, P., and Kosicki, J.Z. (2019). Wintering range of *Pipistrellus nathusii* (Chiroptera) in Central Europe: has the species extended to the north-east using urban heat islands? *Mammalia* 83: 260–271.
- Schwab, N.A. and Mabee, T.J. (2014). Winter acoustic activity of bats in Montana. *Northwest. Nat.* 95: 13–27.
- Sherwin, H.A., Montgomery, W.I., and Lundy, M.G. (2013). The impact and implications of climate change for bats: bats and climate change. *Mammal Rev.* 43: 171–182.
- Siivonen, Y. and Wermundsen, T. (2003). First records of *Myotis dasycneme* and *Pipistrellus pipistrellus* in Finland. *Vespertilio* 7: 177–179.
- Sorte, F.A.L. and Iii, F.R.T. (2007). Poleward shifts in winter ranges of North American birds. *Ecology* 88: 1803–1812.
- Thomas, D.W., Dorais, M., and Bergeron, J.-M. (1990). Winter energy budgets and cost of arousals for hibernating little brown bats, *Myotis lucifugus*. *J. Mammal.* 71: 475–479.
- Thomas, C., Franco, A., and Hill, J. (2006). Range retractions and extinction in the face of climate warming. *Trends Ecol. Evol.* 21: 415–416.
- Thomas, C.D. (2010). Climate, climate change and range boundaries: climate and range boundaries. *Divers. Distrib.* 16: 488–495.
- Tidenberg, E.-M., Liukko, U.-M., and Stjernberg, T. (2019). Atlas of Finnish bats. *Ann. Zool. Fenn.* 56: 207–250.
- Tingley, M.W., Monahan, W.B., Beissinger, S.R., and Moritz, C. (2009). Birds track their Grinnellian niche through a century of climate change. *Proc. Natl. Acad. Sci.* 106: 19637–19643.
- Uhrin, M., Hüttmeir, U., Kipson, M., Estók, P., Sachanowicz, K., Bücs, S., Karapandža, B., Paunović, M., Presetnik, P., Bashta, A.T., et al. 2016. Status of Savi's pipistrelle *Hypsugo savii* (Chiroptera) and range expansion in Central and South-eastern Europe: a review. *Mammal Rev.* 46: 1–16.
- Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B., Roy, D.B., Willis, S.G., and Greatorex-Davies, J.N. (2001). Rapid responses of british butterflies to opposing forces of climate and habitat change. *Nature* 414: 65–68.
- Wermundsen, T. and Siivonen, Y. (2010). Seasonal variation in use of winter roosts by five bat species in South-east Finland. *Open Life Sci.* 5: 262–273.
- Willis, C.K.R. and Brigham, R.M. (2003). New records of the eastern red bat, *Lasiurus borealis*, from Cypress Hills Provincial Park, Saskatchewan: a response to climate change? *Can. Field Nat.* 117: 651–654.
- Zsebok, S., Estók, P., and Görföl, T. (2012). Acoustic discrimination of *Pipistrellus kuhlii* and *Pipistrellus nathusii* and its application to assess changes in species distribution. *Acta Zool. Acad. Sci. Hung.* 58: 199–209.